

List of Tables

Table 1	Carbapenemase genes associated with plasmids and mobile genetic elements (MGE).	15
Table-3.1	Morphological characteristics of collected clinical isolates.	35
Table 3.2	Results of biochemical tests of all isolates.	40
Table 3.3	NCBI BLAST results of the Sanger sequenced amplicons of 16s <i>rRNA</i> .	44
Table 4.1	List of antibiotics used for disk-diffusion method.	52
Table 4.2	List of antibiotics used for categorization of isolates (as per Magiorakos et al., 2011).	54
Table 4.3	Primers used for Multiplex PCR and amplicon size.	56
Table 4.4	Distribution of n=11 primer pairs into three sets and their melting temperature (T_m).	57
Table 4.5	Reaction mixture for multiplex PCR.	58
Table 4.6	Cycle/conditions used for multiplex PCR reaction.	58
Table 4.7	Reaction mixture for DNase treatment of extracted mRNA.	60
Table 4.8	Reaction system-I for cDNA synthesis.	60
Table 4.9	Reaction system-II for cDNA synthesis.	61
Table 4.10	Primers used for qRT-PCR.	61
Table 4.11	Reaction system used for qRT-PCR.	62

Table 4.12	Cycle used for qRT-PCR.	62
Table 4.13	Antimicrobial susceptibility testing using disk diffusion.	69
Table 4.14	MIC determination for tigecycline and colistin.	75
Table 4.15	Results of ESBL and MBL phenotypic assay.	76
Table 4.16	Carbapenemase genes detected using multiplex PCR.	80
Table 4.17	Quality analysis of the sequences.	82
Table 4.18	MLST type of n=8 isolates based on allelic combinations at 7 loci.	83
Table 4.19a	Phenotypic and genomic analysis of the eight <i>Kp</i> isolates.	87
Table 4.19b	Phenotypic and genomic analysis of the eight <i>Kp</i> isolates.	87