

---

## References

- Adu-Gyamfi, J.J., Myaka, F.A., Sakala, W.D., Odgaard, R., Vesterager, J.M., Høgh-Jensen, H., 2007. Biological nitrogen fixation and nitrogen and phosphorus budgets in farmer-managed intercrops of maize-pigeonpea in semi-arid southern and eastern Africa. *Plant Soil*. 295, 127–136. <https://doi.org/10.1007/s11104-007-9270-0>.
- Ae, N., Arihara, J., Okada, K., Yoshihara, T., Johansen, C., 1990. Phosphorus Uptake by Pigeon Pea and Its Role in Cropping Systems of the Indian Subcontinent. *Sci*. 495, 2–5. <https://doi.org/10.1126/science.248.4954.477>.
- Agegnehu, G., Ghizaw, A., & Sinebo, W. 2008. Yield potential and land-use efficiency of wheat and faba bean mixed intercropping. *Agron Sustain Dev*. 28(2), 257-263. <https://doi.org/10.1051/agro:2008012>.
- Ahmad, A., Akram, W., Shahzadi, I., Wang, R., Hu, D., Bashir, Z., Jaleel, W., Ahmed, S., Tariq, W., Li, G. and Wu, T., 2019. Benzenedicarboxylic acid upregulates O48814 and Q9FJQ8 for improved nutritional contents of tomato and low risk of fungal attack. *J. Sci. Food Agric*, 99(14), 6139-6154. <https://doi.org/10.1002/jsfa.9836>.
- Ankati, S., Rani, T.S. and Podile, A.R., 2019. Changes in root exudates and root proteins in groundnut–*Pseudomonas* sp. interaction contribute to root colonization by bacteria and defense response of the host. *J. Plant Growth Regul*. 38(2), 523-538. <https://doi.org/10.1007/s00344-018-9868-x>.
- Badri, D. V., Vivanco, J.M., 2009. Regulation and function of root exudates. *Plant, Cell Environ* 32, 666–681. <https://doi.org/10.1111/j.1365-3040.2009.01926.x>
- Badri, D.V., de-la-Peña, C., Lei, Z., Manter, D.K., Chaparro, J.M., Guimarães, R.L., Sumner, L.W., Vivanco, J.M., 2012. Root secreted metabolites and proteins are involved in the early events of Plant-Plant recognition prior to competition. *PLoS One* 7. <https://doi.org/10.1371/journal.pone.0046640>.
- Bais, H.P., Park, S., Weir, T.L., Callaway, R.M., Vivanco, J.M., 2004. How plants communicate using the underground information superhighway 9, 26–32. <https://doi.org/10.1016/j.tplants.2003.11.008>
- Bais HP, Weir TL, Perry LG, Gilroy S, Vivanco JM. 2006. The role of root exudates in rhizosphere interactions with plants and other organisms. *Annu Rev Plant Biol* 57: 233–266. <https://doi.org/10.1146/annurev.arplant.57.032905.105159>
- Bacic A, Moody SF, Clarke AE (1986) Structural Analysis of Secreted Root Slime from

- Maize (*Zea mays* L.). *Plant Physiol* 80:771–777. <https://doi.org/10.1104/pp.80.3.771>
- Backer, R., Rokem, J.S., Ilangumaran, G., Lamont, J., Praslickova, D., Ricci, E., Subramanian, S., Smith, D.L., 2018. Plant growth-promoting Rhizobacteria : context, mechanisms of action, and roadmap to commercialization of biostimulants for sustainable agriculture. *Front. Plant Sci.* 9, 1–17. <https://doi.org/10.3389/fpls.2018.01473>
  - Bargaz, A., Lyamlouli, K., Chtouki, M., Zeroual, Y., & Dhiba, D. 2018. Soil microbial resources for improving fertilizers efficiency in an integrated plant nutrient management system. *Frontiers in microbiology*, 9, 1606. <https://doi.org/10.3389/fmicb.2018.01606>
  - Beauregard, P.B., 2015. Not Just Sweet Talkers: How Roots Stimulate Their Colonization by Beneficial Bacteria, *Advances in Botanical Research*. Elsevier Ltd. <https://doi.org/10.1016/bs.abr.2015.07.001>
  - Bechtaoui, N., El Alaoui, A., Raklami, A., Benidire, L., Tahiri, A.I., Oufdou, K., 2019. Impact of intercropping and co-inoculation with strains of plant growth-promoting rhizobacteria on phosphorus and nitrogen concentrations and yield of durum wheat (*Triticum durum*) and faba bean (*Vicia faba*). *Crop Pasture Sci.* 70, 649–658. <https://doi.org/10.1071/CP19067>.
  - Bedoussac, L., Journet, E.P., Hauggaard-Nielsen, H., Naudin, C., Corre-Hellou, G., Jensen, E.S., Prieur, L., Justes, E., 2015. Ecological principles underlying the increase of productivity achieved by cereal-grain legume intercrops in organic farming. A review. *Agron Sustain Dev* 35, 911–935. <https://doi.org/10.1007/s13593-014-0277-7>
  - Benyamina, S., Alloing, G., Rouhier, N., Jacquot, J., Puppo, A. and Frenedo, P., 2013. Role of *Sinorhizobium meliloti* glutaredoxins in nitrogen-fixing symbiosis. *BioTechnologia. Journal of Biotechnology Computational Biology and Bionanotechnology*, 94(2). <https://doi:10.1111/j.1462-2920.2012.02835.x>
  - Bertin, C., Yang, X., & Weston, L. A., 2003. The role of root exudates and allelochemicals in the rhizosphere. *Plant and soil*, 256(1), 67-83. <https://doi.org/10.1023/A:1026290508166>
  - Berg, G., Smalla, K., 2009. Plant species and soil type cooperatively shape the structure and function of microbial communities in the rhizosphere. *FEMS Microbiol. Ecol.* 68, 1–13. <http://dx.doi.org/10.1111/j.1574-6941.2009.00654.x>.
  - Bradford, M.M., 1976. A rapid and sensitive method for the quantitation of microgram

- quantities of protein utilizing the principle of protein-dye binding. *Analytical biochemistry*, 72(1-2), pp.248-254.
- Brooker, R. W., Bennett, A. E., Cong, W. F., Daniell, T. J., George, T. S., Hallett, P. D., & White, P. J. 2015. Improving intercropping: a synthesis of research in agronomy, plant physiology and ecology. *New Phytologist*, 206(1), 107-117. <https://doi.org/10.1111/nph.13132>
  - Broughton, W.J., Wong, C.H., Lewin, A., Samrey, U., Myint, H., Meyer, H., Dowling, D.N. and Simon, R. (1986) Identification of *Rhizobium* plasmid sequences involved in recognition of *Psophocarpus*, *Vigna*, and other legumes. *J Cell Biol.* 102, 1173-1182. doi:10.1083/jcb.102.4.1173
  - Broughton, W. J., Zhang, F., Perret, X., and Staehelin, C., 2003. Signals exchanged between legumes and *Rhizobium*: agricultural uses and perspectives. *Plant Soil* 252, 129–137. <https://doi.org/10.1023/A:1024179717780>
  - Calatrava-Morales, N., McIntosh, M., and Soto, M. J., 2018. Regulation Mediated by N-Acyl Homoserine Lactone Quorum Sensing Signals in the *Rhizobium*-Legume Symbiosis. *Genes (Basel)*. 18; 9(5):263. doi: 10.3390/genes9050263
  - Calicioglu, O., Flammini, A., Bracco, S., Bellù, L., & Sims, R., 2019. The future challenges of food and agriculture: An integrated analysis of trends and solutions. *Sustainability*, 11(1), 222. <https://doi.org/10.3390/su11010222>
  - Canarini, A., Kaiser, C., Merchant, A., Richter, A., Wanek, W., 2019. Root exudation of primary metabolites: Mechanisms and their roles in plant responses to environmental stimuli. *Front Plant Sci* 10. <https://doi.org/10.3389/fpls.2019.00157>
  - Canellas, N.O., Olivares, F.L. and Canellas, L.P., 2019. Metabolite fingerprints of maize and sugarcane seedlings: searching for markers after inoculation with plant growth-promoting bacteria in humic acids. *Chem. biol. technol.*, 6(1), pp.1-10. <https://doi.org/10.1186/s40538-019-0153-4>
  - Carranca C (2013) Legumes: properties and symbiosis. In: Camisão AH, Pedroso CC (eds) *Symbiosis: evolution, biology and ecological effects*. Animal science, issues and professions. Nova Science Publishers, New York. ISBN 978-1-62257-211-3
  - Carvalhais, L.C., Dennis, P.G., Fedoseyenko, D., Hajirezaei, M.R., Borriss, R., Von Wirén, N., 2011. Root exudation of sugars, amino acids, and organic acids by maize as

- affected by nitrogen, phosphorus, potassium, and iron deficiency. *J. Plant Nutr. Soil Sci.* 174, 3–11. <https://doi.org/10.1002/jpln.201000085>.
- Cavalcanti, M.I.P., Carvalho, R. De, Ribeiro, D., Elena, I., Escobar, C., Carla, A., Fraiz, R., Pereira, A., Souza, D., Dolores, A., Freitas, S. De, Simão, R., Nóbrega, A., Fernandes-júnior, P.I., 2020. Maize growth and yield promoting endophytes isolated into a legume root nodule by a cross-over approach. *Rhizosphere* 100211. <https://doi.org/10.1016/j.rhisph.2020.100211>.
  - Chagas, F.O., Pessotti, R.D.C., Caraballo-Rodríguez, A.M., Pupo, M.T., 2018. Chemical signaling involved in plant-microbe interactions. *Chem. Soc. Rev.* 47, 1652–1704. <https://doi.org/10.1039/c7cs00343a>.
  - Chang, C., Damiani, I., Puppo, A. and Frenedo, P., 2009. Redox changes during the legume–*Rhizobium* symbiosis. *Molecular Plant*, 2(3), pp.370-377. <https://doi.org/10.1093/mp/ssn090>
  - Chen, J. and Strous, M., 2013. Denitrification and aerobic respiration, hybrid electron transport chains and co-evolution. *Biochimica et Biophysica Acta (BBA)-Bioenergetics*, 1827(2), pp.136-144. <http://dx.doi.org/10.1016/j.bbabi.2012.10.002>
  - Chen, W.M., Prell, J., James, E.K., Sheu, D.S. and Sheu, S.Y., 2012. Biosynthesis of branched-chain amino acids is essential for effective symbioses between betarhizobia and *Mimosa pudica*. *Microbiology*, 158(7), pp.1758-1766. DOI 10.1099/mic.0.058370-0
  - Chen, J., Arafat, Y., Wu, L., Xiao, Z., Li, Q., 2017. Shifts in soil microbial community , soil enzymes and crop yield under peanut / maize intercropping with reduced nitrogen levels. *Appl Soil Ecol* 0–1. <https://doi.org/10.1016/j.apsoil.2017.11.010>
  - Chen, Y., Bonkowski, M., Shen, Y., Griffiths, B.S., Jiang, Y., Wang, X., Sun, B., 2020. Root ethylene mediates rhizosphere microbial community reconstruction when chemically detecting cyanide produced by neighbouring plants. *Microbiome*, 8(1), 1–17. <https://doi.org/10.1186/s40168-019-0775-6>
  - Chimonyo, V. G. P., Snapp, S. S., & Chikowo, R., 2019. Grain legumes increase yield stability in maize based cropping systems. *Crop Science*, 59(3), 1222-1235. <https://doi.org/10.2135/cropsci2018.09.0532>
  - Chong, T.M., Chen, J.W., Seng, W., Too, S., Yu, C.Y., Ang, G.Y., Lim, Y.L., Yin, W.F., Grandclément, C., Faure, D., Dessaux, Y., Chan, K.G., 2017. Phenotypic and genomic

- survey on organic acid utilization profile of *Pseudomonas mendocina* strain S5 . 2, a vineyard soil isolate. *Amb. Express* 7, 138. <https://doi.org/10.1186/s13568-017-0437-7>.
- Cong, W.F., Hoffland, E., Li, L., Janssen, B.H. and van der Werf, W., 2015. Intercropping affects the rate of decomposition of soil organic matter and root litter. *Plant and Soil*, 391(1), 399-411. <https://doi.org/10.1007/s11104-015-2433-5>
  - Contreras, F., Díaz, J., Domenico, A., De, M., Mora, L., 2019. Current Plant Biology Prospecting intercropping between subterranean clover and grapevine as potential strategy for improving grapevine performance. *Curr Plant Biol* 19, 100110. <https://doi.org/10.1016/j.cpb.2019.100110>
  - Coskun, D., Britto, D.T., Shi, W., Kronzucker, H.J., 2017. How Plant Root Exudates Shape the Nitrogen Cycle. *Trends Plant Sci* 22, 661–673. <https://doi.org/10.1016/j.tplants.2017.05.004>
  - Cu, S.T.T., Hutson, J., Schuller, K.A., 2005. Mixed culture of wheat (*Triticum aestivum* L.) with white lupin (*Lupinus albus* L.) improves the growth and phosphorus nutrition of the wheat 143–151. <https://doi.org/10.1007/s11104-004-4336-8>
  - Dai, J., Qiu, W., Wang, N., Wang, T., Nakanishi, H., Zuo, Y., 2019. From leguminosae/gramineae intercropping systems to see benefits of intercropping on iron nutrition. *Front Plant Sci* 10, 1–7. <https://doi.org/10.3389/fpls.2019.00605>
  - Darias, J.A.R., García-Fontana, C., Lugo, A.C., Rico-Jiménez, M., Krell, T., 2014. Qualitative and Quantitative assays for flagellum-mediated chemotaxis. In: Filloux, A., Ramos, J.L. (Eds.), *Pseudomonas Methods and Protocols. Methods in Molecular Biology (Methods and Protocols)*, vol. 1149. Humana Press, New York, NY. [https://doi.org/10.1007/978-1-4939-0473-0\\_10](https://doi.org/10.1007/978-1-4939-0473-0_10)
  - Dall’Agnol, R. F., Ribeiro, R. A., Ormeño-Orrillo, E., Rogel, M. A., Delamuta, J. R. M., Andrade, D. S., (2013). *Rhizobium freirei* sp. nov., a symbiont of *Phaseolus vulgaris* that is very effective at fixing nitrogen. *Int J Syst Evol Microbiol.* 63, 4167–4173. [doi:10.1099/ijms.0.052928-0](https://doi.org/10.1099/ijms.0.052928-0)
  - Dang, K., Gong, X., Zhao, G., Wang, H., Ivanistau, A., 2020. Intercropping Alters the Soil Microbial Diversity and Community to Facilitate Nitrogen Assimilation: A Potential Mechanism for Increasing Proso Millet Grain Yield 11, 1–17. <https://doi.org/10.3389/fmicb.2020.601054>

- Danhorn, T., & Fuqua, C., 2007. Biofilm formation by plant-associated bacteria. *Annual Review of Microbiology*, 61, 401e422.
- da Silva Lima, L., Olivares, F.L., De Oliveira, R.R., Vega, M.R.G., Aguiar, N.O. and Canellas, L.P., 2014. Root exudate profiling of maize seedlings inoculated with *Herbaspirillum seropedicae* and humic acids. *Chemical and Biological Technologies in Agriculture*, 1(1), pp.1-18.<http://www.chembioagro.com/content/1/1/23>
- de Oliveira, I.M., Zanotto-Filho, A., Moreira, J.C.F., Bonatto, D. and Henriques, J.A.P., 2010. The role of two putative nitroreductases, Frm2p and Hbn1p, in the oxidative stress response in *Saccharomyces cerevisiae*. *Yeast*, 27(2), pp.89-102.<https://doi.org/10.1002/yea.1734>
- Dennis, P.G., Miller, A.J., Hirsch, P.R., 2010. Are root exudates more important than other sources of rhizodeposits in structuring rhizosphere bacterial communities? *FEMS Microbiol Ecol* 72, 313–327. <https://doi.org/10.1111/j.1574-6941.2010.00860.x>
- Depuydt, S., 2014. Arguments for and against self and non-self root recognition in plants. *Front Plant Sci* 5, 1–7. <https://doi.org/10.3389/fpls.2014.00614>
- Derrien, D., Marol, C., & Balesdent, J., 2004. The dynamics of neutral sugars in the rhizosphere of wheat. An approach by <sup>13</sup>C pulse-labelling and GC/C/IRMS. *Plant and Soil*, 267(1), 243-253. <https://doi.org/10.1007/s11104-005-5348-8>
- Dhont, C., Castonguay, Y., Nadeau, P., Bélanger, G., Drapeau, R., Laberge, S., Chalifour, F. P., 2006. Nitrogen reserves, spring regrowth and winter survival of field-grown alfalfa (*Medicago sativa*) defoliated in the autumn. *Annals of Botany*, 97(1), 109-120.
- Ding, H., Yip, C.B., Geddes, B.A., Oresnik, I.J. and Hynes, M.F., 2012. Glycerol utilization by *Rhizobium leguminosarum* requires an ABC transporter and affects competition for nodulation. *Microbiology*, 158(5), pp.1369-1378.<https://doi.org/10.1099/mic.0.057281-0>
- Downie, J.A., 2010. The roles of extracellular proteins, polysaccharides and signals in the interactions of rhizobia with legume roots. *FEMS microbiology reviews*, 34(2), pp.150-170. DOI:10.1111/j.1574-6976.2009.00205.x
- Duchene, O., Vian, J., Celette, F., 2017. Agriculture , Ecosystems and Environment Intercropping with legume for agroecological cropping systems : Complementarity and facilitation processes and the importance of soil microorganisms . A review. "Agriculture,

- Ecosyst Environ 240, 148–161. <https://doi.org/10.1016/j.agee.2017.02.019>
- Dunn, M.F., 2015. Key roles of microsymbiont amino acid metabolism in rhizobia-legume interactions. *Critical reviews in microbiology*, 41(4), pp.411-451. DOI: 10.3109/1040841X.2013.856854
  - Dutta, S., & Podile, A. R. 2010. Plant growth promoting rhizobacteria (PGPR): the bugs to debug the root zone. *Critical Reviews in Microbiology*, 36(3), 232e244.
  - Erro, J., Zamarreñ O, A., Yvin, J.C., Garcia-Mina, J.M., 2009. Determination of organic acids in tissues and exudates of maize, lupin, and chickpea by high-performance liquid chromatography-tandem mass spectrometry. *J. Agric. Food Chem.* 57, 4004–4010. <https://doi.org/10.1021/jf804003v>.
  - Faget, M., Nagel, K.A., Walter, A., Herrera, J.M., Jahnke, S., Schurr, U. and Temperton, V.M., 2013. Root–root interactions: extending our perspective to be more inclusive of the range of theories in ecology and agriculture using in-vivo analyses. *Annals of Botany*, 112(2), pp.253-266.<https://doi.org/10.1093/aob/mcs296>
  - Faurobert, M., Pelpoir, E. and Chaïb, J., 2007. Phenol extraction of proteins for proteomic studies of recalcitrant plant tissues. In *Plant Proteomics* (pp. 9-14). Humana Press.
  - Feng, H., Zhang, N., Du, W., Zhang, H., Liu, Y., Fu, R., Shao, J., Zhang, G., Shen, Q., Zhang, R., 2018. Identification of chemotaxis compounds in root exudates and their sensing chemoreceptors in plant growth-promoting rhizobacteria *Bacillus amyloliquefaciens* SQR9. *Mol. Plant Microbe Interact.* 31, 995–1005. <https://doi.org/10.1094/MPMI-01-18-0003-R>
  - Ferraioli, S., Tatè, R., Cermola, M., Favre, R., Iaccarino, M. and Patriarca, E.J., 2002. Auxotrophic mutant strains of *Rhizobium etli* reveal new nodule development phenotypes. *Molecular plant-microbe interactions*, 15(5), pp.501-510. <https://doi.org/10.1094/MPMI.2002.15.5.501>
  - Flemming, H.-C., & Wingender, J., 2010. The biofilm matrix. *Nature Reviews Microbiology*, 8(9), 623e633. <http://dx.doi.org/10.1038/nrmicro2415>
  - Flores-Tinoco, C.E., Tschan, F., Fuhrer, T., Margot, C., Sauer, U., Christen, M. and Christen, B., 2020. Co-catabolism of arginine and succinate drives symbiotic nitrogen fixation. *Molecular systems biology*, 16(6), p.e9419.<https://doi.org/10.15252/msb.20199419>

- Fry, J., Wood, M. and Poole, P.S., 2001. Investigation of myo-inositol catabolism in *Rhizobium leguminosarum* bv. *viciae* and its effect on nodulation competitiveness. *Molecular plant-microbe interactions*, 14(8), pp.1016-1025. <https://doi.org/10.1094/MPMI.2001.14.8.1016>
- Garland, G., Bünenmann, E.K., Oberson, A., 2017. Plant-mediated rhizospheric interactions in maize-pigeon pea intercropping enhance soil aggregation and organic phosphorus storage 37–55. <https://doi.org/10.1007/s11104-016-3145-1>
- Ghaley BB, Hauggaard-Nielsen H, Jensen HH, Jensen ES 2005 Intercropping of wheat and pea as influenced by nitrogen fertilization. *Nutr Cycl Agroecosyst* 73:201–212. <https://doi.org/10.1007/s10705-005-2475-9>
- Glaze-Corcoran, S., Hashemi, M., Sadeghpour, A., Jahanzad, E., Afshar, R.K., Liu, X. and Herbert, S.J., 2020. Understanding intercropping to improve agricultural resiliency and environmental sustainability. In *Advances in Agronomy* (Vol. 162, pp. 199-256). Academic Press. <https://doi.org/10.1016/bs.agron.2020.02.004>
- Glick, B. R., 2012. Plant growth-promoting bacteria: mechanisms and applications. *Scientifica* 2012:15. doi: 10.6064/2012/963401
- Gomes, D.F., da Silva Batista, J.S., Schiavon, A.L., Andrade, D.S. and Hungria, M., 2012. Proteomic profiling of *Rhizobium tropici* PRF 81: identification of conserved and specific responses to heat stress. *BMC microbiology*, 12(1), pp.1-12. <https://doi.org/10.1186/1471-2180-12-84>
- Gómez-Godínez, L.J., Fernandez-Valverde, S.L., Romero, J.C.M. and Martínez-Romero, E., 2019. Metatranscriptomics and nitrogen fixation from the rhizoplane of maize plantlets inoculated with a group of PGPRs. *Systematic and applied microbiology*, 42(4), pp.517-525. <https://doi.org/10.1016/j.syapm.2019.05.003>
- Gonzalez-y-Merchand, J.A., Estrada-Garcia, I., Colston, M.J. and Cox, R.A., 1996. A novel method for the isolation of mycobacterial DNA. *FEMS microbiology letters*, 135(1), pp.71-77. <https://doi.org/10.1111/j.1574-6968.1996.tb07968.x>
- Gordillo, F., Ch´avez, F.P., Jerez, C.A., 2007. Motility and chemotaxis of *Pseudomonas* sp. B4 towards polychlorobiphenyls and chlorobenzoates. *FEMS Microbiol. Ecol.* 60, 322–328. <https://doi.org/10.1111/j.1574-6941.2007.00293.x>
- Gosai, J., Anandhan, S., Bhattacharjee, A., Archana, G., 2019. Elucidation of quorum

- sensing components and their role in regulation of symbiotically important traits in Ensifer nodulating pigeon pea. *Microbiol. Res.* 231, 126354. <https://doi.org/10.1016/j.micres.2019.126354>.
- Grote, J., Krysciak, D., Schorn, A., Dahlke, R.I., Soonvald, L., Müller, J., Hense, B.A., Schwarzfischer, M., Sauter, M., Schmeisser, C. and Streit, W.R., 2014. Evidence of autoinducer-dependent and-independent heterogeneous gene expression in *Sinorhizobium fredii* NGR234. *Applied and environmental microbiology*, 80(18), pp.5572-5582. <http://dx.doi.org/10.1128/AEM.01689-14>.
  - Haichar, F. el Z., Santaella, C., Heulin, T., Achouak, W., 2014. Root exudates mediated interactions belowground. *Soil Biol Biochem* 77, 69–80. <https://doi.org/10.1016/j.soilbio.2014.06.017>
  - Hara, S., Morikawa, T., Wasai, S., Kasahara, Y., Koshiba, T., Yamazaki, K., Fujiwara, T., Tokunaga, T. and Minamisawa, K., 2019. Identification of nitrogen-fixing *Bradyrhizobium* associated with roots of field-grown sorghum by metagenome and proteome analyses. *Frontiers in microbiology*, 10, p.407. <http://dx.doi.org/10.3389/fmicb.2019.00407>
  - Hauggaard-Nielsen, H., Jensen, E.S., 2005. Facilitative root interactions in intercrops. *Plant Soil* 274, 237–250. <https://doi.org/10.1007/s11104-004-1305-1>
  - He, X., Chang, W., Pierce, D. L., Seib, L. O., Wagner, J., and Fuqua, C. 2003. Quorum sensing in *Rhizobium* sp. strain NGR234 regulates conjugal transfer (*tra*) gene expression and influences growth rate. *Journal of Bacteriology*, 185(3), 809-822. Doi: DOI: 10.1128/JB.185.3.809–822.2003
  - Heberle, H., Meirelles, G. V., da Silva, F. R., Telles, G. P., and Minghim, R., (2015). InteractiVenn: a web-based tool for the analysis of sets through Venn diagrams. *BMC bioinformatics*, 16(1), 1-7. <https://doi.org/10.1186/s12859-015-0611-3>
  - Hei, Z., Xiang, H., Zhang, J., Liang, K., Zhong, J., Li, M., Lu, Y., 2021. Rice intercropping with water mimosa ( *Neptunia oleracea* Lour .) can facilitate soil N utilization and alleviate apparent N loss *Agriculture , Ecosystems and Environment*. *Agric Ecosyst Environ* 313, 107378. <https://doi.org/10.1016/j.agee.2021.107378>
  - Hinsinger, P., Bengough, A.G., Vetterlein, D., Young, I.M., 2009. Rhizosphere : biophysics , biogeochemistry and ecological relevance 117–152.

- <https://doi.org/10.1007/s11104-008-9885-9>
- Hinsinger, P., Betencourt, E., Bernard, L., Brauman, A., Plassard, C., Shen, J., Tang, X. and Zhang, F., 2011. P for two, sharing a scarce resource: soil phosphorus acquisition in the rhizosphere of intercropped species. *Plant physiology*, 156(3),1078-1086. <https://doi.org/10.1104/pp.111.175331>
  - Hoagland, D. R., and Arnon, D. I., (1941). Physiological aspects of availability of nutrients for plant growth. *Soil Science*, 51(6), 431-444.
  - Huang, X., Chaparro, J.M., Reardon, K.F., Zhang, R., Shen, Q., Vivanco, J.M., 2014. Rhizosphere interactions: root exudates, microbes, and microbial communities. *Botany* 92, 267–275. <https://doi.org/dx.doi.org/10.1139/cjb-2013-0225>
  - Huang, J., Li, Y., Shi, Y., Wang, L., Id, Q.Z., Huang, X., 2019. Effects of nutrient level and planting density on population relationship in soybean and wheat intercropping populations. *PloS One* 14 (12), e0225810. <https://doi.org/10.1371/journal.pone.0225810>.
  - Hütsch, B. W., Augustin, J., & Merbach, W., 2002. Plant rhizodeposition—an important source for carbon turnover in soils. *Journal of plant nutrition and soil science*, 165(4), 397-407. [https://doi.org/10.1002/1522-2624\(200208\)165:4](https://doi.org/10.1002/1522-2624(200208)165:4)
  - Hunter, P.J., Teakle, G.R., Bending, G.D., 2014. Root traits and microbial community interactions in relation to phosphorus availability and acquisition, with particular reference to Brassica. *Front. Plant Sci.* 5 (27) <https://doi.org/10.3389/fpls.2014.00027>
  - Iyer, B., Rajput, M.S., Jog, R., Joshi, E., Bharwad, K., Rajkumar, S., 2016. Organic acid mediated repression of sugar utilization in rhizobia. *Microbiol. Res.* 192, 211–220. <https://doi.org/10.1016/j.micres.2016.07.006>
  - Jacoby, R.P., Koprivova, A. and Kopriva, S., 2021. Pinpointing secondary metabolites that shape the composition and function of the plant microbiome. *Journal of Experimental Botany*, 72(1), pp.57-69. <https://doi.org/10.1093/jxb/eraa424>
  - Jat, H.S., Datta, A., Choudhary, M., Sharma, P.C., Jat, M.L., 2020. Conservation Agriculture : factors and drivers of adoption and scalable innovative practices in Indo-Gangetic plains of India – a review . *Int J Agric Sustain* 0, 1–16. <https://doi.org/10.1080/14735903.2020.1817655>
  - Jensen, E. S., Carlsson, G., & Hauggaard-Nielsen, H., 2020. Intercropping of grain legumes and cereals improves the use of soil N resources and reduces the requirement for synthetic

- fertilizer N: A global-scale analysis. *Agron. Sustain. Dev.* 40(1), 1-9. <https://doi.org/10.1007/s13593-020-0607-x>
- Jones, D.L., Darrah, P.R., 1995. Influx and efflux of organic acids across the soil-root interface of *Zea mays* L. and its implications in rhizosphere C flow. *Plant Soil* 173, 103–109. <https://doi.org/10.1007/BF00155523>.
  - Jones, D.L., 1998. Organic acids in the rhizosphere - a critical review. *Plant Soil* 205, 25–44. <http://www.jstor.org/stable/42949904>.
  - Jones, D.L., Jones, D.L., Hodge, A., Kuzyakov, Y., 2004. Plant and mycorrhizal regulation of rhizodeposition. *New phytologist*, 163(3), 459-480. <https://doi.org/10.1111/j.1469-8137.2004.01130.x>
  - Juma, G., Thiongo, M., Dutaur, L., Rharrabe, K., Marion-Poll, F., Le Ru, B., Magoma, G., Silvain, J.F. and Calatayud, P.A., 2013. Two sugar isomers influence host plant acceptance by a cereal caterpillar pest. *Bull. Entomol. Res*, 103, pp.20-28. [doi:10.1017/S0007485312000387](https://doi.org/10.1017/S0007485312000387)
  - Kamilova, F., Kravchenko, L.V., Shaposhnikov, A.I., Azarova, T., Makarova, N., Lugtenberg, B., 2006. Organic acids, sugars, and L-tryptophane in exudates of vegetables growing on stonewool and their effects on activities of rhizosphere bacteria. *Mol. Plant Microbe Interact.* 19, 250–256. <https://doi.org/10.1094/MPMI-19-0250>.
  - Kennedy, I. R., Choudhury, A. T. M. A., and Kecskés, M. L., 2004. Non-symbiotic bacterial diazotrophs in crop-farming systems: can their potential for plant growth promotion be better exploited? *Soil Biol. Biochem. J.* 36, 1229–1244. [doi: 10.1016/j.soilbio.2004.04.006](https://doi.org/10.1016/j.soilbio.2004.04.006)
  - Kermah, M., Franke, A.C., Adjei-nsiah, S., Ahiabor, B.D.K., Abaidoo, R.C., Giller, K.E., 2017. Field Crops Research Maize-grain legume intercropping for enhanced resource use efficiency and crop productivity in the Guinea savanna of northern Ghana. *F Crop Res* 213, 38–50. <https://doi.org/10.1016/j.fcr.2017.07.008>
  - Khaosaad, T., Staehelin, C., Steinkellner, S., Hage-Ahmed, K., Ocampo, J.A., Garcia-Garrido, J.M. and Vierheilig, H., 2010. The *Rhizobium* sp. strain NGR234 systemically suppresses arbuscular mycorrhizal root colonization in a split-root system of barley (*Hordeum vulgare*). *Physiologia plantarum*, 140(3), pp.238-245. <https://doi.org/10.1111/j.1399-3054.2010.01396.x>

- King, N. D., Hojnacki, D., and O'Brian, M. R., (2000). The *Bradyrhizobium japonicum* proline biosynthesis gene *proC* is essential for symbiosis. *Appl Environ Microbiol.* 66, 5469–5471. doi:10.1128/AEM.66.12.5469-5471.2000
- Kelly, S.J., Muszyński, A., Kawaharada, Y., Hubber, A.M., Sullivan, J.T., Sandal, N., Carlson, R.W., Stougaard, J. and Ronson, C.W., 2013. Conditional requirement for exopolysaccharide in the *Mesorhizobium*–*Lotus* symbiosis. *Molecular plant-microbe interactions*, 26(3), pp.319-329. <https://doi.org/10.1094/MPMI-09-12-0227-R>
- Khatabi, B., Gharechahi, J., Ghaffari, M.R., Liu, D., Haynes, P.A., McKay, M.J., Mirzaei, M. and Salekdeh, G.H., 2019. Plant–Microbe Symbiosis: What Has Proteomics Taught Us?. *Proteomics*, 19(16), p.1800105. <https://doi.org/10.1002/pmic.201800105>
- Knee, E.M., Gong, F.C., Gao, M., Teplitski, M., Jones, A.R., Foxworthy, A., Mort, A.J., Bauer, W.D., 2001. Root mucilage from pea and its utilization by rhizosphere bacteria as a sole carbon source. *Mol. Plant Microbe Interact.* 14, 775–784. <https://doi.org/10.1094/MPMI.2001.14.6.775>
- Kleijn, D., Bommarco, R., Fijen, T.P.M., Garibaldi, L.A., Potts, S.G., Putten, W.H. Van Der., 2019. Ecological Intensification : Bridging the Gap between Science and Practice. *Trends Ecol Evol* 34, 154–166. <https://doi.org/10.1016/j.tree.2018.11.002>
- Kobayashi, H., Graven, Y.N., Broughton, W.J. and Perret, X., 2004. Flavonoids induce temporal shifts in gene-expression of nod-box controlled loci in *Rhizobium* sp. NGR234. *Molecular microbiology*, 51(2), pp.335-347. <https://doi.org/10.1046/j.1365-2958.2003.03841.x>
- Konkolewska, A., Piechalak, A., Ciszewska, L., Antos-Krzemińska, N., Skrzypczak, T., Hanć, A., Sitko, K., Małkowski, E., Barańkiewicz, D., Małecka, A., 2020. Combined use of companion planting and PGPR for the assisted phytoextraction of trace metals (Zn, Pb, Cd). *Environ. Sci. Pollut. Res.* 27, 13809–13825. <https://doi.org/10.1007/s11356-020-07885-3>
- Krishnappa, R., Hussain, L.S., 2014. Phosphorus acquisition from deficient soil: involvement of organic acids and acid phosphatase in pigeon pea (*Cajanus cajan* L. mills sp). *Indian J. Plant Physiol.* 19 (3), 197–204. <https://doi.org/10.1007/s40502-014-0101-z>.
- Krysciak, D., Schmeisser, C., Preuss, S., Riethausen, J., Quitschau, M., Grond, S. and Streit, W.R., 2011. Involvement of multiple loci in quorum quenching of autoinducer I

- molecules in the nitrogen-fixing symbiont *Rhizobium* (*Sinorhizobium*) sp. strain NGR234. *Applied and environmental microbiology*, 77(15), pp.5089-5099. <https://doi.org/10.1128/AEM.00112-11>
- Krysciak, D., Grote, J., Rodriguez Orbegoso, M., Utpatel, C., Förstner, K.U., Li, L., Schmeisser, C., Krishnan, H.B. and Streit, W.R., 2014. RNA sequencing analysis of the broad-host-range strain *Sinorhizobium fredii* NGR234 identifies a large set of genes linked to quorum sensing-dependent regulation in the background of a *traI* and *ngfI* deletion mutant. *Applied and environmental microbiology*, 80(18), pp.5655-5671. <https://doi.org/10.1128/AEM.01835-14>
  - Kuijken, R.C.P., Snel, J.F.H., Heddes, M.M., Bouwmeester, H.J., Marcelis, L.F.M., 2015. The importance of a sterile rhizosphere when phenotyping for root exudation. *Plant and Soil* 387, 131-142. <https://doi.org/10.1007/s11104-014-2283-6>
  - Kuznetsov, V.V., Rakitin, V.Y. and Zholkevich, V.N., 1999. Effects of preliminary heat-shock treatment on accumulation of osmolytes and drought resistance in cotton plants during water deficiency. *Physiologia Plantarum*, 107(4), pp.399-406.
  - Kuzyakov, Y. and Schneckenberger, K., 2004. Review of estimation of plant rhizodeposition and their contribution to soil organic matter formation. *Archives of Agronomy and Soil Science*, 50(1), pp.115-132. <https://doi.org/10.1080/03650340310001627658>
  - Lamrabet, Y., Bellogín, R.A., Cubo, T., Espuny, R., Gil, A., Krishnan, H.B., Megias, M., Ollero, F.J., Pueppke, S.G., Ruiz-Sainz, J.E. and Spaink, H.P., 1999. Mutation in GDP-fucose synthesis genes of *Sinorhizobium fredii* alters Nod factors and significantly decreases competitiveness to nodulate soybeans. *Molecular plant-microbe interactions*, 12(3), pp.207-217. <https://doi.org/10.1094/MPMI.1999.12.3.207>
  - Latati, M., Bargaz, A., Belarbi, B., Lazali, M., Benlahrech, S., Tellah, S., Kaci, G., Drevon, J.J. and Ounane, S.M., 2016. The intercropping common bean with maize improves the rhizobial efficiency, resource use and grain yield under low phosphorus availability. *European journal of agronomy*, 72, 80-90. <https://doi.org/10.1016/j.eja.2015.09.015>
  - Layek, J., Das, A., Mitran, T., Nath, C., Meena, R. S., Yadav, G., Lal, R., 2018. Cereal+ legume intercropping: an option for improving productivity and sustaining soil health.

- In Legumes for Soil Health and Sustainable Management (pp. 347-386). Springer, Singapore. <https://doi.org/10.1007/978-981-13-0253-4>
- Lei, Z., Huhman, D.V. and Sumner, L.W., 2011. Mass spectrometry strategies in metabolomics. *Journal of Biological Chemistry*, 286(29), pp.25435-25442. <https://doi.org/10.1074/jbc.R111.238691>
  - Lee, H.I., Lee, J.H., Park, K.H., Sangurdekar, D., Chang, W.S., 2012. Effect of soybean coumestrol on *Bradyrhizobium japonicum* nodulation ability, biofilm formation, and transcriptional profile. *Appl. Environ. Microbiol.* 78, 2896–2903. <https://doi.org/10.1128/AEM.07336-11>.
  - Le Strange, K. K., Bender, G. L., Djordjevic, M. A., Rolfe, B. G., & Redmond, J. W., 1990. The Rhizobium strain NGR234 nodD1 gene product responds to activation by the simple phenolic compounds vanillin and isovanillin present in wheat seedling extracts. *Molecular Plant-Microbe Interactions*, 3(4), 214-220.
  - Li, X., Mu, Y., Cheng, Y., Liu, X., Nian, H., 2013a. Effects of intercropping sugarcane and soybean on growth, rhizosphere soil microbes, nitrogen, and phosphorus availability. *Acta Physiol. Plant.* 35, 1113–1119. <https://doi.org/10.1007/s11738-012-1148-y>
  - Li, Y., Tian, C.F., Chen, W.F., Wang, L., Sui, X.H. and Chen, W.X., 2013. High-resolution transcriptomic analyses of *Sinorhizobium* sp. NGR234 bacteroids in determinate nodules of *Vigna unguiculata* and indeterminate nodules of *Leucaena leucocephala*. *PloS one*, 8(8), p.e70531. <https://doi.org/10.1371/journal.pone.0070531>
  - Li, B., Li, Y.-Y., Wu, H.-M., Zhang, F.-F., Li, C.-J., Li, X.-X., Lambers, H., Li, L., 2016. Root exudates drive interspecific facilitation by enhancing nodulation and N<sub>2</sub> fixation. *Proc Natl Acad Sci* 113, 6496–6501. <https://doi.org/10.1073/pnas.1523580113>
  - Li, C., Hoffland, E., Kuyper, T.W., Yu, Y., Zhang, C., Li, H., Zhang, F., van der Werf, W., 2020. Syndromes of production in intercropping impact yield gains. *Nat Plants* 6, 653–660. <https://doi.org/10.1038/s41477-020-0680-9>
  - Li, L., Li, S., Sun, J., Zhou, L., Bao, X., Zhang, H., Zhang, F., 2007. Diversity enhances agricultural productivity via rhizosphere phosphorus facilitation on phosphorus-deficient soils 104, 3–7.
  - Li, Q., Chen, J., Wu, L., Luo, X., Li, N., Arafat, Y., Lin, S., Lin, W., 2018. Belowground interactions impact the soil bacterial community, soil fertility, and crop yield in

- maize/peanut intercropping systems. *Int J Mol Sci* 19, 1–16. <https://doi.org/10.3390/ijms19020622>
- Li, T., Zhan, Z., Lin, Y., Lin, M., Xie, Q., Chen, Y., He, C., Tao, J. and Li, C., 2019. Biosynthesis of Amino Acids in *Xanthomonas oryzae* pv. *oryzae* Is Essential to Its Pathogenicity. *Microorganisms*, 7(12), p.693. <https://doi.org/10.3390/microorganisms7120693>
  - Liang, Y., Cao, Y., Tanaka, K., Thibivilliers, S., Wan, J., Choi, J., ho Kang, C., Qiu, J. and Stacey, G., 2013. Nonlegumes respond to rhizobial Nod factors by suppressing the innate immune response. *Science*, 341(6152), pp.1384-1387. <https://doi.org/10.1126/science.1242736>
  - Ling, N., Raza, W., Ma, J., Huang, Q., Shen, Q., 2011. Identification and role of organic acids in watermelon root exudates for recruiting *Paenibacillus polymyxa* SQR-21 in the rhizosphere. *Eur. J. Soil Biol.* 47, 374–379. <https://doi.org/10.1016/j>
  - Liu, S., Hu, X., Lohrke, S.M., Baker, C.J., Buyer, S., Souza, J.T. De, Roberts, D.P., 2007. Role of *sdhA* and *pfkA* and catabolism of reduced carbon during colonization of cucumber roots by *Enterobacter cloacae*. *Microbiology* 153, 3196–3209. <https://doi.org/10.1099/mic.0.2006/005538-0>.
  - Liu, X, Qiu, W, Rao, B, Cao, Y, Fang, X, Yang, J, Jiang, G, Zhong, Z, Zhu, J 2019a. Bacterioferritin comigratory protein is important in hydrogen peroxide resistance, nodulation, and nitrogen fixation in *Azorhizobium caulinodans*. *Arch Microbiol* 201(6):823–831. <https://doi.org/10.1007/s00203-019-01654-8>
  - Liu, Y., Yin, X., Xiao, J., Tang, L., Zheng, Y., 2019b. Interactive influences of intercropping by nitrogen on flavonoid exudation and nodulation in faba bean. *Sci. Rep.* 9, 4818. <https://doi.org/10.1038/s41598-019-41146-9>.
  - Lithourgidis, A. S., Dordas, C. A., Damalas, C. A., & Vlachostergios, D. 2011. Annual intercrops: an alternative pathway for sustainable agriculture. *Aust. J. Crop. Sci.* 5, 396–410. ISSN: 1835-2693.
  - Livak, K.J. and Schmittgen, T.D., 2001. Analysis of relative gene expression data using real-time quantitative PCR and the 2<sup>-</sup> ΔΔCT method. *methods*, 25(4), pp.402-408. <https://doi.org/10.1006/meth.2001.1262>
  - Lodwig, E. M., Hosie, A. H., Bourdès, A., Findlay, K., Allaway, D., Karunakaran, R.,

- Downie, J. A., Poole, P. S. 2003. Amino-acid cycling drives nitrogen fixation in the legume-*Rhizobium* symbiosis. *Nature*. 422, 722-726. doi: 10.1038/nature01527. PMID: 12700763
- Lugtenberg, B., & Kamilova, F., 2009. Plant-growth-promoting rhizobacteria. *Annual review of microbiology*, 63, 541-556. <https://doi.org/10.1146/annurev.micro.62.081307.162918>
  - Lv, J., Dong, Y., Dong, K., Zhao, Q., Yang, Z. and Chen, L., 2020. Intercropping with wheat suppressed Fusarium wilt in faba bean and modulated the composition of root exudates. *Plant and Soil*, pp.1-12. <https://doi.org/10.1007/s11104-019-04413-2>
  - Macias-benitez, S., Garcia A.M., Jimenez, P.C., Gonzalez, J.M., Moral, M.T., Rubio, J.P., Ryan, P., 2020. Rhizospheric organic acids as Biostimulants : monitoring feedbacks on soil microorganisms and biochemical properties. *Front. Plant Sci.* 11 (633), 1–16. <https://doi.org/10.3389/fpls.2020.00633>.
  - Malézieux, E., Crozat, Y., Dupraz, C., Laurans, M., Makowski, D., Ozier-Lafontaine, H., Valantin-Morison, M. 2009. Mixing plant species in cropping systems: concepts, tools and models: a review. *Sustainable agriculture*, 329-353. [https://doi.org/10.1007/978-90-481-2666-8\\_22](https://doi.org/10.1007/978-90-481-2666-8_22)
  - Maitra, S., Hossain, A., Brestic, M., Skalicky, M., Ondrisik, P., Gitari, H., Brahmachari, K., Shankar, T., Bhadra, P., Palai, J.B., Jena, J., Bhattacharya, U., Duvvada, S.K., Lalichetti, S., Sairam, M., 2021. Intercropping—A Low Input Agricultural Strategy for Food and Environmental Security. *Agronomy* 11, 343. <https://doi.org/10.3390/agronomy11020343>
  - Maitra, S., Palai, J.B., Manasa, P., Kumar, D.P., 2019. Potential of Intercropping System in Sustaining Crop Productivity 12, 39–45. <https://doi.org/10.30954/0974-1712.03.2019.7>
  - Malobane, M.E., Nciizah, A.D., Mudau, F.N., 2020. Tillage , Crop Rotation and Crop Residue Management Effects on Nutrient Availability in a Sweet Sorghum-Based Cropping System in Marginal Soils of South Africa. *Agronomy*, 10(6), 776. <https://doi.org/10.3390/agronomy10060776>
  - Malik, K. A., Mirza, M. S., Hassan, U., Mehnaz, S., Rasul, G., Haurat, J.. 2002. The role of plant-associated beneficial bacteria in rice-wheat cropping system, in *Biofertilisers in Action*, eds I. R. Kennedy and A. Choudhury (Canberra: Rural Industries Research and

- Development Corporation), 73–83.
- Martin-Guay, M.O., Paquette, A., Dupras, J. and Rivest, D., 2018. The new green revolution: sustainable intensification of agriculture by intercropping. *Science of the Total Environment*, 615, 767-772. <https://doi.org/10.1016/j.scitotenv.2017.10.024>.
  - Mathimaran, N., Jegan, S., 2020. Intercropping Transplanted Pigeon Pea With Finger Millet : Arbuscular Mycorrhizal Fungi and Plant Growth Promoting Rhizobacteria Boost Yield While Reducing Fertilizer Input 4, 1–12. <https://doi.org/10.3389/fsufs.2020.00088>
  - Matilla, M.A., Krell, T., 2018. The effect of bacterial chemotaxis on host infection and pathogenicity. *FEMS Microbiol. Rev.* 42, fux052. <https://doi:10.1093/femsre/fux052>.
  - Martínez-Gil, M., Yousef-Coronado, F., & Espinosa-Urgel, M., June 2010. LapF, the second largest *Pseudomonas putida* protein, contributes to plant root colonization and determines biofilm architecture. *Molecular Microbiology*, 77, 549e561.
  - McNear Jr., D. H., 2013 The Rhizosphere - Roots, Soil and Everything In Between. *Nature Education Knowledge* 4(3):1
  - Marschner, H., 1995. *Mineral Nutrition of Higher Plants*. Academic Press, London.
  - Masson-Boivin, C. and Sachs, J.L., 2018. Symbiotic nitrogen fixation by rhizobia—the roots of a success story. *Current opinion in plant biology*, 44, pp.7-15. <https://doi.org/10.1016/j.pbi.2017.12.001>
  - Mendes, R., Garbeva, P., Raaijmakers, J.M., 2013. The rhizosphere microbiome: significance of plant beneficial, plant pathogenic, and human pathogenic microorganisms 37, 634–663. <https://doi.org/10.1111/1574-6976.12028>
  - Meng, L., Zhang, A., Wang, F., Han, X., Wang, D., Li, S., 2015. Arbuscular mycorrhizal fungi and rhizobium facilitate nitrogen uptake and transfer in soybean/maize intercropping system. *Front. Plant Sci.* 6, 1–10. <https://doi.org/10.3389/fpls.2015.00339>.
  - Mommer, L., Hinsinger, P., Prigent-Combaret, C., Visser, E.J.W., 2016a. Advances in the rhizosphere: stretching the interface of life. *Plant Soil* 407, 1–8. <https://doi.org/10.1007/s11104-016-3040-9>
  - Mommer, L., Kirkegaard, J., van Ruijven, J., 2016b. Root-Root Interactions: Towards A Rhizosphere Framework. *Trends Plant Sci* 21, 209–217. <https://doi.org/10.1016/j.tplants.2016.01.009>
  - Nanda, A.K., Andrio, E., Marino, D., Pauly, N. and Dunand, C., 2010. Reactive oxygen

- species during plant-microorganism early interactions. *Journal of integrative plant biology*, 52(2), pp.195-204.<https://doi.org/10.1111/j.1744-7909.2010.00933.x>
- Naudin, C., Werf, H.M.G. Van Der, Jeuffroy, M., 2014. Life cycle assessment applied to pea-wheat intercrops : A new method for handling the impacts of co-products. *J Clean Prod* 73, 80–87. <https://doi.org/10.1016/j.jclepro.2013.12.029>
  - Naveed, M., Mitter, B., Yousaf, S., Pastar, M., Afzal, M., Sessitsch, A., 2013. The endophyte *Enterobacter* sp. FD17: a maize growth enhancer selected based on rigorous testing of plant beneficial traits and colonization characteristics. *Biol. Fertil. Soils* 50, 249–262. <https://doi.org/10.1007/s00374-013-0854-y>
  - Naveed, M., Brown, L.K., Raffan, A.C., George, T.S., Bengough, A.G., Roose, T., Sinclair, I., Koebernick, N., Cooper, L., Hackett, C.A. and Hallett, P.D., 2017. Plant exudates may stabilize or weaken soil depending on species, origin and time. *Eur. J. Soil Sci.*, 68(6), pp.806-816.[doi: 10.1111/ejss.12487](https://doi.org/10.1111/ejss.12487)
  - Neumann, G., Römheld, V., 1999. Root excretion of carboxylic acids and protons in phosphorus-deficient plants. *Plant Soil* 211, 121–130. <https://doi.org/10.1023/A:1004380832118>
  - Noirot-Gros, M., Shinde, S., Larsen E., P., Zerbs, S., Korajczyk J., P., Kemner M., K., Noirot H., P., 2018. Dynamics of aspen roots colonization by pseudomonads reveals strain-specific and mycorrhizal-specific patterns of biofilm formation. *Front. Microbiol.* 9, 853. <https://doi.org/10.3389/fmicb.2018.00853>
  - Nwachukwu, B.C., Ayangbenro, A.S., Babalola, O.O., 2021. Elucidating the rhizosphere associated bacteria for environmental sustainability. *Agric* 11, 1–18. <https://doi.org/10.3390/agriculture11010075>
  - Oburger, E., Dellmour, M., Hann, S., Wieshammer, G., Puschenreiter, M., Wenzel, W.W., 2013. Evaluation of a novel tool for sampling root exudates from soil-grown plants compared to conventional techniques. *Environ. Exp. Bot.* 87, 235–247. <https://doi.org/10.1016/j.envexpbot.2012.11.007>
  - Oburger, E., Jones, D.L., 2018. Sampling root exudates—mission impossible? *Rhizosphere* 6, 116–133. <https://doi.org/10.1016/j.rhisph.2018.06.004>.
  - Oelhermann, M., Echarte, L., Vachon, K., Dubois, C., 2009 The role of complex agroecosystems in sequestering carbon and mitigating global warming. *Earth Environ Sci*

- 6:20–31. IOP Publishing.
- Ogawa, J. and Long, S.R., 1995. The *Rhizobium meliloti* groELc locus is required for regulation of early nod genes by the transcription activator NodD. *Genes & Development*, 9(6), pp.714-729. <https://doi.org/10.1101/gad.9.6.714>
  - Okazaki, S., Hattori, Y. and Saeki, K., 2007. The *Mesorhizobium loti* purB gene is involved in infection thread formation and nodule development in *Lotus japonicus*. *Journal of bacteriology*, 189(22), pp.8347-8352. <https://doi.org/10.1128/JB.00788-07>
  - Orrell, P., Bennett, A.E., 2013. How can we exploit above–belowground interactions to assist in addressing the challenges of food security? *Front Plant Sci* 4, 1–11. <https://doi.org/10.3389/fpls.2013.00432>
  - Patel, K Janki., Archana, G., 2018. Engineered production of 2, 4-diacetylphloroglucinol in the diazotrophic endophytic bacterium *Pseudomonas* sp. WS5 and its beneficial effect in multiple plant-pathogen systems. *Appl. Soil Ecol.* 124, 34–44. <https://doi.org/10.1016/j.apsoil.2017.10.008>
  - Pathak, H., 2010. Mitigating greenhouse gas and nitrogen loss with improved fertilizer management in rice: quantification and economic assessment. *Nutrient cycling in agroecosystems*, 87(3), 443-454. <https://doi.org/10.1007/s10705-010-9350-z>
  - Pawlak, K., & Kołodziejczak, M., 2020. The Role of Agriculture in Ensuring Food Security in Developing Countries : Considerations in the Context of the Problem of Sustainable Food Production. *Sustainability*, 12(13), 5488.<https://doi.org/10.3390/su12135488>
  - Pereg, L., & McMillan, M. 2020. Nitrogen-Cycling Communities in Organically Amended Versus Conventionally Managed Agricultural Soil. In *Carbon and Nitrogen Cycling in Soil* (pp. 377-398). Springer, Singapore. [https://doi.org/10.1007/978-981-13-7264-3\\_11](https://doi.org/10.1007/978-981-13-7264-3_11). <https://doi.org/10.1007/s11103-015-0337-7>
  - Pérez-Jaramillo, J.E., Mendes, R. and Raaijmakers, J.M., 2016. Impact of plant domestication on rhizosphere microbiome assembly and functions. *Plant molecular biology*, 90(6), pp.635-644.
  - Pérez-Montaña, F., Guasch-Vidal, B., González-Barroso, S., López-Baena, F.J., Cubo, T., Ollero, F.J., Gil-Serrano, A.M., Rodríguez-Carvajal, M.Á., Bellogín, R.A. and Espuny, M.R., 2011. Nodulation-gene-inducing flavonoids increase overall production of autoinducers and expression of N-acyl homoserine lactone synthesis genes in

- rhizobia. *Research in microbiology*, 162(7), pp.715-723. <https://doi.org/10.1016/j.resmic.2011.05.002>
- Philippot, L., Raaijmakers, J.M., Lemanceau, P., Van Der Putten, W.H., 2013. Going back to the roots: The microbial ecology of the rhizosphere. *Nat Rev Microbiol* 11, 789–799. <https://doi.org/10.1038/nrmicro3109>
  - Piromyoo, P., Songwattana, P., Greetatorn, T., Okubo, T., Kakizaki, K.C., Prakamhang, J., Tittabutr, P., Boonkerd, N., Teaumroong, N. and Minamisawa, K., 2015. The type III secretion system (T3SS) is a determinant for rice-endophyte colonization by non-photosynthetic *Bradyrhizobium*. *Microbes and environments*, p.ME15080. <https://doi.org/10.1264/jsme2.ME15080>
  - Poole, P.S., Blyth, A., Reid, C.J. and Walters, K., 1994. myo-Inositol catabolism and catabolite regulation in *Rhizobium leguminosarum* bv. *viciae*. *Microbiology*, 140(10), pp.2787-2795.
  - Poole, P., Ramachandran, V. and Terpolilli, J., 2018. Rhizobia: from saprophytes to endosymbionts. *Nature Reviews Microbiology*, 16(5), p.291. [doi:10.1038/nrmicro.2017.171](https://doi.org/10.1038/nrmicro.2017.171)
  - Prell, J., White, J.P., Bourdes, A., Bunnell, S., Bongaerts, R.J. and Poole, P.S., 2009. Legumes regulate *Rhizobium* bacteroid development and persistence by the supply of branched-chain amino acids. *Proceedings of the National Academy of Sciences*, 106(30), pp.12477-12482. <https://doi.org/10.1073/pnas.0903653106>
  - Pueppke, S.G. and Broughton, W.J., 1999. *Rhizobium* sp. strain NGR234 and *R. fredii* USDA257 share exceptionally broad, nested host ranges. *Molecular Plant-Microbe Interactions*, 12(4), pp.293-318. <https://doi.org/10.1094/MPMI.1999.12.4.293>
  - Rajendran, G., Mistry, S., Desai J., A., Archana, G., 2007. Functional expression of *Escherichia coli* *fhuA* gene in *Rhizobium* spp. of *Cajanus cajan* provides growth advantage in the presence of Fe<sup>3+</sup>: ferrichrome as iron source. *Arch. Microbiol.* 187, 257–264. <https://doi.org/10.1007/s00203-006-0191-8>
  - Ramachandran, V.K., East, A.K., Karunakaran, R., Downie, J.A. and Poole, P.S., 2011. Adaptation of *Rhizobium leguminosarum* to pea, alfalfa and sugar beet rhizospheres investigated by comparative transcriptomics. *Genome biology*, 12(10), pp.1-12. <https://doi.org/10.1186/gb-2011-12-10-r106>

- Ramakrishna, W., Yadav, R. and Li, K., 2019. Plant growth promoting bacteria in agriculture: two sides of a coin. *Applied Soil Ecology*, 138, pp.10-18. <https://doi.org/10.1016/j.apsoil.2019.02.019>
- Relic, B., Fellay, R., Lewin, A., Perret, X., Price, N.P.J., Rochepeau, P. and Broughton, W.J., 1993. nod genes and Nod factors of *Rhizobium* species NGR234. In *New horizons in nitrogen fixation* (pp. 183-189). Springer, Dordrecht. [https://doi.org/10.1007/978-94-017-2416-6\\_22](https://doi.org/10.1007/978-94-017-2416-6_22)
- Reddy, P.M., Ladha, J.K., So, R.B., Hernandez, R.J., Ramos, M.C., Angeles, O.R., Dazzo, F.B. and de Bruijn, F.J., 1997. Rhizobial communication with rice roots: induction of phenotypic changes, mode of invasion and extent of colonization. *Plant and soil*, 194(1), pp.81-98. <https://doi.org/10.1023/A:1004243915997>
- Reyes-Pérez, A., del Carmen Vargas, M., Hernández, M., Aguirre-von-Wobeser, E., Pérez-Rueda, E. and Encarnacion, S., 2016. Transcriptomic analysis of the process of biofilm formation in *Rhizobium etli* CFN42. *Archives of microbiology*, 198(9), pp.847-860. <https://doi.org/10.1007/s00203-016-1241-5>
- Rezaei-Chiyaneh, E., Amirnia, R., Amani, M., Machiani, M.A., Javanmard, A., Maggi, F., Morshedloo, M.R., 2019. Intercropping fennel (*Foeniculum vulgare* L.) with common bean (*Phaseolus vulgaris* L.) as affected by PGPR inoculation : a strategy for improving yield, essential oil, and fatty acid composition. *Sci. Hortic. (Amsterdam)* 261, 108951. <https://doi.org/10.1016/j.scienta.2019.108951>
- Rinaudi, L.V. and Giordano, W., 2010. An integrated view of biofilm formation in rhizobia. *FEMS microbiology letters*, 304(1), pp.1-11. <https://doi.org/10.1111/j.1574-6968.2009.01840.x>
- Robertsen, B. K., Aman, P., Darvill, A. G., McNeil, M. & Albersheim, P., (1981). Host-symbiont interactions : V. The structure of acidic extracellular polysaccharides secreted by *Rhizobium leguminosarum* and *Rhizobium trifolii*. *Plant Physiol* 67, 389-400. <https://doi.org/10.1104/pp.67.3.389>
- Rodriguez-Navarro, D.N., Rodriguez-Navarro, D.N., Dardanelli, M.S., Dardanelli, M.S., Ruiz-Sainz, J.E., and Ruiz-Sainz, J.E., (2007) Attachment of bacteria to the roots of higher plants. *FEMS Microbiol Lett* 272: 127–136. <https://doi.org/10.1111/j.1574-6968.2007.00761.x>

- Rodriguez, C., Carlsson, G., Englund, J.E., Flöhr, A., Pelzer, E., Jeuffroy, M.H., Makowski, D., Jensen, E.S., 2020. Grain legume-cereal intercropping enhances the use of soil-derived and biologically fixed nitrogen in temperate agroecosystems. A meta-analysis. *Eur J Agron* 118. <https://doi.org/10.1016/j.eja.2020.126077>
- Romero-Perdomo, F., Abril, J., Camelo, M., Moreno-Galván, A., Pastrana, I., Rojas-Tapias, D., & Bonilla, R., 2017. *Azotobacter chroococcum* as a potentially useful bacterial biofertilizer for cotton (*Gossypium hirsutum*): Effect in reducing N fertilization. *Revista Argentina de microbiología*, 49(4), 377-383.
- Rosenblueth, M., Martínez-Romero, E., 2004. *Rhizobium etli* maize populations and their competitiveness for root colonization. *Arch. Microbiol.* 181, 337–344. <https://doi.org/10.1007/s00203-004-0661-9>
- Ryan, P. R., Delhaize, E., & Jones, D. L. 2001. Function and mechanism of organic anion exudation from plant roots. *Annual review of plant biology*, 52(1), 527-560. <https://doi.org/10.1146/annurev.arplant.52.1.527>
- Sambrook, J., & Russell, D. W., 2001. *Molecular Cloning: a Laboratory Manual*, third ed., vol. 2. Cold Spring Harbor Laboratory Press, New York.
- Saharan, K., Schütz, L., Kahmen, A., Wiemken, A., Boller, T., Mathimaran, N., 2018. Finger millet growth and nutrient uptake is improved in intercropping with pigeon pea through “biofertilization” and “bioirrigation” mediated by arbuscular mycorrhizal fungi and plant growth promoting rhizobacteria. *Front Environ Sci* 6, 1–11. <https://doi.org/10.3389/fenvs.2018.00046>
- Schoebitz, M., Castillo, D., Jorquera, M., Roldan, A., 2020. Responses of microbiological soil properties to intercropping at different planting densities in an acidic Andisol. *Agronomy* 10 (781). <https://doi.org/10.3390/agronomy10060781>.
- Sekaran, U., Loya, J.R., Abagandura, G.O., Subramanian, S., Owens, V., Kumar, S., 2019. Intercropping of kura clover (*Trifolium ambiguum* M . Bieb ) with prairie cordgrass (*Spartina pectinata* link .) enhanced soil biochemical activities and microbial community structure. *Appl Soil Ecol* 103427. <https://doi.org/10.1016/j.apsoil.2019.103427>
- Shao, Z.Q., Zheng, C.C., Postma, J.A., LU, W.L., Qiang, G.A.O., Gao, Y.Z. and Zhang, J.J., 2021. Nitrogen acquisition, fixation and transfer in maize/alfalfa intercrops are increased through root contact and morphological responses to interspecies

- competition. *Journal of Integrative Agriculture*, 20(8), 2240-2254. [https://doi.org/10.1016/S2095-3119\(20\)63330-5](https://doi.org/10.1016/S2095-3119(20)63330-5)
- Sharma, R., Barot, K., Archana, G., 2019. Root colonization by heavy metal resistant *Enterobacter* sp. C1D and its influence on metal-induced oxidative stress on *Cajanus cajan*. *J. Sci. Food Agric.* 100 (4) <https://doi.org/10.1002/jsfa.10161>.
  - Sibhatu KT, Qaim M 2017. Rural food security , subsistence agriculture , and seasonality. *PloS one*, 12(10), e0186406. <https://doi.org/10.1371/journal.pone.0186406>
  - Silva, F.B., Winck, B., Borges, C.S., Santos, F.L., Bataioli, R.D., Backes, T., Bassani, V.L., Borin, J.B.M., Frazzon, A.P.G., S´a, E.L.S., 2020. Native rhizobia from southern Brazilian grassland promote the growth of grasses. *Rhizosphere* 16. <https://doi.org/10.1016/j.rhisph.2020.100240>
  - Skøt, L., and Egsgaard, H., 1984. Identification of ononitol and O-methyl-*scyllo*-inositol in pea root nodules. *Planta*. 161, 32–36. doi:10.1007/BF00951457
  - Solanki, M.K., Wang, F., Li, C., 2020. Impact of Sugarcane – Legume Intercropping on Diazotrophic Microbiome. *Sugar Tech* 22, 52–64. <https://doi.org/10.1007/s12355-019-00755-4>
  - Solanki, M.K., Wang, F., Wang, Z., Li, C., Lan, T., Singh, R.K., Singh, P., Yang, L., Li, Y., 2019. Rhizospheric and endospheric diazotrophs mediated soil fertility intensification in sugarcane-legume intercropping systems 1911–1927.
  - Staehelin, C., Forsberg, L.S., D’Haeze, W., Gao, M.Y., Carlson, R.W., Xie, Z.P., Pellock, B.J., Jones, K.M., Walker, G.C., Streit, W.R. and Broughton, W.J., 2006. Exo-oligosaccharides of *Rhizobium* sp. strain NGR234 are required for symbiosis with various legumes. *Journal of Bacteriology*, 188(17), pp.6168-6178. <https://doi.org/10.1128/JB.00365-06>
  - Stagnari, F., Maggio, A., Galieni, A., Pisante, M., 2017. Multiple benefits of legumes for agriculture sustainability: an overview. *Chem Biol Technol Agric* 4(1), 1-13. <https://doi.org/10.1186/s40538-016-0085-1>.
  - Struik, P.C., & Kuyper, T. W., 2017. Sustainable intensification in agriculture : the richer shade of green . A review. *Agronomy for Sustainable Development*, 37(5), 1-15 <https://doi.org/10.1007/s13593-017-0445-7>
  - Subrahmanyam, G., Sharma, R.K., Kumar, G.N., Archana, G., 2018. *Vigna radiata* var.

- Gm4 plant growth enhancement and root colonization by a multi-metal-resistant plant growth-promoting bacterium *Enterobacter* sp. C1D in Cr(VI)-amended soils. *Pedosphere* 28, 144–156. [https://doi.org/10.1016/S1002-0160\(17\)60448-X](https://doi.org/10.1016/S1002-0160(17)60448-X).
- Sun, Y., Liu, M., Song, M., Tian, Y., Xu, X., 2020. Interactions between neighboring native and alien species are modulated by nitrogen availability. *Rhizosphere* 16, 100242. <https://doi.org/10.1016/j.rhisph.2020.100242>.
  - Szklarczyk, D., Franceschini, A., Wyder, S., Forslund, K., Heller, D., Huerta-Cepas, J., Simonovic, M., Roth, A., Santos, A., Tsafou, K.P. and Kuhn, M., 2015. STRING v10: protein–protein interaction networks, integrated over the tree of life. *Nucleic acids research*, 43(D1), pp.D447-D452. <https://doi.org/10.1093/nar/gku1003>
  - Tang, X., Zhong, R., Jiang, J., He, L., Huang, Z., Shi, G., Wu, H., 2020. Cassava / peanut intercropping improves soil quality via rhizospheric microbes increased available nitrogen contents. *BMC biotechnology*, 20(1), 1–11. <https://doi.org/10.1186/s12896-020-00606-1>
  - Tang, X., Zhang, C., Yu, Y., Shen, J., van der Werf, W. and Zhang, F., 2021. Intercropping legumes and cereals increases phosphorus use efficiency; a meta-analysis. *Plant and Soil*, 460(1), pp.89-104. <https://doi.org/10.1007/s11104-020-04768-x>
  - Taschen, E., Amenc, L., Tournier, E., Deleporte, P., Malagoli, P., Fustec, J., Bru, D., Philippot, L. and Bernard, L., 2017. Cereal-legume intercropping modifies the dynamics of the active rhizospheric bacterial community. *Rhizosphere*, 3, 191-195. <https://doi.org/10.1016/j.rhisph.2017.04.011>
  - Taylor, P., Otani, T., Ae, N., Tanaka, H., 1996. Phosphorus (P) uptake mechanisms of crops grown in soils with low P status (ii . significance of organic acids in root exudates of pigeon pea). *J. Soil Sci. Plant Nutr.* 42 (3), 553–560. <https://doi.org/10.1080/00380768.1996.10416324>.
  - Theunis, M., Kobayashi, H., Broughton, W. J., & Prinsen, E., 2004. Flavonoids, NodD1, NodD2, and nod-box NB15 modulate expression of the y4wEFG locus that is required for indole-3-acetic acid synthesis in *Rhizobium* sp. strain NGR234. *Molecular Plant-Microbe Interactions*, 17(10), 1153-1161.
  - Tian, B., Pei, Y., Huang, W., Ding, J. and Siemann, E., 2021. Increasing flavonoid concentrations in root exudates enhance associations between arbuscular mycorrhizal fungi and an invasive plant. *The ISME Journal*, pp.1-12. <https://doi.org/10.1038/s41396-021->

00894-1

- Thilakarathna, M.S., McElroy, M.S., Chapagain, T., Papadopoulos, Y.A., Raizada, M.N., 2016. Erratum to: Belowground nitrogen transfer from legumes to non-legumes under managed herbaceous cropping systems. A review. *Agron Sustain Dev* 36. <https://doi.org/10.1007/s13593-016-0403-9>
- Tilman, D., Balzer, C., Hill, J., & Befort, B. L., 2011. Global food demand and the sustainable intensification of agriculture. *Proceedings of the national academy of sciences*, 108(50), 20260-20264. <https://doi.org/10.1073/pnas.1116437108>
- Tilman, D., 2020. Benefits of intensive agricultural intercropping. *Nat Plants* 6, 604–605. <https://doi.org/10.1038/s41477-020-0677-4>
- Toukabri, W., Ferchichi, N., Hlel, D., Jadlaoui, M., Kheriji, O., Mhamdi, R. and Trabelsi, D., 2021. Response of intercropped barley and fenugreek to mono-and co-inoculation with *Sinorhizobium meliloti* F42 and *Variovorax paradoxus* F310 under contrasting agroclimatic regions. *Archives of Microbiology*, 203(4), pp.1657-1670. <https://doi.org/10.1007/s00203-020-02180-8>
- Vandermeer, J., van Noordwijk, M., Anderson, J., Ong, C., & Perfecto, I. 1998. Global change and multi-species agroecosystems: concepts and issues. *Agriculture, Ecosystems & Environment*, 67(1), 1-22. [https://doi.org/10.1016/S0167-8809\(97\)00150-3](https://doi.org/10.1016/S0167-8809(97)00150-3)
- van Dam, N.M. and Bouwmeester, H.J., 2016. Metabolomics in the rhizosphere: tapping into belowground chemical communication. *Trends in plant science*, 21(3), pp.256-265. <https://doi.org/10.1016/j.tplants.2016.01.008>
- Van Deynze, A., Zamora, P., Delaux, P.M., Heitmann, C., Jayaraman, D., Rajasekar, S., Graham, D., Maeda, J., Gibson, D., Schwartz, K.D. and Berry, A.M., 2018. Nitrogen fixation in a landrace of maize is supported by a mucilage-associated diazotrophic microbiota. *PLoS biology*, 16(8), p.e2006352. <https://doi.org/10.1371/journal.pbio.2006352>
- Walker, T.S., Bais, H.P., Grotewold, E., Vivanco, J.M., 2003. Update on Root Exudation and Rhizosphere Biology *Root Exudation and Rhizosphere Biology* 1 132, 44–51. <https://doi.org/10.1104/pp.102.019661>.
- Wang, D., Marschner, P., Solaiman, Z., Rengel, Z., 2007. Growth, P uptake, and rhizosphere properties of intercropped wheat and chickpea in soil amended with iron

- phosphate or phytate. *Soil Biol. Biochem.* 39, 249–256. <https://doi.org/10.1016/j.soilbio.2006.07.013>
- Wang, J., Li, R., Zhang, H., Wei, G., Li, Z., 2020a. Beneficial bacteria activate nutrients and promote wheat growth under conditions of reduced fertilizer application. *BMC Microbiol.* 20, 1–12. <https://doi.org/10.1186/s12866-020-1708-z>.
  - Wang, N. Q., Kong, C. H., Wang, P., & Meiners, S. J. 2021b Root exudate signals in plant–plant interactions. *Plant, Cell & Environment*, 44(4), 1044-1058. <https://doi.org/10.1111/pce.13892>
  - Wang, G., Bei, S., Li, J., Bao, X., Zhang, J., Schultz, P. A., Zhang, J. 2021c. Soil microbial legacy drives crop diversity advantage: Linking ecological plant–soil feedback with agricultural intercropping. *Journal of Applied Ecology*, 58(3), 496-506. <https://doi.org/10.1111/1365-2664.13802>
  - Wang, Z., Bao, X., Li, X., Jin, X., 2015. Intercropping maintains soil fertility in terms of chemical properties and enzyme activities on a timescale of one decade. <https://doi.org/10.1007/s11104-015-2428-2>
  - Wieland, G., Neumann, R., Backhaus, H., 2001. Variation of microbial communities in soil, rhizosphere, and rhizoplane in response to crop species, soil type, and crop development. *Appl. Environ. Microbiol.* 67, 5849–5854. doi:<http://dx.doi.org/10.1128/AEM.67.12.5849-5854.2001>.
  - Wezel, A., Casagrande, M., Celette, F., Vian, J., Ferrer, A., Peigné, J., 2014b. Agroecological practices for sustainable agriculture. A review. *Agronomy for sustainable development*, 34(1), 1-20. <https://doi.org/10.1007/s13593-013-0180-7>
  - Wu, F.Y., Chung, A.K.C., Tam, N.F.Y., Wong, M.H., 2012. Root exudates of wetland plants influenced by nutrient status and types of plant cultivation. *Int. J. Phytoremediation* 14, 543–553. <https://doi.org/10.1080/15226514.2011.604691>.
  - Wu, Q., Peng, X., Yang, M., Zhang, W., Dazzo, F.B., Uphoff, N., Jing, Y. and Shen, S., 2018. Rhizobia promote the growth of rice shoots by targeting cell signaling, division and expansion. *Plant molecular biology*, 97(6), pp.507-523. <https://doi.org/10.1007/s11103-018-0756-3>
  - Xia, J. and Wishart, D.S., 2016. Using MetaboAnalyst 3.0 for comprehensive metabolomics data analysis. *Current protocols in bioinformatics*, 55(1), pp.14-10.

- Xu, Y., Qiu, W., Sun, J., Müller, C. and Lei, B., 2019. Effects of wheat/faba bean intercropping on soil nitrogen transformation processes. *Journal of Soils and Sediments*, 19(4), 1724-1734. <https://doi.org/10.1007/s11368-018-2164-3>
- Xu, Z., Li, C., Zhang, C., Yu, Y., van der Werf, W., & Zhang, F. 2020. Intercropping maize and soybean increases efficiency of land and fertilizer nitrogen use; A meta-analysis. *Field Crops Research*, 246, 107661. <https://doi.org/10.1016/j.fcr.2019.107661>
- Xue, Y., Xia, H., Christie, P., Zhang, Z., Li, L., Tang, C., 2016. Crop acquisition of phosphorus, iron and zinc from soil in cereal / legume intercropping systems: A critical review. *Annals of Botany*, 117(3), 363-377. <https://doi.org/10.1093/aob/mcv182>
- Yajiao, Z., Xiaojing, L., Changchun, T., Yong, W., 2020. Effect of root interaction on nodulation and nitrogen fixation ability of alfalfa in the simulated alfalfa / triticale intercropping in pots 1–11. <https://doi.org/10.1038/s41598-020-61234-5>
- Yang, C., Fan, Z., Chai, Q., 2018. Agronomic and Economic Benefits of Pea / Maize Intercropping Systems in Relation to N Fertilizer and Maize Density. *Agronomy*, 8(4), 52. <https://doi.org/10.3390/agronomy8040052>
- Yeh, K.C., Peck, M.C. and Long, S.R., 2002. Luteolin and GroESL modulate in vitro activity of NodD. *Journal of bacteriology*, 184(2), pp.525-530. <https://doi.org/10.1128/JB.184.2.525-530.2002>
- Yin, W., Guo, Y., Hu, F., Fan, Z., Feng, F., Zhao, C., French, R.J., 2018. Wheat-Maize Intercropping With Reduced Tillage and Straw Retention: A Step Towards Enhancing Economic and Environmental Benefits in Arid Areas 9, 1–16. <https://doi.org/10.3389/fpls.2018.01328>
- Yong, T. W., Ping, C. H. E. N., Qian, D. O. N. G., Qing, D. U., Feng, Y. A. N. G., Wang, X. C., Yang, W. Y. 2018. Optimized nitrogen application methods to improve nitrogen use efficiency and nodule nitrogen fixation in a maize-soybean relay intercropping system. *Journal of Integrative Agriculture*, 17(3), 664-676. [https://doi.org/10.1016/S2095-3119\(17\)61836-7](https://doi.org/10.1016/S2095-3119(17)61836-7)
- Yuan, J., Zhao, J., Wen, T., Zhao, M., Li, R., Goossens, P., Huang, Q., Bai, Y., Vivanco, J.M., Kowalchuk, G.A. and Berendsen, R.L., 2018. Root exudates drive the soil-borne legacy of aboveground pathogen infection. *Microbiome*, 6(1), pp.1-12. <https://doi.org/10.1186/s40168-018-0537-x>

- Zhang, N., Wang, D., Liu, Y., Li, S., Shen, Q., Zhang, R., 2014. Effects of different plant root exudates and their organic acid components on chemotaxis, biofilm formation, and colonization by beneficial rhizosphere-associated bacterial strains. *Plant Soil* 374, 689–700. <https://doi.org/10.1007/s11104-013-1915-6>.
- Zhao, M., Jones, C.M., Meijer, J., Lundquist, P., Fransson, P., Carlsson, G., Hallin, S., 2017a. Intercropping affects genetic potential for inorganic nitrogen cycling by root-associated microorganisms in *Medicago sativa* and *Dactylis glomerata*. *Appl Soil Ecol* 119, 260–266. <https://doi.org/10.1016/j.apsoil.2017.06.040>
- Zhao, Y., Liu, X., Tong, C., & Wu, Y. 2020. Effect of root interaction on nodulation and nitrogen fixation ability of alfalfa in the simulated alfalfa/triticale intercropping in pots. *Scientific reports*, 10(1), 1-11. <https://doi.org/10.1038/s41598-020-61234-5>
- Zhi-dan, F.U., Li, Z., Ping, C., Qing, D.U., Ting, P., Chun, S., Xiao-chun, W., 2019. Effects of maize-soybean relay intercropping on crop nutrient uptake and soil bacterial community. *J Integr Agric* 18, 2006–2018. [https://doi.org/10.1016/S2095-3119\(18\)62114-8](https://doi.org/10.1016/S2095-3119(18)62114-8)
- Zhu, S., Vivanco, J.M., Manter, D.K., 2016. Nitrogen fertilizer rate affects root exudation, the rhizosphere microbiome and nitrogen-use-efficiency of maize. *Appl Soil Ecol* 107, 324–333. <https://doi.org/10.1016/j.apsoil.2016.07.009>