References

Allen C.B., Kanda J. and Laws E.A. (1996) New production and photosynthetic rates within and outside a cyclonic mesoscale eddy in the North Pacific subtropical gyre, Deep-Sea Res., 43, 917-936.

Altabet M.A. and McCarthy J.J. (1985) Temporal and spatial variations in the natural abundance of ^{15}N in PON from warm core ring, Deep-Sea Res., Part A, 32, 755-772.

Altabet M.A. and McCarthy J.J. (1986) Vertical patterns in ^{15}N abundance in PON from the surface waters of warm core rings, J Mar. Sci., 44,185-201.

Altabet M.A. (1988) Variations in nitrogen isotopic composition between sinking and suspended particles: Implications for nitrogen cycling and particle transformation in the open ocean, Deep-Sea Res., Part A, 35, 535-554.

Altabet M.A., Deuser W.G., Honjo S. and Stienen C. (1991) Seasonal and depth-related changes in the source of sinking particles in the North Atlantic, Nature, 354, 136-139.

Altabet M.A. and Francois R. (1994) The use of nitrogen isotopic ratio for reconstruction of past changes in the surface ocean nutrient utilization, In: Carbon Cycling in the Glacial Ocean: Constraints on the Ocean's Role in Global Change (Eds) Zahn, R., Kaminski, M.A., Labeyrie, L., and Pederson, T. F., 281-306, Springer-Verlag, New York.

Altabet M.A. (1996) Nitrogen and carbon isotopic tracers of the source and transformation of particles in the Deep-Sea, In: Particle Flux in the Ocean (Eds) Ittekkot, V., Schafer, P., Honjo, S., and Depetris, P.J., 155-184, John Wiley, New York.

Asper V.L. and Smith W.O. (1999) Particle fluxes during austral spring and summer in the southern Ross Sea (Antarctica), J. Geophys. Res., 104, 5345-5360.

Bacon M.P., Huh C.A., Fleer A.P. and Deuser W.G. (1985) Seasonality in the flux of natural radionuclides and plutonium in the deep Sargasso Sea, J. Mar. Res., 44, 185-201.

Banse K. (1984) Overview of the hydrography and associated biological phenomena in the Arabian Sea, off Pakistan, In: Marine geology and oceanography of Arabian Sea and coastal Pakistan (Eds) Haq B.U. and Milliman J.D., 271-303, Van Nostrand Reinhold Co., New York.

Banse K. and McClain (1986) Winter blooms of phytoplankton in the Arabian Sea as observed by the coastal zone color scanner, Mar. Ecol. Progr. Ser., 34, 201-211.

Banse K. (1987) Seasonality of phytoplankton chlorophyll in the central and northern Arabian Sea, Deep-Sea Res. I, 34, 713-723.

Banse K (1992) Grazing, temporal changes of phytoplankton concentrations and the microbial loop in the open sea, In: Primary Productivity and Biogeochemical Cycles, (Eds) Falkowski P. and Woodhead A.D., 409-440, Plenum, New York.

Banse K. and English D.C. (1993) Revision of satellite-based phytoplankton pigment data from the Arabian Sea during the northeast monsoon, Mar. Res., 2, 83-103.

Barber R.T. (1992) Introduction to the WEC88 cruise: An investigation into why the equator is not greener, J. Geophys. Res., 97, 609-610.

Barber R.T., Murray J.W. and McCarthy J.J. (1994) Biogeochemical interactions in the equatorial Pacific, Ambio, 23, 62-66.

Barber R.T., Sanderson M.P., Lindley S.T., Chai F., Newton J., Trees C.C., Foley D.G. and Chavez F.P. (1996) Primary productivity and its regulation in the equatorial pacific during and following the 1991-92 El Nino, Deep-Sea Res. II, 43, 933-969.

Barber R.T., Marra J., Bidigare R.C., Codispoti L.A., Halpern D., Johnson Z, Latasa M., Goericke R. and Smith S.L. (2001) Primary productivity and its regulation in the Arabia Sea during 1995, Deep-Sea Res. II, 48, 1127-1172.

Bender M., Ducklow H., Kiddon J., Marra J. and Martin J. (1992) The carbon balance during the 1989 spring bloom in the North Atlantic Ocean, 47°N, 20°W, Deep-Sea Res., 39, 1707-1725.

Ben-Yaakov S., Ruth E. and Kaplan I.R. (1974) Calcium carbonate saturation in northeastern Pacific: *In situ* determination and geochemical implications, Deep-Sea Res., 21, 229-2243.

Berger W.H., Fisher K., Lai C. and Wu G. (1987) Oceanic productivity and organic carbon flux, Scripps Institution of Oceanography reference 87(30), 1-67.

Bhattathiri P. M. A., Devassy V. P. and Radhakrishna K. (1980) Primary production in the Bay of Bengal during southwest monsoon of 1978, Mahasagar- Bulletin of the National Institute of Oceanography, 13, 315-323.

Bhattathiri P. M. A., Pant A., Sawant S., Gauns M., Matondakar S. G. P. and Mohanraju R. (1996) Phytoplankton production and chlorophyll distribution in eastern and central Arabian Sea in 1994-1995, Curr. Sci., 71, 857-862.

Bird R.E. (1984) A simple, solar spectral model for direct normal and diffuse horizontal irradiance, Solar Energy, 32, 461-471.

Blackmer A.M. and Bremner J.M. (1977) Nitrogen Isotope discrimination in denitrification of nitrate in soils, Soil Biol. Biochem., 9, 73-77.

Brandes J.A., Devol A.H., Yoshinari T., Jayakumar T.A. and Naqvi S.W.A. (1998) Isotopic composition of nitrate in the central Arabian Sea and eastern tropical north Pacific: a tracer for mixing and nitrogen cycles, Limnol. Oceanogr., 43, 1680-1689.

Bremner J.M. and Blackmer, A.M. (1981) Terrestrial nitrification as a source of atmospheric nitrous oxide, In: Denitrification, Nitrification, and atmospheric nitrous oxide (Ed) Delwiche C.C., Wiley, New York.

Brock J.C., McClain C.R., Luther M.E. and Hay W.W. (1991) The phytoplankton bloom in the northwestern Arabian Sea during the southwest monsoon of 1979, J. Geophys. Res., 96,20 623-20642.

Brock J.C. and McClain C.R. (1992) Interannual variability in phytoplankton blooms observed in the northwestern Arabian Sea during the Southwest monsoon, J. Geophys. Res., 97, 733-750.

Brock J., Sathyendranath S. and Platt T. (1993) Modelling the seasonality of subsurface light and primary production in the Arabian Sea, Mar. Ecol. Prog. Ser., 101,209-221.

Brock J., Sathyendranath S. and Platt T. (1994) A model study of seasonal mixed layer primary production in the Arabian Sea, Proc. Ind. Acad. Sci. (Earth Planet. Sci.), 103 (2), 163-176.

Broecker W.S. (1982) Glacial to interglacial changes in ocean chemistry, Progr. Oceanogr., 11, 151-197.

Broecker W.S., Ledwell J.R., Takahashi T., Weiss R., Merlivat L., Memery L., Peng T.H., Jahne B. and Munnich K.O. (1986) Isotopic versus micrometeorologic ocean CO_2 fluxes: A serious conflict, J. Geophys. Res., 91, 10517-10527.

Broenkow W. W. and Cline J.D. (1969) Calorimetric determination of dissolved oxygen at low concentrations, Limnol. Oceanogr., 14,450-14,454.

Bronk D.A., Glibert P.M. and Ward B.B. (1994) Nitrogen uptake, dissolved nitrogen release and new production, Science, 265, 1843-1846.

Buesseler K.O. (1998) The decoupling of production and particulate export in the surface ocean, Global Biogeochem. Cycle, 12, 297-310.

Buesseler K.O. (1991) Do upper-ocean sediment traps provide an accurate record of particle flux? Nature, 353, 420-423.

Buesseler K.O., Bacon M.P., Cochran J.K. and Livingstone H.D. (1992) Carbon and nitrogen export during the JGOFS North Atlantic bloom experiment estimated from ^{234}Th: ^{238}U disequilibria, Deep-Sea Res.,39,115-1137.

Burkill P.H., Mantoura R.F.C. and Owens N.J.P. (1993) Biogeochemical cycling in the northwestern Indian Ocean: a brief overview, Deep-Sea Res. II, 40, 643-649.

Calvert S. E., Nielsen B. and Fontugne M.R. (1992) Evidence from nitrogen isotope ratios for enhanced productivity during formation of eastern Mediterranean sapropels, Nature, 359,223-225.

Capone D.G., Zehr J.P., Paerl H.W., Bergman B. and Carpenter E.J. (1997) *Trichodesmium,* a Globally significant marine Cyanobacterium, Science, 276, 1221-1229.

Capone D.G. (2001) Marine nitrogen fixation: What's the fuss? Current opinion in microbiology, 4, 341-348.

Chauhan P., Nagur C. R. C., Mohan M., Nayak S. and Navalgund, R. R., (2001) Surface Chlorophyll *a* distribution in Arabian Sea and Bay of Bengal using IRS P4 OCM satellite data, Curr. Sci., 80 (2), 127-129.

Cline J.D. and Kaplan I.R. (1975) Isotopic fractionation of dissolved nitrate during denitrification in the eastern tropical North Pacific Ocean, Mar. Chem., 3, 271-299.

Cochlan W.P. and Bronk D.A. (2001) Nitrogen uptake kinetics in Ross Sea, Antarctica, Deep-Sea Res. II, 48, 4127-4153.

Cole J.J., Pace M.L. and Fidlay S. (1988) Bacterial production in fresh and saltwater ecosystems: a cross system overview, Mar. Ecol. Prog. Ser., 43, 1-10.

Collos Y. (1987) Calculation of ^{15}N uptake rates by phytoplankton assimilating one or several nitrogen sources, Int. J. Rad. Appl. Instrum., Part A, 275-282.

Conway H.L. (1977) Interaction of inorganic nitrogen in the uptake and assimilation by marine phytoplankton, Mar. Biol., 39, 221-232.

Cullen J.J. (1991) Hypotheses to explain high nutrient, low chlorophyll conditions in the open sea, Limnol. Oceanogr., 36, 1578-1599.

Das P.K. (1995) The monsoons, 219, National Book Trust, New Delhi.

Datta D.K. (1999) Dissolved phosphate in the Ganges-Brahmaputra-Meghna river system in the Bengal Basin, Bangladesh, Mitteilungen aus dem Geologisch - Palaontologischen Institut der Universitat Hamburg, SCOPE/UNEP Sonderband 82, 131-138.

Delwiche C.C. (1970) The nitrogen Cycle, Scientific American, 223, 136-147.

Delwiche C.C. (1981) The nitrogen cycle and nitrous oxide, In: Denitrification, Nitrification, and atmospheric nitrous oxide (Ed) Delwiche C.C., Wiley, New York.

DeNiro M.J. and Epstein S. (1981) Influence of diet on the distribution of nitrogen isotopes in animals, Geochim. Cosmochim. Acta, 45, 341-351.

DeSousa S. N., Naqvi S. W. A. and Reddy C. V. G. (1981) Distribution of nutrients in the western Bay of Bengal, Ind. J. Mar. Sci., 10, 327-331.

Devol A.H. (1978) Bacterial oxygen uptake kinetics as related to biological processes in oxygen deficient zones of the oceans, Deep-Sea Res., 25, 137-146.

DiTullio G.R. and Laws E.A. (1991) Impact of an atmospheric-oceanic disturbance on phytoplankton community dynamics in the North Pacific Central Gyre, Deep-Sea Res., 38, 1305-1329.

Doerffer R. (1992) Imaging spectroscopy for detection of chlorophyll and suspended matter, In: Imaging Spectroscopy: Fundamentals and Prospective Applications (Eds) Toselli F. and Bodechtel J., 215-257, Kluwer Academic Publisher.

Donald K.M., Joint I., Rees A.P., Woodward E.M.S. and Savidge G. (2001) Uptake of carbon, nitrogen and phosphorus by phytoplankton along the 20°W meridian in the NE Atlantic between 57.5°N and 37°N, Deep-Sea Res. II, 48, 873-897.

Dore J. E. and Karl D. M. (1996) Nitrification in the euphotic zone as a source of nitrite, nitrate, and nitrous oxide at station ALOHA, Limnol. Oceanogr., 41, 1619-1628.

Dortch Q. (1990) The interaction between ammonium and nitrate uptake in phytoplankton, Mar. Ecol. Progr. Ser., 61, 183-201.

Dowling C.B., Poreda R.J. and Basu A.R. (2003) The ground water geochemistry of Bengal Basin: Weathering, chemsorption, and trace metal flux to the oceans, Geochim. Cosmochim. Acta, 67(12), 2117-2136.

Duce et al. (1991) The atmospheric input of trace species to the world ocean, Global Biogeochem. Cycles, 5, 193-259.

Ducklow H.W. and Harris R. (1993) Introduction to the JGOFS North Atlantic Bloom Experiment, Deep-Sea Res. II, 40, 1-8.

Ducklow H.W. (1995) Ocean Biogeochemical fluxes: New production and export of organic matter from the upper ocean, Rev. Geophys., Supplement, 1271-1276.

Ducklow H.W. (1999) The bacterial content of the oceanic euphotic zone, Fems. Microbiol. Ecol., 30,1-10.

Dugdale R.C. (1967) Nutrient limitation in the sea: Dynamics, identification, and significance, Limnol. Oceanogr., 12, 685-695.

Dugdale R.C. and Goering J.J. (1967) Uptake of new and regenerated forms of nitrogen in primary productivity, Limnol. Oceanogr., 12, 196-206.

Dugdale R.C. and Wilkerson F.P. (1986) The use of ^{15}N to measure nitrogen uptake in eutrophic oceans; experimental considerations, Limnol. Oceanogr., 31, 673-689.

Dugdale R.C., Wilkerson F.P., Barber R. T. and Chavez F.P. (1992) Estimating new production in the Equatorial Pacific Ocean at 150° W, J. Geophys. Res., 97, 681-686.

Emerson S.P., Quay P., Stump C., Wilbur S. and Knox M. (1991) O_2, Ar, N_2 and ^{222}Rn in surface waters of subarctic ocean: net biological O_2 production, Global Biogeochem. Cycles, 5, 49-69.

Emerson S., Quay P. and Wheeler P.A. (1993) Biological productivity determined from oxygen mass balance and incubation experiments, Deep-Sea Res., 40, 2351-2358.

Emerson S., Quay P., Karl D., Winn C., Tupas L. and Landry M. (1997) Experimental determination of organic carbon flux from open Ocean surface waters, Nature, 389, 951-954.

Eppley R.W., Sharp J.H., Renger E.H., Perry M.J. and Harrison W.G. (1977) Nitrogen assimilation by phytoplankton and other microorganism in the surface waters of the central North Pacific Ocean, Mar. Biol., 39, 111.

Eppley R.W. and Peterson B.J. (1979) Particulate organic matter flux and planktonic new production in the deep ocean, Nature, 282, 677-680.

Eppley R.W., Renger E.H. and Betzer R.R. (1983) The residence time of particulate organic carbon in the surface layer of the ocean, Deep-Sea Res., 30, 311-323.

Falkowski P.G., Laws E.A., Barber R.T. and Murray J.W. (2003) Phytoplankton and their role in primary new and export production, In: Ocean Biogeochemistry (Ed) Fasham M.J.R., 99-121,Springer, Heideberg.

Farrell J.W., Pederson T.F., Calvert S.E. and Nielsen B. (1995) Glacial-interglacial changes in nutrient utilization in equatorial Pacific Ocean, Nature, 377,514-517.

Field C., Behrenfeld M., Randerson J. and Falkowski P. (1998) Primary production of the biosphere: integrating terrestrial and oceanic components, Science, 281, 237-240.

Findlater J. (1974) The low-level cross equatorial air current of the western Indian Ocean during the northern summer, Weather, 29,411-416.

Fitzwater S.E., Knauer G.A. and Martin J.H. (1982) Metal contamination and its effects on primary production measurements, Limnol. Oceanogr., 27, 544-551.

Flynn K.J., Fasham M. J. R. and Hipkin C.R. (1997) Modelling the interactions between ammonium and nitrate uptake in marine phytoplankton, Philosophical Transactions of London B 352, 1625-1645.

Flynn K.J. (1998) Estimation of kinetic parameters for the transport of nitrate and ammonium into marine phytoplankton, Mar. Ecol. Prog. Ser., 169, 13-28.

Fogel M.L. and Cifuentes L.A. (1993) Isotope fractionation during primary production, In: Organic Geochemistry, (Eds) Engel M.H. and Macko S.A., 73-98, Plenum Press, New York.

Francois R., Altabet M.A. and Burkle L.H. (1992) Glacial to interglacial changes in surface nitrate utilization in the Indian Sector of the Southern Ocean as recorded by sediment $\delta^{15}N$, Paleoceanography, 7, 589-606.

Frost B.W. and Franzen N.C. (1992) Grazing and iron limitation in the control of phytoplankton stock and nutrient concentration: A chemostat analogue of the Pacific equatorial upwelling zone, Mar. Ecol. Prog. Ser., 83, 291-303.

Galloway J.N. et al. (2004) Nitrogen Cycles: past, present and future, Biogeochemistry, 1-73.

George M.D., Kumar M.D., Naqvi S.W.A., Banerjee S., Narvekar P.V., De Sousa S.N. and Jayakumar D.A. (1994) A study of the carbon dioxide system in the northern Indian Ocean during premonsoon, Mar. Chem., 47, 243-254.

Gill A.E. (1982) Atmosphere-Ocean dynamics, Academic Press, New York, 662.

Glibert P.M., Lipschultz F., McCarthy J. J. and Altabet M. A. (1982a) Isotope dilution models of uptake and remineralization of ammonium by marine plankton, Limnol. Oceanogr., 27, 639-650.

Glibert P.M., Biggs D.C. and McCarthy J.J. (1982b) Utilization of ammonium and nitrate during austral summer in the Scotia Sea, Deep-Sea Res., 29, 837-850.

Goldman J.C. and Glibert P.M. (1983) Kinetics of inorganic nitrogen uptake by phytoplankton, In: Nitrogen in the marine environment, (Eds) Carpenter E.J. and Capone D.G., 233-274, Academic.

Gomes H. R., Goes J. I. and Saino T. (2000) Influence of physical processes and fresh water discharge on seasonality of phytoplankton regime in the Bay of Bengal, Cont. Shelf Res., 20, 313-330.

Gordon H.R. and Wang M. (1994) Retrieval of water leaving radiance and aerosol optical thickness over the oceans with SeaWiFS: A preliminary algorithm, Appl. Optics, 33, 443-452.

Haake B., Ittekkot V., Rixen T., Ramaswami V., Nair R. R. and Curry W. B. (1993) Seasonality and interannual variability of particle fluxes to the deep Arabian Sea, Deep-Sea Res. I, 40, 1323-1344.

Handley L.L. and Raven J.H. (1992) The use of natural abundance of nitrogen isotopes in plant physiology and ecology, Plant Cell Environ., 15, 965-985.

Hansell D.A., Whitledge T.E. and Goering J.J. (1993) Patterns of nitrate utilization and new production over the Bering -Chukchi Shelf, Cont. Shelf Res., 13, 601-627.

Hansen J., Johnson D., Lacis A., Lebedeff S., Lee P., Pind D. and Russel G. (1981) Climate impact of increasing atmospheric carbon dioxide, Nature, 213, 957-966.

Harrison W.G. (1980) Nutrient regeneration and primary productivity in the sea, In: Primary productivity in the Sea, (Ed) Falkowski P.G., Plenum, 433-460.

Harrision W.G., Platt T. and Lewis M.R. (1987) f-ratio and its relationship to ambient nitrate concentration in coastal waters, J. Plankton Res., 9, 235-248.

Harrison W.G., Harris L.R. and Irwin B.D. (1996) The kinetics of nitrogen utilization in the oceanic mixed layer: nitrate and ammonium interactions at nanomolar concentrations, Limnol. Oceanogr., 41, 16-32.

Hastenrath S. and Lamb P.J. (1979) Climate Atlas of the Indian Ocean, University of Wisconsin Press.

Hastenrath S. and Greishar L.L. (1989) Climatic Atlas of the Indian Ocean. Part III: Upper Ocean structure (University of Wisconsin press) 247.

Heaton T.H.E. (1986) Isotopic studies of nitrogen pollution in the hydrosphere and atmosphere: a review, Chem. Geol. (Iso. Geo. Sect.), 5, 87-102.

Hedges J.I., Baldock J.A., Gelinas Y., Lee C., Peterson M. and Wakehams S.G. (2001) Evidence for non selective preservation of organic matter in sinking marine particles, Nature, 409, 801-804.

Holland H.D. (1970) Ocean water, nutrients and atmospheric oxygen, Proceedings of the international symposium on hydrogeochemistry and Biogeochemistry, Tokyo, 68-81.

Holm-Hansen O. and Mitchell B.G. (1991) Spatial and temporal distribution of phytoplankton and primary production in the western Bransfield Strait region, Deep-Sea Res., 38, 961-980.

Honjo S., Spencer D.W. and Gardner W.D. (1992) A sediment trap intercomparison experiment in the Panama Basin, Deep-Sea Res., 39, 333-358.

Honjo S. and Manganini S.J. (1993) Annual biogenic particle fluxes to the interior of the North Atlantic Ocean studied at 34°N, 21°W and 48°N, 21°W, Deep-Sea Res. II, 40, 587-607.

Honjo S., Dymond J., Prell W. and Ittekkot V. (1999) Monsoon-controlled export fluxes to the interior of the Arabian Sea, Deep-Sea Res. II, 46, 1859-1902.

Houghton J.T. et al. (2001) Climate change 2001: The scientific basis, Cambridge University Press, New York.

Hubner H. (1986) Isotope effect of nitrogen in the soil and biosphere, In: Handbook of Environmental Isotope Geochemistry, (Eds) Fritz P. and Fontes J.C., The Terrestrial Environment, vol. 2b, 361- 425.

Huntley M., Karl D.M., Niller P. and Holm-Hansen O. (1991) Research on Antarctic Coastal Ecosystem Rates (RACER): An interdisciplinary field experiment, Deep-Sea Res., 38, 911-941.

Ittekkot V. (1991) Particle flux studies in the Indian Ocean, Eos, 72, 527-230.

Ittekkot V., Nair R.R., Honjo S., Ramaswami V., Bartsch M., Manganini S. and Desai

B.N. (1991) Enhanced particle fluxes in the Bay of Bengal induced by injection of freshwater, Nature, 351, 385-387.

Jacques G. (1991) Is the concept of new production - regenerated production valid for the southern Ocean? Mar. Chem., 35,273-286.

Jaffe D.A. (2000) The nitrogen Cycle, In: Earth System Science, 322-340, Academic Press Limited.

Jawed M. (1973) Ammonia excretion by zooplankton and its significance to primary productivity during summer, Mar. Biol., 23,115-120.

JGOFS Report No. 19. (1996) Protocol for the Joint Global Ocean Flux Study (JGOFS) core measurements, 145-150, SCOR, Bergen, Norway.

Jickells T.D. (1998) Nutrient biogeochemistry of the coastal zone, Science, 284,217-222.

Jochem F.J., Pollehne, F. and Zeitschel B. (1993) Productivity regime and phytoplankton size structure in the Arabian Sea, Deep-Sea Res. II, 40, 711-735.

Joint I., Rees A. and Woodward E.M.S. (2001) Primary production and nutrient assimilation in the Iberian upwelling in August 1998, Progr. Oceanogr., 51, 303-320.

Junk G. and Svec H.J. (1958) The absolute abundance of the nitrogen isotopes in the atmosphere and compressed gas from various sources, Geochim. Cosmochim. Acta, 14, 234-243.

Kabanova Y.G. (1968) Primary production of the northern part of the Indian Ocean, Oceanology, 8, 214-225.

Kanda J., Saino T. and Hattori A. (1985) Nitrogen uptake by natural populations of phytoplankton and primary production in the Pacific Ocean: regional variability of uptake capacity, Limnol. Oceanogr., 30, 987-999.

Karl D.M., Tilbrook B.D. and Tien G. (1991) Seasonal coupling of organic matter production and particle flux in the western Bransfield Strait, Antarctica, Deep-Sea Res., 38, 1097-1126.

Karl D.M., Christian J.R., Dore J.E., Hebel D.V., Letelier R.M., Tupas L.M. and Winn C.D. (1996) Seasonal and interannual variability in primary production and particle flux at station ALOHA, Deep-Sea Res. II, 43, 539-568.

Karl D.M., Letelier R., Tupas L., Dore J., Christian J. and Hebel D. (1997) The role of nitrogen fixation in biogeochemical cycling in the subtropical North Pacific Ocean, Nature, 388, 533-538.

Keeling C.D. and Bacastow R.B. (1977) Energy and Climate, 72-95, National Academy of Sciences, Washington, D.C.

Keeling C. D. and Whorf T. P. (2000) Trends: A compendium of data on global change, Carbon dioxide information analysis center, Oak Ridge National Laboratory, Tennessee.

Kendall C. (1998) Tracing nitrogen sources and cycling in catchments, In :Isotope tracers in catchment hydrology (Eds) Kendall C. and McDonnell J.J., 519-576, Elsevier.

Kirchman D.L. (1994) The uptake of inorganic nutrients by heterotrophic bacteria, Microb. Ecol., 28, 255-271.

Kouvarakis G., Mihalopoulos N., Tselepides A. and Stavrakakis S. (2001) On the importance of atmospheric inputs of inorganic nitrogen species on the productivity of the eastern Mediterranean Sea, Global Biogeochem. Cycle, 15(4), 805-817.

Krey J. (1973) Primary production in the Indian Ocean, In: The Biology of the Indian Ocean, (Ed) Zeitschel B., 115-126, Springer, New York.

Kumar M.D., Rajendran A., Somasundar K., Ittekkot V. and Desai B.N. (1992) Processes controlling carbon components in the Arabian Sea, In: Oceanography of the Indian Ocean (Ed) Desai B.N., 313-325, Oxford-IBH, New Delhi.

Kumar M.D., Naqvi S.W.A., George M.D. and Jayakumar D.A. (1996) A sink for atmospheric carbon dioxide in the northeast Indian Ocean, J. Geophys. Res., 101, 18,121-18,125.

Kumar S., Ramesh R., Sardesai S. and Seshshayee S. (2004a) High new production in the Bay of Bengal: possible causes and implications, Geophys. Res. Lett. 31, L18304, doi: 10. 1029/2004GL021005.

LaFond E.C. (1966) Bay of Bengal, In: The Encyclopaedia of Oceanography, (Ed) Fairbridge R.W., Van Nostrand Reinhold Co., New York, 110-118.

LaViolette P.E. (1967) Temperature, salinity and density of the world's seas, Bay of Bengal and Andaman Sea, Informal Report no. 67-57, Naval Oceanographic Office, Washington DC, 81.

Laws E.A., Harrison W.G. and DiTullio G. (1985) A comparison of nitrogen assimilation rates based on ^{15}N uptake and autotrophic protein synthesis, Deep-Sea Res., 32, 85-95.

Laws E.A., Falkowski P.G., Smith W.O., Ducklow H. and McCarthy J.J. (2000) Temperature effects on export production in the open ocean, Global Biogeochem. Cycles, 14, 1231-1246.

Lee et al. (1998) Particulate organic carbon fluxes: compilation of results from the 1995 US JGOFS Arabian Sea Process Study, Deep-Sea Res. II, 45, 2489-2501.

Lee C.M., Jones B.H., Brink K.H. and Fischer K.H. (2000) The upper-Ocean response to monsoonal forcing in the Arabian Sea: seasonal and spatial variability, Deep-Sea Res. II, 47, 1177-1226.

L' Helguen S., LeCorre P., Madec C. and Morin P. (2002) New and regenerated production in the Almeria-Oran front area, eastern Alboran Sea, Deep-Sea Res. I, 49, 83-99.

Liu, K.-K. and Kaplan. I. R. (1989) The eastern tropical Pacific as a source of ^{15}N-enriched nitrate in seawater off southern California, Limnol. Oceanogr., 34, 820-830.

Lohrenz S.E., Knauer G.A., Asper V.L., Tuel M., Michaels A.F. and Knap A.H. (1992) Seasonal variability in primary production and particle flux in the northwestern Sargasso Sea: U.S. JGOFS Bermuda Atlantic Time Series Study, Deep-Sea Res., 39, 1373-1391.

Lourey M.J., Trull T.W. and Sigman D.M. (2003) Sensitivity of $\delta^{15}N$ of nitrate, surface suspended and deep sinking particulate nitrogen to seasonal nitrate depletion in the Southern Ocean, Global Biogeochem. Cycles, 17, 1081, doi: 10.1029/2001GB001973.

MacIsaac J.J. and Dugdale R.C. (1969) The kinetics of nitrate and ammonium uptake by natural populations of marine phytoplankton, Deep-Sea Res., 16, 45-57.

MacIsaac J.J. and Dugdale R.C. (1972) Interactions of light and inorganic nitrogen in controlling nitrogen uptake in the sea, Deep-Sea Res., 19, 209-232.

Madhupratap M., Prasanna Kumar S., Bhattathiri P.M.A., Dileep Kumar M., Raghukumar S., Nair K.K.C. and Ramaiah N. (1996) Mechanism of the biological response to winter cooling in the northeastern Arabian Sea, Nature, 384, 549-552.

Madhupratap M., Gauns M., Ramaiah N., Prasanna Kumar S., Muraleedharan P.M., DeSouza S.N., Sardessai S., Muraleedharan U. (2003) Biogeochemistry of the Bay of Bengal: physical, chemical and primary productivity characteristics of the central and western Bay of Bengal during summer monsoon 2001, Deep-Sea Res. II, 50, 881-896.

Maier-Reimer E., Mikolajewicz U. and Winguth A. (1996) Clim. Dyn., 12, 711.

Malone T.C., Pike S.E. and Conley D.J. (1993) Transient variations in phytoplankton productivity at the JGOFS Bermuda time series station, Deep-Sea Res., 40, 903-924.

Mariotti A., Pierre D., Vedy J.C., Bruckert S. and Guillemot J. (1980) The abundance of natural nitrogen 15 in the organic matter of soils along an altitudinal gradient (Chablais, Haute Savoie, France) Catena, 7, 293-300.

Mariotti A., Lancelot C. and Billen G. (1984) Natural isotopic composition of nitrogen as a tracer of origin for suspended organic matter in the Scheldt estuary, Geochim. Cosmochim. Acta, 48, 549-555.

Martin J. -M. and Whitfield M. (1983) The significance of the river input of chemical elements to the ocean, In: Trace metals in seawater, (Eds) C. S. Wong et al., 265-295, Plenum, New York.

McCarthy J.J. and Eppley R.W. (1972) A comparison of chemical isotopic and enzymatic methods for measuring nitrogen assimilation of marine phytoplankton, Limnol. Oceanogr., 17, 371-382.

McCarthy J.J., Taylor W.R. and Taft T. (1977) Nitrogenous nutrition of the plankton in the Chesapeake Bay: Nutrient availability and phytoplankton preferences, Limnol. Oceanogr., 22, 996-1011.

McCarthy J.J. and Goldman J.C. (1979) Nitrogenous nutrition of marine phytoplankton in nutrient depleted waters, Science, 203, 670-672.

McCarthy J.J. and Carpenter E.J. (1983) Nitrogen cycling in near surface waters of the open ocean, In: Nitrogen in the Marine Environment, (Eds) Carpenter E.J. and Capone D.G., 487-512, Academic Press, New York.

McCarthy J.J., Garside C. and Nevins J.L. (1992) Nitrate supply and phytoplankton uptake kinetics in the euphotic layer of a gulf stream warm core ring, Deep-Sea Res., 39, S393-S403.

McCarthy J.J., Garside C., Nevins J.L. and Barber R.T. (1996) New production along 140°W in the equatorial Pacific during and following the 1992 El Nino event, Deep-Sea Res. II, 43,1065-1093.

McCarthy J.J., Garside C. and Nevins J. (1999) Nitrogen dynamics during the Arabian Sea northeast monsoon, Deep-Sea Res. II, 46, 1623-1664.

McCreary J.P., Kohler K.E., Hood R.R. and Olson D.B. (1996) A four component ecosystem model of biological activity in the Arabian Sea, Progr. Oceanogr., 37, 193-240.

McElroy M.B. (1983) Marine biological controls on atmospheric CO_2 and climate, Nature, 302, 328-329.

McGillicuddy D.J. Jr., McCarthy J.J. and Robinson A.R. (1995) Coupled physical and biological modelling of the spring bloom in the North Atlantic (I): model formulation and one dimensional bloom processes, Deep-Sea Res. I, 42,1313-1357.

Michaels A.F., Siegel D.A., Johnson R.J., Knap A.F. and Galloway J.N. (1993) Episodic inputs of atmospheric nitrogen to the Sargasso Sea: Contribution to new production and phytoplankton blooms, Global Biogeochem. Cycles, 7, 339-351.

Michaels A.F., Bates N.R., Buesseler K.O., Carlson C.A., Knap H.H. (1994) Carbon cycle imbalances in the Sargasso Sea, Nature, 372, 537-540.

Miller C.B., Frost B.W., Wheeler P.A., Landry M.R., Welschmeyer N. and Powell T.M. (1991) Ecological dynamics in the subarctic Pacific, a possibly iron limited ecosystem, Limnol. Oceanogr., 36, 1600-1615.

Miller C.B. (1993) Pelagic production processes in the subarctic Pacific, Progr. Oceanogr., 32, 1-15.

Milliman J.D. and Syvitski J.P.M. (1992) Geomorphic/tectonic control of sediment discharge to the ocean: the importance of small mountainous rivers, The J. Geol., 100,525-100,544.

Minagawa M. and Wada E. (1986) Nitrogen isotope ratios of red tide organisms in the East Chine Sea: A characterization of biological nitrogen fixation, Mar. Chem., 19, 245-259.

Mino Y., Saino T., Suzuki K. and Maranon E. (2002) Isotopic composition of suspended particulate nitrogen ($\delta^{15}N_{sus}$) in surface waters of the Atlantic Ocean from 50°N to 50°S, Global Biogeochem. Cycles, 16, 4, 1059.

Miyake Y. and Wada E. (1967) The abundance ratio of $^{15}N/^{14}N$ in marine environments, Rec. Oceanogr. Works Jpn. 9, 37-53.

Miyake Y. and Wada E. (1971) The isotope effect on the nitrogen in biochemical, oxidation-reduction reactions, Rec. Oceanogr. Works Jpn. 11, 1-6.

Montoya J.P. and McCarthy J.J. (1995) Isotopic fractionation during nitrate uptake by phytoplankton grown in continuous culture, J. Plankton Res., 17, 439-464.

Muller L. J. and Charles C. T (1994) Revised SeaWiFS prelaunch algorithm for the diffuse attenuation coefficient K (490); In case studies for SeaWiFS Calibration and Validation, Part 4, NASA Technical Memorandum, 104566, 28, 18-21.

Mullin M.M., Perry M.J., Renger E.H. and Evans P.M. (1975) Nutrient regeneration by oceanic zooplankton: a comparison of methods, Mar. Sci. Comm., 1, 1-13.

Murray J.W., Downs J.N., Strom S., Wei C.L. and Jannasch H.W. (1989) Nutrient assimilation, export production and ^{234}Th scavenging in the eastern equatorial Pacific, Deep-Sea Res., 36, 1471-1489.

Murray J.W. (1991) The 1988 Black Sea Oceanography Expedition: Introduction and Summary, Deep-Sea Res., 38 (Supplement 2): S655-S661.

Murty V.S.N., Subrahmanyam B., Gangadhara Rao L.V. and Reddy G.V. (1998) Seasonal variation of sea surface temperature in the Bay of Bengal during 1992 as derived from NOAA-AVHRR SST data, Int. J. Remote Sensing, 19, 2361-2371.

Nair P.V.R., Samuel S., Joseph K.J. and Balachandran V.K. (1973) Primary production and potential fishery resources in the seas around India, In: Proceedings of the symposium on Living Resources of the Seas Around India, 1968, special publication, Central Marine Fisheries Research Institute, Cochin, 184-198.

Naqvi S.W.A. (1987) Some aspects of the oxygen deficient conditions and denitrification in the Arabian Sea, J. Mar. Res., 45, 1049-1072.

Naqvi S.W.A., Noronha R.J., Somasundar K. and Sen Gupta R. (1990) Seasonal changes in the denitrification regime of the Arabian Sea, Deep-Sea Res., 37, 593-611.

Naqvi S.W. A (1991) Geographical extent of denitrification in the Arabian Sea in relation to some physical processes, Oceanolog. Acta, 14, 281-290.

Naqvi S.W.A. and Noronha R.J. (1991) Nitrous oxide in the Arabian Sea, Deep-Sea Res., 38, 871-890.

Naqvi S.W.A. and Shailaja M.S. (1993) Activity of the respiratory electron transport system and respiration rates within the oxygen minimum layer of the Arabian Sea, Deep-Sea Res. II, 40, 687-695.

Naqvi S.W.A. (1994) Denitrification processes in the Arabian Sea, Proc. Indian Acad. Sci. (Earth Planet, Sci), 103, 2, 279-300.

Neess J.C., Dugdale R.C. and Goering J.J. (1962) Nitrogen metabolism in lakes, Measurement of nitrogen fixation with ^{15}N, Limnol. Oceanogr., 7, 163-169.

Olson D. B., Hitchcock G. L., Fine R. A. and Warren B. A. (1993) Maintenance of low-oxygen layer in the Central Arabian Sea, Deep-Sea Res. II, 40, 697-709.

O'Reilly J.E., Maritorena S., Mitchell B.G., Siegel D.A., Carder K.L., Garver S.A., Kahru M. and McClain C. (1998) Ocean Colour chlorophyll algorithms for SeaWiFS, J. Geophys. Res., 103, 24937-24953.

Owens N.J.P. (1987) Natural variation in ^{15}N in the marine environment, Adv. Mar. Biol., 24, 390-451.

Owens N.J.P. and Rees A.P. (1989) Determination of nitrogen-15 at sub-microgram levels of nitrogen using automated continuous-flow isotope ratio mass spectrometer, Analyst, 114,1655-1657.

Owens N.J.P., Priddle J. and Whitehouse M.J. (1991) Variation in phytoplanktonic nitrogen assimilation around south Georgia and in the Bransfield Strait (Southern Ocean), Mar. Chem., 35, 287-304.

Owens N.J.P., Burkill P.H., Mantoura R.F.C, Woodward E.M.S., Bellan I.E., Howland R.J.M. and Llewellyn C.A. (1993) Size fractionated primary production and nitrogen assimilation in the northwestern Indian Ocean, Deep-Sea Res. II, 40, 697-709.

Paasche E. and Kristjiansen S. (1982) Nitrogen nutrition of the phytoplankton in the Oslofjord, Estuarine Coastal and Shelf Science, 14, 237-249.

Perry M.J. and Eppley R.W. (1981) Phosphate uptake by phytoplankton in the Central North Pacific Ocean, Deep-Sea Res., 28A, 39-49.

Peterson B.J. and Fry B. (1987) Stable isotopes in ecosystem studies, Ann. Rev. ecol. System., 18, 293-320.

Platt T., Sathyendranath S., Ulloa O., Harrison W.G., Hoepffner N. and Goes J. (1992) Nutrient control of phytoplankton photosynthesis in the western North Atlantic, Nature, 356, 229-231.

Platt T. and Sathyendranath S. (1993) Estimators of primary production for interpretation of remotely sensed data on ocean color, J. Geophys. Res., 98, 14561-14576.

Pollehne F., Cline B. and Zeitzschel B. (1993) Low light adaptation and export production in the deep chlorophyll maximum layer in the northern Indian Ocean, Deep-Sea Res. II, 40, 737-752.

Pomroy A.J. and Joint I.R. (1999) Bacterioplankton activity in the surface mixed layer of the Arabian Sea during and after the 1994 SW monsoon, Deep-Sea Res. II, 46, 767-794.

Prasanna Kumar S., Ramaiah N., Gauns M., Sarma V.V.S.S., Muraleedharan P.M., Raghukumar S., Dileep Kumar M. and Madhupratap M. (2001) Physical forcing of biological productivity in the Northern Arabian Sea during the Northeast Monsoon, Deep-Sea Res., 48, 1115-1126.

Prasanna Kumar S., Muraleedharan P.M., Prasad T.G., Gauns M., Ramaiah N., De Souza S.N., Sardesai S., and Madhupratap M. (2002) Why is the Bay of Bengal less productive during summer monsoon compared to the Arabian Sea? Geophys. Res. Lett., 29 (24), 2235, doi: 10.1029/2002GL016013.

Price N.M., Ahner B.A. and Morel F.M.M. (1994) The equatorial Pacific Ocean: Grazer controlled phytoplankton populations in an iron-limited ecosystem, Limnol. Oceanogr., 39, 520-534.

Qasim S.Z. (1977) Biological productivity of the Indian Ocean, Ind. J. Mar. Sci., 6, 122-137.

Qasim S.Z. (1982) Oceanography of the northern Arabian Sea, Deep-Sea Res., 29, 1041-1068.

Radhakrishna K. (1978) Primary productivity of the Bay of Bengal during March-April, 1975, Ind. J. Mar. Sci., 7, 58-60.

Radhakrishna K., Devassy V.P., Bhargava R.M.S. and Bhattathiri P.M.A. (1978) Primary production in the northern Arabian Sea, Ind. J. Mar. Sci., 7, 271-275.

Ramage C.S. (1971) Monsoon Meteorology, Academic Press, New York, 296.

Ramaiah N., Raghukumar S. and Gauns M. (1996) Bacterial abundance and production in the central and eastern Arabian Sea, Curr. Sci., 71(11), 878-882.

Rau G.H., Sullivan C.W. and Gordon L.I. (1991) $\delta^{13}C$ and $\delta^{15}N$ variations in Weddell Sea particulate organic matter, Mar. Chem., 35, 355-369.

Rau G.H., Low C., Pennington J.T., Buck K.R. and Chavez F.P. (1998) Suspended particulate nitrogen $\delta^{15}N$ versus nitrate utilization: Observations in Montery Bay, CA, Deep-Sea Res. II, 45, 1603-1616.

Redfield A.C. (1934) On the proportions of organic derivatives in seawater and their relation to the composition of plankton, James Johnstone Memorial Volume, Liverpool, 176p.

Rees A.P., Owens N.J.P., Heath M.R., Plummer D.H. and Bellerby R.S. (1995) Seasonal nitrogen assimilation and carbon fixation in a fjordic sea loch, J. Plankton Res., 17(6), 1307-1324.

Rees A. P., Woodward E.M.S., Robinson C., Cummings D. G., Tarran G. A. and Joint I. (2002) Size-fractionated nitrogen uptake and carbon fixation during a developing coccolithophore bloom in the North Sea during June 1999, Deep-Sea Res. II, 49, 2905-2927.

Richards F.A. (1965) Anoxic basins and Fjords; In: Chemical Oceanography I (Eds) J.P. Riley and G. Skirrow, 611-645, Academic Press, New York.

Roman M.R., Dam H.G., Gauzens, A.L. and Napp J.M. (1993) Zooplankton biomass and grazing at the JGOFS Sargasso Sea time series station, Deep-Sea Res., 40, 883-901.

Ryther J.H., Hall J.R., Pease A.K., Bakun A. and Jones M.M. (1966) Primary production in relation to the chemistry and hydrography of the western Indian Ocean, Limnol. Oceanogr., 11, 371-380.

Saino T. and Hattori A. (1980) ^{15}N natural abundance in oceanic suspended particulate matter, Nature, 283, 752-754.

Saino T. and Hattori A. (1985) Variation of ^{15}N natural abundance of suspended organic matter in shallow oceanic water, In: Marine and Estuarine Geochemistry (Eds) A.C. Sigleo and A. Hattori, 1-13, A.F. Lewis, New York.

Saino T. and Hattori A. (1987) Geographical variation of the water column distribution of suspended particulate nitrogen and its ^{15}N natural abundance in the Pacific and its marginal seas, Deep-Sea Res., 34, 807-827.

Sambrotto R.N., Niebauer H.J., Goering J.J. and Iverson R.L. (1986) The relationship among vertical mixing, nitrate uptake, and growth during the spring phytoplankton bloom in the southeast Bering Sea middle shelf, Cont. Shelf Res., 5, 161-198.

Sambrotto R.N. and Lorenzen C.J. (1987) Phytoplankton, phytoplankton production in the coastal, oceanic areas of the gulf of Alaska, In: Hood D.W., Zimerman S.T. (Eds) The Gulf of Alaska: physical environment, biological resources. U.S. Dept. of Commerce, Washington, D.C., 249-282.

Sambrotto R.N., Martin J.H., Broenkow W.W., Carlson C.A. and Fitzwater S.E. (1993) Nitrate utilization in surface waters of the Iceland Basin during spring and summer of 1989, Deep-Sea Res. II, 40, 441-457.

Sambrotto R.N. (2001) Nitrogen production in the northern Arabian Sea during the spring intermonsoon and southwest monsoon seasons, Deep-Sea Res. II, 48, 1173-1198.

Sarin M.M., Krishnaswami S., Dilli K., Somayajulu B.L.K. and Moore W.S. (1989) Major ion chemistry of the Ganga-Brahmaputra river system: weathering processes and fluxes to the Bay of Bengal, Geochim. Cosmochim. Acta, 53, 997-1009.

Sarmiento J.L. and Siegenthaler U. (1992) New production and Global Carbon Cycle in Primary Productivity and Biogeochemical Cycles, In: The Sea, (Eds) Falkowski P. and Woodhead A.D., 317-332, Plenum, NewYork.

Sathyendranath S. and Platt T. (1988) The spectral irradiance field at the surface and in the interior of the ocean: A model for applications in oceanography and remote sensing, J. Geophys. Res., 93, 9270-9280.

Savidge G. and Gilpin L. (1999) Seasonal influences on size fractionated Chlorophyll a concentrations and primary production in the north-west Indian Ocean, Deep-Sea Res. II, 46, 701-723.

Sawant S. and Madhupratap M. (1996) Seasonality and composition of phytoplankton in the Arabian Sea, Curr. Sci., 71, 869-873.

Schafer P., Kreilein H., Muller M. and Gravenhorst G. (1993) Cycling of inorganic nitrogen compounds between atmosphere and ocean in tropical areas off south East Asia, Mitt. Geol.-Palaont. Inst. Univ. Hamburg, 76, 19-36.

Schafer P. and Ittekkot V. (1995) Isotopic biogeochemistry of nitrogen in the northern Indian Ocean, Mitt. Geol. -Palaont. Inst. Univ. Hamburg, 78, 67-93.

Schoenheimer R. and Rittenberg D. (1939) Studies in protein metabolism: I, General consideration in the application of isotopes on the study of protein metabolism, The normal abundance of nitrogen isotopes in amino acids.

Schott F. and McCreary J.P. (2001) The monsoon circulation of the Indian Ocean, Progr. Oceanogr., 51, 1-123.

Schott F., Swallow J.C. and Fieux (1990) The Somali Current at the equator: annual cycles of currents and transport in the upper 1000m and connection with neighbouring latitudes, Deep-Sea Res., 37, 1825-1848.

Shearer G. and Kohl D. (1986) N_2 fixation in field settings, estimation based on natural ^{15}N abundance, Australian J. Plant Physiol., 13, 699-757.

Shetye S.R., Gouveia A.D., Shenoi S.S.C., Sundar D., Michael G. S., Almeida A. M. and Santanam K. (1990) Hydrography and circulation off the west coast of India during the Southwest monsoon 1987, J. Mar. Res., 48, 359-378.

Shetye S.R., Shenoi S.S.C., Gouveia A.D., Michael G. S., Sundar D. and Nampoothiri G. (1991) Wind-driven coastal upwelling along the western boundary of the Bay of Bengal during the southwest monsoon, Cont. Shelf Res., 11, 1397-1408.

Shetye S.R., Gouveia A.D., Shenoi S.S.C., Sundar D., Michael G. S. and Nampoothiri G. (1993) The western boundary current of the seasonal subtropical gyre in the Bay of Bengal, J. Geophys. Res., 98, 945-954.

Shetye S.R., Gouveia A.D., Shankar D., Shenoi S.S.C., Vinaychandran P.N., Sundar D., Michael G. S. and Nampoothiri G. (1996) Hydrography and circulation in the western Bay of Bengal during northeast monsoon, J. Geophys. Res., 101,14011-14025.

Smith R.L. and Bottero J.S. (1977) On upwelling in the Arabian Sea, In: A voyage of discovery (Ed) M. Angel, 291-304, Pergamon Press.

Smith S.L., Codispoti L.A., Morrison, J.M. and Barber R.T. (1998) The 1994-1996 Arabian Sea Expedition: an integrated interdisciplinary investigation of the response of the northwestern Indian Ocean to monsoonal forcing, Deep-Sea Res. II, 45, 1905-1915.

Smith S. L. (2001) Understanding the Arabian Sea: Reflections on the 1994-1996 Arabian Sea Expedition, Deep-Sea Res. II, 48, 1385-1402.

Smith W.O. and Nelson D.M. (1990) Phytoplankton growth and new production in the Weddell Sea marginal ice zone in the austral spring and autumn, Limnol. Oceanogr., 35, 809-821.

Smith W.O. Jr (1995) Primary productivity and new production in the northeast water (Greenland) Polynya during summer 1992, J. Geophys. Res., 100, 4357-4370.

Smith W.O. Jr., Barber R.T., Hiscock M.R. and Marra J. (2000) The seasonal cycle of phytoplankton biomass and primary productivity in the Ross Sea, Antarctica, Deep-Sea Res. II, 47, 3119-3140.

Steinberg D.K., Carlson C A., Bates N.R., Goldthwait S.A. Madin L.P. and Michaels A.F. (2000) Zooplankton vertical migration and the active transport of dissolved organic and inorganic carbon in the Sargasso Sea, Deep-Sea Res. I, 47, 137-158.

Strickland J.D.H. and Parson T.R. (1972) A practical handbook of seawater analysis, 127-130, Fisheries Research Board, Canada.

Subramanian V. (1993) Sediment load of Indian Rivers, Curr. Sci., 64, 928-930.

Sullivan C.W., Arrigo K.R., McClain C.R., Comiso J.C. and Firestone J. (1993) Distribution of phytoplankton blooms in the Southern Ocean, Science, 262, 1832-1837.

Sweeney R.E., Liu K.K. and Kaplan I.R. (1978) Oceanic nitrogen isotopes and their use in determining the source of sedimentary nitrogen, in: Stable isotopes in the earth sciences, edited by Robinson, B.W., DSIR 220 New Zealand dept. Scientific and industrial, Wellington, 9-26.

Takahashi, T. (1989) The carbon dioxide puzzle, Oceanus, 32, 22-29.

Takahashi, T., Olafsson J., Goddard J., Chipman D.W. and Sutherland S.C. (1993) Seasonal variation of CO_2 and nutrients in the high latitude surface oceans: A comparative study, Global Biogeochem. Cycles, 7, 843-878.

Tomczak M. and Godfrey J.S. (1994) Regional Oceanography: an Introduction, 422, Pergamon, Terry town, New York.

Townsend D.W., Keller M.D., Sieracki M.E. and Ackleson S.G. (1992) Spring phytoplankton blooms in the absence of vertical water column stratification, Nature, 360, 59-62.

Treguer P., Legendre L., Rivkin R.T., Ragueneau O. and Dittert N. (2003) Water column biogeochemistry below the euphotic zone, In: Ocean Biogeochemistry (Ed) Fasham M.J.R., 145-156, Springer, Heideberg.

Unger D., Ittekkot V., Schafer P., Tiemann J. and Reschke S. (2003) Seasonality and interannual variability of particle fluxes to the deep Bay of Bengal: influence of riverine input and oceanographic processes, Deep-Sea Res. II, 50, 897-923.

Urey, H.C. (1947) The thermodynamic properties of isotopic substances, J. Chem. Soc., 562-581.

Venkateswaran S.V. (1956) On evaporation from the Indian Ocean, Indian Journal of meteorology and Geophysics, 7, 265-284.

Villareal T.A., Altabet M.A. and Culver-Rymsza K. (1993) Nitrogen transport by vertically migrating diatom mats in the North Pacific Ocean, Nature, 363, 709-712.

Vinaychandran P.N., Murty V.S.N. and Ramesh Babu V. (2002) Observations of barrier layer formation in the Bay of Bengal during summer monsoon, J. Geophy. Res., 107(C12), 8018, doi: 10.1029/2001JC000831.

Vinaychandran P.N. and Mathew S. (2003) Phytoplankton bloom in the Bay of Bengal during the northeast monsoon and its intensification by cyclones, Geophys. Res. Lett., 30 (11), 1572, doi: 10.1029/2002GL016717.

Vitousek P.M., Hattenschwiler S., Olander L. and Allison S. (2002) Nitrogen and nature, Ambio, 31, 97-101.

Wada E., Miyazaki T. and Hattori A. (1971) ^{15}N abundance in nitrogenous compounds in the sea, Proc. 1971 Fall Meet. Oceanogr. Soc. Japan, 227.

Wada E. and Hattori A. (1976) Natural abundance of ^{15}N in particulate organic matter in the North Pacific Ocean, Geochim. Cosmochim. Acta, 40, 249-251.

Wada E. and Hattori A. (1978) Nitrogen isotope effects in the assimilation of inorganic nitrogenous compounds by marine diatoms, Geomicrobiol. J., 1, 85-101.

Wada E. (1980) Nitrogen Isotope fractionation and its significance in biogeochemical processes occurring in marine environments, In :Isotope marine Chemistry, 375-398.

Wada E. and Hattori A. (1991) Nitrogen in the sea: Forms, Abundances, and rate processes, CRC Press, Boca Raton, Fla, 208.

Wafar M.V.M., Wafar S. and Devassy V.P. (1986) Nitrogenous nutrients and primary production in a tropical oceanic environment, Bull. Mar. Sci., 38(2), 273-284.

Wafar M.V.M., Le Corre P. and L'Helguen S. (1995) f-Ratios calculated with and without urea uptake in nitrogen uptake by phytoplankton, Deep-Sea Res., 42,1669-1774.

Walsh J.J., (1981) A carbon budget for overfishing off Peru, Nature, 290, 300-304.

Waser N.A., Yin K., Yu Z., Tada K., Harrison P.J., Turpin D.H. and Calvert S.E. (1998) Nitrogen isotope fractionation during nitrate, ammonium and urea uptake by marine diatoms and coccolithophores under various conditions of N availability, Mar. Ecol. Prog. Ser., 169, 29-41.

Watson A.J. and Orr J.C. (2003) Carbon dioxide fluxes in the global ocean, In: Ocean Biogeochemistry (Ed) Fasham M.J.R., 123-143,Springer, Heideberg.

Watts L.J. and Owens N.J.P. (1999) Nitrogen assimilation and f-ratio in the northwestern Indian Ocean during an Intermonsoon period, Deep-Sea Res. II, 46, 725-743.

Wheeler P.A., Glibert P.M. and McCarthy J.J. (1982) Ammonium uptake and incorporation by Chesapeake Bay phytoplankton: short term uptake kinetics, Limnol. Oceanogr., 27, 1113-1128.

Wheeler P.A. and Kokkinakis S.A. (1990) Ammonium recycling limits nitrate use in the oceanic subarctic Pacific, Limnol. Oceanogr., 35, 1267-1278.

Wiebe P.H., Boyd S. and Cox J.L. (1975) Relationship between zooplankton displacement volume, wet weight, dry weight, and carbon, Fishery Bulletin, 73(4), 777-786.

Wilkerson F.P., Dugdale R.C. and Barber R.T. (1987) Effects of El Nino on new, regenerated, and total production in eastern boundary upwelling systems, J. Geophys. Res., 92, 14347-14353.

Wong C.S., Whitney F.A., Matear R.J. and Iseki K. (1998) Enhancement of new production in the northeast subarctic Pacific Ocean during negative north Pacific index events, Limnol. Oceanogr., 43, 1418-1426.

Woodward E.M.S., Rees A.P. and Stephens J.A. (1999) The influence of south-west monsoon upon the nutrient biogeochemistry of the Arabian Sea, Deep-Sea Res. II, 46, 571-591.

Wyrtki K. (1973) Physical Oceanography of the Indian Ocean, In: The Biology of the Indian Ocean (Eds) Zeitzschel B. and Gerlach S.A., 18-36, Chapman Hall Ltd., London.

Zehr J.P., Waterbury J.B., Turner P.J., Montoya J.P., Omoregie E., Steward G.F., Hansen A. and Karl D.M. (2001) Unicellular cynobacteria fix N_2 in the subtropical North Pacific Ocean, Nature, 412, 635-638.