BIBLIOGRAPHY

BIBLIOGRAPHY

Ahmed, S.M. et al. 1977. Special Publication, Archaean geochemistry, NGRI, Hyderabad.

Allen, P. 1984. The geochemistry of the amphibolite-granulite facies in Central South India. Unpubl. Ph.D. Thesis, New Mexico Institute of Mining and Technology, Socorro, USA.

Aranovich, L, Y. and Podlesskii, K.K.(1989) Geothermobarometry of high grade metapelites : simultaneous operating reactions. In : Saxena S.K. (ed) Kinetics and equilibrium in mineral reactions, Springer Verlag, N.Y., pp. 173-198

Auden, J B. (1931) General Report for 1930, Rec Geol Surv. Ind , LXV, 1· 133 -144.

Balaram, V. (1991) Determination of rare earth elements in geological samples by Inductively coupled plasma - mass spectrometry, Ind. Chem. Soc. v.68, pp 66-603.

Berman, R.G. (1988) Internally consistent thermodynamic data for minerals in the system Na2)-K2O-CaO-MgO-FeO-A12O3-SiO2-TiO2- H2O-CO2. J. Petrol. **52**: 445-522. 29, 495-522.

Bhaskara Rao, Y.J., Warren Beck, Rama Murthy, V., Nirmal Charan, S. and Naqvi, S.M., (1983) Geology, geochemistry and age of metamorphism of the Archaean grey gneisses around Channaraya Patna, Hassan Dist., Karnataka, South India. In : Naqvi and Rogers (Eds) Precambrian of South India, Geol. Soc.India, Mem.4, pp. 309-328.

Bhattacharya, A., (1973) Petrology of leptynites and garnetiferous granite gneisses around Garbham, Srikakulam Dist., Andhra Pradesh. J. Geol. Soc. Ind. v.14, pp. 113-123.

Bhattacharya, A., Krishna Kumar, K.R., Raith, M. and Sen, S.K. (1991) An improved set of a-x parameters for the Fe-Mg-Ca garnets and refinements of the orthopyroxene-garnet thermometer and the orthopyroxene-garnet-plagioclase. J. Petrol. v.32, pp.629-656.

Bhattacharya, A., Mazumdar, A.C. and Sen, S.K. (1988) "Fe-Mg mixing in cordierite : Constraints from natural data and implications for cordierite garnet geothermometry in granulites". Amer. Mineral. v.73, pp.338-344.

Bhattacharya, A., Mohanty. L., Maji, A., Sen, S.K. and Raith, M. (1992) Non-ideal mixing in the phlogopite-annite binary : Constraints from experimentaldata on Fe-Mg partitioning and reformulation of biotite-garnet geothermometer. Contri. Mineral. Petrol., v.111, pp. 87-93.

Bhattacharya, A., Spiering, B., Sen, S.K. , Natarajan, R. and Mazumdar, A.C. (1990) Compositional characteristics and phase equilibria in manganiferous iron formations from high grade terrain near Satnuru, Karnataka, India. J. Met. Geol. 8, 525-538

Bhattacharya, J. Golani, P.R. & reddy, A.B. (1988) Rift related bimodal volcanism and metallogeny in the Delhi fold belt, Rajasthan and Gujarat. Ind. Geol., 60: 191-199.

Binns, R.A. (1962) Metamorphic pyroxenes from the Broken Hill District, New south Wales, Mineral Mag. v.33, pp.320-338.

Binns, R.A. (1964) Zones of progressive regional metamorphism in the Willyam Complex, Broken Hill District, New South Wales, Geol. Soc. Australia J. 11: 283-330.

Biswal, T.K. (1988) Polyphase deformation in Delhi rocks southeast of Amirgarh, Banaskantha Dist., Gujarat. Mem. Geol. Soc. Ind. 7: 267-277.

Biswal, T.K. (1996) Structural and geochemical evidence of a continental Arc in granulite-charnockite rocks of the Delhi Supergroup from from Balaram-Mawal area, Gujarat, Ind. Abs. International Seminar on Charnockites and Granulites, Madras (29-31 August, 1996).

Blundy, J.D. and Holland, T.J.B. (1990) Calcic amphibole equilibria and new amphibole-plagiocalse geothermometer. Contr. Mineral. Petrol. v.104, pp. 208-224.

Bohlen, S.R. (1987) Pressure-temperature-time paths and a tectonic model for the evolution of granulites. J. Geology **95**: 617-632.

Bohlen, S.R. (1991) On the formation of granulites. J. Meta. Geol. 9, 223-229

Bowen, N.L., Schairer, J.F. & Posnjak, E. (1933) The system CaO- FeO-SiO2, Amer. J. Sci. 5th Ser. v.26, p.193.

Brown, M. and Raith (1986) First evidence of ultrahigh - temperature decompression from the granulite province of southern India. J. Geol. Soc. London, 153: 819-822.

Bucher-Nurminen & Droop, Giles. (1983) The metamorphic evolution of garnet-cordierite-sillimanite-gneisses of the Gruf-Complex, Eastern Pennine Alps. Contrib. Mineral. Petrol. 84: 215-227.

Caporuscio, F.A. and Morse, S.A. (1978) Occurrence of sapphirine + quartz at Peekskill, New York. Amer. J. Sci. 278: 1334-1352.

Chakraborty, K.R. & Sen, S.K. (1983) a comparative analyses of geochemical trends in basic granulites, greenstones and recent basalt based on three basic granulite suites from India. Geol. Soc. Ind. Mem. 4. S.M.Naqvi & Rogers (Eds.), 462-487.

Chinner, G.A. (1961) The origin of sillimanite in Glen Clova, Angus, Scotland. J. Petrol. 3: 312-323.

Clemens, J.D. and Vielzeuf, D. (1987) Constraints on melting and magma production in the crust. Earth Planet. Sci. Lett. 86: 287-306.

Condie, K. (1976 c) Trece element geochemistry of Archaean belts, Earth Sci. Rev. v.12, pp. 393-417.

Condie, K. (1981 a) Archaean Greenstone Belts, Elsevier, Amsterdam. 434 pp.

Condie, K. and Allen, P. (1984) Origin of Arcchaean Charnockites from south India. In : A.Kroner, G.N.Hanson and A.M.Goodwith (Eds.) Archaean Geochemistry. pp. 182-203. Springer-Verlag. Berlin Heidelberg.

Cooray, P.G. (1960) Khondalites and charnockites of Laggala Pallegma area, Ceylon. Bull. Mys.Geol.Assn. v.18, p.3-54.

Coulson, A.L.(1933). The Geology of Sirohi State, Rajputana. Mem.

Geol. Surv. India. **6LXIII**(1), 1-166.

Crawford, A.R. (1970) The Precambrian geochronology of Rajasthan and Bundelkhand, Northern India, Can. J. Earth Sci.,v.7(1): 91-110.

Dasgupta, S., Sengupta, P., Guha, D. and Fukuoka, M. (1991) A refined garnet-biotite Fe-Mg exchange : geothermometer and its application to amphibolites and granulites. Contr. Mineral. Petrol. v.109, pp. 130-137.

Dasgupta, S., Sengupta, P., Juergen Ehl, Raith, M. & Bardhan, S. (1995) Reaction textures in a suite od spinel granulites from the Eastern Ghats Belt, India, Evidence for polymetamorphism a partial petrogenetic grid in the system KFMASH and the roles of ZnO and Fe2O3. J.Petrology, 36(2): 453-461.

Deb, M., Talwar, A.K., Anju Tiwary and Banerjee, A.K. (1995) Bimodal volcanism in south Delhi Fold Belt I: A suite of differentiated felsic lava at Jharivav, North gujarat. Mem. Geol. Soc. Ind. No.31: 259-278.

Deb, M., Thorpe, R.I., Cumming, G.L. and Wagner, P.A. (1989) Age, source and stratigraphic implications of Pb isotope data for conformable, sediment-hosted, basemetal deposits in the Proterozoic Aravalli-Delhi orogenic belt, north-western India, precambr. Res. 43: 1-22.

Deer, W.A., Howie, R.A. & Zussman, J. (1974) Rock forming minerals, V.II, Longmans.

Desai, S.J. (1974) Petrologial studies of high grade Metamorphic rocks from an area between Balaram and Kanpura, Banaskantha district, Gujarat, India. Unpublished thesis for DIC, Imperial College of Science and Technology, London, P. 1- 115.

Desai, S.J. & Patel, M.P. (1989) International Seminar on Granitoids, Helsinki. Abs. Vol.

Desai, S.J., Patel, M.P. and Merh, S.S. (1978)Polymetamorphites of Balaram -Abu Road area north Gujarat and southwest Rajasthan. J. Geol. Soc. Ind. v.19(9), p.383-394.

Desai, S.J. and Patel, M.P. (1992). Granulite rocks of the Banas-kantha District of North Gujarat, India. in High grade metamorphics ed.A.Barbo-Kyriakidis. Theophrastus Pub.-S.A., Athens, Greece, p.103-121.

Deshpande, G.G., Powar, K B., Peshwa, V V. & Sukhtankar, R K (1970) Occurrence of hornfels granulite transitional facies in the metasediments of Danta area, Gujarat. J. Ind. Geol. Asso. 12: 72-82.

Devaraju, T.C. and Sadashivaiah, M.S. (1975) Structural geology of granulite area in southern Karnataka. J.Karn. Uni., 20, 99-123.

Dewaard, D. (1964) Mineral assemblages and metamorphic subfacies in the granulite facies terrane of the little Moose mountain syncline, south central Adirondack Highlands, Koninkl. Nederlandse Akad. Weensch. Proc., Amsterdam, B, 67, 344-362.

Dewaard, D. (1965) A proposed subdivision of the granulite facies. Am.J Sci., 263, 455-461.

Droop, G.T.R. (1987) A general equation for estimating Fe3+ concentrations in ferromagnesian silicates andoxides from microprobe analyses, using stoichiometric criteria. Mineral Mag. v.51, pp. 431-435.

Droop, G.T.R. & Bucher-Nurminen, K. (1984) Reaction textures and metamorphic evolution of sapphirine bearing grnulites from the Gruf Complex, Italian Central Alps. J. Petrl., 25(3): 766-803.

Elkins, L.T. and Grove, T.L. (1990) Ternary feldspar experiments and thermodynamic models Am. Mineral. v.75, pp. 544-559.

Ellis D.J. (1980) "Osumilite-sapphirine-quartz granulites from Enderby Land, Antarctica: P-T conditions of metamorphism, implications for garnet-cordierite equilibria and the evolution of the deep crust". Contrib. Miner. and Petrol.,V.74,N.2,p.201-210, 1980.

Ellis, D.J. (1986) Garnet-liquid Fe2+-Mg equilibria and implications for the beginning of melting in the crust and subduction zones. Am. J. sci. v. 286, pp. 765-791.

Ellis, D.J. (1987) Origin and evolution of granulites in normal and thickened crusts. Geology, 15: 167-170.

Ellis, D.J. and Green, D,H. (1979) An experimental study of effect of Ca upon garnet-clinopyroxene exchange equilibria. Contr. Mineral. Petrol. v.71, pp. 13-22.

Ellis, D.J. and Green, D,H. (1985) Garnet forming reactions in mafic granulites from Enderby Land, Antarctica - Implications from geothermometry and geobarometry. J. Petrol., 26, 633-662.

Ellis, D.J., Sheraton, J.W., England, R.N. and Dallwitz, W.B. (1980) Osumilite-Sapphirine-Quartz granulites from Enderby Land, Antarctica : Mineral assemblages and reactions. Contrib. Mineral. Petrol. 72: 123-143.

England, P.C. and Thompson, A. (1984) Pressure-temperature-time paths of regional metamorphism, I. Heat transfer during the evolution of regions of thickened continntal crust. J. Petrol., 25, 894-928.

Eskola, P. (1952) On the granulites of Lapland, Am. J. sci, Bowen Vol., 135-171.

Fareeduddin (1995). Field setting, petrochemistry and P-T regime of the deep crustal rocks to the Northwest of the Aravalli- Delhi mobie belt, North-Central Rajasthan. Mem.Geol.Soc.Ind. **31**: 117-139

Fareeduddin, Sharma, A.K. and Bose, V. (1991) Tectonic slices of high grade rocks in the Delhi fold belt of North central Rajasthan . J.Geol. Soc. Ind. 38, 427-436.

Fermor, L.L. (1936) An attempt at correlation of ancient schistose rocks of Peninsular India. Mem. Geol. Surv. Ind. v.70, Pt.3, pp. 325.

Ferry, J.M. and Spear, F.S. (1978) Experimantal calibration of the patitioning of Fe and Mg between biotite and garnet. Contrib. minera. Petrol. 66 : 113-117.

Field, D., Drury, S.A. and Copper, D.C. (1980) Rare Earth and LIL Element fractionation in high grade charnockitic gneisses. South Norway. Lithos, v.13, pp. 281-289.

Fitzisimons, I.C.W. and Harley, S.L., 1994. Garnet coronas in scapolite-wollastonite calc-silicates from East antactica : the applications and limitations of activity corrected grids. J. Meta. Geol., 12, 761-777.

Fonarev, V.I. and Graphichikov, A.A. (1991) Two pyroxene thermometry : A critical evaluation In : Perchuk, L.L. (Ed.) Progress in metamorphic and magmatic petrology. Cmbridge Univ. Press, pp. 65-92.

Fonarev, V.I. Graphichikov, A.A. and Konilov, A.N. (1991 b) A consistent system of geothermometers for metamorphic complexes. Int. Geol. Rev. v.33, pp. 743-783.

Frisch, T., 1988. Reconnaissance geology of the Precambrianshield of Ellesmere, Devon and Coburg Islands, Canadian Arctic Archipelago. Geology Survey of Canada Mem., 409, 102 pp.

Frost, B.R. and CHacko, T., (1989)Thegranulite uncertainty principle : limitations on thermobarometry in granulites.J. Geol., 97, 435-450.

Fyfe, W. (1973) the granulite facies, partial meltingand the Archaean crust. Phil. Trans. Royal Soc. London. v.A273, pp. 457-461.

Gandhi, S.M., Paliwal, H.V. and Bhtnagar, S.M. (1984) Geology and ore reserve estimates of Rampura-Agucha zic lead deposits, district Bhilwara, Rajasthan, J.Geol. Soc. Ind., 25, 689-705.

Gaeb.-Jahrb. Pet-ol.-85, 248-255

Gangopadhyay, P.K. (1970) Structure and tectonics of Alwar region in Rajasthan with special reference to Precambrian stratigraphy. Proc. Ind. Sci. Cong.

Gangopadhyay, P.K. & Sen, R. (1971) Trend of regional metamorphism, an example of Dehi System of rocks occuring around Bairawas, N E Rajasthan Proc Ind Sci.Cong.

Ghosh, P K (1941) Charnockite series of Bastar State Rec. Geol. Surv. India, 75, paper 15.

Ghosh, S.K. and Naha, K. (1962) recumbent folding in the migmatites in the Archaean basement north of Nathdwara, Rajasthan, Ind. Am. J. Sci. 260: 241-248.

Goscombe Ben (1992) High-grade reworking of Central Australian granulites : Metamorphic evolution of the Arunta Complex. J. Petrol., 33(4): 917-962.

Green, D.H. and Ringwood, A.E. (1967) An experimental investigation of the gabbro to eclogite transfromationnad its petrological applications. Geochem. et. Cosmochimica Acta., 31, 767-833.

Grew, E.S. (1982b) Sapphirine, Kornerupine and sillimanite + Orthopyroxene in the charnockite region of South India. J. Geol. Soc. Ind., 23: 469-505.

Guha, D.B. (1996) Granulites of Sandmata Complex, Rajasthan, India -PT Conditions and evolution. Abs. International Seminar on Charnockites and Granulites, Madras (29-31 August, 1996).

Guha, D.B. and Bhattacharya, A.K. (1995). Metamorphic evolution

and high-grade reworking of Sandmata Complex Granulites. Mem.Geol.Soc.Ind. **31**: 163-198.

Gupta, B.C. (1934) The geology of Central Mewar. Rec. Geol. Surv. Ind. v.65, pt.2.

Gupta, S.N., Mathur, R.K. & Arora, Y.K. (1992). Lithostratigraphy

of Proterozoic rocks of Rajasthan and Gujarat - A review. Rec. Geol. Surv. Ind., **115** (7 & 8) : 63-85.

Gyani, K.C. (1970) Granulitic rocks from BGC of Bandanwara region, Ajmer Distt., Rajasthan. Proc. Sec. Symp. Upper Mantle Proj., Hyderabad. pp. 339-348.

Gyani, K.C. (1971) Petrogenetic studies of amphibolites from Bandanwara region of Central Rajasthan. 58th Sess. Ind. Sci. Cong.Assoc., Abst. p.250.

Gyani, K.C. and Omar, Imad, E.M. (1996) Charnockites from Banded Gneissic Complex of Central Rajasthan : Geochemical Constraints and petrogenesis. ISCGFR, Madras (Abs.), pp. 63- 64.

Halferdahl, L.B. (1961). 'Chloritoid : Its composition, X-ray and optical properties, stability and occurrence', J. Petr., v.2, p.49

Hammarstrom, J.M. and Zen E-An (1986) Aluminium in hornblende : An empericaligneous geobarometer. Amer. Mineral. v.71, pp. 1297-1313.

Harker, A. (1950) Metamorphism a study of the transformation of rock masses (Ed.) Methuen, London, Dutton, Dutton, New York, pp. 362.

Harker, R.I. & Tuttle, O.F. (1956) The experimentally determined univariant PCO2 - T curve for the reaction CaCO3 + SiO2 = CaSiO3 +CO2. Amer. J.Sci. v. 254 : 239

Harley, S.L. (1984) Comparison of the garnet-orthopyroxene geobarometer with recent experimental studies and applications to natural assemblages. J. Petrol., 25, 697-712.

Harley, S.L. (1984b) An experimental study of the partitioning of Fe and Mg between garnet and orthopyroxene. Contrib. Mineral. Petrol. 86: 359-373.

Harley, S. (1989) The origin of granulites: a metamorphic perspective. Geol. Mag. 126: 215-247.

Harley, S.L., Fitzsimans, J.C.W. and Buick, I.S., (1994) Reactions and textures in wollastonite-scapolite granulites and their significance for pressure-temperature-fluid histories of high grade terranes. Precambrian Research, 66, 309-323.

Harley, S. & Buick Ian, S. (1992) Wollastonite-Scapolite assemblages as indicators of granulite P-T-Fluid histories : The Rauer Group, E. Antarctica. J. Petrology, 33(3) : 693- 728.

Harley, S.L. & Gren, D.H. (1982) Garnet-orthopyroxene barometry for granulites and peridotites. Nature., 300: 699-701.

Harris, N.R.W. and Holland, T.J.R. (1984) The significance of cordierite-hypersthene assemblages from the Beitbridge region of the Central Limpopo Belt; evidence for rapid decompression in the Archaean ?. Amer. Mineralogist, 69: 1036-1049.

Haselton, H.T., Jr., Hovis, G.L., Hemingway, B.S. and Robert, R.A. (1983) Calorimetric investigation of the excess entropy of mixing in an albite sanidine solid solution : Lack of evidence for Na, K short-range order and implications for the two feldspar thermometry. Amer. Mineral., v.68, pp. 398- 413.

Henry, D.J. and Medaris, L.G. (1976) Application of pyroxene and olivine-spinel geothermometers to the alpine peridotites in southwestern Oregon Geol. Soc Amer. Abstracts with programs, v.8, pp. 913-914.

Heron, A.M. (1917) The geology of North eastern Rajputana and adjacent districts. Mem. Geol. Surv. Ind. v.44, Pt.1, pp. 28.

Heron, A.M. (1935) Synopsis of the Pre-Vindhyan Geology of Rajputana. Trans. Nat. Inst. Soc. Ind.V.1, No.2, pp. 17-33.

Heron, A.M. (1953) The geology of Central Rajputana, Mem. Geol. Surv. Ind. 72(4) : 367-411.

Heron, A.M. and Ghosh, P.K. (1938) The geology of Palanpur, Danta and part of Idar states. Rec. Geol. Surv. Ind., 79 : 367-412.

Hirschberg, A. and Winkler, A.G.F. (1968) Stabiletatbe Jiehunger zuischen chlorit, cordierit and almandin bei der metamorphose. Contrb. Mineral. Petrol. v.18, pp. 17-42.

Hodges K.V., Spear F.S. (1982) "Geothermometry, geobarometry and the Al2SiO5 triple point at Mt. Moosilauke, New Hempshire". Amer. Mineral.,V.67,N.11/ 12,p.1118-1134,1982

Hodges K.V., Royden L. (1984) "Geologic thermobarometry of retrograded metamorphic rocks: an indication of the uplift trajectory of a portion of the Northern Scandinavian Caledonides". J. of Geophysicaal Research, N.B8,V.89,p.7077- 7090,1984

Holdaway, M.J. and Lee, S.M. (1977) Fe-Mg Cordierite stability in high-grade pelitic rocks based on experimental, theoretical and natural observations. Contrib. Mineral. Petrol. 63: 175- 198.

Holland, T.H. (1900) The charnockite series, a group of Archaean Hypersthenic rocks in Peninsular India. Mem. Geol. Surv. India 28.

Hollister, L.S., Grissom, G.C., Peters, S.K., Stowell, H.H. and Sisson, V.B. (1987) Confirmation of the empirical correlation of Al in hornblende with pressure of the solidification of calc-alkaline plutons. Amer. Mineral. v.72, pp. 231-239.

Huckenbolz, H.G., Lindhurber, W. and Fehr, K.T., 1981. Stability relationships of grossularite + quartz + wollastonite + anorthite I. The effect of andradite and albite. Nues Jahrbuch fur Mineralogie, 42, 223-278.

Irvine, T.N. and Barager, W.R.A. (1971) A guide to the classification of the common volcanic rocks. Can.J.Earth Sc. 8: 523-549.

Jacobs, G.K. and Kerrick, D.M. (1981) Devolatilization equilibria in H2O-CO2 and H2O-CO2-NaCl fluids : and experimental and thermodynamic evaluation at elevated pressures and temperatures. Am. Mineralogist., 66, 1135-1156.

Jahn, B.M. (1990) Origin of granulites : Geochemical constriants from Archaean granulite facies rocks of Sino-Korean Craton, China. In : Vilezeuf., D and Vidal, Ph. (Eds.) Granulites and crustal evolution, Kluwar Academic Pub. pp. 471-492, - Series C. V.311.

Jakes, P. and Taylor, S.R. (1974) Excess Europium content in Precambrian sedimentary rocks and continental evolution. Geochim Cosmochim. Acta. V.38, pp 739-745

Janardhan, A.S., Francis Anto, K. & Sivasubramaniam (1996) Reaction textures and fluid inclusions in calc silicate lithologies of Kambam Valley, Tamil Nadu. Abs. International Seminar on Charnockites and Granulites, Madras (29-31 August, 1996).

Janardhan, A.S., Newton, R.C. and Hansen, E.C. (1982) The transformation of amphibolite facies gneiss to charnockite in southern Karnataka and northern Tamil Nadu, India. Contrib. Moineral. Petrol. 79: 130-149.

Janardhan, A.S. & Sivasubramanian, P. (1996) Metamorphic evolution of the Palni Ranges, Tasmil Nadu, India. Abs. International Seminar on Charnockites and Granulites, Madras (29-31 August, 1996).

Jayananda, M., Peucat, J.J., Mahabaleshwar, B. & Janardhan, A.S. (1995) Magmatic underplating in the deep Archaean crust of Southern India : Implications for the origin of granulites and large scale hybridisation of continental crust.Abs. International Seminar on Charnockites and Granulites, Madras (29-31 August, 1996).

Jhingran, A.G. (1958) Bundelkhand gneiss. Ind. Sci. Cong. Pres. Add. Geol. Section.

Johnson M.C., Rutherford M.J. (1989b) "Experimetal calibration of the aluminium-in-hornblende geobarometer with application to Long Valley Caldera (California) volcanic rocks". J. Geology,V.17,p.837-841,1989 b

Keith, M. and Schairer, J.F. (1952). The stability field of sapphirine in the system MgO-Al2O3-SiO2, J. Geol., v.60, p.181.

Kretz, R (1964) Analysis of equilibrium in garnet-biotite- sillimanite gneisses from Quebec. J. Petrol. v.5, pp. 1-20.

Kretz, R. (1982) Transfer and exchange equilibria in a portion of pyroxene quadrilateral as deduced from natural and experimental data. Geochim. Cosmochim. Acta 46: 411-421.

Krupicka, J. (1973) Geology of Devon Island, Canadian shield.

Lal. R. K., Ackermand, D. & Upadhyay, H. (1987) P-T-X relatioships deduced from corona textures in sapphirine- spinel-quartz assemblages from Paderu, S.India. J.Petrol. 28: 1139-1168.

Langmuier, C.H., Voke, R.D , Hanson, G.N and Hart, S.R (1978) A general mixing equation with application to Icelandic basalts. Earth Planet. Sci. Lett. v.37, pp. 380-392.

Laverentyeva, I,V, and Perchuk, L.L. (1981b) Phase correspondence in the systembiotite-garnet experimental results : Doklady Akad Nauk SSSR, v.260, pp. 731-734.

Lee, H.I. and Ganguly, J. (1988) Equilibrium composition of coexisting orthopyroxene and garnet : experimental determination in the system FeO-MgO-Al2O3-SiO2, and applications. J. Petrol. v.29, pp. 93-113.

Leelanandam, C. (1970) Chemical mineralogy of hornblendes and biotites from the charnockitic rocks of Kondapalli, India. J. Petrology, 11(30: 475-505.

Limaye, M.A. (1992) Geochemistry, structure and metamorphism of rocks around Danta, Banaskantha Dist., N. Gujarat. Unpub. Ph.D. Thesis, M.S.Univ., Baroda.

Limaye, M.A. & Desai, S.J. (1995) Petrological studies of metamorphites of Danta area, N. Gujarat. Proc. Recent Res. Geol. W.India. Oct. 29-31, pp. 87-94.

Lindsey, D.H. (1983) Pyroxene thermometry. Am. Mineral. 68: 477- 493.

Lingadevaru, M., Mahabaleshwar, B. & Shadakshra Swamy, N. (1996) Petrology and geochemistry of mafic granulites from Honur area, Southern Karnataka, India. Abst. International Symposium on Charnockit and Granulite Facies Rocks, Madras, India. pp. 53.

Macdougal, J.D., Gopalan, K., Lugmair, G.W. and Roy, A.B. (1983). The Banded Gneissic Complex of India : Early crust from depleted mantle at 3.5 AE. Trans Amer. Geophys. Union, **64**: 381.

Mahadevan, T.M. (1964) The origin of Charnockite suite of rocks forming part of the Western Ghats in Kerala, Ind. Int. Geol. Cong. XIII, 88-96, New Delhi.

Mahadevan, T.M. 1989. Comments on some major Tectonic lineaments in India, their Geophysical expression and Metallogenic significance. Mem. Geol. Soc. Ind. **12**, 43-48.

Mahadevan, T.M. (1995). Deep continental structure of North- Western and Central Indian peninsular shield - A Review. Mem.Geol.Soc.Ind. **31** : 1-35.

Marston (1978) The geochemistry of the Archaean clastic metasediments in relation to crustal evolution, Northeastern Yilgarn Block, western Australia. Precamb. Res. v.6, pp. 157-175.

Mc Lennan, S.M. and Taylor, S.R. (1984) Archaean sedimentary rocks and their relations to the composition of the Archaean continental crust. In : Kroner, A., Hanson, G.N. and Goodwin, A.M. (Eds.), Archaean Geochemistry. The origin and evolution of the Arrchaean continental crust. Springer- verlag. Berlin-Heidelberg, pp. 47-72.

Mc Lennan, S.M., Taylor, S.R. and Mc Gregor, V.R.(1984) Geochemistry of Archaean metasedimentary rocks from West Greenland. Geochim. Cosmochim. Acta. v.48, pp. 1-13.

Mehnert, K.R. (1968) Migmatites and the origin of granitic rocks. Elsevier Pub. pp. 393.

Mercier J.-C.C. (1980) "Single-pyroxene thermobarometry". Tectonophysics, 70 : 1-37._

Mercier J.-C.C. (1984) "Equilibrium state of diopside-bearing harzburgites from ophiolites: Geobarometric and geodynamic implications." Contrib. Mineral. Petrol., 85(4): 391-403.

Merh, S.S. (1950) On the geology of the area aroundAmbamata, Danta state, N. Gujarat. Quar. J.Min. Met. Soc. Ind. XXII.

Merh, S.S. (1967) Mafic and ultramafic rocks of the Precambrian terrains of Gujarat. Proc. Sym. Upper Mantle Project, Hyderabad. 402-409.

Meschede, M. (1986) A method of discriminating between different types of mid-oceanic ridge basalts and continental tholeiites with Nb-Zr-Y diagram. Chem. Geol. 56: 207-218.

Mitra, A.K. (1971) The petrology of metamorphic rocks around Sheopura-Mewaria-Govindgarh, SW Ajmer Dist., Rajasthan. Proc. 58th Session Ind. Sci. Cong. 3.

Miyashiro, A. (1973) Metamorphism and metamorphic belts. George Allen and Unwin, London, pp. 492.

Miyashiro, A. (1994) Metamorphic petrology. UCL Press Ltd., London. pp. 404.

Moecher, D.P. and Essene, E.J. (1990a) Scapolite phase equilibria : additional constraints on the role of CO2 in granulite genesis. In : Granulites and crustal evolution (eds. Vielzeuf, D. and Vidal, Ph.), Kluwer, Dordrecht, Boston, 385-396.

Moecher, D.P. and Essene, E.J., 1991. Calculation of CO2 activities using scapolite equilibria : constraints on the presence and composition of a fluid phase during high grade metamorphism. Contrb. Mineral. Petrol., 89, 215-225.

Mohanty, M. and Guha, D.B.(1995). Lithotectonic stratigraphy of dismembered Greenstone sequence of the Mangalwar Complex around Lawa- Sardargarh and Parasali areas, Rajsamand District, Rajasthan. Mem.Geol.Soc.Ind. **31**:141-162.

Newton, R.C. (1987) Petrologic aspects of Precambrian granulite facies terrain : bearing on their origins. In Lithospheric evolution. A. Kroner (Ed). Geodynamics Series V.17, AGU- GSA.11-26.

Newton, R.C. (1993) Charnockite alteration evidence for CO2 infiltration in granulite facies metamorphism. J. Metamorphic Geol. 10 : 383-400.

Newton, R.C., Charlu, T.V and Kleppa, O.J. (1974) A calorimetric investigation of the stability of anhydrous magnesium cordierite with application to granulite facies metamorphism. Contrib. Mineral. Petrol., 44: 295-311.

Newton, R.C., Smith, J.V. & Windley, B.F. (1980) Carbonic metamorphism, granulites and crustal growth. Nature, 288 : 45-50.Nichols, G.T., Bery, R.F. and Green, D.H. (1992) Internally consistent gahnitic spinel-cordierite-garnet equilibria in the FMAsHZn systems : geothermobarometry and applications. Contrb. Mineral. Petrol., v.111, pp. 362-377.

Nixon, P.H., Reedman, A.L. and Burns, L.K. (1973) Sapphirine bearing granulites from Labwor, Uganda. Min. Mag., 39, 304, 420-428.

Niyogi, D. (1952) Preliminary structural and tectonic studies of Kishangarh alkaline rocks, Sci. and Cul. 17, 433.

Niyogi, D. (1960) Structural pattern of the Kishangarh rocks. Quar. J. Min. Met. Soc. Ind. 32.

Noguiera, J.R., Choudhuri, A. & Truow, R. (1996) Fluid inclusion studies, P-T-t paths and geotectonic evolution of the granulite facies terrain of Juij De Fora, Southern Brazil. Abs. International Seminar on Charnockites and Granulites, Madras (29-31 August, 1996).

Oldham, R.D. (1893) Manual of Geology of India. Govt. of India Press, Nasik.

Oxburgh, E.R., 1990. Some thermal aspects of granulitic history. In : Granulites and crustal evolution (eds. Vielzeuf, D. and Vidal, Ph.) Kluwer Academic Pub., Netherland. 569-580.

Parras, K. (1958) On the charnockites in the light of a highly metamorphic complex in southwetern Finland. Comm. Geol. Finlnd Bull. 181, pp.

Patel, M.P., Desai, S.J. and Merh, S.S. (1985) Structural and metamorphic studiesof the precambrian rocks of Balaram- Swarupganj area of North Gujarat and SW Rajasthan. Proc. of Fifth session Ind. Geol. Cong. 65-75.

Patel, P.P. (1971) A detailed study of the rocks of Kapasia- Bhatana area of Sirohi-Palanpur (N. Gujarat, Rajasthan) with special reference to the structural and metamorphic history and igneous activities. Unpublished Ph.D. Thesis, M.S.Univ. of Baroda, Baroda.

Pattison, D.R.M. and Newton, R.C. (1989) Reversed experimental calibration of garnet-clinopyroxene Fe-Mg exchange thermometer. Contrb. Mineral. Petrol. v.101, pp. 87-103.

Pearce, J.A. (1983) Role of subcontinental lithosphere in magma genesis at active continental margins. In : Hawkesworth, C.J. and Norry, M.J. (eds.), Continental basalts and mantle xenoliths. Shiva, Nantwich, pp.230-249.

Pearce, J.A. & Norry, M.J. (1979) Petrogenetic implications of Ti, Zr, Y and Nb variations in volcanic rocks. Contrib. Mineral. Petrol., 69 : 33-47.

Perchuk, L.L. (1989) Intercorrelation of Fe-Mg geothermometers using the Nernst Law. Geokhimiya, N.5, pp. 611-622.

Perchuk, L.L. and Lavrent'va, I.V. (1989) Some equilibria involving garnet, orthopyroxene and amphibole as prospective geothermometwers and geobarometers for metamorphic rocks. Experiment-89. (informational volume). Nauka Press, Moscow.

Peterson, J.W. and Newton, R.C., 1989. Reversed experiments on biotite-quartz-feldspar melting in the system KMASH : implications for crustal anatexis. J. Geol., 97, 465-485.

Pigage, L.C. and Greenwood, H.J. (1982) Internally consistent estimates of pressure and temperature : the staurolite problem : Amer. J. Sci. v.282, pp. 943-969.

Powar, K.B. and Patwardhan, A.M. (1984) Tectonic evolution and basemetal mineralisation in the Aravalli-Delhi belt, India. Precamb. Res. 25 : 309-323.

Powell, M. and Powell, R. (1977) Plagioclase-alkali feldspar geothermometry revisited, Mineral. Mag., v.41, pp. 253-256.

Powell, R. (1983) Preocesses in grnulite facies metamorphism. In : Migmatites, melting and metamorphism. Atherton M.P. and Gribble C.D. (Eds.) Shiva Geol. Series, pp. 127-139.

Pride, C. and Muecke, G.K. (1980) Rare earth element geochemistry of the Scourian Complex, N.W> Scotland evidence for the granite - granulite link. Contrb. Mineral. Petrol., v.73, pp. 403-412.

Prider, R.T. (1945) Charnockite and related cordierite bearing rocks from Dangin, Western Australia, Geol. Mag. 82 : 145- 172.

Quensel, P. (1951) The charnockite series of Varberg District on the Southwestern Coast of Sweden. Arkiv. Min. Geol. 1, 10.

Raja Rao, C.S. et al (1971) Precambrian stratigraphy of Rajasthan - A review. Rec. Geol. Surv. Ind. v.101, Pt.2, pp. 60-79.

Ramakrishna, T.S. (1986b) Crustal structure in the Aravalli region. proc. Sem. Crustal Dynamics, IGU, Hyderabad. 134-139.

Raval (1971) Unpub. Ph.D. Thesis, M.S.Univ., Baroda

Ravindra Kumar, G.R. and Chacko, T. (1994) Geothermobarometry of mafic granulite and metapelite from the Palghat gap, south India : petrological evidences for isothermal uplift and rapid cooling. J.Metamorphic Geol., 12(4) : 479-492.

Reddy, A.G.B. and Ramakrishnan, T.S.(1988). Bouguer gravity atlas of Rajasthan-Gujarat shield. Geol. Surv. Ind. Sp. Pub. Reddy, P.R., Rajendra Prasad, B., Vijaya Rao, V.,Prakash Khare,

Kesava Rao, G., Murthy, A.S.N., Sarkar, D., Raju, S., Rao, G.S.P. and Sridher, V. (1995). Deep seismic reflection profiling along Nandsi-Kunjer Section of Nagaur-Jhalawar Transect : Preliminary Results. Mem. Geol. Soc. Ind. **31**: 353-372.

Rietmeijer, F.J. (1988) Pyroxene exsolution in granulites from Fyfe Hills, Enderby Land. Antarctica : Evidence for 1000°C metamorphic temperatures in Archaean continental crust - Discussion. Am. Min. 73:432-433.

Rollinson, H.R. and Windley, B.F. (1980a) An Archaean granulite- grade tonalite-trondhjemite-granite suite from Scouria, NW Scotland : geochemistry and origin. Contrb. Mineral. Petrol., v.72, pp. 265-281.

Roy, A.B. (1988) Stratigraphic and tectonic framework of the Aravalli Mountain Range. In : A.B.Roy (Ed.) Precambrian of the Aravalli Mountain. Mem. Geol. Soc. Ind., v.7, pp.3-31.

Roy, A B., Somani, M K and Sharma, N.K (1981) Aravalli, Pre- Aravalli reelationship : a study from the Bhinder region, southern Rajasthan. Ind. J.Earth Sci., v.8, pp. 119-130.

Roy, A.B. & Rathore, S. (1996) Tectonic setting and model of evolution of granulites in the central part of the Aravalli mountain belt, Rajasthan, India. Abs. International Seminar on Charnockites and Granulites, Madras (29-31 August, 1996).

Saha, A.K. & Ray, S.L., (1984) Early-middle Archaean crustal evolution of the Singhbhum - Orissa craaton, Ind.J.Earth Sci. Sp. Vol. CEISM.

Sandiford, M. & Powell, R. (1986) Pyroxene exsolution in granulites from Fyfe Hills, Enderby Land, Antarctica : Evidence for 1000°C metamorphic temperatures in Archaean continental crust. Am. Mineralogist, 71: 1946-1954.

Santosh, M. (1987) Cordierite gneisses of southern Kerala, India: Petrology and fluid inclusions and implications for crustal uplift history. Contrib. Mineral. Petrol., 96: 343-356.

Sarkar, G., Barman, T.R. and Corfu, F. (1989) Timing of continental arc-type magmatism in North western India : Evidence from U-Pb zircon geochronology. J. Geol., v.97, pp. 607-612.

Sarkar, S.N., Polkanov, A.A., Gerling, E.K. and Chukrov, F.V. (1964) Geochronology of the Precambrians of Peninsular India · A synopsis, Sci. and Cul. V.30, pp. 527-537.

Schenk, V. (1984) Petrology of felsic graanulites, metapelites, metabasites, ultramafics and metacarbonates from southern Calabra (Italy) . prograde metamorphism, uplift and cooling of a former lower crust. J. Petrol., 25, 255-298.

Schreyer, W., Horrocks, P.C. and Abraham, K. (1984) High- magnesium staurolite in a sapphirine-garnet rock from the Limpopo Belt, South Africa. Contrib.Mineral. Petrol. 86: 200-207.

Schmidt, M.W. (1991) Experimental calibration of Al in hornblende geobarometer at 650°C, 3.5-13.0 Kbar. Terra. (Abs.) v.3, pp. 30.

Sen, S. (1959) Mineralogenetic trends in the evolution of metamorphic rocks and origin of granulites of east Manubhùm, India. Proc. Nat. Acad. Scvvi. 25A : 118-138.

Sen, S.K. and Bhatttacharya, A. (1984) An orthopyroxene-garnet thermometer and its application to the Madras charnockites. Contrib. Minerl. Petrol. 88: 64-71.

Sengupta, P., Dasgupta, S., Bhattacharya, P.K., Fukuoka, M., Chakraborty, S. & Bhowmick, S. (1990) Petro-tectonic imprints in the sapphirine granulites from Anantagiri, Eastern Ghats Mobile Belt,, India. J. Petrol., 33: 971-996.

Sengupta, P., Dasgupta, S., Bhui, U.K., jurgen, E. & Fukuoka, M. (1996) Magmatic evolution of mafic granulites from Anakapalle, Eastern Ghats, India : implications for tectonic setting of a Precambrian high-grade terrain. J. Southeast Asian Earth Sciences, 14: 185-198.

Sharma, N.L., Nandy, N.C. (1936) A note on the petrological classification of the basic intrusives of Danta State (N.Gujarat) Proc. Ind. Acad. Sci. 3(B) : 388-376.

Sharma, R.S. (1988) Patterns of metamorphism in the Aravalli mountain belt. Mem. Geol. Soc. Ind. 7: 33-75.

Sharma, R.S. (1995). An evolutionary model for the Precambrian crust of Rajasthan : Some petrological and geochrono-logical considerations. Mem.Geol.Soc.Ind. **31**: 91-115.

Singh, S (1966) Orthopyroxene bearing rocks of charnockitic affinities in South Savanna Kanuku Complex of British Guiana. J. Petrology. 7(2): 171-192

Sinha-Roy, S., Guha, D.B. and Bhattacharya A.K. (1992) Polymetamorphic granulite facies gneisses of the Precambrian Sandmata Complex, Rajasthan, India. Indian minerals, 46(1): 1-12.

Sinha-Roy, S.,Malhotra, G. and Guha, D.B. (1995). A transect across Rajasthan Precambrian terrain in relation to geology, tectonics and crustal evolution of South-Central Rajasthan. Mem. Geol. Soc. Ind. **31**: 63-89.

Sivasbramanian, P., Natarajan, R. and Janardhan, A.S. (1991) Sapphirine bearing assemblage from Perumalmalai, Palni Hills, Tamil Nadu. J. Geol. Soc. Ind., 38: 532-537.

Sivasbramanian, P., Natarajan, R and Janardhan, A S. (1992) Sapphirine bearing assemblage from Perumalmalai, Palni Hills, Tamil Nadu. (Reply) J. Geol Soc. Ind., 40: 287-288.

Siva Kumar Babu, E.V.S. (1996) High temperature assemblages and stability of osumilite in the granulite-facies metapelites around the Chimakurti pluton, India. Abs. International Seminar on Charnockites and Granulites, Madras (29-31 August, 1996).

Slavinsky, V.V. (1980) Partitioning of the thee components among coexisting minerals and the geothermometry of garnet-two pyroxene equilibria : Izvestiya Akad. Nauk SSSR, Ser. Geol., No.1, pp. 72-85.

Slavinskiy V.V. (1983) Two-pyroxene geothermometry. Mineralogicheskii zhournal, N 6, p. 29-38, 1983.(in Russian)

Spear, F.S. (1981) An experimental study of hornblende stability and compositional variability in amphibole. Amer. J. Sci., 281, 697-734.

Spear, F.S. (1991) On the interpretation of peak metamorphic temperatures in light of garnet diffusion during cooling. J. Meta. Geol., 379-388.

Srikarni, C., Desai, S.J. and Patel, M.P. (1996). Tectono- stratigraphic significance of granulites of North Gujarat, W.India. Abs. International Seminar on Charnockites and Granulites, Madras (29-31 August, 1996).

Srinivasulu, S., Rajasekharan, K.C., Ram Mohan, V. & Balasubramanian, S. (1996) Geochemical evolution of metapelites of Kirnur, Tamil Nadu, S.India. Abs. International Seminar on Charnockites and Granulites, Madras (29-31 August, 1996).

Stromer, J.C. Jr. (1975) A practical two-feldspar geothermometer. Amer. Mineral., v.60, pp. 667-674.

Stromer, J.C. Jr. and Whitney, J.A. (1977) Two-feldspar geothremometry in granulite facies metamorphic rocks : Sappherine granulites from Brazil. Contrb. Mineral. Petrol., v.65, pp. 123-133.

Subba Rao, D.V., , Niraml Charan, S., Narayana, B.L. and Naqvi, S.M. (1995) Symplectites in high-grade pelitic gneisses of Usilampatti, Tamil Nadu : P-T conditions and geochemistry. J. Geol. Soc. Ind., 46: 37-45.

Sutton, J. and Watson, J. (1951) The Torridonian metamorphic history of the Loch Torridon and Scourie areas in the Nw Highlands and its bearing on the chronological clasification of the Lewisian. Quart. J. Geol. Soc. London 106 : 241-308.

Sutton, J., Watson, J. & James, C.T. (1954) A study of metamorphic rocks of Krema and Kungwe Bay, Western Tanganyika, Geol. Suv. Dept., Tanganyika, Bull. 22.

Sycanthovong, S.P.H. and Merh, S.S. (1984) Proto Plate Tectonics : the energetic model for the structural, metamorphic and igneous evolution of the Precambrian rocks of northwest peninsular India. Special Pu. Geol. Surv. Ind. 12 : 419-457.

Tewari, H.C., Vijaya Rao, V., Dixit, M.M., Rajendra Prasad, B., Madhava Rao, N., Venkateswarlu, N., Prakash Khare, Kesava Rao, G., Raju, S. and Kaila, K.L. (1995). Deep crustal reflection studiesacross the Delhi-Aravalli Fold Belt, Results from the North-Western part : by Controlled Source Seismology Group. Mem.Geol.Soc.Ind. **31**: 383-402.

Tewari, H.C., Rajendra Prasad, B., Vijaya Rao, V.,P.R.Reddy, Dixit, M.M. and Madhava Rao, N.(1997). Crustal reflectivity parameter fordeciphering the evolutionary processes across the proterozoic Aravalli-Delhi Fold Belt. J. Geol. Soc. Ind. 50: 779-785

Thompson, A.B. (1976) Mineral reactions in pelitic rocks : prediction of P-T-X (Fe-Mg) phase relations. Am.J.Sci. v.276, pp. 401-424.

Touret, J. (1971) Lithos 4: 423-436.

Touret, J.L.R. (1981) Fluid inclusions in high grade rocks. In: Hollister, L.S.and Crawford, M.J. (eds.) Short course in fluid inclusions: Applications to petrology: 182-208. Mineral. Ass. Canada, Short course handbook, vol. 6.

Tracy, R.J. and McLellan, E.L. (1985) A natural example of the kinetic controls of compositional and textural equilibration. In : Thompson, A.B. and David C. Rubie "Metamorphic Reactions -Kinetics, Textures, and Deformation" - Advances in Physical Geochemistry V.4 : 118-137, Springer- Verlag, N.Y.,

Turner, F.J. and Verhoogen, J. (1960) Igneous and metamorphic petrology, McGraw Hill Publ. p.694.

Turner, F.J. (1968) Metamorphic Petrology Mineralogical and Field Aspects. McGraw Hill Publ. p. 403.

Turnock, A.C. (1962) Phase diagram of the join diopside- hedenbergite. Carnegie Inst. Washington, Ann. Rep. Dir. Geophys. Lab., 1961-62, p. 81.

Vielzuef, D. and Holloway, J.R. (1988) Experimental determination of the fluid absent melting reactions in the pelitic system. Consequences for crustal differentiation. Contrib. Mineral. Petrol., 98: 257-276.

Vielzeuff, D. and Vidal, Ph. (1990) Granulites and crustal evolution. NATO ASI Series C: Mathematical and Physical Sciences, Vol.311. Kluwer Acad. Publ., Dordrecht, Holland.

Wang, K.Y., Yan, Y.H., Yang, R.Y. and Chen, Y.F., 1985. REE geochemistry of Early Precambrian charnockites and tonalitic-granonodioritic gneisses of Qianan Region, eastern Hebei, North China, Precambrian Res., 27: 63-84.

Watson, J. (1949) The Geology of Lewisian of the Scourie area, Unpublished Ph D Thesis, University of London

Watson, J. (1973) The effects of reworking on high grade gneiss complexes. Phil. Trans. R. Soc. London, A.273: 443-455.

Weaver, B.L. (1980) Rare Earth Element geochemistry of Madras granulites. Contrb. Mineral Petrol., 71: 271-297.

Weaver, B.L. and Tarney, J. (1980) Rare earth geochemistry of Lewisian granulite facies gneisses, northwest Scotland : implications for the petrogenesis of the Archaean lower contineal crust. Earth Planet. Sci. Lett., v.51, pp. 279-296

Wells, P.R.A. (1979) Chemical and thermal evolution nof Archaean sialic crust, southwest Greenland. J. Petrology., v.20, pp. 187-226.

White, A.J.R. (1971) Granulites - results of discussion II. N.Jb.Miner.Mh., 4, p.139.

Whitney, J.A. and Stromer, J.C. (1977) Two-feldspar geothermometry in mesozonal granitic intrusions : Three examples from the Piedmont of Georgia. Contrb Mineral. Petrol., 63, 51-69.

Wilson, A.F. (1952) The charnoickite problem in Australia, Sir Douglas Mawson Anniv. University of Adelaide 205-224.

Wilson, A.F. (1978) Comparison of some of the geochemical features and tectonic settings of Archaean and Proterozoic granulites, with reference to Australia. In : B.F.Windley and S.M.Naqvi (Eds.) Archaean Geochemistry, Elsevier, Amsterdam, pp. 241-267

Windley, B.F. (1972) Regional geology of early pre-Cambrian high-grademetamorphic rocks in West Greenland. The Geol. Surv. Greenland, 46 : 5-46.

Windley, B.F. (1984) The evolving continents. Wiley & Sons (Publ.) 2nd Edition. 399pp.

Winkler, H.G.F. (1979) Petrogenesis of metamorphic rocks. Springer-Verlag, New YorkInc, p.344.

Wood, D.A. (1980) The application of Th-Hf-Ta diagram to problems of tectonomagmatic classification and to establishing the nature of crustal contamination of basaltic lavas of British tertiary volcanic province Erath Plan Sci. lett 50. 11-30.

Wood, B.J and Banno, S. (1973) Garnet-orhtopyroxene and orthopyroxene-clinopyroxene relationships in simple and complex systems. Contrib. Mineral. Petrol. 42: 109-124

Wynne-Edwards, H.R. (1967) West Port map ara, Ontario with special emphasis on the Precambrian rocks, Geol Surv Can Mem 346 · 1-142.

Young, E.D. (1989) Petrology of biotite-cordierite-garnet gneiss of the McCullough Range, Nevada II. P-T-aH2O path and growth of cordierite during late stages of low-P granulite-grade metamorphism. J.Petrol. 30-1: 61-78.

Young, E.D., Anderson, J.L., Clarke, S.H. and Thomas, W.M. (1989) Petrology of biotite-cordierite-garnet gneiss of McCullough Range, Nevada. 1. evidence for Proterozoic low pressure fluid absent granulite grade metamorphism in the southern Cordillera. J.Petrol.30: 39-60.

* References quoted in computer programme by Fonarev, V.I., Graphicikov, A.A. and Konilov, A.N. (1991). A Consistent system of geothermometers for metamorphic complexes.