

## List of Tables

Table 1.1	The theoretical yield from each step of denitrification.....	4
Table 1.2	Distribution of denitrifiers among bacteria and Archaea.....	8
Table 1.3	Functional importance of EPS in biofilms.....	13
Table 2.1	Characteristics of the activated sludge samples.....	35
Table 2.2	PCR primers used in the study.....	38
Table 2.3	Percentage and ratio of the denitrification genes in the activated sludge samples.....	42
Table 2.4	Nitrate reduction test results of the isolates.....	44
Table 2.5	Determination of the denitrifying bacterial isolates by partial 16S rRNA gene sequence similarity.....	47
Table 2.6	Calculated diversity estimates of <i>nosZ</i> gene in the sludge samples...	51
Table 2.7	Calculated rates during denitrification deduced from Fig. 2.9.....	53
Table 2.8	Nitrate and Nitrite reduction rates by resting-state cells.....	53
Table 2.9	Nitrate reduction rates during denitrification at increasing nitrate concentrations.....	54
Table 2.10	Denitrification and growth of the isolates at high nitrate concentrations.....	57
Table 3.1	Plackett-Burman design matrix.....	80
Table 3.2	Variables with media components and their concentrations.....	81
Table 3.3	Structural parameters of biofilm formed in 12 different media of the Plackett-Burman experiment.....	92
Table 3.4	Statistical analysis for influence of nutritional components on biofilm structural parameters from the P-B design results.....	93
Table 3.5	The cellular fatty acids analysed by FAME.....	104
Table 4.1	Percentage of <i>nosZ</i> gene as analysed from Fig. 4.4 data.....	127

Table 4.2	Diversity indices calculated from the DGGE band patterns.....	132
Table 4.3	16S rRNA gene similarity to the closely related organisms from the eluted bands of DGGE.....	133