

Bibliography

- Abbamondi, G.R., Suner, S., Cutignano, A., Grauso, L., Nicolaus, B., Toksoy Oner, E., Tommonaro, G., 2016. Identification of N-Hexadecanoyl-L-homoserine lactone (C16-AHL) as signal molecule in halophilic bacterium *Halomonas smyrnensis* AAD6. *Ann. Microbiol.* <https://doi.org/10.1007/s13213-016-1206-6>
- Acosta-Jurado, S., Alías-Villegas, C., Almozara, A., Espuny, M.R., Vinardell, J.M., Pérez-Montaño, F., 2020. Deciphering the symbiotic significance of quorum sensing systems of *Sinorhizobium fredii* HH103. *Microorganisms* 8. <https://doi.org/10.3390/microorganisms8010068>
- Alami, Y., Achouak, W., Marol, C., Heulin, T., 2000. Rhizosphere soil aggregation and plant growth promotion of sunflowers by an exopolysaccharide-producing *Rhizobium* sp. strain isolated from sunflower roots. *Appl. Environ. Microbiol.* <https://doi.org/10.1128/AEM.66.8.3393-3398.2000>
- Altschul, S.F., Gish, W., Miller, W., Myers, E.W., Lipman, D.J., 1990. Basic local alignment search tool. *J. Mol. Biol.* [https://doi.org/10.1016/S0022-2836\(05\)80360-2](https://doi.org/10.1016/S0022-2836(05)80360-2)
- Amaya-Gómez, C. V, Hirsch, A.M., Soto, M.J., 2015. Biofilm formation assessment in *Sinorhizobium meliloti* reveals interlinked control with surface motility. *BMC Microbiol.* 15, 58. <https://doi.org/10.1186/s12866-015-0390-z>
- Aneja, P., Zachertowska, A., & Charles, T. C. (2005). Comparison of the symbiotic and competition phenotypes of *Sinorhizobium meliloti* PHB synthesis and degradation pathway mutants. *Canadian Journal of Microbiology*.
- Archana, G. (2010). Engineering nodulation competitiveness of rhizobial bioinoculants in soils. In *Microbes for legume improvement* (p. 157).
- Arnold, M.F.F., Penterman, J., Shabab, M., Chen, E.J., Walker, G.C., 2018. Important late stage symbiotic role of the *Sinorhizobium meliloti* exopolysaccharide succinoglycan. *J. Bacteriol.* JB.00665-17. <https://doi.org/10.1128/JB.00665-17>
- Atkinson, S., Chang, C.-Y., Sockett, R.E., Camara, M., Williams, P., 2006. Quorum sensing in *Yersinia enterocolitica* controls swimming and swarming motility. *J. Bacteriol.* 188, 1451–1461. <https://doi.org/10.1128/JB.188.4.1451-1461.2006>
- Aufrecht, J. A., Timm, C. M., Bible, A., Morrell-Falvey, J. L., Pelletier, D. A., Doktycz, M. J., & Retterer, S. T. (2018). Quantifying the Spatiotemporal Dynamics of Plant Root Colonization by Beneficial Bacteria in a Microfluidic Habitat. *Advanced Biosystems*, 1800048, 1800048.
- Bahlawane, C., McIntosh, M., Krol, E., Becker, A., 2008. *Sinorhizobium meliloti* Regulator MucR Couples Exopolysaccharide Synthesis and Motility. *Mol. Plant-Microbe Interact.* 21, 1498–1509. <https://doi.org/10.1094/MPMI-21-11-1498>
- Barnett, M.J., Long, S.R., 2017. Novel genes and regulators that influence production of cell surface exopolysaccharides in *Sinorhizobium meliloti*. *J. Bacteriol.* JB.00501-17. <https://doi.org/10.1128/JB.00501-17>

- Baumgardt, K., Šmidová, K., Rahn, H., Lochnit, G., Robledo, M., Evgenieva-Hackenberg, E., 2015. The stress-related, rhizobial small RNA RcsR1 destabilizes the autoinducer synthase encoding mRNA *sini* in *Sinorhizobium meliloti*. *RNA Biol.* <https://doi.org/10.1080/15476286.2015.1110673>
- Becana, M., Matamoros, M. A., Udvardi, M., & Dalton, D. A. (2010). Recent insights into antioxidant defenses of legume root nodules. *New Phytologist*.
- Benizri, E., Baudoin, E., Guckert, A., 2001. Root colonization by inoculated plant growth-promoting rhizobacteria. *Biocontrol Sci. Technol.* 11, 557–574. <https://doi.org/10.1080/09583150120076120>
- Bernabéu-Roda, L., Calatrava-Morales, N., Cuéllar, V., & Soto, M. J. (2015). Characterization of surface motility in *Sinorhizobium meliloti*: regulation and role in symbiosis. *Symbiosis*, 67(1–3), 79–90.
- Bogino, P., Nievas, F., Banchio, E., & Giordano, W. (2011). Increased competitiveness and efficiency of biological nitrogen fixation in peanut via in-furrow inoculation of rhizobia. *European Journal of Soil Biology*, 47(3), 188–193.
- Bogino, P.C., Nievas, F.L., Giordano, W., 2015. A review: Quorum sensing in *Bradyrhizobium*. *Appl. Soil Ecol.* 94, 49–58. <https://doi.org/10.1016/J.APSOIL.2015.04.016>
- Boyer, M., Wisniewski-Dyé, F., 2009. Cell-cell signalling in bacteria: Not simply a matter of quorum. *FEMS Microbiol. Ecol.* <https://doi.org/10.1111/j.1574-6941.2009.00745.x>
- Brockwell, J., Roughley, R.J., Herridge, D.F., 1987. Population dynamics of *Rhizobium japonicum* strains used to inoculate three successive crops of soybean. *Aust. J. Agric. Res.* <https://doi.org/10.1071/AR9870061>
- Caetano-Anollés, G., Wrobel-Boerner, E., Bauer, W.D., 1992. Growth and movement of spot inoculated *Rhizobium meliloti* on the root surface of alfalfa. *Plant Physiol.* 98, 1181–1189. <https://doi.org/10.1104/pp.98.3.1181>
- Calatrava-Morales, N., McIntosh, M., Soto, M., 2018. Regulation mediated by N-acyl homoserine lactone quorum sensing signals in the *Rhizobium*-legume symbiosis. *Genes*. 9, 263. <https://doi.org/10.3390/genes9050263>
- Calatrava-Morales, N., Nogales, J., Amezttoy, K., van Steenberg, B., & Soto, M. J. (2017b). The NtrY/NtrX System of *Sinorhizobium meliloti* GR4 Regulates Motility, EPS I Production, and Nitrogen Metabolism but Is Dispensable for Symbiotic Nitrogen Fixation. *Molecular Plant-Microbe Interactions*, 30(7), MPMI-01-17-0021.
- Calvert, H. E., Pence, M. K., Pierce, M., Malik, N. S. A., & Bauer, W. D. (1984). Anatomical analysis of the development and distribution of *Rhizobium* infections in soybean roots. *Canadian Journal of Botany*, 62(11), 2375–2384.
- Cataldi, T.R.I., Bianco, G., Abate, S., 2007. Profiling of N-acyl-homoserine lactones by liquid chromatography coupled with electrospray ionization and a hybrid quadrupole linear ion-trap and Fourier-transform ion-cyclotron-resonance mass spectrometry (LC-ESI-LTQ-FTICR-MS). *J. Mass Spectrom.* 43, 82–96. <https://doi.org/10.1002/jms.1275>

- Cesco, S., Neumann, G., Tomasi, N., Pinton, R., & Weiskopf, L. (2010). Release of plant-borne flavonoids into the rhizosphere and their role in plant nutrition. *Plant and Soil*.
- Cha, C., Gao, P., Chen, Y.-C., Shaw, P.D., Farrand, S.K., 1998. Production of acyl-homoserine lactone quorum-sensing signals by Gram-negative plant-associated bacteria. *Mol. Plant-Microbe Interact.* 11, 1119–1129. <https://doi.org/10.1094/MPMI.1998.11.11.1119>
- Chanda Venkata, S. K., Nadigatla Veera Prabha Rama, G. R., Saxena, R. K., Saxena, K., Upadhyaya, H. D., Siambi, M., ... Varshney, R. K. (2018). Pigeonpea improvement: An amalgam of breeding and genomic research. *Plant Breeding*, (November 2017), 1–10. <https://doi.org/10.1111/pbr.12656>
- Charoenpanich, P., Meyer, S., Becker, A., McIntosh, M., 2013. Temporal expression program of quorum sensing-based transcription regulation in *Sinorhizobium meliloti*. *J. Bacteriol.* 195, 3224–3236. <https://doi.org/10.1128/JB.00234-13>
- Chen, H., Teplitski, M., Robinson, J.B., Rolfe, B.G., Bauer, W.D., 2003. Proteomic analysis of wild-type *Sinorhizobium meliloti* responses to N-acyl homoserine lactone quorum-sensing signals and the transition to stationary phase. *J. Bacteriol.* 185, 5029–5036. <https://doi.org/10.1128/jb.185.17.5029-5036.2003>
- Chi, F., Yang, P., Han, F., Jing, Y., Shen, S., 2010. Proteomic analysis of rice seedlings infected by *Sinorhizobium meliloti* 1021. *Proteomics*. <https://doi.org/10.1002/pmic.200900694>
- Chilton, M.D., Currier, T.C., Farrand, S.K., Bendich, a J., Gordon, M.P., Nester, E.W., 1974. *Agrobacterium tumefaciens* DNA and PS8 bacteriophage DNA not detected in crown gall tumors. *Proc. Natl. Acad. Sci. U. S. A.* 71, 3672–3676.
- Choo, J.H., Rukayadi, Y., Hwang, J.-K., 2006. Inhibition of bacterial quorum sensing by vanilla extract. *Lett. Appl. Microbiol.* 42, 637–641. <https://doi.org/10.1111/j.1472-765X.2006.01928.x>
- Chun, J., Lee, J.-H., Jung, Y., Kim, M., Kim, S., Kim, B.K., Lim, Y.-W., 2007. EzTaxon: a web-based tool for the identification of prokaryotes based on 16S ribosomal RNA gene sequences. *Int. J. Syst. Evol. Microbiol.* 57, 2259–2261.
- Cooper, J.E., 2007. Early interactions between legumes and rhizobia : disclosing complexity in a molecular dialogue. *J. Appl. Microbiol.* 103, 1355–1365. <https://doi.org/10.1111/j.1365-2672.2007.03366.x>
- Cummings, S. P. (2005). The role and future potential of nitrogen fixing bacteria to boost productivity in organic and low-input sustainable farming systems. *Biotechnology*, (1), 1–10.
- Deaker, R., Roughley, R.J., Kennedy, I.R., 2004. Legume seed inoculation technology - A review. *Soil Biol. Biochem.* 36, 1275–1288. <https://doi.org/10.1016/j.soilbio.2004.04.009>
- Doyle, J. J. (2011). Phylogenetic Perspectives on the Origins of Nodulation. *Molecular Plant-Microbe Interactions*, 24(11), 1289–1295.
- Dworkin, M., Foster, J.W., 1958. Experiments with some microorganisms which utilize ethane and hydrogen. *J. Bacteriol.* 75, 592–603.

- El Msehli, S., Lambert, A., Hopkins, J., Boncompagni, E., Smiti-Aschi, S., Hérouart, D., & Frendo, P. (2019). Physiological and genetic changes during natural senescence of *Medicago truncatula* root nodules. *Journal of Plant Nutrition and Soil Science*, 182(3), 385–392.
- Ferguson, B. J., Indrasumunar, A., Hayashi, S., Lin, M., Lin, Y., Reid, D. E., & Gresshoff, P. M. (2010). Molecular Analysis of Legume Nodule Development and Autoregulation. *Journal of Integrative Plant Biology*, 52(1), 61–76.
- Ferguson, B. J., Mens, C., Hastwell, A. H., Zhang, M., Su, H., Jones, C. H., ... Gresshoff, P. M. (2019). Legume nodulation: The host controls the party. *Plant, Cell & Environment*, 42(1), 41–51.
- Ferraioli, S., Patriarca, E. J., & Tate, R. (2004). Organogenesis of Legume Root Nodules. *Review Literature And Arts Of The Americas*, 234, 201–262.
- Finnie, C., Hartley, N. M., Findlay, K. C., & Downie, J. A. (1997). The *Rhizobium leguminosarum* *prsDE* genes are required for secretion of several proteins, some of which influence nodulation, symbiotic nitrogen fixation and exopolysaccharide modification . *Molecular Microbiology*.
- Fossou, R.K., Ziegler, D., Zézé, A., Barja, F., Perret, X., 2016. Two major clades of *Bradyrhizobia* dominate symbiotic interactions with pigeonpea in fields of Côte d'Ivoire. *Front. Microbiol.* 7, 1–11. <https://doi.org/10.3389/fmicb.2016.01793>
- Foucher, F., & Kondorosi, E. (2000). Cell cycle regulation in the course of nodule organogenesis in *Medicago*. *Plant Molecular Biology*, 43(5–6), 773–786.
- Frayse, N., Couderc, F., & Poinso, V. (2003). Surface polysaccharide involvement in establishing the rhizobium-legume symbiosis. *European Journal of Biochemistry*.
- Frederix, M., Edwards, A., Mcanulla, C., Downie, J.A., 2011. Co-ordination of quorum-sensing regulation in *Rhizobium leguminosarum* by induction of an anti-repressor. *Mol. Microbiol.* 81, 994–1007. <https://doi.org/10.1111/j.1365-2958.2011.07738.x>
- Frederix, M., Edwards, A., Swiderska, A., Stanger, A., Karunakaran, R., Williams, A., Abbruscato, P., Sanchez-Contreras, M., Poole, P.S., Downie, J.A., 2014. Mutation of *praR* in *R. hizobium leguminosarum* enhances root biofilms, improving nodulation competitiveness by increased expression of attachment proteins. *Mol. Microbiol.* 93, 464–478. <https://doi.org/10.1111/mmi.12670>
- Fujishige, N.A., Kapadia, N.N., De Hoff, P.L., Hirsch, A.M., 2006. Investigations of *Rhizobium* biofilm formation. *FEMS Microbiol. Ecol.* 56, 195–206. <https://doi.org/10.1111/j.1574-6941.2005.00044.x>
- Gage, D.J., Bobo, T., Long, S.R., 1996. Use of Green Fluorescent Protein To Visualize the Early Events of Symbiosis between *Rhizobium meliloti* and Alfalfa (*Medicago sativa*) 178, 7159–7166.
- Gao, M., Barnett, M.J., Long, S.R., Teplitski, M., 2010. Role of the *Sinorhizobium meliloti* global regulator *hfq* in gene regulation and symbiosis. *Mol. Plant-Microbe Interact.* 23, 355–365. <https://doi.org/10.1094/MPMI-23-4-0355>

- Gao, M., Chen, H., Eberhard, A., Gronquist, M.R., Robinson, J.B., Rolfe, B.G., Bauer, W.D., 2005. *sinI*- and *expR*-dependent quorum sensing in *Sinorhizobium meliloti*. *J. Bacteriol.* 187, 7931–7944. <https://doi.org/10.1128/JB.187.23.7931-7944.2005>
- García-Fraile, P., Carro, L., Robledo, M., Ramírez-Bahena, M.-H., Flores-Félix, J.-D., Fernández, M. T., ... Velázquez, E. (2012). Rhizobium promotes non-legumes growth and quality in several production steps: towards a biofertilization of edible raw vegetables healthy for humans. *PloS One*, 7(5), e38122.
- Garg, B., Dogra, R. C., & Sharma, P. K. (1999). High-efficiency transformation of *Rhizobium leguminosarum* by electroporation. *Applied and Environmental Microbiology*, 65(6), 2802–2804.
- Gates, M. F. (2014). Putting women and girls at the center of development. *Science* (New York, N.Y.), 345(6202), 1273–5. <https://doi.org/10.1126/science.1258882>
- Gaunt, M.W., Turner, S.L., Rigottier-Gois, L., Lloyd-Macgilp, S. a, Young, J.P., 2001. Phylogenies of *atpD* and *recA* support the small subunit rRNA-based classification of rhizobia. *Int. J. Syst. Evol. Microbiol.* 51, 2037–2048.
- Gibson, K.E., Kobayashi, H., Walker, G.C., 2008. Molecular determinants of a symbiotic chronic infection. *Annu. Rev. Genet.* 42, 413–41. <https://doi.org/10.1146/annurev.genet.42.110807.091427>
- González, J.E., Marketon, M.M., 2003. Quorum sensing in nitrogen-fixing rhizobia. *Microbiol. Mol. Biol. Rev.* 67, 574–592. <https://doi.org/10.1128/MMBR.67.4.574>
- Gosai, J., Anandhan, S., Bhattacharjee, A., & Archana, G. (2020). Elucidation of quorum sensing components and their role in regulation of symbiotically important traits in *Ensifer nodulating pigeon pea*. *Microbiological Research*, 231, 126354.
- Gotz, R., Limmer, N., Ober, K., Schmitt, R., 1982. Motility and Chemotaxis in Two Strains of *Rhizobium* with Complex Flagella. *J. Gen. Microbiol.* 128, 789–798. <https://doi.org/10.1099/00221287-128-4-789>
- Gould, T.A., Herman, J., Krank, J., Murphy, R.C., Churchill, M.E.A., 2006. Specificity of acyl-homoserine lactone synthases examined by mass spectrometry. *J. Bacteriol.* 188, 773–783. <https://doi.org/10.1128/JB.188.2.773-783.2006>
- Graham, P. H., & Vance, C. P. (2003). Legumes : Importance and Constraints to Greater Use. *Plant Physiology*, 131(March), 872–877.
- Gurich, N., & González, J. E. (2009). Role of quorum sensing in *Sinorhizobium meliloti*-Alfalfa symbiosis. *Journal of Bacteriology*, 191(13), 4372–82 . <https://doi.org/10.1128/JB.00376-09>
- Heidstra, R., Nilsen, G., Martinez-Abarca, F., Van Kammen, A., & Bisseling, T. (1997). Nod factor-induced expression of leghemoglobin to study the mechanism of NH₄NO₃ inhibition on root hair deformation. *Molecular Plant-Microbe Interactions*, 10(2), 215–220.

- Hense, B. a, Kuttler, C., Müller, J., Rothballer, M., Hartmann, A., & Kreft, J.-U. (2007). Does efficiency sensing unify diffusion and quorum sensing? *Nature Reviews. Microbiology*, 5(3), 230–239.
- Herridge, D. F., Peoples, M. B., & Boddey, R. M. (2008). Global inputs of biological nitrogen fixation in agricultural systems. *Plant and Soil*, 311(1–2), 1–18. <https://doi.org/10.1007/s11104-008-9668-9673>
- Hirsch, A. M., Lum, M. R., & Downie, J. A. (2001). What makes the rhizobia-legume symbiosis so special? *Plant Physiology*, 127(4), 1484–92.
- Hoang, H.H., Becker, A., González, J.E., 2004. The LuxR Homolog ExpR, in combination with the sin quorum sensing system, plays a central role in *Sinorhizobium meliloti* gene expression. *J. Bacteriol.* 186, 5460–5472. <https://doi.org/10.1128/JB.186.16.5460-5472.2004>
- Hoang, H.H., Gurich, N., González, J.E., 2008. Regulation of motility by the ExpR/Sin quorum-sensing system in *Sinorhizobium meliloti*. *J. Bacteriol.* 190, 861–871. <https://doi.org/10.1128/JB.01310-07>
- Huang, Y., Zeng, Y., Yu, Z., 2013. A novel degenerated primer pair detects diverse genes of acyl homoserine lactone synthetase in Rhizobiaceae family. *Curr. Microbiol.* 67, 183–187. <https://doi.org/10.1007/s00284-013-0351-4>
- Iyer, B., Rajput, M. S., Jog, R., Joshi, E., Bharwad, K., & Rajkumar, S. (2016). Organic acid mediated repression of sugar utilization in rhizobia. *Microbiological Research*, 192, 211–220. <https://doi.org/10.1016/J.MICRES.2016.07.006>
- Janczarek, M. (2011). Environmental Signals and Regulatory Pathways That Influence Exopolysaccharide Production in Rhizobia, 7898–7933.
- Janczarek, M., Skorupska, A., 2011. Modulation of *rosR* expression and exopolysaccharide production in *Rhizobium leguminosarum* bv. *trifolii* by phosphate and clover root exudates. *Int. J. Mol. Sci.* <https://doi.org/10.3390/ijms12064132>
- Jensen, J.B., Peters, N.K., Bhuvaneshwari, T. V, 2002. Redundancy in periplasmic binding protein-dependent transport systems for trehalose, sucrose, and maltose in *Sinorhizobium meliloti*. *J. Bacteriol.* 184, 2978–2986. <https://doi.org/10.1128/JB.184.11.2978-2986.2002>
- Jiménez-Zurdo, J. I., Frugier, F., Crespi, M. D., & Kondorosi, A. (2000). Expression profiles of 22 novel molecular markers for organogenetic pathways acting in alfalfa nodule development. *Molecular Plant-Microbe Interactions*, 13(1), 96–106.
- Jimenez-Zurdo, J. I., Van Dillewijn, P., Soto, M. J., De Felipe, M. R., Olivares, J., & Toro, N. (1995). Characterization of a *Rhizobium meliloti* proline dehydrogenase mutant altered in nodulation efficiency and competitiveness on alfalfa roots. *Molecular Plant-Microbe Interactions*.
- Jitackson, S., Sadowsky, M.J., 2008. Nodulation gene regulation and quorum sensing control density-dependent suppression and restriction of nodulation in the *Bradyrhizobium japonicum*-soybean symbiosis. *Appl. Environ. Microbiol.* 74, 3749–56. <https://doi.org/10.1128/AEM.02939-07>

- Johnston, A.W.B., Behringer, J.E., 1975. Identification of the rhizobium strains in pea root nodules using genetic markers. *J. Gen. Microbiol.* <https://doi.org/10.1099/00221287-87-2-343>
- Jones, K. M., Kobayashi, H., Davies, B. W., Taga, M. E., & Walker, G. C. (2007). How rhizobial symbionts invade plants: The Sinorhizobium - Medicago model. *Nature Reviews Microbiology*, 5(8), 619–633.
- Knox, O. G. G., Killham, K., Artz, R. R. E., Mullins, C., & Wilson, M. (2004). Effect of nematodes on rhizosphere colonization by seed-applied bacteria. *Applied and Environmental Microbiology*, 70(8), 4666–4671.
- Kovach, M.E., Elzer, P.H., Steven Hill, D., Robertson, G.T., Farris, M.A., Roop, R.M., Peterson, K.M., 1995. Four new derivatives of the broad-host-range cloning vector pBBR1MCS, carrying different antibiotic-resistance cassettes. *Gene* 166, 175–176. [https://doi.org/10.1016/0378-1119\(95\)00584-1](https://doi.org/10.1016/0378-1119(95)00584-1)
- Krishnan, H. B. (2002). NoIX of Sinorhizobium fredii USDA257, a type III-secreted protein involved in host range determination, is localized in the infection threads of cowpea (*Vigna unguiculata* [L.] Walp) and soybean (*Glycine max* [L.] Merr.) nodules. *Journal of Bacteriology*, 184(3), 831–839.
- Krol, E., & Becker, A. (2014). Rhizobial homologs of the fatty acid transporter FadL facilitate perception of long-chain acyl-homoserine lactone signals. *Proceedings of the National Academy of Sciences of the United States of America*.
- Krysciak, D., Grote, J., Orbegoso, M. R., Utpatel, C., Förstner, K. U., Li, L., ... Streit, W. R. (2014). RNA sequencing analysis of the broad-host-range strain sinorhizobium fredii NGR234 identifies a large set of genes linked to quorum sensing-dependent regulation in the background of a traI and ngrI deletion mutant. *Applied and Environmental Microbiology*, 80(18), 5655–5671.
- Krysciak, D., Grote, J., Rodriguez Orbegoso, M., Utpatel, C., Förstner, K.U., Li, L., Schmeisser, C., Krishnan, H.B., Streit, W.R., 2014. RNA sequencing analysis of the broad-host-range strain Sinorhizobium fredii NGR234 identifies a large set of genes linked to quorum sensing-dependent regulation in the background of a traI and ngrI deletion mutant. *Appl. Environ. Microbiol.* 80, 5655–5671. <https://doi.org/10.1128/AEM.01835-14>
- Kumar, S., Stecher, G., Li, M., Knyaz, C., Tamura, K., 2018. MEGA X: Molecular evolutionary genetics analysis across computing platforms. *Mol. Biol. Evol.* 35, 1547–1549. <https://doi.org/10.1093/molbev/msy096>
- Kuzyakov, Y., Razavi, B.S., 2019. Rhizosphere size and shape: Temporal dynamics and spatial stationarity. *Soil Biol. Biochem.* 135, 343–360. <https://doi.org/10.1016/j.soilbio.2019.05.011>
- Lardi, M., de Campos, S. B., Purtschert, G., Eberl, L., & Pessi, G. (2017). Competition experiments for legume infection identify Burkholderia phymatum as a highly competitive β -rhizobium. *Frontiers in Microbiology*.
- Lee, D.J., Jo, A.R., Jang, M.C., Nam, J., Choi, H.J., Choi, G.-W., Sung, H.Y., Bae, H., Ku, Y.-G., Chi, Y.-T., 2018. Analysis of two quorum sensing-deficient isolates of Pseudomonas aeruginosa. *Microb. Pathog.* 119, 162–169. <https://doi.org/10.1016/J.MICPATH.2018.04.014>

- Lee, Y. W., Jeong, S. Y., In, Y. H., Kim, K. Y., So, J. S., & Chang, W. S. (2010). Lack of O-polysaccharide enhances biofilm formation by *Bradyrhizobium japonicum*. *Letters in Applied Microbiology*.
- Lei, M. J., Wang, Q., Li, X., Chen, A., Luo, L., Xie, Y., ... Wang, Y. Z. (2015). The small Gtpase ROP10 of *Medicago truncatula* is required for both tip growth of root hairs and nod factor-induced root hair deformation. *Plant Cell*, 27(3), 806–822.
- Li, D. M., & Alexander, M. (1986). Bacterial growth rates and competition affect nodulation and root colonization by *Rhizobium meliloti*. *Applied and Environmental Microbiology*, 52(4), 807–811.
- Libault, M., Farmer, A., Brechenmacher, L., Drnevich, J., Langley, R. J., Bilgin, D. D., ... Stacey, G. (2010). Complete Transcriptome of the Soybean Root Hair Cell, a Single-Cell Model, and Its Alteration in Response to *Bradyrhizobium japonicum* Infection. *Plant Physiology*, 152(2), 541–552.
- Lithgow, J.K., Wilkinson, A., Hardman, A., Rodelas, B., Wisniewski-Dyé, F., Williams, P., Downie, J.A., 2000. The regulatory locus *cinRI* in *Rhizobium leguminosarum* controls a network of quorum-sensing loci. *Mol. Microbiol.* <https://doi.org/10.1046/j.1365-2958.2000.01960.x>
- Liu, Y., Guan, D., Jiang, X., Ma, M., Li, L., Cao, F., ... Li, J. (2015). Proteins involved in nodulation competitiveness of two *Bradyrhizobium diazoefficiens* strains induced by soybean root exudates. *Biology and Fertility of Soils*.
- Liu, Y., Jiang, X., Guan, D., Zhou, W., Ma, M., Zhao, B., ... Li, J. (2017). Transcriptional analysis of genes involved in competitive nodulation in *Bradyrhizobium diazoefficiens* at the presence of soybean root exudates. *Scientific Reports*, 7(1), 10946.
- Livak, K.J., Schmittgen, T.D., 2001. Analysis of relative gene expression data using real-time quantitative PCR and the 2- $\Delta\Delta$ CT method. *Methods* 25, 402–408. <https://doi.org/10.1006/meth.2001.1262>
- Lodwig, E., & Poole, P. (2003, January). *Metabolism of Rhizobium bacteroids*. *Critical Reviews in Plant Sciences*. Taylor & Francis Group.
- Loper, J.E., Schroth, M.N., 1986. Influence of Bacterial Sources of Indole-3-acetic Acid on Root Elongation of Sugar Beet. *Phytopathology* 76, 386. <https://doi.org/10.1094/Phyto-76-386>
- López-García, S. L., Peticari, A., Piccinetti, C., Ventimiglia, L., Arias, N., De Battista, J. J., ... Lodeiro, A. R. (2009). In-Furrow Inoculation and Selection for Higher Motility Enhances the Efficacy of Nodulation. *Agronomy Journal*, 101(2), 357.
- Lopez-Garcia, S.L., Vazquez, T.E.E., Favelukes, G., Lodeiro, A.R., 2002. Rhizobial position as a main determinant in the problem of competition for nodulation in soybean. *Environ. Microbiol.* 4, 216–224. <https://doi.org/10.1046/j.1462-2920.2002.00287.x>
- Luo, Z.Q., Su, S., Farrand, S.K., 2003. In situ activation of the quorum-sensing transcription factor TraR by cognate and noncognate acyl-homoserine lactone ligands: Kinetics and consequences. *J. Bacteriol.* 185, 5665–5672. <https://doi.org/10.1128/JB.185.19.5665-5672.2003>

- Ma, J., Campbell, A., Karlin, S., 2002. Correlations between Shine-Dalgarno Sequences and Gene Features Such as Predicted Expression Levels and Operon Structures Correlations between Shine-Dalgarno Sequences and Gene Features Such as Predicted Expression Levels and Operon Structures. <https://doi.org/10.1128/JB.184.20.5733>
- MacNeil, D.J., Gewain, K.M., Ruby, C.L., Dezeny, G., Gibbons, P.H., MacNeil, T., 1992. Analysis of *Streptomyces avermitilis* genes required for avermectin biosynthesis utilizing a novel integration vector. *Gene*. [https://doi.org/10.1016/0378-1119\(92\)90603-M](https://doi.org/10.1016/0378-1119(92)90603-M)
- Marketon, M.M., Glenn, S.A., Eberhard, A., Gonzalez, J.E., 2003. Quorum Sensing Controls Exopolysaccharide Production in *Sinorhizobium meliloti*. *J. Bacteriol.* 185, 325–331. <https://doi.org/10.1128/JB.185.1.325-331.2003>
- Marketon, M.M., González, J.E., 2002. Identification of two quorum-sensing systems in *Sinorhizobium meliloti*. *J. Bacteriol.* 184, 3466–3475. <https://doi.org/10.1128/jb.184.13.3466-3475.2002>
- Marketon, M.M., Gronquist, M.R., Eberhard, A., Gonzalez, J.E., 2002. Characterization of the *Sinorhizobium meliloti* *sinR/sinI* Locus and the Production of Novel N-Acyl Homoserine Lactones. *J. Bacteriol.* 184, 5686–5695. <https://doi.org/10.1128/JB.184.20.5686-5695.2002>
- Massalha, H., Korenblum, E., Malitsky, S., Shapiro, O. H., & Aharoni, A. (2017). Live imaging of root–bacteria interactions in a microfluidics setup. *Proceedings of the National Academy of Sciences*, 114(17), 4549–4554. <https://doi.org/10.1073/pnas.1618584114>
- Mateos, P. F., Baker, D. L., Petersen, M., Velázquez, E., Jiménez-Zurdo, J. I., Martínez-Molina, E., ... Dazzo, F. B. (2001). Erosion of root epidermal cell walls by *Rhizobium* polysaccharide-degrading enzymes as related to primary host infection in the *Rhizobium*-legume symbiosis. *Canadian Journal of Microbiology*, 47(6), 475–87.
- Mathesius, U., Charon, C., Rolfe, B. G., Kondorosi, A., & Crespi, M. (2000). Temporal and spatial order of events during the induction of cortical cell divisions in white clover by *Rhizobium leguminosarum* bv. *trifolii* Inoculation or localized cytokinin addition. *Molecular Plant-Microbe Interactions*, 13(6), 617–628.
- Matse, D.T., Huang, C.H., Huang, Y.M., Yen, M.Y., 2020. Effects of coinoculation of *Rhizobium* with plant growth promoting rhizobacteria on the nitrogen fixation and nutrient uptake of *Trifolium repens* in low phosphorus soil. *J. Plant Nutr.* <https://doi.org/10.1080/01904167.2019.1702205>
- Mazumder, R., Phelps, T.J., Krieg, N.R., Benoit, R.E., 1999. Determining chemotactic responses by two subsurface microaerophiles using a simplified capillary assay method. *J. Microbiol. Methods* 37, 255–263. [https://doi.org/10.1016/S0167-7012\(99\)00072-X](https://doi.org/10.1016/S0167-7012(99)00072-X)
- McIntosh, M., Meyer, S., & Becker, A. (2009). Novel *Sinorhizobium meliloti* quorum sensing positive and negative regulatory feedback mechanisms respond to phosphate availability. *Molecular Microbiology*, 74(5), 1238–1256. <https://doi.org/10.1111/j.1365-2958.2009.06930.x>

- McIntosh, M., Serrania, J., Lacanna, E., 2019. A novel LuxR-type solo of *Sinorhizobium meliloti*, NurR, is regulated by the chromosome replication coordinator, DnaA and activates quorum sensing. *Mol. Microbiol.* <https://doi.org/10.1111/mmi.14312>
- Meade, J., Higgins, P., Gara, F.O., 1985. Studies on the Inoculation and Competitiveness of a *Rhizobium leguminosarum* Strain in Soils Containing Indigenous Rhizobia 49, 899–903.
- Mergaert, P., Uchiumi, T., Alunni, B., Evanno, G., Cheron, A., Catrice, O., ... Kondorosi, E. (2006). Eukaryotic control on bacterial cell cycle and differentiation in the *Rhizobium*-legume symbiosis. *Proceedings of the National Academy of Sciences of the United States of America*, 103(13), 5230–5235.
- Miao, J., Zhang, N., Liu, H., Wang, H., Zhong, Z., & Zhu, J. (2018). Soil commensal rhizobia promote *Rhizobium etli* nodulation efficiency through CinR-mediated quorum sensing. *Archives of Microbiology*, 200(5), 685–694.
- Miller, D. D., Klooster, H. B. L., & Emons, A. M. C. (2000). Lipochito-Oligosaccharide Nodulation Factors Stimulate Cytoplasmic Polarity with Longitudinal Endoplasmic Reticulum and Vesicles at the Tip in Vetch Root Hairs. *Molecular Plant-Microbe Interactions*, 13(12), 1385–1390.
- Miller, L. D., Yost, C. K., Hynes, M. F., & Alexandre, G. (2007). The major chemotaxis gene cluster of *Rhizobium leguminosarum* bv. *viciae* is essential for competitive nodulation, 63(December 2006), 348–362.
- Miller, M. B., & Bassler, B. L. (2001). Quorum Sensing in Bacteria. *Annual Review of Microbiology*, 55(1), 165–199.
- Mithöfer, A., Bhagwat, A. A., Keister, D. L., & Ebel, J. (2001). *Bradyrhizobium japonicum* mutants defective in cyclic β -glucan synthesis show enhanced sensitivity to plant defense responses. *Zeitschrift Fur Naturforschung - Section C Journal of Biosciences*.
- Mongiardini, E. J., Pérez-Giménez, J., Althabegoiti, M. J., Covelli, J., Quelas, J. I., López-García, S. L., & Lodeiro, A. R. (2009). Overproduction of the rhizobial adhesin RapA1 increases competitiveness for nodulation. *Soil Biology and Biochemistry*, 41(9), 2017–2020. <https://doi.org/10.1016/j.soilbio.2009.07.016>
- Moriya, N., Minamino, T., Imada, K., Namba, K., 2011. Genetic analysis of the bacterial hook-capping protein FlgD responsible for hook assembly. *Microbiology* 157, 1354–1362. <https://doi.org/10.1099/mic.0.047100-0>
- Mueller, K., Gonzalez, J.E., 2011. Complex Regulation of Symbiotic Functions Is Coordinated by MucR and Quorum Sensing in *Sinorhizobium meliloti*. *J. Bacteriol.* 193, 485–496. <https://doi.org/10.1128/JB.01129-10>
- Naseer, I., Ahmad, M., Nadeem, S.M., Ahmad, I., Najm-ul-Seher, Zahir, Z.A., 2019. Rhizobial Inoculants for Sustainable Agriculture: Prospects and Applications. https://doi.org/10.1007/978-3-030-18933-4_11
- Nealson, K. H., Platt, T., & Hastings, J. W. (1970). Cellular control of the synthesis and activity of the bacterial luminescent system. *Journal of Bacteriology*, 104(1), 313–22.

- Niehaus, K., Lagares, A., & Pühler, A. (1998). A *Sinorhizobium meliloti* lipopolysaccharide mutant induces effective nodules on the host plant *Medicago sativa* (Alfalfa) but fails to establish a symbiosis with *Medicago truncatula*. *Molecular Plant-Microbe Interactions*.
- Nievas, F., Bogino, P., Sorroche, F., & Giordano, W. (2012). Detection, characterization, and biological effect of quorum-sensing signaling molecules in Peanut-nodulating bradyrhizobia. *Sensors*, 12(3), 2851–2873.
- Obranić, S., Babić, F., & Maravić-Vlahoviček, G. (2013). Improvement of pBBR1MCS plasmids, a very useful series of broad-host-range cloning vectors. *Plasmid*. <https://doi.org/10.1016/j.plasmid.2013.04.001>
- Ortori, C.A., Halliday, N., Cámara, M., Williams, P., Barrett, D.A., 2014. LC-MS/MS quantitative analysis of quorum sensing signal molecules, in: Filloux, A., Ramos, J.-L. (Eds.), *Methods in Molecular Biology*. Springer New York, New York, NY, pp. 255–270. https://doi.org/10.1007/978-1-4939-0473-0_21
- Papenfort, K., & Bassler, B. L. (2016). Quorum sensing signal-response systems in Gram-negative bacteria. *Nature Reviews Microbiology*, 14(9), 576–588. <https://doi.org/10.1038/nrmicro.2016.89>
- Parke, J. L. (1991). Root colonization by indigenous and introduced microorganisms. In *The Rhizosphere and Plant Growth* (pp. 33–42). Dordrecht: Springer Netherlands. https://doi.org/10.1007/978-94-011-3336-4_4
- Parsek, M. R., & Greenberg, E. P. (2000). Acyl-homoserine lactone quorum sensing in Gram-negative bacteria: A signaling mechanism involved in associations with higher organisms. *Proceedings of the National Academy of Sciences of the United States of America*.
- Patankar, A. V., & Gonzalez, J. E. (2009). An Orphan LuxR Homolog of *Sinorhizobium meliloti* Affects Stress Adaptation and Competition for Nodulation. *Applied and Environmental Microbiology*, 75(4), 946–955. <https://doi.org/10.1128/AEM.01692-08>
- Patankar, A. V., & González, J. E. (2009b). Orphan LuxR regulators of quorum sensing. *FEMS Microbiology Reviews*, 33(4), 739–756.
- Patriarca, E. J., Tatè, R., Ferraioli, S., & Iaccarino, M. (2004). Organogenesis of Legume Root Nodules. *International Review of Cytology*, 234, 201–262.
- Pellock, B. J., Teplitski, M., Boinay, R. P., Bauer, W. D., & Walker, G. C. (2002). A LuxR homolog controls production of symbiotically active extracellular polysaccharide II by *Sinorhizobium meliloti*. *Journal of Bacteriology*, 184(18), 5067–5076. <https://doi.org/10.1128/JB.184.18.5067-5076.2002>
- Pellock, B.J., Cheng, H.P., Walker, G.C., 2000. Alfalfa root nodule invasion efficiency is dependent on *Sinorhizobium meliloti* polysaccharides. *J. Bacteriol.* 182, 4310–4318. <https://doi.org/10.1128/JB.182.15.4310-4318.2000>
- Pereira, C. S., Thompson, J. A., & Xavier, K. B. (2013). AI-2-mediated signalling in bacteria. *FEMS Microbiology Reviews*.
- Philippot, L., Raaijmakers, J. M., Lemanceau, P., & Van Der Putten, W. H. (2013). Going back to the roots: The microbial ecology of the rhizosphere. *Nature Reviews Microbiology*.

- Pierre, O., Hopkins, J., Combiér, M., Baldacci, F., Engler, G., Brouquisse, R., ... Boncompagni, E. (2014). Involvement of papain and legumain proteinase in the senescence process of *Medicago truncatula* nodules. *New Phytologist*, 202(3), 849–863.
- Pillai, P., Archana, G., 2008. Hide depilation and feather disintegration studies with keratinolytic serine protease from a novel *Bacillus subtilis* isolate. *Appl. Microbiol. Biotechnol.* 78, 643–650. <https://doi.org/10.1007/s00253-008-1355-z>
- Pini, F., East, A. K., Appia-Ayme, C., Tomek, J., Karunakaran, R., Mendoza-Suárez, M., ... Poole, P. S. (2017). Bacterial biosensors for in vivo spatiotemporal mapping of root secretion. *Plant Physiology*, 174(3), 1289–1306.
- Ponnusamy, K., Paul, D., Kweon, J.H., 2009. Inhibition of quorum sensing mechanism and *Aeromonas hydrophila* biofilm formation by vanillin. *Environ. Eng. Sci.* 26, 1359–1363. <https://doi.org/10.1089/ees.2008.0415>
- Poole, P. S. (2017). Title : Bacterial biosensors for in vivo spatiotemporal mapping of root secretion, (0).
- Poole, P., Ramachandran, V., & Terpolilli, J. (2018). Rhizobia: from saprophytes to endosymbionts. *Nature Reviews Microbiology*, 16(5), 291–303. <https://doi.org/10.1038/nrmicro.2017.171>
- Prell, J., & Poole, P. (2006). Metabolic changes of rhizobia in legume nodules. *Trends in Microbiology*, 14(4), 161–8.
- Rachwał, K., Lipa, P., Wojda, I., Vinardell, J.M., Janczarek, M., 2017. Regulatory elements located in the upstream region of the *Rhizobium leguminosarum* rosR global regulator are essential for its transcription and mRNA stability. *Genes (Basel)*. 8. <https://doi.org/10.3390/genes8120388>
- Rachwał, K., Matczyńska, E., Janczarek, M., 2015. Transcriptome profiling of a *Rhizobium leguminosarum* bv. *trifolii* rosR mutant reveals the role of the transcriptional regulator RosR in motility, synthesis of cell-surface components, and other cellular processes. *BMC Genomics* 16, 1111. <https://doi.org/10.1186/s12864-015-2332-4>
- Raina, J.B., Fernandez, V., Lambert, B., Stocker, R., Seymour, J.R., 2019. The role of microbial motility and chemotaxis in symbiosis. *Nat. Rev. Microbiol.* <https://doi.org/10.1038/s41579-019-0182-9>
- Ravn, L., Beck, A., Molin, S., Givskov, M., 2001. Methods for detecting acylated homoserine lactones produced by Gram-negative bacteria and their application in studies of AHL-production kinetics. *Image (Rochester, N.Y.)* 239–251.
- Redfield, R. J. (2002). Is quorum sensing a side effect of diffusion sensing? *Trends in Microbiology*.
- Ren, B., Wang, X., Duan, J., & Ma, J. (2019). Rhizobial tRNA-derived small RNAs are signal molecules regulating plant nodulation. *Science (New York, N.Y.)*, 365(6456), 919–922.
- Rinaudi, L. V, Gonzalez, J.E., 2009. The Low-Molecular-Weight Fraction of Exopolysaccharide II from *Sinorhizobium meliloti* Is a Crucial Determinant of

- Biofilm Formation. *J. Bacteriol.* 191, 7216–7224. <https://doi.org/10.1128/JB.01063-09>
- Rinaudi, L. V., Giordano, W., 2010. An integrated view of biofilm formation in rhizobia. *FEMS Microbiol. Lett.* 304, 1–11. <https://doi.org/10.1111/j.1574-6968.2009.01840.x>
- Rinaudi-Marron, L. V., González, J.E., 2015. Regulation of symbiotically important functions by quorum sensing in the *Sinorhizobium meliloti* -Alfalfa interaction, in: *Biological Nitrogen Fixation*. John Wiley & Sons, Inc, Hoboken, NJ, USA, pp. 381–388. <https://doi.org/10.1002/9781119053095.ch37>
- Rivas, R., García-Fraile, P., & Velázquez, E. (2009). Taxonomy of Bacteria Nodulating Legumes. *Microbiology Insights*, 2, MBIS3137.
- Rivilla, R., Martín, M., & Lloret, J. (2017). What makes rhizobia rhizosphere colonizers? *Environmental Microbiology*, 0. <https://doi.org/10.1111/1462-2920.13917>
- Rodelas, B., Lithgow, J.K., Wisniewski-Dye, F., Hardman, a, Wilkinson, a, Economou, a, Williams, P., Downie, J. a, 1999. Analysis of quorum-sensing-dependent control of rhizosphere-expressed (rhi) genes in *Rhizobium leguminosarum* bv. *viciae*. *J. Bacteriol.* 181, 3816–23.
- Roumiantseva, M. L., & Muntyan, V. S. (2015). Root nodule bacteria *Sinorhizobium meliloti*: Tolerance to salinity and bacterial genetic determinants. *Microbiology*, 84(3), 303–318.
- Rubio-Canalejas, A., Celador-Lera, L., Cruz-González, X., Menéndez, E., & Rivas, R. (2016). *Rhizobium* as potential biofertilizer of *Eruca Sativa*. In *Biological Nitrogen Fixation and Beneficial Plant-Microbe Interaction* (pp. 213–220). Springer.
- Rusconi, R., Garren, M., & Stocker, R. (2014). Microfluidics Expanding the Frontiers of Microbial Ecology. *Annual Review of Biophysics*, 43(1), 65–91.
- Russo, D.M., Williams, A., Edwards, A., Posadas, D.M., Finnie, C., Dankert, M., Downie, J.A., Zorreguieta, A., 2006. Proteins exported via the PrsD-PrsE type I secretion system and the acidic exopolysaccharide are involved in biofilm formation by *Rhizobium leguminosarum*. *J. Bacteriol.* <https://doi.org/10.1128/JB.00246-06>
- Sanchez-Contreras, M., Bauer, W.D., Gao, M., Robinson, J.B., Allan Downie, J., 2007. Quorum-sensing regulation in rhizobia and its role in symbiotic interactions with legumes. *Philos. Trans. R. Soc. Lond. B. Biol. Sci.* 362, 1149–1163. <https://doi.org/10.1098/rstb.2007.2041>
- Santaella, C., Schue, M., Berge, O., Heulin, T., Achouak, W., 2008. The exopolysaccharide of *Rhizobium* sp. YAS34 is not necessary for biofilm formation on *Arabidopsis thaliana* and *Brassica napus* roots but contributes to root colonization. *Environ. Microbiol.* 10, 2150–2163. <https://doi.org/10.1111/j.1462-2920.2008.01650.x>
- Saraf, M., Pandya, U., Thakkar, A., 2014. Role of allelochemicals in plant growth promoting rhizobacteria for biocontrol of phytopathogens. *Microbiol. Res.* 169, 18–29. <https://doi.org/10.1016/J.MICRES.2013.08.009>

- Saurav, K., Burgsdorf, I., Teta, R., Esposito, G., Bar-Shalom, R., Costantino, V., Steindler, L., 2016. Isolation of marine *Paracoccus* sp. Ss63 from the sponge *Sarcotragus* sp. and characterization of its quorum-sensing chemical-signaling molecules by LC-MS/MS Analysis. *Isr. J. Chem.* 56, 330–340. <https://doi.org/10.1002/ijch.201600003>
- Savka, M. A., Dessaux, Y., Oger, P., & Rossbach, S. (2002). Engineering Bacterial Competitiveness and Persistence in the Phytosphere. *Society*, 15(9), 866–874.
- Saxena, K. B., Kumar, R. V., & Sultana, R. (2010). Quality nutrition through pigeonpea—a review. *Health*, 2(11), 1335–1344. <https://doi.org/10.4236/health.2010.211199>
- Saxena, R. K., Saxena, K. B., & Varshney, R. K. (2019). Pigeonpea (*Cajanus cajan* L. Millsp.): An Ideal Crop for Sustainable Agriculture. In *Advances in Plant Breeding Strategies: Legumes* (pp. 409-429). Springer, Cham.
- Scharf, B. E., Hynes, M. F., & Alexandre, G. M. (2016). Chemotaxis signaling systems in model beneficial plant – bacteria associations. *Plant Molecular Biology*.
- Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., ... Cardona, A. (2012). Fiji: An open-source platform for biological-image analysis. *Nature Methods*. <https://doi.org/10.1038/nmeth.2019>
- Schmidt, H., Nunan, N., Höck, A., Eickhorst, T., Kaiser, C., Wobken, D., Raynaud, X., 2018. Recognizing patterns: Spatial analysis of observed microbial colonization on root surfaces. *Front. Environ. Sci.* 6, 1–12. <https://doi.org/10.3389/fenvs.2018.00061>
- Schroeder, B.K., House, B.L., Mortimer, M.W., Yurgel, S.N., Maloney, S.C., Ward, K.L., Kahn, M.L., 2005. Development of a functional genomics platform for *Sinorhizobium meliloti*: Construction of an ORFeome. *Appl. Environ. Microbiol.* <https://doi.org/10.1128/AEM.71.10.5858-5864.2005>
- Schwember, A. R., Schulze, J., del Pozo, A., & Cabeza, R. A. (2019). Regulation of Symbiotic Nitrogen Fixation in Legume Root Nodules. *Plants*, 8(9), 333.
- Schwyn, B., Neilands, J.B., 1987. Universal chemical assay for the detection and determination of siderophores. *Anal. Biochem.* 160, 47–56. [https://doi.org/10.1016/0003-2697\(87\)90612-9](https://doi.org/10.1016/0003-2697(87)90612-9)
- Selma, M. V., Martínez-Sánchez, A., Allende, A., Ros, M., Hernández, M. T., & Gil, M. I. (2010). Impact of organic soil amendments on phytochemicals and microbial quality of rocket leaves (*Eruca sativa*). *Journal of Agricultural and Food Chemistry*.
- Sessitsch, A., Howieson, J. G., Perret, X., Antoun, H., & Martínez-Romero, E. (2002). *Advances in Rhizobium Research. Critical Reviews in Plant Sciences*, 21(4), 323–378.
- Smit, G., Kijne, J. W., & Lugtenberg, B. J. (1987). Involvement of both cellulose fibrils and a Ca²⁺-dependent adhesin in the attachment of *Rhizobium leguminosarum* to pea root hair tips. *Journal of Bacteriology*.
- Shaw, P.D., Ping, G., Daly, S.L., Cha, C., Cronan, J.E., Rinehart, K.L., Farrand, S.K., 1997. Detecting and characterizing N-acyl-homoserine lactone signal

- molecules by thin-layer chromatography. *Proc. Natl. Acad. Sci.* 94, 6036–6041. <https://doi.org/10.1073/pnas.94.12.6036>
- Sorroche, F., Bogino, P., Russo, D. M., Zorreguieta, A., Nievas, F., Morales, G. M., ... Giordano, W. (2018). Cell autoaggregation, biofilm formation, and plant attachment in a sinorhizobium meliloti lpsB mutant. *Molecular Plant-Microbe Interactions*.
- Sorroche, F., Walch, M., Zou, L., Rengel, D., Maillet, F., Gibelin-Viala, C., ... Garnerone, A. M. (2019). Endosymbiotic Sinorhizobium meliloti modulate Medicago root susceptibility to secondary infection via ethylene. *New Phytologist*, 223(3), 1505–1515.
- Sourjik, V., Muschler, P., Scharf, B., & Schmitt, R. (2000). VisN and VisR Are Global Regulators of Chemotaxis, Flagellar, and Motility Genes in Sinorhizobium (Rhizobium) meliloti. *Journal of Bacteriology*, 182(3), 782–788.
- Sourjik, V., Schmitt, R., 1996. Different roles of CheY1 and CheY2 in the chemotaxis of Rhizobium meliloti. *Mol. Microbiol.* 22, 427–436. <https://doi.org/10.1046/j.1365-2958.1996.1291489.x>
- Spini, G., Decorosi, F., Cerboneschi, M., Tegli, S., Mengoni, A., Viti, C., Giovannetti, L., 2016. Effect of the plant flavonoid luteolin on Ensifer meliloti 3001 phenotypic responses. *Plant Soil*. <https://doi.org/10.1007/s11104-015-2659-2>
- Stacey, G. (2007). The Rhizobium-Legume Nitrogen-Fixing Symbiosis. In *Biology of the Nitrogen Cycle* (pp. 147–163). Elsevier B.V.
- Stanfield, S.W., Ielpi, L., O'Brochta, D., Helinski, D.R., Ditta, G.S., 1988. The ndvA gene product of Rhizobium meliloti is required for beta-(1---2)glucan production and has homology to the ATP-binding export protein HlyB. *J. Bacteriol.* 170, 3523–3530. <https://doi.org/10.1128/jb.170.8.3523-3530.1988>
- Steindler, L., Venturi, V., 2007. Detection of quorum-sensing N-acyl homoserine lactone signal molecules by bacterial biosensors. *FEMS Microbiol. Lett.* 266, 1–9. <https://doi.org/10.1111/j.1574-6968.2006.00501.x>
- Sturme, M. H. J., Kleerebezem, M., Nakayama, J., Akkermans, A. D. L., Vaughan, E. E., & De Vos, W. M. (2002). Cell to cell communication by autoinducing peptides in gram-positive bacteria. *Antonie van Leeuwenhoek, International Journal of General and Molecular Microbiology*.
- Sugawara, M., & Sadowsky, M. J. (2014). Enhanced Nodulation and Nodule Development by nolR Mutants of Sinorhizobium medicae on Specific Medicago Host Genotypes. *Molecular Plant-Microbe Interactions*, 27(4), 328–335.
- Suzuki, T., & Nishida, H. (2019). Autoregulation of Legume Nodulation by Sophisticated Transcriptional Regulatory Networks. *Molecular Plant*, 12(9), 1179–1181.
- Tambalo, D.D., Del Bel, K.L., Bustard, D.E., Greenwood, P.R., Steedman, A.E., Hynes, M.F., 2010. Regulation of flagellar, motility and chemotaxis genes in Rhizobium leguminosarum by the VisN/R-Rem cascade. *Microbiology* 156, 1673–1685. <https://doi.org/10.1099/mic.0.035386-0>

- Taylor, P., Sessitsch, A., Howieson, J. G., Perret, X., & Antoun, H. (2002). Please scroll down for article. Online, (909277354).
- Tecon, R., & Or, D. (2017). Biophysical processes supporting the diversity of microbial life in soil. *FEMS Microbiology Reviews*, 41(5), 599–623.
- Teplitski, M., Eberhard, A., Gronquist, M.R., Gao, M., Robinson, J.B., Bauer, W.D., 2003. Chemical identification of N -acyl homoserine lactone quorum-sensing signals produced by *Sinorhizobium meliloti* strains in defined medium. *Arch. Microbiol.* 180, 494–497. <https://doi.org/10.1007/s00203-003-0612-x>
- Thies, J. E., Singleton, P. W., & Bohlool, B. B. (1991). Modeling symbiotic performance of introduced rhizobia in the field by use of indices of indigenous population size and nitrogen status of the soil. *Applied and Environmental Microbiology*.
- Tkacz, A., Cheema, J., Chandra, G., Grant, A., & Poole, P. S. (2015). Stability and succession of the rhizosphere microbiota depends upon plant type and soil composition. *The ISME Journal*, 9(11), 2349–2359.
- Toro, N. (1996). Nodulation competitiveness in the R / VzoMwwlegume symbiosis, 157–162.
- Turgeon, B. G., & Bauer, W. D. (1982). Early events in the infection of soybean by *Rhizobium japonicum* . Time course and cytology of the initial infection process. *Canadian Journal of Botany*, 60(2), 152–161.
- Turner, T. R., Ramakrishnan, K., Walshaw, J., Heavens, D., Alston, M., Swarbreck, D., ... Poole, P. S. (2013). Comparative metatranscriptomics reveals kingdom level changes in the rhizosphere microbiome of plants. *ISME Journal*.
- Vanderlinde, E. M., Muszyński, A., Harrison, J. J., Koval, S. F., Foreman, D. L., Ceri, H., ... Yost, C. K. (2009). *Rhizobium leguminosarum* biovar *viciae* 3841, deficient in 27-hydroxyoctacosanoate-modified lipopolysaccharide, is impaired indeseccation tolerance, biofilm formation and motility. *Microbiology*, 155(9), 3055–3069.
- Varshney, R. K., Chen, W., Li, Y., Bharti, A. K., Saxena, R. K., Schlueter, J. A., ... Jackson, S. A. (2012). Draft genome sequence of pigeonpea (*Cajanus cajan*), an orphan legume crop of resource-poor farmers. *Nature Biotechnology*, 30(1), 83–89. <https://doi.org/10.1038/nbt.2022>
- Vincent, J.M., 1970. A Manual for the Practical Study of Root-nodule Bacteria. [Published for the] International Biological Programme [by] Blackwell Scientific.
- Vlassak, K. M., & Vanderleyden, J. (1997). Factors Influencing Nodule Occupancy by Inoculant Rhizobia. *Critical Reviews in Plant Sciences*, 16(2), 163–229.
- Wadisirisuk, P., Danso, S. K., Hardarson, G., & Bowen, G. D. (1989). Influence of *Bradyrhizobium japonicum* Location and Movement on Nodulation and Nitrogen Fixation in Soybeans. *Applied and Environmental Microbiology*, 55(7), 1711–1716.
- Walker, S. A., & Downie, J. A. (2000). Entry of *Rhizobium leguminosarum* bv. *viciae* into root hairs requires minimal nod factor specificity, but subsequent

- infection thread growth requires nodO or nodE. *Molecular Plant-Microbe Interactions*, 13(7), 754–762.
- Wang, Q., Garrity, G.M., Tiedje, J.M., Cole, J.R., 2007. Naive Bayesian classifier for rapid assignment of rRNA sequences into the new bacterial taxonomy. *Appl. Environ. Microbiol.* 73, 5261–5267. <https://doi.org/10.1128/AEM.00062-07>
- Webb, B. A., Hildreth, S., Helm, R. F., & Scharf, B. E. (2014). Sinorhizobium meliloti chemoreceptor McpU mediates chemotaxis toward host plant exudates through direct proline sensing. *Applied and Environmental Microbiology*, 80(11), 3404–15.
- Wei, X., & Bauer, W. D. (1998). Starvation-induced changes in motility, chemotaxis, and flagellation of *Rhizobium meliloti*. *Applied and Environmental Microbiology*.
- Weir, B.S. (2016) The current taxonomy of rhizobia. NZ Rhizobia website. <https://www.rhizobia.co.nz/taxonomy/rhizobia> Last updated: X Jan, 2016
- West, S. A., Winzer, K., Gardner, A., & Diggle, S. P. (2012). Quorum sensing and the confusion about diffusion. *Trends in Microbiology*.
- Wheatley, R.M., Poole, P.S., 2018. Mechanisms of bacterial attachment to roots. *FEMS Microbiol. Rev.* 448–461. <https://doi.org/10.1093/femsre/fuy014>
- Whitehead, N. A., Barnard, A. M. L., Slater, H., Simpson, N. J. L., & Salmond, G. P. C. (2001). Quorum-sensing in Gram-negative bacteria. *FEMS Microbiology Reviews.* 25, 365–404. [https://doi.org/10.1016/S0168-6445\(01\)00059-6](https://doi.org/10.1016/S0168-6445(01)00059-6)
- Wielbo, J., Marek-Kozaczuk, M., Kubik-Komar, A., & Skorupska, A. (2007). Increased metabolic potential of *Rhizobium* spp. is associated with bacterial competitiveness. *Canadian Journal of Microbiology*.
- Williams, A., Wilkinson, A., Krehenbrink, M., Russo, D.M., Zorreguieta, A., Downie, J.A., 2008. Glucosaminan-mediated attachment of *Rhizobium leguminosarum* to pea root hairs is required for competitive nodule infection. *J. Bacteriol.* 190, 4706–4715. <https://doi.org/10.1128/JB.01694-07>
- Willis, L. B., & Walker, G. C. (1998). The phbC (poly- β -hydroxybutyrate synthase) gene of *Rhizobium* (*Sinorhizobium*) *meliloti* and characterization of phbC mutants. *Canadian Journal of Microbiology*.
- Wilson, K., 2001. Preparation of Genomic DNA from Bacteria. *Curr. Protoc. Mol. Biol.* 56, 2.4.1-2.4.5. <https://doi.org/10.1002/0471142727.mb0204s56>
- Wisniewski-Dyé, F., Downie, J.A., 2002. Quorum-sensing in *Rhizobium*. *Antonie van Leeuwenhoek, Int. J. Gen. Mol. Microbiol.* 81, 397–407. <https://doi.org/10.1023/A:1020501104051>
- Wisniewski-Dyé, F., Jones, J., Chhabra, S.R., Downie, J.A., 2002. *raiIR* genes are part of a quorum-sensing network controlled by *cinI* and *cinR* in *Rhizobium leguminosarum*. *J. Bacteriol.* 184, 1597–1606. <https://doi.org/10.1128/JB.184.6.1597-1606.2002>
- Wood, S. M., & Newcomb, W. (1989). Nodule morphogenesis: the early infection of Alfalfa (*Medicago sativa*) root hairs by *Rhizobium meliloti*. *Canadian Journal of Botany*, 67(10), 3108–3122.

- Yates, R. J., Howieson, J. G., Reeve, W. G., & O'Hara, G. W. (2011). A re-appraisal of the biology and terminology describing rhizobial strain success in nodule occupancy of legumes in agriculture. *Plant and Soil*, 348(1–2), 255–267.
- Peter, J., Young, W., & Haukka, K. E. (1996). Diversity and phylogeny of rhizobia. *New phytologist*, 133(1), 87-94.
- Zaim, S., Bekkar, A. A., & Belabid, L. (2017). Rhizobium as a Crop Enhancer and Biofertilizer for Increased Non-legume Production. In *Rhizobium Biology and Biotechnology* (pp. 25–37). Springer.
- Zatakia, H.M., Nelson, C.E., Syed, U.J., Scharf, B.E., 2014. ExpR coordinates the expression of symbiotically important, bundle-forming Flp pili with quorum sensing in *Sinorhizobium meliloti*. *Appl. Environ. Microbiol.* 80, 2429–2439. <https://doi.org/10.1128/AEM.04088-13>
- Zgad Zaj, R., Garrido-Oter, R., Jensen, D. B., Koprivova, A., Schulze-Lefert, P., & Radutoiu, S. (2016). Root nodule symbiosis in *Lotus japonicus* drives the establishment of distinctive rhizosphere, root, and nodule bacterial communities. *Proceedings of the National Academy of Sciences*, 113(49), E7996–E8005.
- Zheng, H., Mao, Y., Teng, J., Zhu, Q., Ling, J., & Zhong, Z. (2015). Flagellar-dependent motility in *Mesorhizobium tianshanense* is involved in the early stage of plant host interaction: Study of an flgE mutant. *Current Microbiology*.
- Zheng, H., Mao, Y., Zhu, Q., Ling, J., Zhang, N., Naseer, N., Zhong, Z., Zhu, J., 2015. The quorum sensing regulator CinR hierarchically regulates two other quorum sensing pathways in ligand-dependent and -independent fashions in *Rhizobium etli*. *J. Bacteriol.* 197, 1573–1581. <https://doi.org/10.1128/JB.00003-15>.