

# **Bibliography**



## 6. Bibliography

- Aist, J.R., and P.H. Williams. 1972. Ultrastructure and time course of mitosis in the fungus *Fusarium oxysporum*. *J. Cell Biol.* 55:368-389.
- Asbury, C.L., D.R. Gestaut, A.F. Powers, A.D. Franck, and T.N. Davis. 2006. The Dam1 kinetochore complex harnesses microtubule dynamics to produce force and movement. *Proc. Natl. Acad. Sci. U S A.* 103:9873-9878.
- Biggins, S. 2013. The composition, functions, and regulation of the budding yeast kinetochore. *Genetics.* 194:817-846.
- Boettcher, B., and Y. Barral. 2013. The cell biology of open and closed mitosis. *Nucleus.* 4:160-165.
- Burrack, L.S., S.E. Applen, and J. Berman. 2011. The requirement for the Dam1 complex is dependent upon the number of kinetochore proteins and microtubules. *Curr. Biol.* 21:889-896.
- Buttrick, G.J., T.C. Lancaster, J.C. Meadows, and J.B. Millar. 2012. Plo1 phosphorylates Dam1 to promote chromosome bi-orientation in fission yeast. *J. Cell Sci.* 125:1645-1651.
- Cai, X., X. Zhang, X. Li, M. Liu, X. Liu, X. Wang, H. Zhang, X. Zheng, and Z. Zhang. 2017. The Atypical Guanylate Kinase MoGuk2 Plays Important Roles in Asexual/Sexual Development, Conidial Septation, and Pathogenicity in the Rice Blast Fungus. *Front. Microbiol.* 8:2467.
- Cheeseman, I.M. 2014. The kinetochore. *Cold Spring Harb. Perspect. Biol.* 6:a015826.
- Cheeseman, I.M., S. Anderson, M. Jwa, E.M. Green, J. Kang, J.R. Yates, 3rd, C.S. Chan, D.G. Drubin, and G. Barnes. 2002. Phospho-regulation of kinetochore-microtubule attachments by the Aurora kinase Ipl1p. *Cell.* 111:163-172.
- Cheeseman, I.M., C. Brew, M. Wolyniak, A. Desai, S. Anderson, N. Muster, J.R. Yates, T.C. Huffaker, D.G. Drubin, and G. Barnes. 2001a. Implication of a novel multiprotein Dam1p complex in outer kinetochore function. *J. Cell Biol.* 155:1137-1145.
- Cheeseman, I.M., M. Enquist-Newman, T. Muller-Reichert, D.G. Drubin, and G. Barnes. 2001b. Mitotic spindle integrity and kinetochore function linked by the Duo1p/Dam1p complex. *J. Cell Biol.* 152:197-212.
- Chen, J., W. Zheng, S. Zheng, D. Zhang, W. Sang, X. Chen, G. Li, G. Lu, and Z. Wang. 2008. Rac1 is required for pathogenicity and Chm1-dependent conidiogenesis in rice fungal pathogen *Magnaporthe grisea*. *PLoS Pathog.* 4:e1000202.
- Chen, Y., X. Le, Y. Sun, M. Li, H. Zhang, X. Tan, D. Zhang, Y. Liu, and Z. Zhang. 2017. MoYcp4 is required for growth, conidiogenesis and pathogenicity in *Magnaporthe oryzae*. *Mol. Plant Pathol.* 18:1001-1011.
- Chen, Y., S. Zhai, Y. Sun, M. Li, Y. Dong, X. Wang, H. Zhang, X. Zheng, P. Wang, and Z. Zhang. 2015. MoTup1 is required for growth,

- conidiogenesis and pathogenicity of *Magnaporthe oryzae*. *Mol. Plant Pathol.* 16:799-810.
- Chen, Y., S. Zhai, H. Zhang, R. Zuo, J. Wang, M. Guo, X. Zheng, P. Wang, and Z. Zhang. 2014a. Shared and distinct functions of two Gti1/Pac2 family proteins in growth, morphogenesis and pathogenicity of *Magnaporthe oryzae*. *Environ. Microbiol.* 16:788-801.
- Chen, Y., R. Zuo, Q. Zhu, Y. Sun, M. Li, Y. Dong, Y. Ru, H. Zhang, X. Zheng, and Z. Zhang. 2014b. MoLys2 is necessary for growth, conidiogenesis, lysine biosynthesis, and pathogenicity in *Magnaporthe oryzae*. *Fungal Genet. Biol.* 67:51-57.
- Choi, J., K.S. Kim, H.S. Rho, and Y.H. Lee. 2011. Differential roles of the phospholipase C genes in fungal development and pathogenicity of *Magnaporthe oryzae*. *Fungal Genet. Biol.* 48:445-455.
- Choi, J., Y. Kim, S. Kim, J. Park, and Y.H. Lee. 2009. MoCRZ1, a gene encoding a calcineurin-responsive transcription factor, regulates fungal growth and pathogenicity of *Magnaporthe oryzae*. *Fungal Genet. Biol.* 46:243-254.
- Choi, W., and R.A. Dean. 1997. The adenylate cyclase gene MAC1 of *Magnaporthe grisea* controls appressorium formation and other aspects of growth and development. *Plant Cell.* 9:1973-1983.
- Ciferri, C., A. Musacchio, and A. Petrovic. 2007. The Ndc80 complex: hub of kinetochore activity. *FEBS Lett.* 581:2862-2869.
- Couch, B.C., and L.M. Kohn. 2002. A multilocus gene genealogy concordant with host preference indicates segregation of a new species, *Magnaporthe oryzae*, from *M. grisea*. *Mycologia.* 94:683-693.
- Dawe, A.L., V.C. McMains, M. Panglao, S. Kasahara, B. Chen, and D.L. Nuss. 2003. An ordered collection of expressed sequences from *Cryphonectria parasitica* and evidence of genomic microsynteny with *Neurospora crassa* and *Magnaporthe grisea*. *Microbiology.* 149:2373-2384.
- De Souza, C.P., S.B. Hashmi, T. Nayak, B. Oakley, and S.A. Osmani. 2009. Mlp1 acts as a mitotic scaffold to spatially regulate spindle assembly checkpoint proteins in *Aspergillus nidulans*. *Mol. Biol. Cell.* 20:2146-2159.
- De Souza, C.P., and S.A. Osmani. 2007. Mitosis, not just open or closed. *Eukaryot. Cell.* 6:1521-1527.
- Deng, Y.Z., Z. Qu, Y. He, and N.I. Naqvi. 2012. Sorting nexin Snx41 is essential for conidiation and mediates glutathione-based antioxidant defense during invasive growth in *Magnaporthe oryzae*. *Autophagy.* 8:1058-1070.
- Deng, Y.Z., M. Ramos-Pamplona, and N.I. Naqvi. 2009. Autophagy-assisted glycogen catabolism regulates asexual differentiation in *Magnaporthe oryzae*. *Autophagy.* 5:33-43.
- Dhatchinamoorthy, K., M. Shivaraju, J.J. Lange, B. Rubinstein, J.R. Unruh, B.D. Slaughter, and J.L. Gerton. 2017. Structural plasticity of the living kinetochore. *J. Cell Biol.* 216:3551-3570.

- Ding, R., K.L. McDonald, and J.R. McIntosh. 1993. Three-dimensional reconstruction and analysis of mitotic spindles from the yeast, *Schizosaccharomyces pombe*. *J. Cell Biol.* 120:141-151.
- Dong, Y., Q. Zhao, X. Liu, X. Zhang, Z. Qi, H. Zhang, X. Zheng, and Z. Zhang. 2015. MoMyb1 is required for asexual development and tissue-specific infection in the rice blast fungus *Magnaporthe oryzae*. *BMC Microbiol.* 15:37.
- Du, Y., L. Hong, W. Tang, L. Li, X. Wang, H. Ma, Z. Wang, H. Zhang, X. Zheng, and Z. Zhang. 2014. Threonine deaminase Mollv1 is important for conidiogenesis and pathogenesis in the rice blast fungus *Magnaporthe oryzae*. *Fungal Genet. Biol.* 73:53-60.
- Du, Y., Y. Shi, J. Yang, X. Chen, M. Xue, W. Zhou, and Y.L. Peng. 2013. A serine/threonine-protein phosphatase PP2A catalytic subunit is essential for asexual development and plant infection in *Magnaporthe oryzae*. *Curr. Genet.* 59:33-41.
- Enquist-Newman, M., I.M. Cheeseman, D. Van Goor, D.G. Drubin, P.B. Meluh, and G. Barnes. 2001. Dad1p, third component of the Duo1p/Dam1p complex involved in kinetochore function and mitotic spindle integrity. *Mol. Biol. Cell.* 12:2601-2613.
- Fernandez, J., and K. Orth. 2018. Rise of a Cereal Killer: The Biology of *Magnaporthe oryzae* Biotrophic Growth. *Trends Microbiol.* 26:582-597.
- Franco, A., J.C. Meadows, and J.B. Millar. 2007. The Dam1/DASH complex is required for the retrieval of unclustered kinetochores in fission yeast. *J. Cell Sci.* 120:3345-3351.
- Freitag, M. 2016. The kinetochore interaction network (KIN) of ascomycetes. *Mycologia.* 108:485-505.
- Freitag, M., P.C. Hickey, N.B. Raju, E.U. Selker, and N.D. Read. 2004. GFP as a tool to analyze the organization, dynamics and function of nuclei and microtubules in *Neurospora crassa*. *Fungal Genet. Biol.* 41:897-910.
- Fukagawa, T. 2017. Critical histone post-translational modifications for centromere function and propagation. *Cell Cycle.* 16:1259-1265.
- Funabiki, H., I. Hagan, S. Uzawa, and M. Yanagida. 1993. Cell cycle-dependent specific positioning and clustering of centromeres and telomeres in fission yeast. *J. Cell Biol.* 121:961-976.
- Gao, H.M., X.G. Liu, H.B. Shi, J.P. Lu, J. Yang, F.C. Lin, and X.H. Liu. 2013. MoMon1 is required for vacuolar assembly, conidiogenesis and pathogenicity in the rice blast fungus *Magnaporthe oryzae*. *Res. Microbiol.* 164:300-309.
- Gao, Q., T. Courtheoux, Y. Gachet, S. Tournier, and X. He. 2010. A non-ring-like form of the Dam1 complex modulates microtubule dynamics in fission yeast. *Proc Natl Acad Sci U S A.* 107:13330-13335.
- Goh, J., K.S. Kim, J. Park, J. Jeon, S.Y. Park, and Y.H. Lee. 2011. The cell cycle gene MoCDC15 regulates hyphal growth, asexual development and plant infection in the rice blast pathogen *Magnaporthe oryzae*. *Fungal Genet. Biol.* 48:784-792.

- Goshima, G., S. Saitoh, and M. Yanagida. 1999. Proper metaphase spindle length is determined by centromere proteins Mis12 and Mis6 required for faithful chromosome segregation. *Genes Dev.* 13:1664-1677.
- Goshima, G., and J.M. Scholey. 2010. Control of mitotic spindle length. *Annu. Rev. Cell Dev. Biol.* 26:21-57.
- Grishchuk, E.L., A.K. Efremov, V.A. Volkov, I.S. Spiridonov, N. Gudimchuk, S. Westermann, D. Drubin, G. Barnes, J.R. McIntosh, and F.I. Ataullakhanov. 2008a. The Dam1 ring binds microtubules strongly enough to be a processive as well as energy-efficient coupler for chromosome motion. *Proc. Natl. Acad. Sci. U S A.* 105:15423-15428.
- Grishchuk, E.L., I.S. Spiridonov, V.A. Volkov, A. Efremov, S. Westermann, D. Drubin, G. Barnes, F.I. Ataullakhanov, and J.R. McIntosh. 2008b. Different assemblies of the DAM1 complex follow shortening microtubules by distinct mechanisms. *Proc. Natl. Acad. Sci. U S A.* 105:6918-6923.
- Guo, M., F. Gao, X. Zhu, X. Nie, Y. Pan, and Z. Gao. 2015. MoGrr1, a novel F-box protein, is involved in conidiogenesis and cell wall integrity and is critical for the full virulence of *Magnaporthe oryzae*. *Appl. Microbiol. Biotechnol.* 99:8075-8088.
- Guo, M., L. Tan, X. Nie, and Z. Zhang. 2017. A class-II myosin is required for growth, conidiation, cell wall integrity and pathogenicity of *Magnaporthe oryzae*. *Virulence*:1-20.
- Hagan, I.M., A. Grallert, and V. Simanis. 2016. Analysis of the *Schizosaccharomyces pombe* Cell Cycle. *Cold Spring Harb. Protoc.* 2016.
- Hamer, J.E., B. Valent, and F.G. Chumley. 1989. Mutations at the smo genetic locus affect the shape of diverse cell types in the rice blast fungus. *Genetics.* 122:351-361.
- Han, J.H., J.H. Shin, Y.H. Lee, and K.S. Kim. 2018. Distinct roles of the YPEL gene family in development and pathogenicity in the ascomycete fungus *Magnaporthe oryzae*. *Sci. Rep.* 8:14461.
- Harris, S.D. 2008. Branching of fungal hyphae: regulation, mechanisms and comparison with other branching systems. *Mycologia.* 100:823-832.
- Harris, S.D. 2019. Hyphal branching in filamentous fungi. *Dev Biol.* 451:35-39.
- Hartwell, L.H., J. Culotti, and B. Reid. 1970. Genetic control of the cell-division cycle in yeast. I. Detection of mutants. *Proc. Natl. Acad. Sci. U S A.* 66:352-359.
- He, M., M.J. Kershaw, D.M. Soanes, Y. Xia, and N.J. Talbot. 2012. Infection-associated nuclear degeneration in the rice blast fungus *Magnaporthe oryzae* requires non-selective macro-autophagy. *PLoS One.* 7:e33270.
- He, Y., Y.Z. Deng, and N.I. Naqvi. 2013. Atg24-assisted mitophagy in the foot cells is necessary for proper asexual differentiation in *Magnaporthe oryzae*. *Autophagy.* 9:1818-1827.
- Hofmann, C., I. Cheeseman, B. Goode, K.L. McDonald, G. Barnes, and D.G. Drubin. 1998. *Saccharomyces cerevisiae* Duo1p and Dam1p, Novel

- Proteins Involved in Mitotic Spindle Function. *J. Cell Biol.* 143:1029-1040.
- Hou, H., Z. Zhou, Y. Wang, J. Wang, S.P. Kallgren, T. Kurchuk, E.A. Miller, F. Chang, and S. Jia. 2012. Csi1 links centromeres to the nuclear envelope for centromere clustering. *J. Cell Biol.* 199:735-744.
- Huang, L., S. Zhang, Z. Yin, M. Liu, B. Li, H. Zhang, X. Zheng, P. Wang, and Z. Zhang. 2017. MoVrp1, a putative verprolin protein, is required for asexual development and infection in the rice blast fungus *Magnaporthe oryzae*. *Sci. Rep.* 7:41148.
- Jacob, S., A.J. Foster, A. Yemelin, and E. Thines. 2015. High osmolarity glycerol (HOG) signalling in *Magnaporthe oryzae*: Identification of MoYPD1 and its role in osmoregulation, fungicide action, and pathogenicity. *Fungal Biol.* 119:580-594.
- Janke, C., J. Ortiz, T.U. Tanaka, J. Lechner, and E. Schiebel. 2002. Four new subunits of the Dam1-Duo1 complex reveal novel functions in sister kinetochore biorientation. *EMBO J.* 21:181-193.
- Jaqaman, K., J.F. Dorn, E. Marco, P.K. Sorger, and G. Danuser. 2007. Phenotypic clustering of yeast mutants based on kinetochore microtubule dynamics. *Bioinformatics.* 23:1666-1673.
- Jenkinson, C.B., K. Jones, J. Zhu, S. Dorhmi, and C.H. Khang. 2017. The appressorium of the rice blast fungus *Magnaporthe oryzae* remains mitotically active during post-penetration hyphal growth. *PLoS Genet.* 98:35-38.
- Jeon, J., J. Goh, S. Yoo, M.H. Chi, J. Choi, H.S. Rho, J. Park, S.S. Han, B.R. Kim, S.Y. Park, S. Kim, and Y.H. Lee. 2008. A putative MAP kinase kinase, MCK1, is required for cell wall integrity and pathogenicity of the rice blast fungus, *Magnaporthe oryzae*. *Mol. Plant Microbe Interact.* 21:525-534.
- Jeon, J., H. Rho, S. Kim, K.S. Kim, and Y.H. Lee. 2014. Role of MoAND1-mediated nuclear positioning in morphogenesis and pathogenicity in the rice blast fungus, *Magnaporthe oryzae*. *Fungal Genet. Biol.* 69:43-51.
- Jones, K., C.B. Jenkinson, M. Borges Araujo, J. Zhu, R.Y. Kim, D.W. Kim, and C.H. Khang. 2016. Mitotic stopwatch for the blast fungus *Magnaporthe oryzae* during invasion of rice cells. *Fungal Genet. Biol.* 93:46-49.
- Jones, M.H., J.B. Bachant, A.R. Castillo, T.H. Giddings, Jr., and M. Winey. 1999. Yeast Dam1p is required to maintain spindle integrity during mitosis and interacts with the Mps1p kinase. *Mol. Biol. Cell.* 10:2377-2391.
- Jones, M.H., X. He, T.H. Giddings, and M. Winey. 2001. Yeast Dam1p has a role at the kinetochore in assembly of the mitotic spindle. *Proc. Natl. Acad. Sci. U S A.* 98:13675-13680.
- Jorgensen, P., N.P. Edgington, B.L. Schneider, I. Rupes, M. Tyers, and B. Futcher. 2007. The size of the nucleus increases as yeast cells grow. *Mol. Biol. Cell.* 18:3523-3532.

- Kang, J., I.M. Cheeseman, G. Kallstrom, S. Velmurugan, G. Barnes, and C.S. Chan. 2001. Functional cooperation of Dam1, Ipl1, and the inner centromere protein (INCENP)-related protein Sli15 during chromosome segregation. *J. Cell Biol.* 155:763-774.
- Keating, P., N. Rachidi, T.U. Tanaka, and M.J. Stark. 2009. Ipl1-dependent phosphorylation of Dam1 is reduced by tension applied on kinetochores. *J. Cell Sci.* 122:4375-4382.
- Kershaw, M.J., M. Basiewicz, D.M. Soanes, X. Yan, L.S. Ryder, M. Csukai, M. Oses-Ruiz, B. Valent, and N.J. Talbot. 2019. Conidial Morphogenesis and Septin-Mediated Plant Infection Require Smo1, a Ras GTPase-Activating Protein in *Magnaporthe oryzae*. *Genetics.* 211:151-167.
- Khan, I.A., G. Ning, X. Liu, X. Feng, F. Lin, and J. Lu. 2015. Mitochondrial fission protein MoFis1 mediates conidiation and is required for full virulence of the rice blast fungus *Magnaporthe oryzae*. *Microbiol Res.* 178:51-58.
- Kiermaier, E., S. Woehrer, Y. Peng, K. Mechtler, and S. Westermann. 2009. A Dam1-based artificial kinetochore is sufficient to promote chromosome segregation in budding yeast. *Nat. Cell Biol.* 11:1109-1115.
- Kim, S., S.Y. Park, K.S. Kim, H.S. Rho, M.H. Chi, J. Choi, J. Park, S. Kong, J. Park, J. Goh, and Y.H. Lee. 2009. Homeobox transcription factors are required for conidiation and appressorium development in the rice blast fungus *Magnaporthe oryzae*. *PLoS Genet.* 5:e1000757.
- Kozubowski, L., V. Yadav, G. Chatterjee, S. Sridhar, M. Yamaguchi, S. Kawamoto, I. Bose, J. Heitman, and K. Sanyal. 2013. Ordered kinetochore assembly in the human-pathogenic basidiomycetous yeast *Cryptococcus neoformans*. *mBio.* 4:e00614-00613.
- Lai, J., S.K. Ng, F.F. Liu, R.N. Patkar, Y. Lu, J.R. Chan, A. Suresh, N. Naqvi, and G. Jedd. 2010. Marker fusion tagging, a new method for production of chromosomally encoded fusion proteins. *Eukaryot. Cell.* 9:827-830.
- Lampert, F., P. Hornung, and S. Westermann. 2010. The Dam1 complex confers microtubule plus end-tracking activity to the Ndc80 kinetochore complex. *J. Cell Biol.* 189:641-649.
- Lara-Gonzalez, P., F.G. Westhorpe, and S.S. Taylor. 2012. The spindle assembly checkpoint. *Curr. Biol.* 22:R966-980.
- Lau, G.W., and J.E. Hamer. 1998. Acropetal: a genetic locus required for conidiophore architecture and pathogenicity in the rice blast fungus. *Fungal Genet. Biol.* 24:228-239.
- Lee, K., P. Singh, W.C. Chung, J. Ash, T.S. Kim, L. Hang, and S. Park. 2006. Light regulation of asexual development in the rice blast fungus, *Magnaporthe oryzae*. *Fungal Genet. Biol.* 43:694-706.
- Legal, T., J. Zou, A. Sochaj, J. Rappsilber, and J.P. Welburn. 2016. Molecular architecture of the Dam1 complex-microtubule interaction. *Open Biol.* 6: 150237.

- Li, C., S. Cao, C. Zhang, Y. Zhang, Q. Zhang, J.R. Xu, and C. Wang. 2016a. MoCDC14 is important for septation during conidiation and appressorium formation in *Magnaporthe oryzae*. *Mol. Plant Pathol.*
- Li, C., J. Yang, W. Zhou, X.L. Chen, J.G. Huang, Z.H. Cheng, W.S. Zhao, Y. Zhang, and Y.L. Peng. 2014. A spindle pole antigen gene MoSPA2 is important for polar cell growth of vegetative hyphae and conidia, but is dispensable for pathogenicity in *Magnaporthe oryzae*. *Curr. Genet.* 60:255-263.
- Li, J.M., Y. Li, and S.J. Elledge. 2005. Genetic analysis of the kinetochore DASH complex reveals an antagonistic relationship with the ras/protein kinase A pathway and a novel subunit required for Ask1 association. *Mol. Cell Biol.* 25:767-778.
- Li, M., X. Liu, Z. Liu, Y. Sun, M. Liu, X. Wang, H. Zhang, X. Zheng, and Z. Zhang. 2016b. Glycoside Hydrolase MoGls2 Controls Asexual/Sexual Development, Cell Wall Integrity and Infectious Growth in the Rice Blast Fungus. *PLoS One.* 11:e0162243.
- Li, Y., J. Bachant, A.A. Alcasabas, Y. Wang, J. Qin, and S.J. Elledge. 2002. The mitotic spindle is required for loading of the DASH complex onto the kinetochore. *Genes Dev.* 16:183-197.
- Li, Y., Y. Que, Y. Liu, X. Yue, X. Meng, Z. Zhang, and Z. Wang. 2015. The putative Ggamma subunit gene MGG1 is required for conidiation, appressorium formation, mating and pathogenicity in *Magnaporthe oryzae*. *Curr. Genet.* 61:641-651.
- Li, Y., X. Zheng, M. Zhu, M. Chen, S. Zhang, F. He, X. Chen, J. Lv, M. Pei, Y. Zhang, Y. Zhang, W. Wang, J. Zhang, M. Wang, Z. Wang, G. Li, and G. Lu. 2019. MoIVD-Mediated Leucine Catabolism Is Required for Vegetative Growth, Conidiation and Full Virulence of the Rice Blast Fungus *Magnaporthe oryzae*. *Front Microbiol.* 10:444.
- Li, Y., J. Zhu, J. Hu, X. Meng, Q. Zhang, K. Zhu, X. Chen, X. Chen, G. Li, Z. Wang, and G. Lu. 2016c. Functional characterization of electron-transferring flavoprotein and its dehydrogenase required for fungal development and plant infection by the rice blast fungus. *Sci. Rep.* 6:24911.
- Lim, Y.J., K.T. Kim, and Y.H. Lee. 2018. SUMOylation is required for fungal development and pathogenicity in the rice blast fungus *Magnaporthe oryzae*. *Mol. Plant Pathol.* 19:2134-2148.
- Liu, H., F. Liang, F. Jin, and Y. Wang. 2008. The coordination of centromere replication, spindle formation, and kinetochore-microtubule interaction in budding yeast. *PLoS Genet.* 4:e1000262.
- Liu, H., A. Suresh, F.S. Willard, D.P. Siderovski, S. Lu, and N.I. Naqvi. 2007. Rgs1 regulates multiple Galpha subunits in *Magnaporthe* pathogenesis, asexual growth and thigmotropism. *EMBO J.* 26:690-700.
- Liu, W., S. Xie, X. Zhao, X. Chen, W. Zheng, G. Lu, J.R. Xu, and Z. Wang. 2010. A homeobox gene is essential for conidiogenesis of the rice blast fungus *Magnaporthe oryzae*. *Mol. Plant Microbe Interact.* 23:366-375.

- Liu, X.H., G.A. Ning, L.Y. Huang, Y.H. Zhao, B. Dong, J.P. Lu, and F.C. Lin. 2016. Calpains are involved in asexual and sexual development, cell wall integrity and pathogenicity of the rice blast fungus. *Sci. Rep.* 6:31204.
- Lu, Q., J.P. Lu, X.D. Li, X.H. Liu, H. Min, and F.C. Lin. 2008. *Magnaporthe oryzae* MTP1 gene encodes a type III transmembrane protein involved in conidiation and conidial germination. *J Zhejiang Univ Sci. B.* 9:511-519.
- Maiato, H., P.J. Hergert, S. Moutinho-Pereira, Y. Dong, K.J. Vandenbeldt, C.L. Rieder, and B.F. McEwen. 2006. The ultrastructure of the kinetochore and kinetochore fiber in *Drosophila* somatic cells. *Chromosoma.* 115:469-480.
- Matar, K.A.O., X. Chen, D. Chen, W.M. Anjago, J. Norvienyeku, Y. Lin, M. Chen, Z. Wang, D.J. Ebbole, and G.D. Lu. 2017. WD40-repeat protein MoCreC is essential for carbon repression and is involved in conidiation, growth and pathogenicity of *Magnaporthe oryzae*. *Curr. Genet.* 63:685-696.
- Matheis, S., A. Yemelin, D. Scheps, K. Andresen, S. Jacob, E. Thines, and A.J. Foster. 2017. Functions of the *Magnaporthe oryzae* Flb3p and Flb4p transcription factors in the regulation of conidiation. *Microbiol. Res.* 196:106-117.
- McKinley, K.L., and I.M. Cheeseman. 2016. The molecular basis for centromere identity and function. *Nat. Rev. Mol. Cell Biol.* 17:16-29.
- Mehrabi, R., S. Ding, and J.R. Xu. 2008. MADS-box transcription factor mig1 is required for infectious growth in *Magnaporthe grisea*. *Eukaryot. Cell.* 7:791-799.
- Meraldi, P., A.D. McAinsh, E. Rheinbay, and P.K. Sorger. 2006. Phylogenetic and structural analysis of centromeric DNA and kinetochore proteins. *Genome Biol.* 7:R23.
- Miranda, J.J., P. De Wulf, P.K. Sorger, and S.C. Harrison. 2005. The yeast DASH complex forms closed rings on microtubules. *Nat. Struct. Mol. Biol.* 12:138-143.
- Miranda, J.J., D.S. King, and S.C. Harrison. 2007. Protein arms in the kinetochore-microtubule interface of the yeast DASH complex. *Mol. Biol. Cell.* 18:2503-2510.
- Mohanam, V.C., P.M. Chandarana, B.B. Chattoo, R.N. Patkar, and J. Manjrekar. 2017. Fungal Histidine Phosphotransferase Plays a Crucial Role in Photomorphogenesis and Pathogenesis in *Magnaporthe oryzae*. *Front. Chem.* 5.
- Musacchio, A. 2015. The Molecular Biology of Spindle Assembly Checkpoint Signaling Dynamics. *Curr. Biol.* 25:R1002-1018.
- Musacchio, A., and A. Desai. 2017. A Molecular View of Kinetochore Assembly and Function. *Biology (Basel).* 6.
- Neumann, F.R., and P. Nurse. 2007. Nuclear size control in fission yeast. *J. Cell Biol.* 179:593-600.
- Nishimura, M., N. Hayashi, N.S. Jwa, G.W. Lau, J.E. Hamer, and A. Hasebe. 2000. Insertion of the LINE retrotransposon MGL causes a

- conidiophore pattern mutation in *Magnaporthe grisea*. *Mol. Plant Microbe Interact.* 13:892-894.
- Norvienenye, J., Z. Zhong, L. Lin, X. Dang, M. Chen, X. Lin, H. Zhang, W.M. Anjago, L. Lin, W. Abdul, and Z. Wang. 2017. Methylmalonate-semialdehyde dehydrogenase mediated metabolite homeostasis essentially regulate conidiation, polarized germination and pathogenesis in *Magnaporthe oryzae*. *Environ. Microbiol.* 19:4256-4277.
- Nurse, P., P. Thuriaux, and K. Nasmyth. 1976. Genetic control of the cell division cycle in the fission yeast *Schizosaccharomyces pombe*. *Mol. Gen. Genet.* 146:167-178.
- O'Toole, E.T., M. Winey, and J.R. McIntosh. 1999. High-voltage electron tomography of spindle pole bodies and early mitotic spindles in the yeast *Saccharomyces cerevisiae*. *Mol. Biol. Cell.* 10:2017-2031.
- Odenbach, D., B. Breth, E. Thines, R.W. Weber, H. Anke, and A.J. Foster. 2007. The transcription factor Con7p is a central regulator of infection-related morphogenesis in the rice blast fungus *Magnaporthe grisea*. *Mol. Microbiol.* 64:293-307.
- Oses-Ruiz, M., W. Sakulkoo, G.R. Littlejohn, M. Martin-Urdiroz, and N.J. Talbot. 2017. Two independent S-phase checkpoints regulate appressorium-mediated plant infection by the rice blast fungus *Magnaporthe oryzae*. *Proc. Natl. Acad. Sci. U S A.* 114:E237-E244.
- Palmer, D.K., K. O'Day, H.L. Trong, H. Charbonneau, and R.L. Margolis. 1991. Purification of the centromere-specific protein CENP-A and demonstration that it is a distinctive histone. *Proc. Natl. Acad. Sci. U S A.* 88:3734-3738.
- Park, S.Y., J. Choi, S.E. Lim, G.W. Lee, J. Park, Y. Kim, S. Kong, S.R. Kim, H.S. Rho, J. Jeon, M.H. Chi, S. Kim, C.H. Khang, S. Kang, and Y.H. Lee. 2013. Global expression profiling of transcription factor genes provides new insights into pathogenicity and stress responses in the rice blast fungus. *PLoS Pathog.* 9:e1003350.
- Patkar, R.N., A. Suresh, and N.I. Naqvi. 2010. MoTea4-mediated polarized growth is essential for proper asexual development and pathogenesis in *Magnaporthe oryzae*. *Eukaryot. Cell.* 9:1029-1038.
- Peng, H., Y. Feng, X. Zhu, X. Lan, M. Tang, J. Wang, H. Dong, and B. Chen. 2011. MoDUO1, a Duo1-like gene, is required for full virulence of the rice blast fungus *Magnaporthe oryzae*. *Curr. Genet.* 57:409-420.
- Pesenti, M.E., J.R. Weir, and A. Musacchio. 2016. Progress in the structural and functional characterization of kinetochores. *Curr. Opin. Struct. Biol.* 37:152-163.
- Prakash, C., J. Manjrekar, and B.B. Chattoo. 2016. Skp1, a component of E3 ubiquitin ligase, is necessary for growth, sporulation, development and pathogenicity in rice blast fungus (*Magnaporthe oryzae*). *Mol. Plant Pathol.* 17:903-919.
- Qi, Z., M. Liu, Y. Dong, Q. Zhu, L. Li, B. Li, J. Yang, Y. Li, Y. Ru, H. Zhang, X. Zheng, P. Wang, and Z. Zhang. 2016. The syntaxin protein

- (MoSyn8) mediates intracellular trafficking to regulate conidiogenesis and pathogenicity of rice blast fungus. *New Phytol.* 209:1655-1667.
- Ramanujam, R., and N.I. Naqvi. 2010. PdeH, a high-affinity cAMP phosphodiesterase, is a key regulator of asexual and pathogenic differentiation in *Magnaporthe oryzae*. *PLoS Pathog.* 6:e1000897.
- Redemann, S., J. Baumgart, N. Lindow, M. Shelley, E. Nazockdast, A. Kratz, S. Prohaska, J. Bruges, S. Furthauer, and T. Muller-Reichert. 2017. *C. elegans* chromosomes connect to centrosomes by anchoring into the spindle network. *Nat. Commun.* 8:15288.
- Reza, M.H., H. Shah, J. Manjrekar, and B.B. Chattoo. 2016. Magnesium Uptake by CorA Transporters Is Essential for Growth, Development and Infection in the Rice Blast Fungus *Magnaporthe oryzae*. *PLoS One.* 11:e0159244.
- Rieder, C.L., and H. Maiato. 2004. Stuck in division or passing through: what happens when cells cannot satisfy the spindle assembly checkpoint. *Dev. Cell.* 7:637-651.
- Roy, B., L.S. Burrack, M.A. Lone, J. Berman, and K. Sanyal. 2011. CaMtw1, a member of the evolutionarily conserved Mis12 kinetochore protein family, is required for efficient inner kinetochore assembly in the pathogenic yeast *Candida albicans*. *Mol. Microbiol.* 80:14-32.
- Sabnam, N., and S. Roy Barman. 2017. WISH, a novel CFEM GPCR is indispensable for surface sensing, asexual and pathogenic differentiation in rice blast fungus. *Fungal Genet. Biol.* 105:37-51.
- Saitoh, H., A. Hirabuchi, S. Fujisawa, C. Mitsuoka, R. Terauchi, and Y. Takano. 2014. MoST1 encoding a hexose transporter-like protein is involved in both conidiation and mycelial melanization of *Magnaporthe oryzae*. *FEMS Microbiol. Lett.* 352:104-113.
- Sambrook, J., and D.W. Russell. 2001. Molecular Cloning A Laboratory Manual. *Cold Spring Harb. Lab. Press.* Edition III.
- Sanchez-Perez, I., S.J. Renwick, K. Crawley, I. Karig, V. Buck, J.C. Meadows, A. Franco-Sanchez, U. Fleig, T. Toda, and J.B. Millar. 2005. The DASH complex and Klp5/Klp6 kinesin coordinate bipolar chromosome attachment in fission yeast. *EMBO J.* 24:2931-2943.
- Saunders, D.G., S.J. Aves, and N.J. Talbot. 2010a. Cell cycle-mediated regulation of plant infection by the rice blast fungus. *Plant Cell.* 22:497-507.
- Saunders, D.G., Y.F. Dagdas, and N.J. Talbot. 2010b. Spatial uncoupling of mitosis and cytokinesis during appressorium-mediated plant infection by the rice blast fungus *Magnaporthe oryzae*. *Plant Cell.* 22:2417-2428.
- Sazer, S., M. Lynch, and D. Needleman. 2014. Deciphering the evolutionary history of open and closed mitosis. *Curr. Biol.* 24:R1099-1103.
- Schumacher, M.A. 2012. Bacterial plasmid partition machinery: a minimalist approach to survival. *Curr. Opin. Struct. Biol.* 22:72-79.

- Sesma, A., and A.E. Osbourn. 2004. The rice leaf blast pathogen undergoes developmental processes typical of root-infecting fungi. *Nature*. 431:582-586.
- Sullivan, K.F., M. Hechenberger, and K. Masri. 1994. Human CENP-A contains a histone H3 related histone fold domain that is required for targeting to the centromere. *J. Cell Biol.* 127:581-592.
- Talbot, N.J. 2003. On the trail of a cereal killer: Exploring the biology of *Magnaporthe grisea*. *Annu. Rev. Microbiol.* 57:177-202.
- Tanaka, K., E. Kitamura, Y. Kitamura, and T.U. Tanaka. 2007. Molecular mechanisms of microtubule-dependent kinetochore transport toward spindle poles. *J. Cell. Biol.* 178:269-281.
- Thakur, J., and K. Sanyal. 2011. The essentiality of the fungus-specific Dam1 complex is correlated with a one-kinetochore-one-microtubule interaction present throughout the cell cycle, independent of the nature of a centromere. *Eukaryot. Cell.* 10:1295-1305.
- Tien, J.F., N.T. Umbreit, D.R. Gestaut, A.D. Franck, J. Cooper, L. Wordeman, T. Gonen, C.L. Asbury, and T.N. Davis. 2010. Cooperation of the Dam1 and Ndc80 kinetochore complexes enhances microtubule coupling and is regulated by aurora B. *J. Cell Biol.* 189:713-723.
- Tooley, J., and P.T. Stukenberg. 2011. The Ndc80 complex: integrating the kinetochore's many movements. *Chromosome Res.* 19:377-391.
- van Hooff, J.J.E., B. Snel, and G. Kops. 2017. Unique Phylogenetic Distributions of the Ska and Dam1 Complexes Support Functional Analogy and Suggest Multiple Parallel Displacements of Ska by Dam1. *Genome Biol. Evol.* 9:1295-1303.
- Veneault-Fourrey, C., M. Barooah, M. Egan, G. Wakley, and N.J. Talbot. 2006. Autophagic fungal cell death is necessary for infection by the rice blast fungus. *Science*. 312:580-583.
- Villalba, F., J. Collemare, P. Landraud, K. Lambou, V. Brozek, B. Cirer, D. Morin, C. Bruel, R. Beffa, and M.H. Lebrun. 2008. Improved gene targeting in *Magnaporthe grisea* by inactivation of MgKU80 required for non-homologous end joining. *Fungal Genet. Biol.* 45:68-75.
- Waldo, J., and M. Scherrer. 2008. Production and initial characterization of Dad1p, a component of the Dam1-DASH kinetochore complex. *PLoS One*. 3:e3888.
- Wang, Y., D. He, Y. Chu, Y.S. Zuo, X.W. Xu, X.L. Chen, W.S. Zhao, Y. Zhang, J. Yang, and Y.L. Peng. 2016. MoCps1 is important for conidiation, conidial morphology and virulence in *Magnaporthe oryzae*. *Curr Genet.* 62:861-871.
- Westermann, S., A. Avila-Sakar, H.W. Wang, H. Niederstrasser, J. Wong, D.G. Drubin, E. Nogales, and G. Barnes. 2005. Formation of a dynamic kinetochore- microtubule interface through assembly of the Dam1 ring complex. *Mol. Cell.* 17:277-290.
- Westermann, S., H.W. Wang, A. Avila-Sakar, D.G. Drubin, E. Nogales, and G. Barnes. 2006. The Dam1 kinetochore ring complex moves processively on depolymerizing microtubule ends. *Nature*. 440:565-569.

- Wilson, R.A., J.M. Jenkinson, R.P. Gibson, J.A. Littlechild, Z.Y. Wang, and N.J. Talbot. 2007. Tps1 regulates the pentose phosphate pathway, nitrogen metabolism and fungal virulence. *EMBO J.* 26:3673-3685.
- Wilson, R.A., and N.J. Talbot. 2009. Under pressure: investigating the biology of plant infection by *Magnaporthe oryzae*. *Nat. Rev. Microbiol.* 7:185-195.
- Winey, M., C.L. Mamay, E.T. O'Toole, D.N. Mastronarde, T.H. Giddings, Jr., K.L. McDonald, and J.R. McIntosh. 1995. Three-dimensional ultrastructural analysis of the *Saccharomyces cerevisiae* mitotic spindle. *J. Cell Biol.* 129:1601-1615.
- Xu, J.R., C.J. Staiger, and J.E. Hamer. 1998. Inactivation of the mitogen-activated protein kinase Mps1 from the rice blast fungus prevents penetration of host cells but allows activation of plant defense responses. *Proc. Natl. Acad. Sci. U S A.* 95:12713-12718.
- Yan, X., Y. Li, X. Yue, C. Wang, Y. Que, D. Kong, Z. Ma, N.J. Talbot, and Z. Wang. 2011a. Two novel transcriptional regulators are essential for infection-related morphogenesis and pathogenicity of the rice blast fungus *Magnaporthe oryzae*. *PLoS Pathog.* 7:e1002385.
- Yan, X., W.B. Ma, Y. Li, H. Wang, Y.W. Que, Z.H. Ma, N.J. Talbot, and Z.Y. Wang. 2011b. A sterol 14 $\alpha$ -demethylase is required for conidiation, virulence and for mediating sensitivity to sterol demethylation inhibitors by the rice blast fungus *Magnaporthe oryzae*. *Fungal Genet Biol.* 48:144-153.
- Yang, F., and N.I. Naqvi. 2014. Sulfonylurea resistance reconstitution as a novel strategy for ILV2-specific integration in *Magnaporthe oryzae*. *Fungal Genet. Biol.* 68:71-76.
- Yang, J., L. Liu, Y. Wang, C. Wang, J. Yan, Y. Liu, C. Wang, and C. Li. 2017. Overexpression of BAS1 in rice blast fungus can promote blast fungus growth, sporulation and virulence in planta. *Saudi J. Biol. Sci.* 24:1884-1893.
- Yang, J., X. Zhao, J. Sun, Z. Kang, S. Ding, J.R. Xu, and Y.L. Peng. 2010. A novel protein Com1 is required for normal conidium morphology and full virulence in *Magnaporthe oryzae*. *Mol. Plant Microbe Interact.* 23:112-123.
- Yang, L., L. Ukil, A. Osmani, F. Nahm, J. Davies, C.P. De Souza, X. Dou, A. Perez-Balaguer, and S.A. Osmani. 2004. Rapid production of gene replacement constructs and generation of a green fluorescent protein-tagged centromeric marker in *Aspergillus nidulans*. *Eukaryot. Cell.* 3:1359-1362.
- Ye, W., X. Chen, Z. Zhong, M. Chen, L. Shi, H. Zheng, Y. Lin, D. Zhang, G. Lu, G. Li, J. Chen, and Z. Wang. 2014. Putative RhoGAP proteins orchestrate vegetative growth, conidiogenesis and pathogenicity of the rice blast fungus *Magnaporthe oryzae*. *Fungal Genet. Biol.* 67:37-50.
- Yi, M., M.H. Chi, C.H. Khang, S.Y. Park, S. Kang, B. Valent, and Y.H. Lee. 2009. The ER chaperone LHS1 is involved in asexual development

- and rice infection by the blast fungus *Magnaporthe oryzae*. *Plant Cell*. 21:681-695.
- Yi, M., J.H. Park, J.H. Ahn, and Y.H. Lee. 2008. MoSNF1 regulates sporulation and pathogenicity in the rice blast fungus *Magnaporthe oryzae*. *Fungal Genet. Biol.* 45:1172-1181.
- Zhang, Y., H. Shi, S. Liang, G. Ning, N. Xu, J. Lu, X. Liu, and F. Lin. 2015a. MoARG1, MoARG5,6 and MoARG7 involved in arginine biosynthesis are essential for growth, conidiogenesis, sexual reproduction, and pathogenicity in *Magnaporthe oryzae*. *Microbiol. Res.* 180:11-22.
- Zhang, Z., J. Wang, R. Chai, H. Qiu, H. Jiang, X. Mao, Y. Wang, F. Liu, and G. Sun. 2015b. An S-(hydroxymethyl)glutathione dehydrogenase is involved in conidiation and full virulence in the rice blast fungus *Magnaporthe oryzae*. *PLoS One*. 10:e0120627.
- Zhou, W., W. Shi, X.W. Xu, Z.G. Li, C.F. Yin, J.B. Peng, S. Pan, X.L. Chen, W.S. Zhao, Y. Zhang, J. Yang, and Y.L. Peng. 2017. Glutamate synthase MoGlt1-mediated glutamate homeostasis is important for autophagy, virulence and conidiation in the rice blast fungus. *Mol. Plant Pathol.*
- Zhou, X., W. Liu, C. Wang, Q. Xu, Y. Wang, S. Ding, and J.R. Xu. 2011. A MADS-box transcription factor MoMcm1 is required for male fertility, microconidium production and virulence in *Magnaporthe oryzae*. *Mol. Microbiol.* 80:33-53.
- Zhu, X., T. Zhou, L. Chen, S. Zheng, S. Chen, D. Zhang, G. Li, and Z. Wang. 2016. Arf6 controls endocytosis and polarity during asexual development of *Magnaporthe oryzae*. *FEMS Microbiol. Lett.* 363.

