

**Characterisation of genes involved in the development and  
virulence of pathogenic fungi**

Thesis submitted to  
The Maharaja Sayajirao University of Baroda  
For the award of the Degree of  
**DOCTOR OF PHILOSOPHY**

By  
**HIRAL SHAH**

**Department of Microbiology and Biotechnology Centre**  
Faculty of Science  
The Maharaja Sayajirao University of Baroda  
Vadodara – 390 002 Gujarat  
INDIA

**DECEMBER 2019**

## TABLE OF CONTENTS

I. List of tables .....	3
II. List of figures .....	3
III. List of abbreviations .....	5
Abstract.....	7
1. Chapter 1: Review of literature .....	13
1.1 Introduction .....	13
1.1.1 The Cell Cycle and Mitosis .....	13
1.1.2 The Kinetochore .....	17
1.2 The Dam1 Complex (also known as the DASH Complex) in yeasts .....	20
1.2.1 DASH Complex proteins in association with spindle and kinetochore .....	22
1.2.2 Structure of the DASH complex.....	26
1.2.3 Regulation of DASH complex proteins.....	29
1.2.4 The DASH complex, MT Network and MAPs .....	31
1.3 The cell cycle in fungi .....	33
1.3.1 Development in <i>M. oryzae</i> and its regulation by the cell cycle.....	35
1.4 Rice Blast Disease .....	38
2. Chapter 2: PRESENT STUDY- Characterisation of genes involved in the development and virulence of pathogenic fungi- Introduction, Objectives and Materials and Methods .....	45
2.1. INTRODUCTION .....	45
2.2. OBJECTIVES.....	48
2.3. MATERIALS AND METHODS .....	48
2.3.1. Fungal Strains and culture conditions .....	48
2.3.2. Bacterial strains and culture conditions.....	49
2.3.3. Bioinformatics analysis of DASH Complex proteins.....	50
2.3.4. Nucleic acid manipulation.....	50
2.3.5. Bacterial and fungal transformation .....	53

Protoplast transformation of <i>M. oryzae</i> .....	54
2.3.6. Pathogenicity assays .....	57
2.3.7. Microscopy .....	58
2.3.7.1 Conidiation .....	59
2.3.7.2 Hyphal growth .....	60
3. Chapter 3: Generation of strains for study of kinetochore genes ..	65
3.1. Analysis of DASH complex proteins in <i>M. oryzae</i> .....	65
3.1.1 Homology based identification of Dam1 complex proteins using <i>Magnaporthe oryzae</i> genome sequence .....	65
3.2. Development of strains for study of kinetochore proteins in <i>M.</i> <i>oryzae</i> .....	73
3.2.1 Tagging of genes for localisation studies .....	73
3.2.2 Generation of mutant strains.....	87
4. Chapter 4: The role of kinetochore proteins in the development and virulence of <i>Magnaporthe oryzae</i> .....	99
4.1 Mis12 is associated with the nucleus during interphase.....	101
4.2 Outer kinetochore DASH complex proteins Dam1 and Ask1 associate with the nucleus during mitosis.....	106
4.3 Dam1 is crucial to proper spindle structure and segregation of chromosomes .....	110
4.4 Polarised growth is dependent on proper DASH complex function	116
4.5.1 Dam1 function is crucial for proper conidiation .....	128
4.5.1 Conidium development displays a convergence of distinct cell division patterns.....	138
4.6 DASH complex protein Dam1 is required for pathogenic development and virulence.....	143
5. Chapter 5: Summary and Conclusion.....	151
6. Bibliography .....	159
7. Publications, oral and poster presentations.....	173

## ABSTRACT

Chromosome segregation is a critical process in eukaryotic cell development, which is brought about by the multi-protein kinetochore complex along spindle microtubules (MTs). The outer kinetochore DASH complex proteins maintain spindle structure and chromosome-MT association to bring about high-fidelity chromosome segregation during anaphase. While the role and sub-cellular localisation of DASH complex proteins varies among yeasts, their role in filamentous fungi had not been investigated in detail thus far. Pathogenic filamentous fungi cycle through diverse developmental stages associated with changes in cell polarity. Pathogenic development in the rice blast fungus *Magnaporthe oryzae* requires an isotropic distension of the polarised germ tube to form the appressorium (infection structure). These developmental transitions are precisely regulated progression of nuclear segregation and migration. Vegetative growth, on the other hand, proceeds by hyphal extension and lateral branching. Here, I studied the dynamics of GFP-tagged middle (Mis12) and outer (Dam1 and Ask1) kinetochore proteins and their role in *M. oryzae* vegetative and pathogenic development. The Mis12 foci constitutively associated with the nucleus throughout the cell cycle, clustered into a single patch during interphase, and de-clustered during mitosis; however, Dam1 and Ask1 showed a mitosis-specific nuclear association. Absence of Dam1 function, (*dam1* $\Delta$ ) led to atypical mitotic progression with sluggish nuclear movement and difficulty in segregation. Dam1 was involved in the three consecutive rounds of mitosis required for development of the three celled conidia such that loss of Dam1 led to defects in conidial development and impaired appressorium formation. The *dam1* $\Delta$  mutant also showed reduced hyphal growth and morphological defects in vegetative hyphae, indicating altered cell polarity during filamentous growth in *M. oryzae*. The *ask1* $\Delta$  strain showed all the defects seen in the *dam1* $\Delta$  mutant. Interestingly, both Dam1 and Ask1

localised to the hyphal tip during vegetative growth and to the germ tube tip during pathogenic development, and oscillated back and forth from the tip. Thus, the findings indicate that *Magnaporthe* Dam1 and Ask1, in addition to their function at mitotic spindle, likely perform a broader role in regulating microtubule dynamics and/or maintenance of cell polarity even during interphase. Aberrant appressorial (infection structure) development and impaired host penetration by the *dam1* $\Delta$  mutant suggest that the fungus-specific DASH complex proteins Dam1 and Ask1 could constitute attractive targets for development of novel antifungal strategies.

## **2. Chapter 2: PRESENT STUDY- Characterisation of genes involved in the development and virulence of pathogenic fungi- Introduction, Objectives and Materials and Methods**

### **2.1. INTRODUCTION**

Fungi form an extremely diverse group of eukaryotic organisms, including unicellular yeasts, dimorphic yeasts that can undergo hyphal transition and multi-cellular filamentous fungi. Fungi are suited to lifestyles in a wide range of niches that call for efficient adaptation in the acquisition of nutrients, response to environmental triggers and stresses and maintenance of basal homeostasis and growth. Excellent examples of such adaptability are found in the class of fungal pathogens that have managed to infect plant, animal and human hosts, where successful colonisation is often achieved through the development of several distinct morphological stages with specialised function. Development of resistant conidia, elongation of germ tube, specialised infection structures that bring about chemical and physical disruption of host tissue, hyphal extension and branching, and substratum attachment are all important for dispersal and virulence in filamentous fungi. The rice blast fungus *Magnaporthe oryzae* is one such filamentous fungus which passes through multiple developmental stages during its infection cycle. Pathogenic development starts when the conidium germinates on the host surface and extends a polarised germ tube that swells at the tip to form the dome-shaped appressorium. The appressorium then extends a penetration peg which ruptures the leaf surface to enter the host and establish infection. These developmental transitions are regulated by the cell cycle. Though a good deal is known about mitosis in *M. oryzae*, as far as kinetochore proteins and the process of chromosome segregation are concerned, little work has been done. *M. oryzae* infects rice and several other cereal crops in India and across the world, negatively impacting global food security. The millions of hectares

of crops lost every year to rice blast necessitate a better understanding of the disease. A detailed investigation of the role of fungus-specific components in development and virulence will suggest ways for overcoming this devastating pathogen without adversely affecting the crop. The microtubule (MT)-associated Dam1/DASH complex proteins are fungus-specific outer kinetochore components whose role has remained relatively unknown in filamentous fungi. Further components of the Dam1 complex are prospective antifungal targets due to their absence from plant genomes. MoDuo1 is the only gene investigated hence far within the group of filamentous fungi, albeit with very little scrutiny of its function at the kinetochore during mitosis or subcellular dynamics through the cell cycle.

The DAM/DASH complex is important for spindle structure and dynamics in yeasts. However, features of the DASH complex vary in *S. cerevisiae*, *S. pombe* and *C. albicans* in sub-cellular localisation, cell-cycle dependent nuclear association, oligomeric states and requirement for survival. As discussed earlier, it has been suggested that the requirement of the DASH complex depends on the number of MTs bound to each kinetochore (Burrack *et al.*, 2011, Thakur and Sanyal, 2011). Whether this is true can only be answered when studies are extended to other fungi beyond the few model yeasts. Therefore, ‘Is *DAM1* essential in filamentous fungi?’ is still an open question in fungal biology. In the filamentous fungus *N. crassa*, not all DASH complex members may be essential (Freitag, 2016) suggesting an outer KT system in filamentous fungi, different from what has been observed in the various yeasts. A better understanding of the function of Dam1 in filamentous fungi can be obtained by studying its role in poleward movement during anaphase, nuclear transport and microtubule organisation. Understanding Dam1 localisation patterns during the cell cycle with respect to chromosomes and other KT proteins will provide information about kinetochore assembly in fungi. Yeasts

show key differences in KT clustering through the cell cycle as well as in assembly of sub-complexes. In *C. neoformans*, there is hierarchical KT assembly of inner followed by middle and subsequently outer layer, and perturbing the inner KT leads to mis-localisation of the outer proteins (Kozubowski *et al.*, 2013). Interestingly, in *C. albicans* perturbing the outer KT proteins also destabilises the inner layer (Roy *et al.*, 2011). Localisation studies of proteins of different layers of the KT in filamentous fungi will provide information about the interaction and assembly of different layers. Co-localisation studies with other DASH complex members will address whether Dam1 localisation is dependent on other subunits, provide information about DASH complex interactions and functions unique to certain proteins. Further, the localisation with MTs or MT perturbation studies will provide information about the dependence of DASH complex function on MT dynamics. Though early studies have termed the DASH proteins as a kinetochore complex, recent studies are beginning to show a broader spectrum of microtubule association, opening up the possibility that, more than just a kinetochore complex, the DASH complex proteins are likely regulators of microtubule dynamics, while also playing a role at the kinetochore. However, the studies are few and will require more support from studies in diverse organisms. Whether Dam1 localisation is restricted to the mitotic spindle MTs, or extends to the interphase and cytoplasmic MTs, will shed further light on its role in filamentous fungal development. I addressed some of these questions about KT proteins and their roles, in the filamentous fungus *M. oryzae* in the present study.

Here, I particularly investigated the function of components of the DASH complex in chromosome segregation during the developmental switches in *M. oryzae*. Based on protein sequence homology, I was able to identify all 10 members of the DASH complex in *M. oryzae*. Using gene deletion and GFP-tagged strains, I investigated the role of DASH complex proteins in mitosis, particularly spindle structure and dynamics, nuclear migration and

its effect on the development of *M. oryzae*. The role of middle KT protein Mis12 and outer KT proteins Dam1 and Ask1 was addressed in *M. oryzae* development and virulence. I looked at the localisation pattern of the proteins during different developmental stages of the filamentous fungus *M. oryzae*, specifically during mitosis. In order to do so, multiple tagging constructs and tagged fungal strains were developed. Further, I generated mutants of *DAMI* and *ASK1* and studied their effects on fungal development, pathogenicity and nuclear division. I suggest a role for Dam1 in KT-MT attachment, proper poleward movement of chromosomes during anaphase and in nuclear transport. I show that although Dam1 and Ask1 are not essential for viability, they are involved in proper conidiation and polarised hyphal growth and pathogenic development. This work shows that beyond its role in proper segregation of chromosomes during mitosis, DASH complex proteins Dam1 and Ask1 are involved in polarised growth likely through association and regulation of cytoplasmic MT dynamics at the hyphal tip.

## **2.2. OBJECTIVES**

### **A. Characterisation of kinetochore genes in *M. oryzae***

1. Study of localisation of kinetochore proteins
2. Generation of mutants of selected kinetochore genes
3. Phenotypic characterisation of mutants

## **2.3. MATERIALS AND METHODS**

### **2.3.1. Fungal Strains and culture conditions**

This study was carried out in a rice infection isolate of *Magnaporthe oryzae* from our laboratory (Kachroo et. al., 1994), namely, strain B157 (MTCC accession number 12236, international race IC9). The fungus was

maintained on prune agar (PA). Fungal biomass was cultured in complete medium broth (Dawe et al., 2003) for isolation of nucleic acids, etc. Colony diameter was used as readout of vegetative hyphal growth. Dry weight was determined for fungal biomass grown in liquid CM. For collection of conidia, fungal cultures were maintained on PA for 3 days in the dark and subjected to constant photo-induction till 9-10 days post inoculation when the conidia were collected.

### **Media composition (1L) for fungal culture**

**Prune Agar-** 40 ml prune juice, 1 g yeast extract, 2.5 g lactose, 2.5 g sucrose, 20 g agar. pH was set to 6.5 with 10M NaOH

**YEG Agar-** 2 g yeast extract, 10 g dextrose, 20 g agar

**Complete medium for *Magnaporthe oryzae*-** 10g Dextrose, 5 g Peptone, 1 g Yeast extract, 1 g CAA, 6 g NaNO<sub>3</sub>, 0.5 g KCl, 0.5 g MgSO<sub>4</sub>, 1.5 g KH<sub>2</sub>PO<sub>4</sub>, pH 6.5

**Basal Medium-** 1.7g Yeast Nitrogen Base without amino acids or ammonium sulphate, 2g asparagines, 1g NH<sub>4</sub>NO<sub>3</sub>, 10g Glucose. pH 6.0 was set with 1M Na<sub>2</sub>HPO<sub>4</sub>.

### **2.3.2. Bacterial strains and culture conditions**

#### ***Escherichia coli (E. coli)***

*E. coli* strain DH5 $\alpha$  (F<sup>-</sup>, *endA1*, *hsdR17* (rk<sup>-</sup>, mk<sup>+</sup>), *supE44*, *thi-1*,  $\lambda$ , *recA1*, *gyrA96*, *relA1* $\Delta$  (*lacZYA-argF*) U169 *deoR* ( $\phi$ 80*dlac* $\Delta$  (*lacZ*) M15; Bethesda Research Laboratories) was used for bacterial transformation and plasmid propagation. The *E. coli* strain was grown at 37<sup>0</sup>C on Luria Bertani (LB) agar medium. *E. coli* transformants carrying plasmid vectors

were grown on LB agar containing either ampicillin or kanamycin at a concentration of 100  $\mu\text{g ml}^{-1}$  or 50  $\mu\text{g ml}^{-1}$ , respectively, as required. Long term preservation was done in 25% glycerol solution at  $-80^{\circ}\text{C}$ .

#### ***Agrobacterium tumefaciens* (*A. tumefaciens*)**

*A. tumefaciens* strain LBA4404/pSB1 was used for *Agrobacterium tumefaciens* mediated transformation (ATMT). The culture was grown at 28/30 $^{\circ}\text{C}$  on LB agar medium. Plasmids were transformed into *Agrobacterium* by triparental mating or electroporation. Transformants carrying plasmid vectors were grown on LB agar containing 50  $\mu\text{g ml}^{-1}$  kanamycin. Long term preservation was done in 25% glycerol solution at  $-80^{\circ}\text{C}$ .

#### **2.3.3. Bioinformatics analysis of DASH Complex proteins**

Middle (Mis12) and outer (DASH complex) kinetochore proteins were identified from the *M. oryzae* 70-15 genome sequence, previously provided by the Broad Institute (Cambridge, USA) ([http://www.broadinstitute.org/annotation/genome/magnaporthe\\_grisea/MultiHome.html](http://www.broadinstitute.org/annotation/genome/magnaporthe_grisea/MultiHome.html)), by a BLAST\_P search using the full length *S. cerevisiae* and *S. pombe* protein sequence. Fungal Dam1 and Ask1 protein sequences were obtained from Uniprot database. The Multiple sequence alignment and phylogenetic analysis was done using COBALT and CLUSTAL OMEGA. The conserved domain prediction was done using Pfam tool. NCOILS was used to determine the coiled coil domains of DASH complex proteins. WoLFPSORT tool was used for predicting localisation of the proteins.

#### **2.3.4. Nucleic acid manipulation**

##### **Genomic DNA extraction from *M. oryzae***

*M. oryzae* B157 was grown in 40 ml complete medium for 2.5 days, harvested on filter paper discs, dried and frozen in liquid nitrogen and

ground to a fine powder. The powdered biomass was transferred to 700  $\mu$ l DNA extraction buffer (pre-heated at 68<sup>0</sup>C) and mixed properly. Next, 70  $\mu$ l of 20% SDS was added and the contents were mixed properly. The samples were then kept at 68<sup>0</sup>C for 10 min. This was followed by addition of 350  $\mu$ l of 5M potassium acetate and incubation on ice for 30 min. The cell debris was separated by centrifugation at 8,000 rpm for 20 min. The supernatant was transferred to a fresh centrifuge tube and 500  $\mu$ l of ice cold 100% isopropyl alcohol was added. The solutions were mixed gently and the DNA was precipitated at -20<sup>0</sup>C for 30 min. The DNA was collected by centrifugation (10000 rpm, 15 min). The DNA pellet was allowed to air dry at 37<sup>0</sup>C and resuspended in 500  $\mu$ l of TE buffer at 68<sup>0</sup>C for 10 min. 20  $\mu$ l of 1 mg ml<sup>-1</sup> RNase was added to this solution and further incubated at 68<sup>0</sup>C for 30 min. This was mixed with an equal volume of C:I (chloroform:Isoamyl alcohol - 24:1) and centrifuged at 10,000 rpm for 15 min. The upper aqueous phase was collected and to this was added 1/10<sup>th</sup> volume of 3M sodium acetate pH 5.0 and 0.6 volume of ice cold 100% isopropanol and incubated at -20<sup>0</sup>C for 1-2 hours. The DNA pellet was recovered by centrifugation and washed once with 70% ethanol. The pellet was then air dried at 37<sup>0</sup>C and dissolved in autoclaved milliQ water. The DNA quality of the DNA prep was assessed by agarose gel electrophoresis and concentration was determined using nanodrop spectrophotometer.

**DNA extraction buffer for *M. oryzae*** (150 ml)- 15 ml 1M Tris HCl pH 8.0, 15 ml 0.5M EDTA pH 8.0, 18.75 ml 4M NaCl. After autoclaving 105  $\mu$ l  $\beta$ -mercaptoethanol was added.

### **Isolation of plasmid DNA**

Overnight grown culture (1.5 ml) was centrifuged in a microfuge tube and the bacterial pellet was resuspended in 300  $\mu$ l of STET buffer. To this suspension, 10  $\mu$ l of 50 mg ml<sup>-1</sup> lysozyme was added, mixed by vortexing

and incubated for 2 min at room temperature (RT), followed by boiling in a water bath for 45 sec. This was followed by centrifugation at 13,000 rpm for 10 min. The resulting spot was removed with a tooth pick and 20 µl of 5% CTAB was added followed by centrifugation for 10 min. To the pellet, 300 µl of 1.2M NaCl was added and vortexed. To this 180 µl of 100% isopropanol was added. The DNA pellet was obtained by centrifugation for 10 min; washed with 70% ethanol and resuspended in 50 µl of TE.

**STET Buffer** (100 ml)- 8% Sucrose, 50mM Tris HCl pH 8, 50mM EDTA pH 8.0. After autoclaving 0.1% TritonX-100 was added.

### **Restriction Digestion**

Restriction enzyme digestion of DNA was carried out in reaction buffers, supplied with the corresponding enzymes, as recommended by the manufacturer (New England Biolabs, USA). Ligations were carried out at 22<sup>0</sup>C, as recommended (Thermo Fisher, USA).

### **Southern Hybridization**

For Southern blot analysis, 30 µg of genomic DNA was digested with restriction enzymes and the digested products were run on 0.8% agarose gel at 3V/cm. After denaturation and neutralisation of the digested products in the gel, capillary blot transfer was done for 16-20 hours to transfer the DNA fragments onto a nylon membrane (Hybond N+, Amersham, UK). The transferred DNA on the membrane was crosslinked by UV crosslinker (Spectrolinker, XL-1000, USA) at preset optimum cross-linking programme. The labelling of the probes was done according to the manufacturer's instructions (Amersham, Buckinghamshire, UK). The probe was hybridised with DNA on the membrane for 16 hours. The membrane was washed and substrate was added for 5 min. X-ray film was exposed for 4-16 hours with the membrane in Hypercassette (Amersham,

Buckinghamshire, UK). X-ray film was developed and targeted integration in knockout and tagging was confirmed.

### **2.3.5. Bacterial and fungal transformation**

#### **Competent cell preparation and transformation of *E. coli***

A single colony of *E. coli* DH5 $\alpha$  was inoculated in 3 ml of LB broth and grown overnight. 100 ml of fresh LB broth was inoculated with 1.0 ml of overnight grown culture and grown at 37 $^{\circ}$ C till 0.3 to 0.4 O.D <sub>$\lambda=600$</sub> . The cells were collected by centrifugation at 5,000 rpm for 4 min at 4 $^{\circ}$ C and resuspended in 10 ml chilled 0.1M CaCl<sub>2</sub>. The cell suspension was incubated on ice for 30 min. The cells were centrifuged at 4000 rpm for 5 min at 4 $^{\circ}$ C. The supernatant was discarded and the cells were re-suspended in 1.0 ml of ice cold 0.1M CaCl<sub>2</sub> + 20% glycerol. The culture was stored at -80 $^{\circ}$ C in 100  $\mu$ l aliquots. For transformation, a 100  $\mu$ l aliquot of competent cell suspension was mixed with the plasmid DNA (50-100 ng) or ligation mixture and incubated at 4 $^{\circ}$ C for 30 minutes. Cells were subjected to heat shock at 42 $^{\circ}$ C for 90 sec, followed by 5 min incubation on ice. 1 ml of LB broth was added to the above suspension and incubated at 37 $^{\circ}$ C with 200rpm shaking for 45 minutes. 100  $\mu$ l of transformation mixture was spread on LB agar medium containing either Ampicillin (100  $\mu$ g ml $^{-1}$ ) or kanamycin (50  $\mu$ g ml $^{-1}$ ).

*Agrobacterium tumefaciens* mediated transformation (ATMT) (Mullins et al., 2001) or protoplast transformation was used for fungal transformation of tagging and gene deletion constructs. The transformants were selected on YEGA or basal medium supplemented with the appropriate antibiotic at concentrations listed below. The antibiotic resistant transformants were screened by locus specific PCR and/or microscopy. Single targeted integration was confirmed by Southern hybridisation. Virulence of transformants was assessed by barley leaf spot inoculation assay. The transformants were stored as filter stocks at -20 $^{\circ}$ C.

### **Protoplast transformation of *M. oryzae***

*M. oryzae* B157 was inoculated in 30 ml of complete medium broth and grown for 2.5 days. The biomass was then filtered with Miracloth (Calbiochem, Darmstadt, Germany), washed with sterile water and the mycelia were resuspended in 30 ml of 1M sorbitol containing 30 mg (1mg ml<sup>-1</sup>) lysing enzyme (Sigma, St. Louis, USA). The culture was incubated at 28<sup>0</sup>C at 100 rpm for 12-16 hours for protoplasting. Next day, the protoplasts were filtered through miracloth, washed and resuspended in 10 ml 1M sorbitol at 4,000 rpm for 5 min at 4<sup>0</sup>C twice. The protoplasts were resuspended in 10 ml STC buffer, washed and resuspended again in 1 ml STC buffer. The protoplasts were counted using a haemocytometer (Marienfeld Superior, Lauda-Konigshofen, Germany) and a density of 10<sup>8</sup> cells per ml was maintained. For every transformation 200 µl of protoplasts was mixed with 2-10 µg of DNA and incubated on ice for 15 min. 1 ml of PTC buffer was added to the protoplast-DNA mixture and incubated at 28<sup>0</sup>C for 30 min. The whole mixture was transferred to sugar tube containing 3 ml of YEGS and was incubated at 28<sup>0</sup>C at 100 rpm for 12-14 hours. After incubation, 10 ml of molten regeneration medium was added, mixed well and poured on YEG agar containing appropriate antibiotics (Hygromycin/ Zeocin/ Geneticin). Appropriate selection antibiotic was also added to the molten medium. Appearance of colonies was monitored within 2-3 days of incubation at 28<sup>0</sup>C. *M. oryzae* B157 protoplasts (not treated with DNA) were kept as a control which did not grow on selection medium. The colonies were transferred to selection medium, for secondary selection to eliminate any stray growth. The resistant transformants were then screened by PCR, microscopy and Southern hybridisation.

### **Media and Reagents for Protoplast transformation**

**STC Buffer** (100 mL)- 18.2 g D-Sorbitol (1M), 5 ml of 1M Tris HCl pH 8.0 (50mM Tris HCl pH 8.0), 0.735 g CaCl<sub>2</sub>.2H<sub>2</sub>O (50mM)

**PTC Buffer** (100 mL)- 40 g Polyethylene glycol 3350 (40% w/v), 18.2 g D-Sorbitol (1M), 5 ml of 1M Tris HCl pH 8.0 (50mM Tris HCl pH 8.0), 0.735 g CaCl<sub>2</sub>.2H<sub>2</sub>O (50mM)

**YEGS Solution** (100mL)- 0.2g Yeast extract, 1g Dextrose, 18.2 g D-Sorbitol

**Molten Regeneration Medium** (100mL)- 0.2g Yeast extract, 1g Dextrose, 0.4 g agarose (0.4% w/v)

### Antibiotics

Antibiotic	Stock Concentration (mg ml <sup>-1</sup> )	Working Concentration (µg ml <sup>-1</sup> )
<b>Bacterial Selection</b>		
Ampicillin	100	100
Kanamycin	100	50
Rifampicin	5	5
Tetracycline	10	10
Cefotaxime	250	250
<b>Fungal Selection</b>		
Hygromycin B	100	300
Zeocin	100	300
Chlorimuron Ethyl	100 in DMF	100
Glufosinate Ammonium	50	50

### *Agrobacterium tumefaciens* mediated transformation (ATMT) of *M. oryzae*

The *Agrobacterium* strain LBA4404/pSB1 (Komari *et al.*, 1996) was first transformed with respective vectors via triparental mating (Helper plasmid pRK2013; Helper strain *E. coli* DH5α) or electroporation. For triparental mating, a single colony each from the freshly grown recipient *Agrobacterium* LBA4404/pSB1, helper plasmid pRK2013 and the donor strain carrying the respective vector were mixed together on YEP agar (0.2% yeast extract, 0.5% peptone, 2% agar) and incubated at 28<sup>0</sup>C for 18-

20 hours. The mixture was scraped and resuspended in 0.9% NaCl. 100  $\mu$ l of serial dilutions of this bacterial suspension was plated onto selection plates and incubated at 28<sup>0</sup>C for 2-3 days till the appearance of colonies. The *Agrobacterium* harbouring T-DNA vector was then used to transform *M. oryzae*. *A. tumefaciens* mediated transformation (ATMT) was performed as described previously (Mullins *et al.*, 2001). Briefly, *M. oryzae* spores were collected from 9-10 days old PA plates and spore concentration was adjusted to 1x10<sup>6</sup> spores/ml. *Agrobacterium* carrying the *HPT/BAR* gene expression cassette was grown at 28/30<sup>0</sup>C overnight in LB supplemented with appropriate antibiotics. The *A. tumefaciens* cells were diluted to OD<sub>600</sub> = 0.15 in induction medium (IM) and grown for another 6 hours, both in the presence (IM+ACS) and absence (IM-ACS) of 200 $\mu$ M acetosyringone (ACS) before mixing them with an equal volume of a conidial suspension from *M. oryzae*. This mix (200  $\mu$ l per plate) was spotted on a 0.45  $\mu$ m pore size, 45 mm diameter nitrocellulose filter (Millipore, Bangalore, India) placed on co-cultivation medium in presence of 200  $\mu$ M ACS. Following incubation at 28<sup>0</sup>C for 2 days, the filter was transferred to YEG containing Hygromycin B (Invitrogen Life Technologies, CA) to a final concentration of 300  $\mu$ g ml<sup>-1</sup> (or any other appropriate antibiotic) as a selection agent for transformants and 250  $\mu$ g ml<sup>-1</sup> cefotaxime to eliminate the *A. tumefaciens* cells. Untransformed *M. oryzae* was kept as a control which did not grow on hygromycin containing medium. The transformants were maintained as mono-conidial isolates.

### **Media and Reagents for ATMT**

#### **Induction medium (IM) for *Agrobacterium***

10 ml K salts, 20 ml M salts, 2.5 ml 20% NH<sub>4</sub>NO<sub>3</sub>, 0.5 ml 1% CaCl<sub>2</sub>, 10 mM Glucose, 40mM MES, 0.5% Glycerol (v/v), 10 ml 0.01% FeSO<sub>4</sub>, 5 ml Z salts.

0.01 % FeSO<sub>4</sub> and Z-salts solution were prepared separately and autoclaved; they were added after autoclaving the medium, just before use. After autoclaving they were stored at 4°C.

1.8 g or 1.98 g glucose and 7.8 g of MES (2-(N-Morpholino) ethanesulfonic acid) were added to give a final concentration of 10 and 40 mM respectively. 200µM acetosyringone (ACS) was added just before pouring.

#### **Composition for salt solutions for IM**

K Salts Solution : K<sub>2</sub>HPO<sub>4</sub> 20.5%, KH<sub>2</sub>PO<sub>4</sub> 14.5%

M Salts Solution : MgSO<sub>4</sub>.7H<sub>2</sub>O 3%, NaCl 1.5% , (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub> 2.5%

Z Salts Solution : ZnSO<sub>4</sub>.7H<sub>2</sub>O 0.01% , CuSO<sub>4</sub>. 5H<sub>2</sub>O 0.01% , ZnSO<sub>4</sub>. H<sub>2</sub>O 0.01%, H<sub>3</sub>BO<sub>3</sub> 0.01%, Na<sub>2</sub>MbO<sub>4</sub>.2H<sub>2</sub>O 0.01%

#### **Co-cultivation medium for *Agrobacterium* (1L)**

10 ml K salts, 20 ml M salts, 2.5 ml 20% NH<sub>4</sub>NO<sub>3</sub>, 0.5 ml 1% CaCl<sub>2</sub>, 0.5 ml 10mM Glucose, 40mM MES, 0.05% Glycerol (v/v), 2% agar.

1.g or 1.98 g glucose and 7.8 g of MES were added to give a final concentration of 10mM and 40mM respectively. 200µM ACS was added just before pouring.

**1X Phosphate Buffer Saline (1L PBS)**- 8.0 g NaCl (137mM), 2.68 g Na<sub>2</sub>HPO<sub>4</sub>.7H<sub>2</sub>O (100mM), 0.2 g KCl (2.7mM), 0.27 g KH<sub>2</sub>PO<sub>4</sub> (2mM), pH 7.4

#### **2.3.6. Pathogenicity assays**

For assessment of appressorium forming ability, freshly prepared conidial suspension ( $5 \times 10^4$  conidia ml<sup>-1</sup> in sterile water with Streptomycin) obtained from 10-day-old prune agar cultures were applied in aliquots of 20 µl on hydrophobic cover glass. The coverglasses were incubated at room temperature under high humidity for 24 h post inoculation. Conidial germination and appressorium formation were examined by microscopy.

The percentage of conidia that germinated and formed appressoria was calculated.

To determine penetration ability of the fungus, rice leaf sheath inoculation was performed with conidial suspensions (Kankanala et al., 2007) and penetration events and spread of invasive hyphae were assessed 36-40 hpi by microscopy, using a 100X oil immersion lens. For assessment of virulence, detached leaf assay was performed with 8-12 days old barley leaves. The leaves were inoculated with three drops of 10µl each of conidial suspension in 0.2% gelatin and maintained on water agar supplemented with 2 mg l<sup>-1</sup> kinetin. The disease outcome was recorded 5 days post inoculation.

For *in-planta* infection assays, susceptible, three-four weeks old (3<sup>rd</sup> – 4<sup>th</sup> leaf stage) rice seedlings of CO-39 cultivar were sprayed with 10<sup>5</sup> ml<sup>-1</sup> of WT and mutant conidia in 0.2% gelatin and disease symptoms were recorded 5 dpi.

### **2.3.7. Microscopy**

Brightfield microscopy and widefield fluorescence microscopy were carried out on the Olympus BX51 (Olympus, Japan) with the 40x ELWD or 100x/1.40 oil immersion objective. The long working distance of the 40X ELWD objective lens allowed observation of conidiophores as well as cultures grown on agar medium in their native conditions. This was also useful for live cell imaging. The suitable filter set was selected based on the fluorescent protein or stain being imaged. The Nikon Eclipse80i (Nikon, Japan) microscope with 20X or 40X objective lens was also used for brightfield microscopy of fungal structures like appressoria and hyphae. The images were captured using a DS-Qi2 camera operated by the Nikon Elements software. Laser scanning confocal microscopy was used to study sub-cellular localisation of tagged proteins (Model: LSM700, Carl

Zeiss Inc., USA) using either the 40X/1.30 EC Plan-Neofluar or a 63X/1.40 Plan-Apochromat oil immersion objective lenses. The microscope was operated using the ZEN 2010 software. Fungal structures were determined by differential interference contrast (DIC) microscopy. GFP and mCherry tagged proteins were imaged in the 488 nm and 555 nm laser channels respectively. For live-cell imaging of fungal structures, cultures were grown on glass-bottom petridishes and viewed with an inverted confocal microscope. To span the entire thickness of fungal structures, images were captured as z-stacks taken at regular intervals ranging from 0.3 - 1µm. To study protein dynamics, time series programs were used with intervals ranging from 5 seconds to 10 minutes, based on the speed of the process being study. Zeiss AxioCam MR Camera was used for acquisition of images. ImageJ tool was used for analysis of images (<https://imagej.nih.gov/ij/download.html>). Adobe Photoshop CS6 was used for preparation of figures.

#### **2.3.7.1 Conidiation**

Conidia were harvested as described previously (Patkar et al., 2012). Briefly, fungal conidia were harvested by scraping the biomass grown on prune agar plates with a sterile loop, resuspending in sterile water and purifying by passing through two layers of Miracloth (Calbiochem, Darmstadt, Germany). Quantification of conidia was done using a haemocytometer (Marienfeld Superior, Lauda-Konigshofen, Germany). Morphology and cell number of the WT, *dam1Δ* and *ask1Δ* conidia was determined by staining with CFW. Cell wall and septa (chitin-based structures) of conidia and vegetative hyphae were stained using 3 µg ml<sup>-1</sup> Calcofluor White (Whitener 28, Sigma-Aldrich, USA). Arrangement of conidia on conidiophores was assessed in WT, *dam1Δ* and *ask1Δ* strains. A thin mycelial plug was inoculated on 1% agarose and incubated under moist conditions. The conidiophore arrangement was monitored using brightfield microscopy 1 or 2 days post inoculation. Number of conidia

per conidiophore was counted for 100 conidiophores. Bar graph showing the frequency of each cluster size (1 to 8 conidia) was plotted to depict the distribution for a given strain.

### 2.3.7.2 Hyphal growth

Hyphal morphology of *M. oryzae* was studied by adapting methods developed for analysis of hyphal growth and branching patterns in *N. crassa* (Riquelme et al, 1998, Riquelme and Bartnicki-Garcia, 2004). *M. oryzae* WT and mutant hyphal plugs were inoculated on 35 mm plates filled with 3-5mm thick basal medium (with agarose as the solidifying agent) The fungal colony was allowed to reach a diameter of 2 cm before imaging. The actively growing cells at the colony periphery were then observed using a 40X or 100X objective lens. The hyphal and branch lengths and branching angles were determined using ImageJ. For hyphal morphology studies, 200 images of colony edge (100 hyphae x 2 plates) were captured for WT and *dam1Δ* strain using ProgResC5 Software on Olympus BX51. The distance from the hyphal tip to the point at which branches emerge and branch lengths were analysed for 100 – 130 images using Image J. A scatter plot was used to depict actual distances of 1<sup>st</sup>, 2<sup>nd</sup> and 3<sup>rd</sup> branch form hyphal tip. Branching frequency was plotted as percentage of hyphae that produced a certain number of branches (zero, 1, 2 or 3) within lengths of 300 μm or 500 μm from the growing tip of the primary hypha. Branching angles were measured with reference to the primary hypha from which the branch had emerged in the WT and *dam1Δ* strain for 100 hyphae and frequency of branch angles greater than 60° was determined. All the data shown are from three independent experiments. For fluorescence microscopy of hyphal structures, the fungal strains were inoculated on 2mm thick basal medium with agarose.

To study whether GFP-Dam1 dynamics were dependent on the microtubule network, germinating conidia were treated with 0.5 μM

nocodazole (microtubule inhibitor) and the process was followed for 15 mins capturing images every 5 min.

## 5. Chapter 5: Summary and Conclusion

1. Strains were developed to study chromosome/ nuclear and microtubule organisation during the cell cycle in *M. oryzae*. The nucleus was marked with a histone H1-mCherry tag in the B157 strain using the SRR strategy by ATMT. A separate  $\beta$ -tubulin-sGFP tagged B157 strain was generated by protoplast transformation in order to observe the microtubule network.
2. The hH1-mCherry tagged B157 was used to develop the hH1-mCherry,  $\beta$ -tubulin-sGFP double tagged strain. One round of nuclear division during the process of appressorium formation is critical to pathogenic development. This process was observed by time-lapse laser-scanning confocal/ fluorescence microscopy using the above RGB12 strain to obtain a WT reference for the subsequent analysis of kinetochore mutants.
3. Gene tagging constructs were developed for *DAM1*, *ASK1* and *MIS12* to replace the native copy of the gene with a GFP/RFP-tagged one. Dam1 was tagged on the N-terminus, while Ask1 and Mis12 were tagged on the C-terminus in the hH1-mCherry strain to give hH1-mCherry;GFP-Dam1, hH1-mCherry;Ask1-GFP and hH1-mCherry;Mis12-GFP strains. These strains were generated by ATMT.
4. All transformants generated in the study were screened by PCR and/or fluorescence microscopy. Southern blotting was used to identify transformants with single integration and confirm locus-specific incorporation. Virulence of strains was assessed by detached leaf spot assay. Strains with phenotype comparable to WT were used for further study.

5. Time-lapse laser-scanning confocal/fluorescence microscopy was used to monitor localisation of the kinetochore proteins both during mitosis and interphase at different stages of fungal development. Mis12-GFP-marked *M. oryzae* kinetochores were clustered during interphase, de-clustered during mitosis and re-clustered during telophase.
6. Mis12 was associated with the nucleus even during interphase while Dam1 and Ask1 showed more dynamic localisation patterns and associated with the nucleus specifically during mitosis. Importantly, multiple GFP-Dam1 and Ask1-GFP spots associated with the chromosomes at the start of mitosis and condensed into one large spot per nucleus during segregation and nuclear migration. This pattern was constant irrespective of developmental stage including, conidia, appressoria and vegetative hyphae.
7. Additional cytoplasmic Dam1 and Ask1 spots were seen in the vegetative hyphae, germ tube and appressorium. Further, Dam1 spots also vary in size.
8. Intense dynamic GFP-Dam1 and Ask1-GFP was also observed as multiple dynamic spots or streaks at the growing tips of vegetative hyphae and germ tubes and in the form of punctae moving back and forth from the hyphal tip cell. The dynamic localisation pattern was dependent on the microtubule network.
9. Conidiation involves three rounds of mitosis. The first mitosis occurs in the conidiophore stalk and the site of septation, which occurs at the base of the conidium, is uncoupled from the site of mitosis.
10. Representative genes, *CENPC/MIF2* from the inner, *MIS12/MTW1* from the middle and *DAM1* and *ASK1* from the outer kinetochore

complex were selected for study. Gene knock-down constructs were generated in pSilentDual-1 (pSD1) to study the role of *CENPC*, *MIS12* and *DAMI* in *M. oryzae*. The pSD1-CenpC, pSD1-Mis12 and pSD1-Dam1 plasmids were moved into *M. oryzae* RGB12 strain (hH1:mCherry,  $\beta$ -tubulin:sGFP) by protoplast transformation. However, most of the transformants were unstable and difficult to revive from stocks and were not used for further study.

11. A *DAMI* knock-out cassette was generated. This construct was transformed into WT strain B157 as well as the nuclear-microtubule dual tagged strain RGB12 by ATMT/ protoplast transformation. The *dam1* $\Delta$  mutant in RGB12 was used to study the effect of loss of *DAMI* on nuclear and spindle organisation, mitosis and nuclear migration. Fungal growth, development and virulence were studied in the *dam1* $\Delta$  mutant in the B157 background. Similarly, *ask1* $\Delta$  mutant was also generated and analysed for fungal development.
12. *dam1* $\Delta$  and *ask1* $\Delta$  showed reduced hyphal growth. The mutants produced flat colonies with reduced aerial hyphae and melanisation. The vegetative hyphae of *dam1* $\Delta$  and *ask1* $\Delta$  showed aberrant morphology with smaller and irregular cell compartments and frequent branching pattern with altered branching angles.
13. *dam1* $\Delta$  and *ask1* $\Delta$  display a decrease in conidium count to ~10% of the WT. Most of the *dam1* $\Delta$  and *ask1* $\Delta$  spores are 1 or 2 celled and oval-shaped, with only ~10-15% three celled conidia. Most of the *dam1* $\Delta$  conidiophores bear only 1-2 conidia, with only 6% conidiophores showing more than 2 conidia.
14. While 81% WT conidia formed appressoria, only 15% of the *dam1* $\Delta$  conidia were able to do so. Around half of the *dam1* $\Delta$  conidia failed to

germinate while others formed irregular germ tubes or were stuck at the hooking stage. In the rice sheath assay, only 28% *dam1* $\Delta$  mutant appressoria showed normal host penetration and invasion compared to 85% of WT appressoria. Most of the conidia failed to form normal functional appressoria even on the host surface, either failing to show penetration or displaying restricted invasive hyphae.

15. In whole plant infection assay, *dam1* $\Delta$  produced smaller and fewer lesions than the WT. Defective development of appressoria and impaired infection in the *dam1* $\Delta$  mutant raises the possibility of novel anti-fungal strategies directed towards these fungus-specific Dam1 complex proteins.
16. The length of the mitotic spindle was altered in the absence of Dam1. The nuclear and microtubule organization was affected by the loss of Dam1.
17. In the absence of Dam1, mitosis was prolonged due to delayed anaphase onset, with slow nuclear migration and improper segregation, both during vegetative and pathogenic development. Thus, Dam1 plays a critical role in the poleward segregation of chromosomes during anaphase and in nuclear migration.

## **Conclusion**

The outer kinetochore DASH complex proteins Dam1 and Ask1 display subcellular dynamics different from the inner kinetochore protein Mis12, such that they are associated with the nucleus during mitosis. Further, Dam1 plays a role in proper mitotic progression and chromosome segregation, likely through the establishment of correct spindle structure and KT-MT interactions. Dam1 is important for the development of

conidia, appressoria and host invasion in *M. oryzae*. Thus, the fungus-specific complex can be targeted for development of antifungal treatment of plant pathogens. During interphase, DASH complex proteins associate with cytoplasmic microtubules, accumulate at the tip and play a role in proper vegetative hyphal growth. Thus, in *M. oryzae*, and likely other filamentous fungi, the DASH complex may play additional roles in cell polarity and MT dynamics beyond its role at the mitotic spindle.

### **Future Directions**

What is the role of cytoplasmic DASH complex proteins during interphase?

What are the different oligomeric assemblies of the DASH complex observed *in vivo* in filamentous fungi?

Do different oligomeric DASH assemblies affect MT dynamics differently?

Does DASH interact with other MAPs while regulating MT dynamics?

Are all components of the DASH complex essential in *M. oryzae* and in other filamentous fungi?

Do all components of the DASH complex show similar sub-cellular localisation patterns?

Is there a difference in DASH complex roles in uninucleate and multi-nucleate filamentous fungi?

## 6. Bibliography

- Aist, J.R., and P.H. Williams. 1972. Ultrastructure and time course of mitosis in the fungus *Fusarium oxysporum*. *J. Cell Biol.* 55:368-389.
- Asbury, C.L., D.R. Gestaut, A.F. Powers, A.D. Franck, and T.N. Davis. 2006. The Dam1 kinetochore complex harnesses microtubule dynamics to produce force and movement. *Proc. Natl. Acad. Sci. U S A.* 103:9873-9878.
- Biggins, S. 2013. The composition, functions, and regulation of the budding yeast kinetochore. *Genetics.* 194:817-846.
- Boettcher, B., and Y. Barral. 2013. The cell biology of open and closed mitosis. *Nucleus.* 4:160-165.
- Burrack, L.S., S.E. Applen, and J. Berman. 2011. The requirement for the Dam1 complex is dependent upon the number of kinetochore proteins and microtubules. *Curr. Biol.* 21:889-896.
- Buttrick, G.J., T.C. Lancaster, J.C. Meadows, and J.B. Millar. 2012. Plo1 phosphorylates Dam1 to promote chromosome bi-orientation in fission yeast. *J. Cell Sci.* 125:1645-1651.
- Cai, X., X. Zhang, X. Li, M. Liu, X. Liu, X. Wang, H. Zhang, X. Zheng, and Z. Zhang. 2017. The Atypical Guanylate Kinase MoGuk2 Plays Important Roles in Asexual/Sexual Development, Conidial Septation, and Pathogenicity in the Rice Blast Fungus. *Front. Microbiol.* 8:2467.
- Cheeseman, I.M. 2014. The kinetochore. *Cold Spring Harb. Perspect. Biol.* 6:a015826.
- Cheeseman, I.M., S. Anderson, M. Jwa, E.M. Green, J. Kang, J.R. Yates, 3rd, C.S. Chan, D.G. Drubin, and G. Barnes. 2002. Phospho-regulation of kinetochore-microtubule attachments by the Aurora kinase Ipl1p. *Cell.* 111:163-172.
- Cheeseman, I.M., C. Brew, M. Wolyniak, A. Desai, S. Anderson, N. Muster, J.R. Yates, T.C. Huffaker, D.G. Drubin, and G. Barnes. 2001a. Implication of a novel multiprotein Dam1p complex in outer kinetochore function. *J. Cell Biol.* 155:1137-1145.
- Cheeseman, I.M., M. Enquist-Newman, T. Muller-Reichert, D.G. Drubin, and G. Barnes. 2001b. Mitotic spindle integrity and kinetochore function linked by the Duo1p/Dam1p complex. *J. Cell Biol.* 152:197-212.
- Chen, J., W. Zheng, S. Zheng, D. Zhang, W. Sang, X. Chen, G. Li, G. Lu, and Z. Wang. 2008. Rac1 is required for pathogenicity and Chm1-dependent conidiogenesis in rice fungal pathogen *Magnaporthe grisea*. *PLoS Pathog.* 4:e1000202.
- Chen, Y., X. Le, Y. Sun, M. Li, H. Zhang, X. Tan, D. Zhang, Y. Liu, and Z. Zhang. 2017. MoYcp4 is required for growth, conidiogenesis and pathogenicity in *Magnaporthe oryzae*. *Mol. Plant Pathol.* 18:1001-1011.
- Chen, Y., S. Zhai, Y. Sun, M. Li, Y. Dong, X. Wang, H. Zhang, X. Zheng, P. Wang, and Z. Zhang. 2015. MoTup1 is required for growth,

- conidiogenesis and pathogenicity of *Magnaporthe oryzae*. *Mol. Plant Pathol.* 16:799-810.
- Chen, Y., S. Zhai, H. Zhang, R. Zuo, J. Wang, M. Guo, X. Zheng, P. Wang, and Z. Zhang. 2014a. Shared and distinct functions of two Gti1/Pac2 family proteins in growth, morphogenesis and pathogenicity of *Magnaporthe oryzae*. *Environ. Microbiol.* 16:788-801.
- Chen, Y., R. Zuo, Q. Zhu, Y. Sun, M. Li, Y. Dong, Y. Ru, H. Zhang, X. Zheng, and Z. Zhang. 2014b. MoLys2 is necessary for growth, conidiogenesis, lysine biosynthesis, and pathogenicity in *Magnaporthe oryzae*. *Fungal Genet. Biol.* 67:51-57.
- Choi, J., K.S. Kim, H.S. Rho, and Y.H. Lee. 2011. Differential roles of the phospholipase C genes in fungal development and pathogenicity of *Magnaporthe oryzae*. *Fungal Genet. Biol.* 48:445-455.
- Choi, J., Y. Kim, S. Kim, J. Park, and Y.H. Lee. 2009. MoCRZ1, a gene encoding a calcineurin-responsive transcription factor, regulates fungal growth and pathogenicity of *Magnaporthe oryzae*. *Fungal Genet. Biol.* 46:243-254.
- Choi, W., and R.A. Dean. 1997. The adenylate cyclase gene MAC1 of *Magnaporthe grisea* controls appressorium formation and other aspects of growth and development. *Plant Cell.* 9:1973-1983.
- Ciferri, C., A. Musacchio, and A. Petrovic. 2007. The Ndc80 complex: hub of kinetochore activity. *FEBS Lett.* 581:2862-2869.
- Couch, B.C., and L.M. Kohn. 2002. A multilocus gene genealogy concordant with host preference indicates segregation of a new species, *Magnaporthe oryzae*, from *M. grisea*. *Mycologia.* 94:683-693.
- Dawe, A.L., V.C. McMains, M. Panglao, S. Kasahara, B. Chen, and D.L. Nuss. 2003. An ordered collection of expressed sequences from *Cryphonectria parasitica* and evidence of genomic microsynteny with *Neurospora crassa* and *Magnaporthe grisea*. *Microbiology.* 149:2373-2384.
- De Souza, C.P., S.B. Hashmi, T. Nayak, B. Oakley, and S.A. Osmani. 2009. Mlp1 acts as a mitotic scaffold to spatially regulate spindle assembly checkpoint proteins in *Aspergillus nidulans*. *Mol. Biol. Cell.* 20:2146-2159.
- De Souza, C.P., and S.A. Osmani. 2007. Mitosis, not just open or closed. *Eukaryot. Cell.* 6:1521-1527.
- Deng, Y.Z., Z. Qu, Y. He, and N.I. Naqvi. 2012. Sorting nexin Snx41 is essential for conidiation and mediates glutathione-based antioxidant defense during invasive growth in *Magnaporthe oryzae*. *Autophagy.* 8:1058-1070.
- Deng, Y.Z., M. Ramos-Pamplona, and N.I. Naqvi. 2009. Autophagy-assisted glycogen catabolism regulates asexual differentiation in *Magnaporthe oryzae*. *Autophagy.* 5:33-43.
- Dhatchinamoorthy, K., M. Shivaraju, J.J. Lange, B. Rubinstein, J.R. Unruh, B.D. Slaughter, and J.L. Gerton. 2017. Structural plasticity of the living kinetochore. *J. Cell Biol.* 216:3551-3570.

- Ding, R., K.L. McDonald, and J.R. McIntosh. 1993. Three-dimensional reconstruction and analysis of mitotic spindles from the yeast, *Schizosaccharomyces pombe*. *J. Cell Biol.* 120:141-151.
- Dong, Y., Q. Zhao, X. Liu, X. Zhang, Z. Qi, H. Zhang, X. Zheng, and Z. Zhang. 2015. MoMyb1 is required for asexual development and tissue-specific infection in the rice blast fungus *Magnaporthe oryzae*. *BMC Microbiol.* 15:37.
- Du, Y., L. Hong, W. Tang, L. Li, X. Wang, H. Ma, Z. Wang, H. Zhang, X. Zheng, and Z. Zhang. 2014. Threonine deaminase Mollv1 is important for conidiogenesis and pathogenesis in the rice blast fungus *Magnaporthe oryzae*. *Fungal Genet. Biol.* 73:53-60.
- Du, Y., Y. Shi, J. Yang, X. Chen, M. Xue, W. Zhou, and Y.L. Peng. 2013. A serine/threonine-protein phosphatase PP2A catalytic subunit is essential for asexual development and plant infection in *Magnaporthe oryzae*. *Curr. Genet.* 59:33-41.
- Enquist-Newman, M., I.M. Cheeseman, D. Van Goor, D.G. Drubin, P.B. Meluh, and G. Barnes. 2001. Dad1p, third component of the Duo1p/Dam1p complex involved in kinetochore function and mitotic spindle integrity. *Mol. Biol. Cell.* 12:2601-2613.
- Fernandez, J., and K. Orth. 2018. Rise of a Cereal Killer: The Biology of *Magnaporthe oryzae* Biotrophic Growth. *Trends Microbiol.* 26:582-597.
- Franco, A., J.C. Meadows, and J.B. Millar. 2007. The Dam1/DASH complex is required for the retrieval of unclustered kinetochores in fission yeast. *J. Cell Sci.* 120:3345-3351.
- Freitag, M. 2016. The kinetochore interaction network (KIN) of ascomycetes. *Mycologia.* 108:485-505.
- Freitag, M., P.C. Hickey, N.B. Raju, E.U. Selker, and N.D. Read. 2004. GFP as a tool to analyze the organization, dynamics and function of nuclei and microtubules in *Neurospora crassa*. *Fungal Genet. Biol.* 41:897-910.
- Fukagawa, T. 2017. Critical histone post-translational modifications for centromere function and propagation. *Cell Cycle.* 16:1259-1265.
- Funabiki, H., I. Hagan, S. Uzawa, and M. Yanagida. 1993. Cell cycle-dependent specific positioning and clustering of centromeres and telomeres in fission yeast. *J. Cell Biol.* 121:961-976.
- Gao, H.M., X.G. Liu, H.B. Shi, J.P. Lu, J. Yang, F.C. Lin, and X.H. Liu. 2013. MoMon1 is required for vacuolar assembly, conidiogenesis and pathogenicity in the rice blast fungus *Magnaporthe oryzae*. *Res. Microbiol.* 164:300-309.
- Gao, Q., T. Courtheoux, Y. Gachet, S. Tournier, and X. He. 2010. A non-ring-like form of the Dam1 complex modulates microtubule dynamics in fission yeast. *Proc Natl Acad Sci U S A.* 107:13330-13335.
- Goh, J., K.S. Kim, J. Park, J. Jeon, S.Y. Park, and Y.H. Lee. 2011. The cell cycle gene MoCDC15 regulates hyphal growth, asexual development and plant infection in the rice blast pathogen *Magnaporthe oryzae*. *Fungal Genet. Biol.* 48:784-792.

- Goshima, G., S. Saitoh, and M. Yanagida. 1999. Proper metaphase spindle length is determined by centromere proteins Mis12 and Mis6 required for faithful chromosome segregation. *Genes Dev.* 13:1664-1677.
- Goshima, G., and J.M. Scholey. 2010. Control of mitotic spindle length. *Annu. Rev. Cell Dev. Biol.* 26:21-57.
- Grishchuk, E.L., A.K. Efremov, V.A. Volkov, I.S. Spiridonov, N. Gudimchuk, S. Westermann, D. Drubin, G. Barnes, J.R. McIntosh, and F.I. Ataullakhanov. 2008a. The Dam1 ring binds microtubules strongly enough to be a processive as well as energy-efficient coupler for chromosome motion. *Proc. Natl. Acad. Sci. U S A.* 105:15423-15428.
- Grishchuk, E.L., I.S. Spiridonov, V.A. Volkov, A. Efremov, S. Westermann, D. Drubin, G. Barnes, F.I. Ataullakhanov, and J.R. McIntosh. 2008b. Different assemblies of the DAM1 complex follow shortening microtubules by distinct mechanisms. *Proc. Natl. Acad. Sci. U S A.* 105:6918-6923.
- Guo, M., F. Gao, X. Zhu, X. Nie, Y. Pan, and Z. Gao. 2015. MoGrr1, a novel F-box protein, is involved in conidiogenesis and cell wall integrity and is critical for the full virulence of *Magnaporthe oryzae*. *Appl. Microbiol. Biotechnol.* 99:8075-8088.
- Guo, M., L. Tan, X. Nie, and Z. Zhang. 2017. A class-II myosin is required for growth, conidiation, cell wall integrity and pathogenicity of *Magnaporthe oryzae*. *Virulence*:1-20.
- Hagan, I.M., A. Grallert, and V. Simanis. 2016. Analysis of the *Schizosaccharomyces pombe* Cell Cycle. *Cold Spring Harb. Protoc.* 2016.
- Hamer, J.E., B. Valent, and F.G. Chumley. 1989. Mutations at the smo genetic locus affect the shape of diverse cell types in the rice blast fungus. *Genetics.* 122:351-361.
- Han, J.H., J.H. Shin, Y.H. Lee, and K.S. Kim. 2018. Distinct roles of the YPEL gene family in development and pathogenicity in the ascomycete fungus *Magnaporthe oryzae*. *Sci. Rep.* 8:14461.
- Harris, S.D. 2008. Branching of fungal hyphae: regulation, mechanisms and comparison with other branching systems. *Mycologia.* 100:823-832.
- Harris, S.D. 2019. Hyphal branching in filamentous fungi. *Dev Biol.* 451:35-39.
- Hartwell, L.H., J. Culotti, and B. Reid. 1970. Genetic control of the cell-division cycle in yeast. I. Detection of mutants. *Proc. Natl. Acad. Sci. U S A.* 66:352-359.
- He, M., M.J. Kershaw, D.M. Soanes, Y. Xia, and N.J. Talbot. 2012. Infection-associated nuclear degeneration in the rice blast fungus *Magnaporthe oryzae* requires non-selective macro-autophagy. *PLoS One.* 7:e33270.
- He, Y., Y.Z. Deng, and N.I. Naqvi. 2013. Atg24-assisted mitophagy in the foot cells is necessary for proper asexual differentiation in *Magnaporthe oryzae*. *Autophagy.* 9:1818-1827.
- Hofmann, C., I. Cheeseman, B. Goode, K.L. McDonald, G. Barnes, and D.G. Drubin. 1998. *Saccharomyces cerevisiae* Duo1p and Dam1p, Novel

- Proteins Involved in Mitotic Spindle Function. *J. Cell Biol.* 143:1029-1040.
- Hou, H., Z. Zhou, Y. Wang, J. Wang, S.P. Kallgren, T. Kurchuk, E.A. Miller, F. Chang, and S. Jia. 2012. Csi1 links centromeres to the nuclear envelope for centromere clustering. *J. Cell Biol.* 199:735-744.
- Huang, L., S. Zhang, Z. Yin, M. Liu, B. Li, H. Zhang, X. Zheng, P. Wang, and Z. Zhang. 2017. MoVrp1, a putative verprolin protein, is required for asexual development and infection in the rice blast fungus *Magnaporthe oryzae*. *Sci. Rep.* 7:41148.
- Jacob, S., A.J. Foster, A. Yemelin, and E. Thines. 2015. High osmolarity glycerol (HOG) signalling in *Magnaporthe oryzae*: Identification of MoYPD1 and its role in osmoregulation, fungicide action, and pathogenicity. *Fungal Biol.* 119:580-594.
- Janke, C., J. Ortiz, T.U. Tanaka, J. Lechner, and E. Schiebel. 2002. Four new subunits of the Dam1-Duo1 complex reveal novel functions in sister kinetochore biorientation. *EMBO J.* 21:181-193.
- Jaqaman, K., J.F. Dorn, E. Marco, P.K. Sorger, and G. Danuser. 2007. Phenotypic clustering of yeast mutants based on kinetochore microtubule dynamics. *Bioinformatics.* 23:1666-1673.
- Jenkinson, C.B., K. Jones, J. Zhu, S. Dorhmi, and C.H. Khang. 2017. The appressorium of the rice blast fungus *Magnaporthe oryzae* remains mitotically active during post-penetration hyphal growth. *PLoS Genet.* 98:35-38.
- Jeon, J., J. Goh, S. Yoo, M.H. Chi, J. Choi, H.S. Rho, J. Park, S.S. Han, B.R. Kim, S.Y. Park, S. Kim, and Y.H. Lee. 2008. A putative MAP kinase kinase, MCK1, is required for cell wall integrity and pathogenicity of the rice blast fungus, *Magnaporthe oryzae*. *Mol. Plant Microbe Interact.* 21:525-534.
- Jeon, J., H. Rho, S. Kim, K.S. Kim, and Y.H. Lee. 2014. Role of MoAND1-mediated nuclear positioning in morphogenesis and pathogenicity in the rice blast fungus, *Magnaporthe oryzae*. *Fungal Genet. Biol.* 69:43-51.
- Jones, K., C.B. Jenkinson, M. Borges Araujo, J. Zhu, R.Y. Kim, D.W. Kim, and C.H. Khang. 2016. Mitotic stopwatch for the blast fungus *Magnaporthe oryzae* during invasion of rice cells. *Fungal Genet. Biol.* 93:46-49.
- Jones, M.H., J.B. Bachant, A.R. Castillo, T.H. Giddings, Jr., and M. Winey. 1999. Yeast Dam1p is required to maintain spindle integrity during mitosis and interacts with the Mps1p kinase. *Mol. Biol. Cell.* 10:2377-2391.
- Jones, M.H., X. He, T.H. Giddings, and M. Winey. 2001. Yeast Dam1p has a role at the kinetochore in assembly of the mitotic spindle. *Proc. Natl. Acad. Sci. U S A.* 98:13675-13680.
- Jorgensen, P., N.P. Edgington, B.L. Schneider, I. Rupes, M. Tyers, and B. Futcher. 2007. The size of the nucleus increases as yeast cells grow. *Mol. Biol. Cell.* 18:3523-3532.

- Kang, J., I.M. Cheeseman, G. Kallstrom, S. Velmurugan, G. Barnes, and C.S. Chan. 2001. Functional cooperation of Dam1, Ipl1, and the inner centromere protein (INCENP)-related protein Sli15 during chromosome segregation. *J. Cell Biol.* 155:763-774.
- Keating, P., N. Rachidi, T.U. Tanaka, and M.J. Stark. 2009. Ipl1-dependent phosphorylation of Dam1 is reduced by tension applied on kinetochores. *J. Cell Sci.* 122:4375-4382.
- Kershaw, M.J., M. Basiewicz, D.M. Soanes, X. Yan, L.S. Ryder, M. Csukai, M. Oses-Ruiz, B. Valent, and N.J. Talbot. 2019. Conidial Morphogenesis and Septin-Mediated Plant Infection Require Smo1, a Ras GTPase-Activating Protein in *Magnaporthe oryzae*. *Genetics.* 211:151-167.
- Khan, I.A., G. Ning, X. Liu, X. Feng, F. Lin, and J. Lu. 2015. Mitochondrial fission protein MoFis1 mediates conidiation and is required for full virulence of the rice blast fungus *Magnaporthe oryzae*. *Microbiol Res.* 178:51-58.
- Kiermaier, E., S. Woehrer, Y. Peng, K. Mechtler, and S. Westermann. 2009. A Dam1-based artificial kinetochore is sufficient to promote chromosome segregation in budding yeast. *Nat. Cell Biol.* 11:1109-1115.
- Kim, S., S.Y. Park, K.S. Kim, H.S. Rho, M.H. Chi, J. Choi, J. Park, S. Kong, J. Park, J. Goh, and Y.H. Lee. 2009. Homeobox transcription factors are required for conidiation and appressorium development in the rice blast fungus *Magnaporthe oryzae*. *PLoS Genet.* 5:e1000757.
- Kozubowski, L., V. Yadav, G. Chatterjee, S. Sridhar, M. Yamaguchi, S. Kawamoto, I. Bose, J. Heitman, and K. Sanyal. 2013. Ordered kinetochore assembly in the human-pathogenic basidiomycetous yeast *Cryptococcus neoformans*. *mBio.* 4:e00614-00613.
- Lai, J., S.K. Ng, F.F. Liu, R.N. Patkar, Y. Lu, J.R. Chan, A. Suresh, N. Naqvi, and G. Jedd. 2010. Marker fusion tagging, a new method for production of chromosomally encoded fusion proteins. *Eukaryot. Cell.* 9:827-830.
- Lampert, F., P. Hornung, and S. Westermann. 2010. The Dam1 complex confers microtubule plus end-tracking activity to the Ndc80 kinetochore complex. *J. Cell Biol.* 189:641-649.
- Lara-Gonzalez, P., F.G. Westhorpe, and S.S. Taylor. 2012. The spindle assembly checkpoint. *Curr. Biol.* 22:R966-980.
- Lau, G.W., and J.E. Hamer. 1998. Acropetal: a genetic locus required for conidiophore architecture and pathogenicity in the rice blast fungus. *Fungal Genet. Biol.* 24:228-239.
- Lee, K., P. Singh, W.C. Chung, J. Ash, T.S. Kim, L. Hang, and S. Park. 2006. Light regulation of asexual development in the rice blast fungus, *Magnaporthe oryzae*. *Fungal Genet. Biol.* 43:694-706.
- Legal, T., J. Zou, A. Sochaj, J. Rappsilber, and J.P. Welburn. 2016. Molecular architecture of the Dam1 complex-microtubule interaction. *Open Biol.* 6: 150237.

- Li, C., S. Cao, C. Zhang, Y. Zhang, Q. Zhang, J.R. Xu, and C. Wang. 2016a. MoCDC14 is important for septation during conidiation and appressorium formation in *Magnaporthe oryzae*. *Mol. Plant Pathol.*
- Li, C., J. Yang, W. Zhou, X.L. Chen, J.G. Huang, Z.H. Cheng, W.S. Zhao, Y. Zhang, and Y.L. Peng. 2014. A spindle pole antigen gene MoSPA2 is important for polar cell growth of vegetative hyphae and conidia, but is dispensable for pathogenicity in *Magnaporthe oryzae*. *Curr. Genet.* 60:255-263.
- Li, J.M., Y. Li, and S.J. Elledge. 2005. Genetic analysis of the kinetochore DASH complex reveals an antagonistic relationship with the ras/protein kinase A pathway and a novel subunit required for Ask1 association. *Mol. Cell Biol.* 25:767-778.
- Li, M., X. Liu, Z. Liu, Y. Sun, M. Liu, X. Wang, H. Zhang, X. Zheng, and Z. Zhang. 2016b. Glycoside Hydrolase MoGls2 Controls Asexual/Sexual Development, Cell Wall Integrity and Infectious Growth in the Rice Blast Fungus. *PLoS One.* 11:e0162243.
- Li, Y., J. Bachant, A.A. Alcasabas, Y. Wang, J. Qin, and S.J. Elledge. 2002. The mitotic spindle is required for loading of the DASH complex onto the kinetochore. *Genes Dev.* 16:183-197.
- Li, Y., Y. Que, Y. Liu, X. Yue, X. Meng, Z. Zhang, and Z. Wang. 2015. The putative Ggamma subunit gene MGG1 is required for conidiation, appressorium formation, mating and pathogenicity in *Magnaporthe oryzae*. *Curr. Genet.* 61:641-651.
- Li, Y., X. Zheng, M. Zhu, M. Chen, S. Zhang, F. He, X. Chen, J. Lv, M. Pei, Y. Zhang, Y. Zhang, W. Wang, J. Zhang, M. Wang, Z. Wang, G. Li, and G. Lu. 2019. MoIVD-Mediated Leucine Catabolism Is Required for Vegetative Growth, Conidiation and Full Virulence of the Rice Blast Fungus *Magnaporthe oryzae*. *Front Microbiol.* 10:444.
- Li, Y., J. Zhu, J. Hu, X. Meng, Q. Zhang, K. Zhu, X. Chen, X. Chen, G. Li, Z. Wang, and G. Lu. 2016c. Functional characterization of electron-transferring flavoprotein and its dehydrogenase required for fungal development and plant infection by the rice blast fungus. *Sci. Rep.* 6:24911.
- Lim, Y.J., K.T. Kim, and Y.H. Lee. 2018. SUMOylation is required for fungal development and pathogenicity in the rice blast fungus *Magnaporthe oryzae*. *Mol. Plant Pathol.* 19:2134-2148.
- Liu, H., F. Liang, F. Jin, and Y. Wang. 2008. The coordination of centromere replication, spindle formation, and kinetochore-microtubule interaction in budding yeast. *PLoS Genet.* 4:e1000262.
- Liu, H., A. Suresh, F.S. Willard, D.P. Siderovski, S. Lu, and N.I. Naqvi. 2007. Rgs1 regulates multiple Galpha subunits in *Magnaporthe* pathogenesis, asexual growth and thigmotropism. *EMBO J.* 26:690-700.
- Liu, W., S. Xie, X. Zhao, X. Chen, W. Zheng, G. Lu, J.R. Xu, and Z. Wang. 2010. A homeobox gene is essential for conidiogenesis of the rice blast fungus *Magnaporthe oryzae*. *Mol. Plant Microbe Interact.* 23:366-375.

- Liu, X.H., G.A. Ning, L.Y. Huang, Y.H. Zhao, B. Dong, J.P. Lu, and F.C. Lin. 2016. Calpains are involved in asexual and sexual development, cell wall integrity and pathogenicity of the rice blast fungus. *Sci. Rep.* 6:31204.
- Lu, Q., J.P. Lu, X.D. Li, X.H. Liu, H. Min, and F.C. Lin. 2008. *Magnaporthe oryzae* MTP1 gene encodes a type III transmembrane protein involved in conidiation and conidial germination. *J Zhejiang Univ Sci. B.* 9:511-519.
- Maiato, H., P.J. Hergert, S. Moutinho-Pereira, Y. Dong, K.J. Vandenbeldt, C.L. Rieder, and B.F. McEwen. 2006. The ultrastructure of the kinetochore and kinetochore fiber in *Drosophila* somatic cells. *Chromosoma.* 115:469-480.
- Matar, K.A.O., X. Chen, D. Chen, W.M. Anjago, J. Norvienyeku, Y. Lin, M. Chen, Z. Wang, D.J. Ebbole, and G.D. Lu. 2017. WD40-repeat protein MoCreC is essential for carbon repression and is involved in conidiation, growth and pathogenicity of *Magnaporthe oryzae*. *Curr. Genet.* 63:685-696.
- Matheis, S., A. Yemelin, D. Scheps, K. Andresen, S. Jacob, E. Thines, and A.J. Foster. 2017. Functions of the *Magnaporthe oryzae* Flb3p and Flb4p transcription factors in the regulation of conidiation. *Microbiol. Res.* 196:106-117.
- McKinley, K.L., and I.M. Cheeseman. 2016. The molecular basis for centromere identity and function. *Nat. Rev. Mol. Cell Biol.* 17:16-29.
- Mehrabi, R., S. Ding, and J.R. Xu. 2008. MADS-box transcription factor mig1 is required for infectious growth in *Magnaporthe grisea*. *Eukaryot. Cell.* 7:791-799.
- Meraldi, P., A.D. McAinsh, E. Rheinbay, and P.K. Sorger. 2006. Phylogenetic and structural analysis of centromeric DNA and kinetochore proteins. *Genome Biol.* 7:R23.
- Miranda, J.J., P. De Wulf, P.K. Sorger, and S.C. Harrison. 2005. The yeast DASH complex forms closed rings on microtubules. *Nat. Struct. Mol. Biol.* 12:138-143.
- Miranda, J.J., D.S. King, and S.C. Harrison. 2007. Protein arms in the kinetochore-microtubule interface of the yeast DASH complex. *Mol. Biol. Cell.* 18:2503-2510.
- Mohanar, V.C., P.M. Chandarana, B.B. Chattoo, R.N. Patkar, and J. Manjrekar. 2017. Fungal Histidine Phosphotransferase Plays a Crucial Role in Photomorphogenesis and Pathogenesis in *Magnaporthe oryzae*. *Front. Chem.* 5.
- Musacchio, A. 2015. The Molecular Biology of Spindle Assembly Checkpoint Signaling Dynamics. *Curr. Biol.* 25:R1002-1018.
- Musacchio, A., and A. Desai. 2017. A Molecular View of Kinetochore Assembly and Function. *Biology (Basel).* 6.
- Neumann, F.R., and P. Nurse. 2007. Nuclear size control in fission yeast. *J. Cell Biol.* 179:593-600.
- Nishimura, M., N. Hayashi, N.S. Jwa, G.W. Lau, J.E. Hamer, and A. Hasebe. 2000. Insertion of the LINE retrotransposon MGL causes a

- conidiophore pattern mutation in *Magnaporthe grisea*. *Mol. Plant Microbe Interact.* 13:892-894.
- Norvienenku, J., Z. Zhong, L. Lin, X. Dang, M. Chen, X. Lin, H. Zhang, W.M. Anjago, L. Lin, W. Abdul, and Z. Wang. 2017. Methylmalonate-semialdehyde dehydrogenase mediated metabolite homeostasis essentially regulate conidiation, polarized germination and pathogenesis in *Magnaporthe oryzae*. *Environ. Microbiol.* 19:4256-4277.
- Nurse, P., P. Thuriaux, and K. Nasmyth. 1976. Genetic control of the cell division cycle in the fission yeast *Schizosaccharomyces pombe*. *Mol. Gen. Genet.* 146:167-178.
- O'Toole, E.T., M. Winey, and J.R. McIntosh. 1999. High-voltage electron tomography of spindle pole bodies and early mitotic spindles in the yeast *Saccharomyces cerevisiae*. *Mol. Biol. Cell.* 10:2017-2031.
- Odenbach, D., B. Breth, E. Thines, R.W. Weber, H. Anke, and A.J. Foster. 2007. The transcription factor Con7p is a central regulator of infection-related morphogenesis in the rice blast fungus *Magnaporthe grisea*. *Mol. Microbiol.* 64:293-307.
- Oses-Ruiz, M., W. Sakulkoo, G.R. Littlejohn, M. Martin-Urdiroz, and N.J. Talbot. 2017. Two independent S-phase checkpoints regulate appressorium-mediated plant infection by the rice blast fungus *Magnaporthe oryzae*. *Proc. Natl. Acad. Sci. U S A.* 114:E237-E244.
- Palmer, D.K., K. O'Day, H.L. Trong, H. Charbonneau, and R.L. Margolis. 1991. Purification of the centromere-specific protein CENP-A and demonstration that it is a distinctive histone. *Proc. Natl. Acad. Sci. U S A.* 88:3734-3738.
- Park, S.Y., J. Choi, S.E. Lim, G.W. Lee, J. Park, Y. Kim, S. Kong, S.R. Kim, H.S. Rho, J. Jeon, M.H. Chi, S. Kim, C.H. Khang, S. Kang, and Y.H. Lee. 2013. Global expression profiling of transcription factor genes provides new insights into pathogenicity and stress responses in the rice blast fungus. *PLoS Pathog.* 9:e1003350.
- Patkar, R.N., A. Suresh, and N.I. Naqvi. 2010. MoTea4-mediated polarized growth is essential for proper asexual development and pathogenesis in *Magnaporthe oryzae*. *Eukaryot. Cell.* 9:1029-1038.
- Peng, H., Y. Feng, X. Zhu, X. Lan, M. Tang, J. Wang, H. Dong, and B. Chen. 2011. MoDUO1, a Duo1-like gene, is required for full virulence of the rice blast fungus *Magnaporthe oryzae*. *Curr. Genet.* 57:409-420.
- Pesenti, M.E., J.R. Weir, and A. Musacchio. 2016. Progress in the structural and functional characterization of kinetochores. *Curr. Opin. Struct. Biol.* 37:152-163.
- Prakash, C., J. Manjrekar, and B.B. Chattoo. 2016. Skp1, a component of E3 ubiquitin ligase, is necessary for growth, sporulation, development and pathogenicity in rice blast fungus (*Magnaporthe oryzae*). *Mol. Plant Pathol.* 17:903-919.
- Qi, Z., M. Liu, Y. Dong, Q. Zhu, L. Li, B. Li, J. Yang, Y. Li, Y. Ru, H. Zhang, X. Zheng, P. Wang, and Z. Zhang. 2016. The syntaxin protein

- (MoSyn8) mediates intracellular trafficking to regulate conidiogenesis and pathogenicity of rice blast fungus. *New Phytol.* 209:1655-1667.
- Ramanujam, R., and N.I. Naqvi. 2010. PdeH, a high-affinity cAMP phosphodiesterase, is a key regulator of asexual and pathogenic differentiation in *Magnaporthe oryzae*. *PLoS Pathog.* 6:e1000897.
- Redemann, S., J. Baumgart, N. Lindow, M. Shelley, E. Nazockdast, A. Kratz, S. Prohaska, J. Bruges, S. Furthauer, and T. Muller-Reichert. 2017. *C. elegans* chromosomes connect to centrosomes by anchoring into the spindle network. *Nat. Commun.* 8:15288.
- Reza, M.H., H. Shah, J. Manjrekar, and B.B. Chattoo. 2016. Magnesium Uptake by CorA Transporters Is Essential for Growth, Development and Infection in the Rice Blast Fungus *Magnaporthe oryzae*. *PLoS One.* 11:e0159244.
- Rieder, C.L., and H. Maiato. 2004. Stuck in division or passing through: what happens when cells cannot satisfy the spindle assembly checkpoint. *Dev. Cell.* 7:637-651.
- Roy, B., L.S. Burrack, M.A. Lone, J. Berman, and K. Sanyal. 2011. CaMtw1, a member of the evolutionarily conserved Mis12 kinetochore protein family, is required for efficient inner kinetochore assembly in the pathogenic yeast *Candida albicans*. *Mol. Microbiol.* 80:14-32.
- Sabnam, N., and S. Roy Barman. 2017. WISH, a novel CFEM GPCR is indispensable for surface sensing, asexual and pathogenic differentiation in rice blast fungus. *Fungal Genet. Biol.* 105:37-51.
- Saitoh, H., A. Hirabuchi, S. Fujisawa, C. Mitsuoka, R. Terauchi, and Y. Takano. 2014. MoST1 encoding a hexose transporter-like protein is involved in both conidiation and mycelial melanization of *Magnaporthe oryzae*. *FEMS Microbiol. Lett.* 352:104-113.
- Sambrook, J., and D.W. Russell. 2001. Molecular Cloning A Laboratory Manual. *Cold Spring Harb. Lab. Press.* Edition III.
- Sanchez-Perez, I., S.J. Renwick, K. Crawley, I. Karig, V. Buck, J.C. Meadows, A. Franco-Sanchez, U. Fleig, T. Toda, and J.B. Millar. 2005. The DASH complex and Klp5/Klp6 kinesin coordinate bipolar chromosome attachment in fission yeast. *EMBO J.* 24:2931-2943.
- Saunders, D.G., S.J. Aves, and N.J. Talbot. 2010a. Cell cycle-mediated regulation of plant infection by the rice blast fungus. *Plant Cell.* 22:497-507.
- Saunders, D.G., Y.F. Dagdas, and N.J. Talbot. 2010b. Spatial uncoupling of mitosis and cytokinesis during appressorium-mediated plant infection by the rice blast fungus *Magnaporthe oryzae*. *Plant Cell.* 22:2417-2428.
- Sazer, S., M. Lynch, and D. Needleman. 2014. Deciphering the evolutionary history of open and closed mitosis. *Curr. Biol.* 24:R1099-1103.
- Schumacher, M.A. 2012. Bacterial plasmid partition machinery: a minimalist approach to survival. *Curr. Opin. Struct. Biol.* 22:72-79.

- Sesma, A., and A.E. Osbourn. 2004. The rice leaf blast pathogen undergoes developmental processes typical of root-infecting fungi. *Nature*. 431:582-586.
- Sullivan, K.F., M. Hechenberger, and K. Masri. 1994. Human CENP-A contains a histone H3 related histone fold domain that is required for targeting to the centromere. *J. Cell Biol.* 127:581-592.
- Talbot, N.J. 2003. On the trail of a cereal killer: Exploring the biology of *Magnaporthe grisea*. *Annu. Rev. Microbiol.* 57:177-202.
- Tanaka, K., E. Kitamura, Y. Kitamura, and T.U. Tanaka. 2007. Molecular mechanisms of microtubule-dependent kinetochore transport toward spindle poles. *J. Cell. Biol.* 178:269-281.
- Thakur, J., and K. Sanyal. 2011. The essentiality of the fungus-specific Dam1 complex is correlated with a one-kinetochore-one-microtubule interaction present throughout the cell cycle, independent of the nature of a centromere. *Eukaryot. Cell.* 10:1295-1305.
- Tien, J.F., N.T. Umbreit, D.R. Gestaut, A.D. Franck, J. Cooper, L. Wordeman, T. Gonen, C.L. Asbury, and T.N. Davis. 2010. Cooperation of the Dam1 and Ndc80 kinetochore complexes enhances microtubule coupling and is regulated by aurora B. *J. Cell Biol.* 189:713-723.
- Tooley, J., and P.T. Stukenberg. 2011. The Ndc80 complex: integrating the kinetochore's many movements. *Chromosome Res.* 19:377-391.
- van Hooff, J.J.E., B. Snel, and G. Kops. 2017. Unique Phylogenetic Distributions of the Ska and Dam1 Complexes Support Functional Analogy and Suggest Multiple Parallel Displacements of Ska by Dam1. *Genome Biol. Evol.* 9:1295-1303.
- Veneault-Fourrey, C., M. Barooah, M. Egan, G. Wakley, and N.J. Talbot. 2006. Autophagic fungal cell death is necessary for infection by the rice blast fungus. *Science*. 312:580-583.
- Villalba, F., J. Collemare, P. Landraud, K. Lambou, V. Brozek, B. Cirer, D. Morin, C. Bruel, R. Beffa, and M.H. Lebrun. 2008. Improved gene targeting in *Magnaporthe grisea* by inactivation of MgKU80 required for non-homologous end joining. *Fungal Genet. Biol.* 45:68-75.
- Waldo, J., and M. Scherrer. 2008. Production and initial characterization of Dad1p, a component of the Dam1-DASH kinetochore complex. *PLoS One*. 3:e3888.
- Wang, Y., D. He, Y. Chu, Y.S. Zuo, X.W. Xu, X.L. Chen, W.S. Zhao, Y. Zhang, J. Yang, and Y.L. Peng. 2016. MoCps1 is important for conidiation, conidial morphology and virulence in *Magnaporthe oryzae*. *Curr Genet.* 62:861-871.
- Westermann, S., A. Avila-Sakar, H.W. Wang, H. Niederstrasser, J. Wong, D.G. Drubin, E. Nogales, and G. Barnes. 2005. Formation of a dynamic kinetochore- microtubule interface through assembly of the Dam1 ring complex. *Mol. Cell.* 17:277-290.
- Westermann, S., H.W. Wang, A. Avila-Sakar, D.G. Drubin, E. Nogales, and G. Barnes. 2006. The Dam1 kinetochore ring complex moves processively on depolymerizing microtubule ends. *Nature*. 440:565-569.

- Wilson, R.A., J.M. Jenkinson, R.P. Gibson, J.A. Littlechild, Z.Y. Wang, and N.J. Talbot. 2007. Tps1 regulates the pentose phosphate pathway, nitrogen metabolism and fungal virulence. *EMBO J.* 26:3673-3685.
- Wilson, R.A., and N.J. Talbot. 2009. Under pressure: investigating the biology of plant infection by *Magnaporthe oryzae*. *Nat. Rev. Microbiol.* 7:185-195.
- Winey, M., C.L. Mamay, E.T. O'Toole, D.N. Mastronarde, T.H. Giddings, Jr., K.L. McDonald, and J.R. McIntosh. 1995. Three-dimensional ultrastructural analysis of the *Saccharomyces cerevisiae* mitotic spindle. *J. Cell Biol.* 129:1601-1615.
- Xu, J.R., C.J. Staiger, and J.E. Hamer. 1998. Inactivation of the mitogen-activated protein kinase Mps1 from the rice blast fungus prevents penetration of host cells but allows activation of plant defense responses. *Proc. Natl. Acad. Sci. U S A.* 95:12713-12718.
- Yan, X., Y. Li, X. Yue, C. Wang, Y. Que, D. Kong, Z. Ma, N.J. Talbot, and Z. Wang. 2011a. Two novel transcriptional regulators are essential for infection-related morphogenesis and pathogenicity of the rice blast fungus *Magnaporthe oryzae*. *PLoS Pathog.* 7:e1002385.
- Yan, X., W.B. Ma, Y. Li, H. Wang, Y.W. Que, Z.H. Ma, N.J. Talbot, and Z.Y. Wang. 2011b. A sterol 14 $\alpha$ -demethylase is required for conidiation, virulence and for mediating sensitivity to sterol demethylation inhibitors by the rice blast fungus *Magnaporthe oryzae*. *Fungal Genet Biol.* 48:144-153.
- Yang, F., and N.I. Naqvi. 2014. Sulfonylurea resistance reconstitution as a novel strategy for ILV2-specific integration in *Magnaporthe oryzae*. *Fungal Genet. Biol.* 68:71-76.
- Yang, J., L. Liu, Y. Wang, C. Wang, J. Yan, Y. Liu, C. Wang, and C. Li. 2017. Overexpression of BAS1 in rice blast fungus can promote blast fungus growth, sporulation and virulence in planta. *Saudi J. Biol. Sci.* 24:1884-1893.
- Yang, J., X. Zhao, J. Sun, Z. Kang, S. Ding, J.R. Xu, and Y.L. Peng. 2010. A novel protein Com1 is required for normal conidium morphology and full virulence in *Magnaporthe oryzae*. *Mol. Plant Microbe Interact.* 23:112-123.
- Yang, L., L. Ukil, A. Osmani, F. Nahm, J. Davies, C.P. De Souza, X. Dou, A. Perez-Balaguer, and S.A. Osmani. 2004. Rapid production of gene replacement constructs and generation of a green fluorescent protein-tagged centromeric marker in *Aspergillus nidulans*. *Eukaryot. Cell.* 3:1359-1362.
- Ye, W., X. Chen, Z. Zhong, M. Chen, L. Shi, H. Zheng, Y. Lin, D. Zhang, G. Lu, G. Li, J. Chen, and Z. Wang. 2014. Putative RhoGAP proteins orchestrate vegetative growth, conidiogenesis and pathogenicity of the rice blast fungus *Magnaporthe oryzae*. *Fungal Genet. Biol.* 67:37-50.
- Yi, M., M.H. Chi, C.H. Khang, S.Y. Park, S. Kang, B. Valent, and Y.H. Lee. 2009. The ER chaperone LHS1 is involved in asexual development

- and rice infection by the blast fungus *Magnaporthe oryzae*. *Plant Cell*. 21:681-695.
- Yi, M., J.H. Park, J.H. Ahn, and Y.H. Lee. 2008. MoSNF1 regulates sporulation and pathogenicity in the rice blast fungus *Magnaporthe oryzae*. *Fungal Genet. Biol.* 45:1172-1181.
- Zhang, Y., H. Shi, S. Liang, G. Ning, N. Xu, J. Lu, X. Liu, and F. Lin. 2015a. MoARG1, MoARG5,6 and MoARG7 involved in arginine biosynthesis are essential for growth, conidiogenesis, sexual reproduction, and pathogenicity in *Magnaporthe oryzae*. *Microbiol. Res.* 180:11-22.
- Zhang, Z., J. Wang, R. Chai, H. Qiu, H. Jiang, X. Mao, Y. Wang, F. Liu, and G. Sun. 2015b. An S-(hydroxymethyl)glutathione dehydrogenase is involved in conidiation and full virulence in the rice blast fungus *Magnaporthe oryzae*. *PLoS One*. 10:e0120627.
- Zhou, W., W. Shi, X.W. Xu, Z.G. Li, C.F. Yin, J.B. Peng, S. Pan, X.L. Chen, W.S. Zhao, Y. Zhang, J. Yang, and Y.L. Peng. 2017. Glutamate synthase MoGlt1-mediated glutamate homeostasis is important for autophagy, virulence and conidiation in the rice blast fungus. *Mol. Plant Pathol.*
- Zhou, X., W. Liu, C. Wang, Q. Xu, Y. Wang, S. Ding, and J.R. Xu. 2011. A MADS-box transcription factor MoMcm1 is required for male fertility, microconidium production and virulence in *Magnaporthe oryzae*. *Mol. Microbiol.* 80:33-53.
- Zhu, X., T. Zhou, L. Chen, S. Zheng, S. Chen, D. Zhang, G. Li, and Z. Wang. 2016. Arf6 controls endocytosis and polarity during asexual development of *Magnaporthe oryzae*. *FEMS Microbiol. Lett.* 363.

## 7.1 Publications

1. **Shah, H.**, Rawat, K., Ashar, H., Patkar, R. and Manjrekar, J. (2019) Dual role for fungal-specific outer kinetochore proteins during cell cycle progression in *Magnaporthe oryzae*. *Journal of Cell Science*. 132, 224147. doi: 10.1242/jcs.224147.

First Person Interview with JCS alongwith above article published at <https://jcs.biologists.org/content/132/12/jcs234971>

2. Manjrekar, J. and **Shah, H.** (2019) Protein-based inheritance. *Seminars in Cell and Developmental Biology*. doi:10.1016/j.semcdb.2019.07.007

3. Reza M. H., **Shah H.**, Manjrekar J., and Chattoo B. B. (2016). Magnesium Uptake by CorA Transporters is Essential for Growth, Development and Infection in the Rice Blast Fungus *Magnaporthe oryzae*. *PLoS ONE* 11(7) 1-30 e0159244.

## **7.2 Oral and poster presentation at conferences**

1. Oral presentation on 'Understanding the role of kinetochore protein Dam1 in a plant fungal pathogen' at Seminar on 'Plant-Microbe Interactions in plant health, disease and biocontrol' on 4<sup>th</sup> October 2017 at Dept. of Microbiology and Biotechnology Centre, The Maharaja Sayajirao University of Baroda.
2. Poster presentation entitled "Dynamics of a kinetochore Protein Dam1 during development of the filamentous fungus *Magnaporthe oryzae*" at 14<sup>th</sup> European Conference on Fungal Genetics held at Haifa, Israel from 25<sup>th</sup> to 28<sup>th</sup> February 2018.
3. Best Poster Award for presentation of "Dynamics of a kinetochore Protein Dam1 during development of the filamentous fungus *Magnaporthe oryzae*" at 10<sup>th</sup> Bangalore Microscopy Course held at NCBS-TIFR, Bangalore, from 16<sup>th</sup> to 23<sup>th</sup> September 2018.
4. Poster presentation entitled "A non-canonical role of DASH proteins during interphase in *Magnaporthe oryzae* development" at 30<sup>th</sup> Fungal Genetics Conference held at Pacific Grove, California, USA, from 12<sup>th</sup> to 17<sup>th</sup> March 2019.