
8. REFERENCES

- Adachi, K. & Hamer, J.E (1998) Divergent cAMP signalling pathways regulate growth and pathogenesis in the rice blast fungus *Magnaporthe grisea*. *Plant Cell* **10**: 1361–1374.
- Adams, D.J. (2004) Fungal cell wall chitinases and glucanases. *Microbiology* **150**: 2029–2035.
- Alex, L.A., Korch C., Seltrennikoff, C.P. & Simon, M.I. (1998) COS1, a two-component histidine kinase that is involved in hyphal development in the opportunistic pathogen *Candida albicans*. *Proc Natl Acad Sci USA* **95**: 7069–7073.
- Backerman, J.L. & Ebbole, D.J. (1996) MPG1, a gene encoding fungal hydrophobin of *Magnaporthe grisea*, is involved in surface recognition. *Mol Plant-Microbe Interact* **9**: 450–456.
- Bahn, Y-S. (2008) Master and commander in fungal pathogens: the two-component system and the HOG signalling pathway. *Eukaryotic cell* **7**: 2017–2036.
- Bahn, Y-S. Kojima, K., Cox, G.M. & Heitman J (2006) A Unique Fungal Two-Component System Regulates Stress Responses, Drug Sensitivity, Sexual Development, and Virulence of *Cryptococcus neoformans*. *Mol biol cell* **17**: 3122–3135.
- Beckerman, J.L. & Ebbole, D.J. (1996) MPG1, a gene encoding a fungal hydrophobin of *Magnaporthe grisea*, is involved in surface recognition. *Mol. Plant Microbe Interact.* **9**:450–456
- Bhambra, G.K., Wang, Z.-Y., Soanes, D.M., Wakley, G.E. & Talbot, N.J. (2006) Peroxisomal carnitine acetyl transferase is required for elaboration of penetration hyphae during plant infection by *Magnaporthe grisea*. *Mol. Microbiol.* **61**: 46–60.
- Bonman, J.M., Estrada, B.A. Kim, C.K. & Lee E. J. (1991) Assessment of blast disease and yield loss in susceptible and partially resistant rice cultivars in two irrigated lowland environments. *Plant Disease.* **75**: 462–466.
- Bourett, T.M., & Howard, R.J. (1990) In vitro development of penetration structures in the rice blast fungus *Magnaporthe grisea*. *Can J Bot* **68**: 329–342.

- Catlett, N.L., Yoder, O.C. & Turgeon, B.G. (2003) Whole-Genome Analysis of Two-Component Signal Transduction Genes in Fungal Pathogens. *Eukaryotic cell*. **2**: 1151–1161.
- Chihiro Tanaka & Kosuke Izumitsu (2010). Two-Component Signaling System in Filamentous Fungi and the Mode of Action of Dicarboximide and Phenylpyrrole Fungicides, Fungicides, Odile Carisse (Ed.), ISBN: 978-953-307-266-1, InTech, DOI: 10.5772/13774.
- Choi, W. & Dean, R. A. (1997). The adenylate cyclase gene MAC1 of *Magnaporthe grisea* controls appressorium formation and other aspects of growth and development. *Plant Cell* **9**: 1973–1983.
- Dellaporta, S.L., Wood, J. & Hicks, J.B. (1983). *PMB Reporter* **4**: 19-21
- Desjardins, A.E. & Hohn, T.M. (1997) Mycotoxins in Plant Pathogenesis. *Mol. Plant-Microbe Interact.* **10**: 147-152.
- Divon, H.H. & Fluhr, R. (2007) Nutrition acquisition strategies during fungal infection of plants. *FEMS Microbiol Lett.* **266**: 65-74.
- Dixon, K.P., Xu, J.R., Smirnoff, N. & Talbot, N.J. (1999). Independent signalling pathways regulate cellular turgor during hyperosmotic stress and appressorium-mediated plant infection by *Magnaporthe grisea*. *Plant Cell* **11**: 2045– 2058.
- Dry, I.B., Yuan, K.H. & Hutton, D.G. (2004). Dicarboximide resistance in field isolates of *Alternaria alternata* is mediated by a mutation in a two-component histidine kinase gene. *Fungal Genet.Biol.* **41**: 102– 108.
- Ebbole, D.J. (2007) *Magnaporthe* as a model for understanding host-pathogen interactions. *Ann rev of phytopathol* **45**: 437-456.
- Egger, L.A., Park, H. & Inouye, M. (1997). Signal transduction via the histidyl-aspartyl phospho- relay histidyl-aspartyl phosphorelay. *Genes to Cells.* **2**: 167–184.
- Enjalbert, B., Smith, D.A., Cornell, M.J., Alam, I., Nicholls, S., Brown, A.J. & Quinn, J. (2006) Role of the Hog1 stress-activated protein kinase in the global transcriptional response to stress in the fungal pathogen *Candida albicans*. *Mol biol cell* **17**: 1018-1032.
- Fassler, J.S. & West, A.H. (2013) Histidine phosphotransfer proteins in fungal two-component signal transduction pathways. *Eukaryotic cell* **12**: 1052-1060.

- Fuchs, B.B. & Mylonakis, E. (2009) Our Paths Might Cross: the Role of the Fungal Cell Wall Integrity Pathway in Stress Response and Cross Talk with Other Stress Response Pathways. *Eukaryotic cell* **8**: 1616–1625.
- Furukawaa, K., Randhawa, A., Kaur, H., Mondal, A.K. & Hohmann, S. (2012) Fungal fludioxonil sensitivity is diminished by a constitutively active form of the group III histidine kinase. *FEBS letters* **586**: 2417-2422.
- Gilbert, R.D., Johnson, A.M. & Dean, R.A. (1996) Chemical signals responsible for appressorium formation in the rice blast fungus *Magnaporthe grisea*. *Physiol Mol Plant Pathol* **48**: 335–346.
- Gowda, M., Shirke, M. D., Mahesh, H. B., Chandarana, P., Rajamani, A., & Chattoo, B. B. (2015). Genome analysis of rice-blast fungus *Magnaporthe oryzae* field isolates from southern India. *Genomics Data*, **5**: 284–291.
- Gruber, S. & Seidl-Seiboth, V. (2012) Self versus non-self: fungal cell wall degradation in *Trichoderma*. *Microbiology* **158**: 26-34.
- Guo, M., Chen, Y., Du, Y., Dong, Y., Guo, W., Zhai, S., Zhang, H., Dong, S., Zhang, Z., Wang, Y., Wang, P. & Zheng, X. (2011) The bZIP transcription factor MoAP1 mediates the oxidative stress response and is critical for pathogenicity of the rice blast fungus *Magnaporthe oryzae*. *PLoS pathogens* **7**
- Hohmann, S., (2002). Osmotic Stress Signalling and Osmoadaptation in Yeasts Osmotic Stress Signalling and Osmoadaptation in Yeasts. *Microbiol Mol Biol Rev* **66**:2 : 300-372.
- Ikner, A., Shiozaki, K., (2005). Yeast signalling pathways in the oxidative stress response. *Mutat. Res.* **569**: 13–27.
- Iyer, G. & Chattoo, B.B. (2003) Purification and characterization of laccase from the rice blast fungus, *Magnaporthe grisea*. *FEMS Microbiol Lett* **227**: 121-126.
- Jacob, S., Foster, A.J., Yemelin, A. & Thines, E. (2015) High osmolarity glycerol (HOG) signalling in *Magnaporthe oryzae*: Identification of MoYPD1 and its role in osmoregulation, fungicide action, and pathogenicity. *Fungal Biology* **119**: 580-594.
- Jelitto, T.C., Page, H.A. & Read, N.D. (1994) Role of external signals in regulating the pre-penetration phase of infection by the rice blast fungus *Magnaporthe grisea*. *Planta* **194**: 471–477.

- Kaba, H.E.J., Nimtz, M., Müller, P.P. & Bilitewski, U., (2013). Involvement of the mitogen activated protein kinase Hog1p in the response of *Candida albicans* to iron availability. *BMC Microbiol.* **13**: 1-16.
- Lacal, J., Busch, A., Guazzaroni, M.E., Krell, T. & Ramos, J.L. (2006) The TodS-TodT two-component regulatory system recognizes a wide range of effectors and works with DNA-bending proteins. *Proc Natl Acad Sci USA* **103**: 8191-8196.
- Lamb, T.M., Goldsmith, C.S., Bennett, L., Finch, K.E. & Bell-Pedersen, D. (2011) Direct transcriptional control of a p38 MAPK pathway by the circadian clock in *Neurospora crassa*. *PLoS One* **6**.
- Lee, K., Singh, P., Chung, W.C., Ash, J., Kim, T.S., Hang, L. & Park, S. (2006) Light regulation of asexual development in the rice blast fungus, *Magnaporthe oryzae*. *Fungal Genet.Biol.***43**: 694-706.
- Lee, Y.H. & Dean, R.A. (1994) Hydrophobicity of contact surface induces appressorium formation in *Magnaporthe grisea*. *FEMS Microbiol Lett* **115**: 71–75.
- Lee, Y.H. & Dean, R. A. (1993) cAMP Regulates Infection Structure Formation in the Plant Pathogenic Fungus *Magnaporthe grisea*. *Plant Cell* **5**: 693–700.
- Li, D., Agrellos, O.A. & Calderone, R., (2010) Histidine kinases keep fungi safe and vigorous. *Curr.Opin.Microbiol.***13**: 424– 430.
- Lu, J.M-Y., Deschenes, R.J. & Fassler, J.S. (2003) *Saccharomyces cerevisiae* Histidine Phosphotransferase Ypd1p Shuttles between the Nucleus and Cytoplasm for SLN1-Dependent Phosphorylation of Ssk1p and Skn7p. *Eukaryotic cell* **2**: 1304–1314.
- Maeda, T., Wurgler- Murphy, S. M. & Saito, H.. (1994). A two - component system that regulates an osmosensing MAP kinase cascade in yeast. *Nature* **369**: 242–245.
- Mendgen, K., Hahn, M. & Deising, H. (1996) Morphogenesis and mechanisms of penetration by plant pathogenic fungi. *Annu. Rev. Phytopathol.* **34**: 367–86.
- Mew, T.M. (1991) Rice diseases. Chapter 5 In: Luh BS (ed.) Rice production 2nd Edition Van Nostrand Reinhold, New York. **1**: 187-236
- Millar, J.B., Buck, V. & Wilkinson, M.G., (1995). Pyp1 and Pyp2 PTPases dephosphorylate an osmosensing MAP kinase controlling cell size at division in fission yeast. *Genes Dev.* **9**: 2117–2130.
- Missall, T. A., Moran, J. M., Corbett, J. A., and Lodge, J. K. (2005). Distinct stress responses of two functional laccases in *Cryptococcus neoformans* are revealed in the absence of the thiol-specific antioxidant Tsa1. *Eukaryot. Cell* **4**, 202–208.

- Mizuno, T. (2004) Plant response regulators implicated in signal transduction and circadian rhythm. *Curr opin plant biol* **7**: 499-505.
- Motoyama, T., Kadokura, K., Ohira, T., Ichiishi, A., Fujimura, M., Yamaguchi, I. & Kudo, T. (2005) A two-component histidine kinase of the rice blast fungus is involved in osmotic stress response and fungicide action. *Fungal genet biol* **42**: 200-212.
- Motoyama, T., Ochiai, N., Morita, M., Iida, Y., Usami, R. & Kudo, T. (2008) Involvement of putative response regulator genes of the rice blast fungus *Magnaporthe oryzae* in osmotic stress response, fungicide action, and pathogenicity. *Curr genet* **54**: 185-195.
- Motoyama, T., Ohira, T., Kadokura, K., Ichiishi, A., Fujimura, M., Yamaguchi, I. & Kudo, T. (2005) An Os-1 family histidine kinase from a filamentous fungus confers fungicide-sensitivity to yeast. *Curr genet* **47**: 298-306.
- Mullins, E.D., Chen, X., Romaine, P., Raina, R., Geiser, D.M. & Kang, S. (2001) Agrobacterium-Mediated Transformation of *Fusarium oxysporum*: An Efficient Tool for Insertional Mutagenesis and Gene Transfer. *Phytopathology* **91**: 173-180.
- Nakayashiki, H., Hanada, S., Nguyen, B.Q., Kadotani, N., Tosa, Y. & Mayama, S. (2005) RNA silencing as a tool for exploring gene function in ascomycete fungi. *Fungal Genet. Biol.* **42**: 275-283.
- Oide, S., Liu, J., Yun, S.H., Wu, D., Michev, A., Choi, M.Y., Horwitz, B.A. & Turgeon, B.G. (2010) Histidine Kinase Two-Component Response Regulator Proteins Regulate Reproductive Development, Virulence, and Stress Responses of the Fungal Cereal Pathogens *Cochliobolus heterostrophus* and *Gibberella zeae*. *Eukaryotic cell* **9**: 1867-1880.
- Oshima, M., Fujimura, M., Banno, S., Hashimoto, C., Motoyama, T., Ichiishi, A. & Yamaguchi, I., (2002) A point mutation in the two-component histidine kinase BcOS-1 gene confers dicarboximide resistance in Weld isolates of *Botrytis cinerea*. *Phytopathology* **92**: 75– 80.
- Ota, I. M. & Varshavsky, A. (1993) A Yeast Protein Similar to Bacterial Two-Component Regulators. *Science* **262**: 566-569.
- Ou, S.H. (1985) Rice Diseases, 2nd ed. Commonwealth Mycological Institute, Kew, Surrey, England.

- Park, G., Xue, C., Zhao, X., Kim, Y., Orbach, M. & Xu, J.-R. (2006) Multiple upstream signals converge on the adaptor protein Mst50 in *Magnaporthe grisea*. *Plant Cell* **18**: 2822–2835.
- Patkar, R. N., and Chattoo, B. B. (2006). Transgenic indica Rice Expressing ns-LTP-Like Protein Shows Enhanced Resistance to Both Fungal and Bacterial Pathogens. *Mol. Breed.* **17**, 159–171. doi:10.1007/s11032-005-4736-3.
- Posas, F., Wurgler-Murphy, S.M., Maeda, T., Witten, E. a, Thai, T.C. & Saito, H. (1996) Yeast HOG1 MAP kinase cascade is regulated by a multistep phosphorelay mechanism in the SLN1-YPD1-SSK1 “two-component” osmosensor. *Cell* **86**: 865–875.
- Pott, G.B., Miller, T.K., Bartlett, J.A., Palas, J.S. & Selitrennikoff, C.P. (2000) The isolation of FOS-1, a gene encoding a putative two-component histidine kinase from *Aspergillus fumigatus*. *Fungal Genet. Biol.* **31**: 55-67.
- Qi, Z., Wang, Q., Dou, X., Wang, W., Zhao, Q., Lv R, Zhang, H., Zheng, X., Wang, P. & Zhang, Z. (2012) MoSwi6, an APSES family transcription factor, interacts with MoMps1 and is required for hyphal and conidial morphogenesis, appressorial function and pathogenicity of *Magnaporthe oryzae*. *Mol plant pathol* **13**: 677-689.
- Rahme, G.L., Mindrinos, M.N. & Panopoulos, N.J. (1992) Plant and environmental sensory signals control the expression of hrp genes in *Pseudomonas syringae* pv. phaseolicola. *J Bacteriol* **174**: 3499-3507.
- Ram, A.F.J., A. Wolters, S.S.C. Brekelmans, and F. M. Klis. (1994) A new approach for isolating cell wall mutants of *Saccharomyces cerevisiae* by screening for hypersensitivity of calcoflour white. *Yeast* **10**: 1019-1030
- Ramos-Pamplona, M. & Naqvi, N.I., (2006) Host invasion during rice-blast disease requires carnitine-dependent transport of peroxisomal acetyl-CoA. *Mol. Microbiol.* **61**: 61–75.
- Rispail, N. & Pietro, A.D.I., (2010) The two-component histidine kinase Fhk1 controls stress adaptation and virulence of *Fusarium oxysporum*. *Mol. Plant. Pathol.* **11**: 395–407.
- Saito, H. & Posas, F., (2012) Response to Hyperosmotic stress. *Genetics* **192**: 289–318
- Schmoll, M., Esquivel-Naranjo, E.U. & Herrera-Estrella, A. (2010) *Trichoderma* in the light of day--physiology and development. *Fungal Genet.Biol.* **47**: 909-916.

- Schmoll, M., Franchi, L. & Kubicek, C. P. (2005) Envoy, a PAS/LOV Domain Protein of *Hypocrea jecorina* (Anamorph *Trichoderma reesei*), Modulates Cellulase Gene Transcription in Response to Light. *Eukaryotic cell* **4**: 1998–2007.
- Schmoll, M., Tian, C., Sun, J., Tisch, D. & Glass, N.L. (2012) Unravelling the molecular basis for light modulated cellulase gene expression - the role of photoreceptors in *Neurospora crassa*. *BMC genomics* **13**: 127.
- Schulte, R. & Bonas, U., (1992). A *Xanthomonas* pathogenicity locus is induced by sucrose and sulphur-containing amino acids. *Plant Cell* **4**: 79- 86.
- Sesma A& Osbourn AE (2004) The rice leaf blast pathogen undergoes developmental processes typical of root-infecting fungi. *Nature* **431**: 582-586.
- Shinohara, M.L., Correa, A., Bell-Pedersen, D., Dunlap, J.C. & Loros, J.J. (2002) *Neurospora* Clock-Controlled Gene 9 (cgc-9) Encodes Trehalose Synthase: Circadian Regulation of Stress Responses and Development. *Eukaryotic cell* **1**[1]: 33-43.
- Smith, D., Morgan, B. & Quinn, J. (2010) Stress signalling to fungal stress-activated protein kinase pathways. *FEMS Microbiol.Lett.* **306**: 1–8.
- Snoeiijers, S.S., P´erez-Garc´ia, A., Joosten, M. H. A. J. & DeWit, P.J.G.M. (2000) The effect of nitrogen on disease development and gene expression in bacterial and fungal plant pathogens. *Eur J Plant Path* **106**: 493–506.
- Solomon, P.S., Tan, K-C. & Oliver, R.P. (2003) The nutrient supply of pathogenic fungi; a fertile field for study. *Mol Plant Pathol* **4**[3]: 203- 210.
- Song, H.K., Lee, J.Y., Lee, M.G., Min, J.M.K., Yang, J.K. & Suh, S.W. (1999) Insights into Eukaryotic Multistep Phosphorelay Signal transduction revealed by the crystal structure of Ypd1 p From *Saccharomyces cerevisiae*. *J Mol Biol* **293**: 753- 761.
- Su, C., Lu, Y. & Liu, H., (2013) Reduced TOR signalling sustains hyphal development in *Candida albicans* by lowering Hog1 basal activity. *Mol. Biol. Cell* **24**: 385–97.
- Talbot, N. J., Kershaw, M. J., Wakley, G. E., Vries, O. M. H. De, Wessels, J. G. H., & Hamer, J. E. (1996). MPCI Encodes a Fungal Hydrophobin Involved in Surface Interactions during Infection-Related Development of *Magnaporthe grisea*. *Plant cell* **8**: 985–999.
- Talbot, N.J. (2003) On the trail of a cereal killer: Exploring the biology of *Magnaporthe grisea*. *Annu. Rev. Microbiol.* **57**: 177–202.

- Talbot, N.J., Ebbole, D.J. & Hamer, J.E. (1993) Identification and characterization of MPG1, a gene involved in pathogenicity from the rice blast fungus *Magnaporthe grisea*. *Plant Cell* **5**: 1575–1590.
- Talbot, N.J., McCafferty, H.R.K., Ma, M., Moore, K. & Hamer, J.E. (1997) Nitrogen starvation of the rice blast fungus *Magnaporthe grisea* may act as an environmental cue for disease symptom expression. *Physiol.Mol Plant Pathol* **50**: 179–195.
- Tanaka, C. and Izumitsu, K. (2010) Two-component signaling system in filamentous fungi and the mode of action of dicarboximide and phenylpyrrole fungicides. In: *Fungicides* (Carisse, O., ed.), pp. 523–538. Rijeka: InTech
- Taylor, B.L. & Zhulin, I.B. (1999) PAS Domains: Internal Sensors of Oxygen, Redox Potential, and Light. *Microbiol Mol Biol Rev* **63**: 479–506.
- TeBeest, D.O., Guerber C.& Ditmore M. (2007) Rice Blast. *Plant Health Instructor*.
- Thines, E., Weber, R.W., & Talbot, N.J. (2000) MAP kinase and protein kinase A-dependent mobilization of triacylglycerol and glycogen during appressorium turgor generation by *Magnaporthe grisea*. *Plant Cell* **12**: 1703–1718.
- Tucker, S.L. & Talbot, N.J. (2001) Surface attachment and pre-penetration stage development by plant pathogenic fungi. *Annu. Rev. Phytopathol.* **39**, 385–417.
- Viaud, M., Fillinger, S., Liu, W., Polepalli, J.S., Le Pêcheur, P., Kunduru, A.R., Leroux, P. & Legendre, L. (2006) A class III histidine kinase acts as a novel virulence factor in *Botrytis cinerea*. *Mol. Plant. Microbe. Interact.* **19**: 1042–1050.
- Wang, Z.-Y., Jenkinson, J.M., Holcombe, L.J., Soanes, D.M., Veneault-Fourrey, C., Bhambra, G.K. & Talbot, N.J. (2005).The molecular biology of appressorium turgor generation by the rice blast fungus *Magnaporthe grisea*. *Biochem. Soc. Trans.* **33**: 384–388.
- Wood PJ, Fulcher RG (1983) Dye interactions, a basis for specific detection and histochemistry of polysaccharides. *J Histochem Cytochem* **31**:823–826
- Xu, J-R., Staiger, C.J. & Hamer, J.E. (1998) Inactivation of the mitogen-activated protein kinase Mps1 from the rice blast fungus prevents penetration of host cells but allows activation of plant defense responses. *Proc Natl Acad Sci USA* **95**: 12713–12718.
- Xu, Q. & West, A.H. (1999) Conservation of structure and function among histidine-containing phosphotransfer (HPT) domains as revealed by the crystal structure of YPD1. *J Mol Biol* **292**: 1039-1050.

- Zhang, H., Liu, K., Zhang, X., Song, W., Zhao, Q., Dong, Y., Guo, M., Zheng, X. & Zhang, Z. (2010) A two-component histidine kinase, MoSLN1, is required for cell wall integrity and pathogenicity of the rice blast fungus, *Magnaporthe oryzae*. *Curr genet* **56**: 517-528.
- Zhao, X., Mehrabi, R. & Xu, J.-R. (2007) Mitogen-activated protein kinase pathways and fungal pathogenesis. *Eukaryotic Cell* **6**: 1701-14.

