

- Adas, M. (2001). *Agricultural and pastoral societies in ancient and classical history* (Vol. 79). Temple University Press.
- Akiba, T., Abe, Y., Kitada, S., Kusaka, Y., Ito, A., Ichimatsu, T., & Ohba, M. (2009). Crystal structure of the parasporin-2 *Bacillus thuringiensis* toxin that recognizes cancer cells. *Journal of molecular biology*, 386(1), 121-133.
- Aly, C., & Mulla, M. S. (1987). Effect of two microbial insecticides on aquatic predators of mosquitoes. *Journal of Applied Entomology*, 103(1-5), 113-118.
- Alzate, O., Hemann, C. F., Osorio, C., Hille, R., & Dean, D. H. (2009). Ser170 of *Bacillus thuringiensis* Cry1Ab  $\delta$ -endotoxin becomes anchored in a hydrophobic moiety upon insertion of this protein into *Manduca sexta* brush border membranes. *BMC biochemistry*, 10(1), 25.
- Alzate, O., Osorio, C., Florez, A. M., & Dean, D. H. (2010). Participation of valine 171 in  $\alpha$ -helix 5 of *Bacillus thuringiensis* Cry1Ab  $\delta$ -endotoxin in translocation of toxin into *Lymantria dispar* midgut membranes. *Applied and environmental microbiology*, 76(23), 7878-7880.
- Angus TA. Studies of *Bacillus* spp. pathogenic for silkworm. Progress Report 1953; 9:Forest Biology Division, Canada Department of Science Service 6
- Anjali, C. H., Sharma, Y., Mukherjee, A., & Chandrasekaran, N. (2012). Neem oil (*Azadirachta indica*) nanoemulsion—a potent larvicidal agent against *Culex quinquefasciatus*. *Pest management science*, 68(2), 158-163.
- Aven, M., Hasui, H., & Motoyoshi, M. (2002). *U.S. Patent No. 6,387,848*. Washington, DC: U.S. Patent and Trademark Office.
- Aziz, A., Trotel-Aziz, P., Dhuicq, L., Jeandet, P., Couderchet, M., & Vernet, G. (2006). Chitosan oligomers and copper sulfate induce grapevine defense reactions and resistance to gray mold and downy mildew. *Phytopathology*, 96(11), 1188-1194.

- Babu, R. M., Sajeena, A., Seetharaman, K., & Reddy, M. S. (2003). Advances in genetically engineered (transgenic) plants in pest management—an over view. *Crop protection*, 22(9), 1071-1086.
- Baig DN, Mehnaz S (2010) Determination and distribution of *cry*-type genes in halophilic *Bacillus thuringiensis* isolates of Arabian Sea sedimentary rocks. *Microbiol Res* 165: 376-383.
- Bakar, N. A., Salleh, M. M., Umar, A. A., & Yahaya, M. (2011). The detection of pesticides in water using ZnCdSe quantum dot films. *Advances in Natural Sciences: Nanoscience and Nanotechnology*, 2(2), 025011.
- Becker, (2003) Ice granules containing endotoxins of microbial agents for the control of mosquito larvae--a new application technique." *Journal of the American Mosquito Control Association* 19, no. 1 (63-66).
- Becker, N M. Zgomba, M. Ludwig, D. Petric, F. Rettich (1992) Factors influencing the efficacy of the microbial control agent *Bacillus thuringiensis israelensis* J. Am. Mosq. Control Assoc., 8 (1992), 285–289
- Behle, R. W., McGuire, M. R., & Shasha, B. S. (1996). Extending the residual toxicity of *Bacillus thuringiensis* with casein-based formulations. *Journal of economic entomology*, 89(6), 1399-1405.
- Behle, R. W., McGuire, M. R., & Shasha, B. S. (1997). Effects of sunlight and simulated rain on residual activity of *Bacillus thuringiensis* formulations. *Journal of economic entomology*, 90(6), 1560-1566.
- Benhamou, N., & Thériault, G. (1992). Treatment with chitosan enhances resistance of tomato plants to the crown and root rot pathogen *Fusarium oxysporum* f. sp. *radicis-lycopersici*. *Physiological and Molecular Plant Pathology*, 41(1), 33-52.

- Ben-Shalom, N., Ardi, R., Pinto, R., Aki, C., & Fallik, E. (2003). Controlling gray mould caused by *Botrytis cinerea* in cucumber plants by means of chitosan. *Crop Protection*, 22(2), 285-290.
- Bernhard K, Jarrett P, Meadows M (1997) Natural isolates of *Bacillus thuringiensis*: Worldwide distribution, characterization, and activity against insect pests. *J Invertebr Pathol* 70: 59–68.
- Bhagat, D., Samanta, S. K., & Bhattacharya, S. (2013). Efficient management of fruit pests by pheromone nanogels. *Scientific reports*, 3, 1294.
- Bizzarri MF, Bishop AH (2008) The ecology of *Bacillus thuringiensis* on the phylloplane: Colonization from soil, plasmid transfer, and interaction with larvae of *Pieris brassicae*. *Microbial Ecol* 56: 133-139.
- Boehm, A. L, Martinon, I., Zerrouk, R., Rump, E., Fessi, H. (2003) Nanoprecipitation technique for the encapsulation of agrochemical active ingredients. *J. Microencapsulation*, 20, 433–441.
- Boehm, A. L. L. R., Zerrouk, R., & Fessi, H. (2000). Poly epsilon-caprolactone nanoparticles containing a poorly soluble pesticide: formulation and stability study. *Journal of microencapsulation*, 17(2), 195-205..
- Bok, S. H., Lee, H. W., Son, K. H., Kim, S. U., Lee, J. W., Kim, D. Y., & Kwon, Y. K. (1993) *U.S. Patent No. 5,273,749*. Washington, DC: U.S. Patent and Trademark Office.
- Boonlertnirun, S., Sarobol, E. D., Meechoui, S., & Sooksathan, I. (2007). Drought recovery and grain yield potential of rice after chitosan application. *Kasetsart Journal: Natural Science*, 41, 1-6.
- Bordes, P., Pollet, E., & Avérous, L. (2009). Nano-biocomposites: biodegradable polyester/nanoclay systems. *Progress in Polymer Science*, 34(2), 125-155.

- Brookes G, Barfoot P. (2013) The global income and production effects of genetically modified (GM) crops 1996–2011, *GM Crops and Food*; (4)1:74-83.
- Brookes, G., & Barfoot, P. (2015). Environmental impacts of genetically modified (GM) crop use 1996–2013: Impacts on pesticide use and carbon emissions. *GM crops & food*, 6(2), 103-133.
- Brookes, G., & Barfoot, P. (2016). Global income and production impacts of using GM crop technology 1996–2014. *GM crops & food*, 7(1), 38-77.
- Burges, H. D. (2012). Formulation of microbial biopesticides: beneficial microorganisms, nematodes and seed treatments. Springer Science & Business Media.
- Castagnola, A. S., & Jurat-Fuentes, J. L. (2012). Bt crops: past and future. In *Bacillus thuringiensis Biotechnology* (pp. 283-304). Springer Netherlands.
- Chandrkrachang, S. (2002). The application of chitin and chitosan in agriculture in Thailand. *Advances in chitin science*, 5(1), 458-462.
- .
- Chegâsaki Y. Aoeber (1915) die pathogenität der sog. sotto *Bacillen* (Ishiwata) bei seidenrauen. *Mitt Med Fak Kais*; 13:419
- Chen, C., Han, D., Cai, C., & Tang, X. (2010). An overview of liposome lyophilization and its future potential. *Journal of Controlled Release*, 142(3), 299-311.
- Chen, H., & Yada, R. (2011). Nanotechnologies in agriculture: New tools for sustainable development. *Trends in Food Science & Technology*, 22(11), 585-594.
- Chen, J. F., Wang, J. X., Liu, R. J., Shao, L., & Wen, L. X. (2004). Synthesis of porous silica structures with hollow interiors by templating nanosized calcium carbonate. *Inorganic Chemistry Communications*, 7(3), 447-449.

- Chen, J., Hua, G., Jurat-Fuentes, J. L., Abdullah, M. A., & Adang, M. J. (2007). Synergism of *Bacillus thuringiensis* toxins by a fragment of a toxin-binding cadherin. *Proceedings of the National Academy of Sciences*, 104(35), 13901-13906.
- chitosan two postharvest pathogens of strawberry fruits. *Phytopathology*, 82(4),
- Clive J.. (2015). 20th Anniversary (1996 to 2015) of the global commercialization of biotech crops and biotech crop highlights in 2015. *ISAAA Brief*, 51.
- coated microbial pesticides and pesticides producedtherefrom. US Patent 5,273,749;
- Cohen, S., Dym, O., Albeck, S., Ben-Dov, E., Cahan, R., Firer, M., & Zaritsky, A. (2008). High-resolution crystal structure of activated Cyt2Ba monomer from *Bacillus thuringiensis* subsp. israelensis. *Journal of molecular biology*, 380(5), 820-827.
- Cote , J. C., Vincent, C., Son, K. H., & Bok, S. H. (2001). Persistence of insecticidal activity of novel bio-encapsulated formulations of *Bacillus thuringiensis* var. kurstaki against *Choristoneura rosaceana* [Lepidoptera: Tortricidae]. *Phytoprotection*, 82(2), 73-82.
- Crickmore N, Zeigler DR, Feitelson J et al. (1998) Revision of the nomenclature for the *Bacillus thuringiensis* pesticidal crystal proteins. *Microbiol Mol Biol Rev* 62:807–813
- Cucchi, A., & Sanchez de Rivas, C. (1998). SASP (small, acid-soluble spore proteins) and spore properties in *Bacillus thuringiensis* israelensis and *Bacillus sphaericus*. *Current microbiology*, 36(4), 220-225.
- Damgaard PH, Abdel-Hameed A, Eilenberg J, Smits PH (1998) Natural occurrence of *Bacillus thuringiensis* on grass foliage. *World J Microbiol Biotechnol* 14: 239-242.

- de Escudero, I. R., Banyuls, N., Bel, Y., Maeztu, M., Escriche, B., Muñoz, D., ... & Ferré, J. (2014). A screening of five *Bacillus thuringiensis* Vip3A proteins for their activity against lepidopteran pests. *Journal of invertebrate pathology*, *117*, 51-55.
- de Maagd, R. A., Bravo, A., Berry, C., Crickmore, N., & Schnepf, H. E. (2003). Structure, diversity, and evolution of protein toxins from spore-forming entomopathogenic bacteria. *Annual review of genetics*, *37*(1), 409-433.
- Du, C., & Nickerson, K. W. (1996). *Bacillus thuringiensis* HD-73 spores have surface-localized Cry1Ac toxin: physiological and pathogenic consequences. *Applied and environmental microbiology*, *62*(10), 3722-3726.
- Dulmage, H. T., & Aizawa, K. (1982). Distribution of *Bacillus thuringiensis* in nature. *Microbial and viral pesticides*, 209-237.
- Dunkle, R. L., & Shasha, B. S. (1988). Starch-encapsulated *Bacillus thuringiensis*: a potential new method for increasing environmental stability of entomopathogens. *Environmental Entomology*, *17*(1), 120-126.
- El Ghaouth, A., Arul, J., Asselin, A., & Benhamou, N. (1992). Antifungal activity of chitosan on post-harvest pathogens: induction of morphological and cytological alterations in *Rhizopus stolonifer*. *Mycological research*, *96*(9), 769-779.
- Esmaeili, A., & Saremnia, B. (2012). Preparation of extract-loaded nanocapsules from *Onopordon leptolepis* DC. *Industrial Crops and Products*, *37*(1), 259-263.
- Esmaeili, A.; Saremnia, B. Preparation of extract-loaded nanocapsules from *Onopordon leptolepis* DC. *Ind. Crops Prod.* 2012, *37*, 259–263.
- Ferré, J., & Van Rie, J. (2002). Biochemistry and Genetics of Insect Resistance to *Bacillus thuringiensis*. *Annual review of entomology*, *47*(1), 501-533.
- Frankenhuyzen, K. (2009). Insecticidal activity of *Bacillus thuringiensis* crystal proteins. *Journal of invertebrate pathology*, *101*(1), 1-16.

- Fuller, D. Q. (2006). Agricultural origins and frontiers in South Asia: a working synthesis. *Journal of World Prehistory*, 20(1), 1-86.
- Gangal, K., Sarson, G. R., & Shukurov, A. (2014). The Near-Eastern Roots of the Neolithic in South Asia. *PloS one*, 9(5), e95714.
- Glare, T.R.; O'Callaghan, M. (2000). *Bacillus thuringiensis: biology, ecology and safety*. Chichester: John Wiley & Sons, 350.
- Gornik, K., Grzesik, M., & Romanowska-Duda, B. (2008). The effect of chitosan on rooting of grapevine cuttings and on subsequent plant growth under drought and temperature stress. *Journal of Fruit and Ornamental Plant Research*, 16, 333-343.
- Grochulski, P., Masson, L., Borisova, S., Pusztai-Carey, M., Schwartz, J. L., Brousseau, R., & Cygler, M. (1995). *Bacillus thuringiensis* CryIA (a) Insecticidal Toxin: Crystal Structure and Channel Formation. *Journal of molecular biology*, 254(3), 447-464.
- .
- Gunning, R. V., Dang, H. T., Kemp, F. C., Nicholson, I. C., & Moores, G. D. (2005). New resistance mechanism in *Helicoverpa armigera* threatens transgenic crops expressing *Bacillus thuringiensis* Cry1Ac toxin. *Applied and environmental microbiology*, 71(5), 2558-2563.
- Gupta, A. K. (2004). Origin of agriculture and domestication of plants and animals linked to early Holocene climate amelioration. *Current science-Bangalore-*, 87, 54-59.
- Handford, C. E., Dean, M., Henchion, M., Spence, M., Elliott, C. T., & Campbell, K. (2014). Implications of nanotechnology for the agri-food industry: Opportunities, benefits and risks. *Trends in Food Science & Technology*, 40(2), 226-241.
- Hannay, C. L. (1953). Crystalline inclusions in aerobic spore forming bacteria. *Nature*, 172, 1004.

- Hannay, C. L., & Fitz-James, P. (1955). The protein crystals of *Bacillus thuringiensis* Berliner. *Canadian journal of microbiology*, 1(8), 694-710.
- Harris, D. R. (1996). *Origins and spread of agriculture and pastoralism in Eurasia*. Smithsonian Institution Press.
- Hernández-Martínez, P., Navarro-Cerrillo, G., Caccia, S., de Maagd, R. A., Moar, W. J., Ferré, J.Herrero, S. (2010). Constitutive activation of the midgut response to *Bacillus thuringiensis* in Bt-resistant *Spodoptera exigua*. *PLoS One*, 5(9), e12795.
- Hwang, I. C., Kim, T. H., Bang, S. H., Kim, K. S., Kwon, H. R., Seo, M. J. (2011). Insecticidal effect of controlled release formulations of etofenprox based on nano-bio technique. 56(1), 33-40.
- Ifoulis, A. A., & Savopoulou-Soultani, M. (2004). Biological control of *Lobesia botrana* (Lepidoptera: Tortricidae) larvae by using different formulations of *Bacillus thuringiensis* in 11 vine cultivars under field conditions. *Journal of economic entomology*, 97(2), 340-343.
- Ikezawa, H. (1991). Bacterial PIPLCs-unique properties and usefulness in studies on GPI anchors. *Cell biology international reports*, 15(11), 1115-1131.
- Jindal V, Dhaliwal GS, Opende K. Pest Management in 21st Century (2013) Roadmap for future. *Biopesticides International* 9(1):1-22.
- Jindal, Vikas., Dhaliwal, G. S., & Koul, opende (2013). Pest management in 21st century: roadmap for future. *Biopest. Int*, 9(1), 1-22.
- Jurat-Fuentes, J. L., Gahan, L. J., Gould, F. L., Heckel, D. G., & Adang, M. J. (2004). The HevCaLP protein mediates binding specificity of the Cry1A class of *Bacillus thuringiensis* toxins in *Heliothis virescens*. *Biochemistry*, 43(44), 14299-14305.

- Kaelin, P., Morel, P., & Gadani, F. (1994). Isolation of *Bacillus thuringiensis* from stored tobacco and *Lasioderma serricorne* (F.). *Applied and environmental microbiology*, 60(1), 19-25.
- Kashyap, P. L., Xiang, X., & Heiden, P. (2015). Chitosan nanoparticle based delivery systems for sustainable agriculture. *International journal of biological macromolecules*, 77, 36-51.
- Kleter, G. A., Bhula, R., Bodnaruk, K., Carazo, E., Felsot, A. S., Harris, C. A. & Shevah, Y. (2007). Altered pesticide use on transgenic crops and the associated general impact from an environmental perspective. *Pest management science*, 63(11), 1107-1115.
- Knapp, O., Stiles, B., & Popoff, M. R. (2010). The aerolysin-like toxin family of cytolytic, pore-forming toxins. *The open toxinology Journal*, 3(1).
- Knight, P. J., Crickmore, N., & Ellar, D. J. (1994). The receptor for *Bacillus thuringiensis* CryIA (c) delta-endotoxin in the brush border membrane of the lepidopteran *Manduca sexta* is aminopeptidase N. *Molecular microbiology*, 11(3), 429-436.
- Knowles, R. G., Palacios, M., Palmer, R. M., & Moncada, S. (1989). Formation of nitric oxide from L-arginine in the central nervous system: a transduction mechanism for stimulation of the soluble guanylate cyclase. *Proceedings of the National Academy of Sciences*, 86(13), 5159-5162.
- Kookana, R. S.; Boxall, A. B. A.; Reeves, P. T.; Ashauer, R.; Beulke, S.; Chaudhry, Q.; Cornelis, G.; Fernandes, T. F.; Gan, J.; Kah, M.; Lynch, I.; Ranville, J.; Sinclair, C.; Spurgeon, D.; Tiede, K.; Van den Brink, P. J. Nanopesticides: guiding principles for regulatory evaluation of environmental risks. *J. Agric. Food Chem.* 2014, 62, 4227–4240.

- Krebs, R. E., & Krebs, C. A. (2003). *Groundbreaking scientific experiments, inventions, and discoveries of the ancient world*. Greenwood Publishing Group.
- Lao, S. B., Zhang, Z. X., Xu, H. H., & Jiang, G. B. (2010). Novel amphiphilic chitosan derivatives: synthesis, characterization and micellar solubilization of rotenone. *Carbohydrate Polymers*, 82(4), 1136-1142.
- Lew CW, Branly K, Gaytan J (1997). Encapsulation with water soluble polymer. United States Patent 5,643,351;
- Li, C., Wang, C., Wang, C., & Hu, S. (2006). Development of a parathion sensor based on molecularly imprinted nano-TiO<sub>2</sub> self-assembled film electrode. *Sensors and Actuators B: Chemical*, 117(1), 166-171.
- Li, J., Carroll, J. and Ellar, D. J. (1991) Crystal structure of insecticidal delta-endotoxin from *Bacillus thuringiensis* at 2.5 Å resolution. *Nature* 353, 815-821
- Liu, J., Wang, F. H., Wang, L. L., Xiao, S. Y., Tong, C. Y., Tang, D. Y., & Liu, X. M. (2008). Preparation of fluorescence starch-nanoparticle and its application as plant transgenic vehicle. *Journal of Central South University of Technology*, 15(6), 768-773.
- LiuF, XuZ, Zhu YC, HuangF, WangY, LiH, etal, Evidence of field evolved resistance to Cry1Ac-expressing Bt cotton in *Helicoverpa armigera* (Lepidoptera: Noctuidae) in northern China. *Pest Manag Sci* 66:155–161 (2010).
- Ma, G., Rahman, M. M., Grant, W., Schmidt, O., & Asgari, S. (2012). Insect tolerance to the crystal toxins Cry1Ac and Cry2Ab is mediated by the binding of monomeric toxin to lipophorin glycolipids causing oligomerization and sequestration reactions. *Developmental & Comparative Immunology*, 37(1), 184-192.

- Maldonado, B. M., Galán, W. L., Rodríguez, P. C., & Quiroz, M. H. (2002). Evaluation of polymer-based granular formulations of *Bacillus thuringiensis israelensis* against larval *Aedes aegypti* in the laboratory. *J Am Mosq Control Assoc*, *18*, 352-358.
- Malthankar, P. A., & Gujar, G. T. (2016). Toxicity of *Bacillus thuringiensis* Cry2Ab and the inheritance of Cry2Ab resistance in the Pink bollworm, *Pectinophora gossypiella* (Saunders).
- Martin, P. A., & Travers, R. S. (1989). Worldwide abundance and distribution of *Bacillus thuringiensis* isolates. *Applied and Environmental Microbiology*, *55*(10), 2437-2442.
- McGuire MR, Shasha BS, Eastman CE, Oloumi-Sadeghi H (1996) Starch and flour based sprayable formulations: effect on rainfastness and solar stability of *Bacillus thuringiensis*. *J Econ Entomol* 89:863–869.
- Meadows, M. P., Ellis, D. J., Butt, J., Jarrett, P., & Burges, H. D. (1992). Distribution, frequency, and diversity of *Bacillus thuringiensis* in an animal feed mill. *Applied and Environmental Microbiology*, *58*(4), 1344-1350.
- Memarizadeh, N., Ghadamyari, M., Adeli, M., & Talebi, K. (2014). Preparation, characterization and efficiency of nanoencapsulated imidacloprid under laboratory conditions. *Ecotoxicology and environmental safety*, *107*, 77-83.
- Milani, N., McLaughlin, M. J., Stacey, S. P., Kirby, J. K., Hettiarachchi, G. M., Beak, D. G., & Cornelis, G. (2012). Dissolution kinetics of macronutrient fertilizers coated with manufactured zinc oxide nanoparticles. *Journal of agricultural and food chemistry*, *60*(16), 3991-3998.
- Mizuki, E., Maeda, M., Tanaka, R., Lee, D. W., Hara, M., Akao, T., ... & Ohba, M. (2001). *Bacillus thuringiensis*: a common member of microflora in activated sludges of a sewage treatment plant. *Current microbiology*, *42*(6), 422-425.

- Mora-Huertas, C. E., Fessi, H., & Elaissari, A. (2010). Polymer-based nanocapsules for drug delivery. *International journal of pharmaceutics*, 385(1), 113-142.
- Morse, R. J., Yamamoto, T., & Stroud, R. M. (2001). Structure of Cry2Aa suggests an unexpected receptor binding epitope. *Structure*, 9(5), 409-417.
- Nair, R., Varghese, S. H., Nair, B. G., Maekawa, T., Yoshida, Y., & Kumar, D. S. (2010). Nanoparticulate material delivery to plants. *Plant science*, 179(3), 154-163.
- Nguyen, H. M., Hwang, I. C., Park, J. W., & Park, H. J. (2012). Photoprotection for deltamethrin using chitosan-coated beeswax solid lipid nanoparticles. *Pest management science*, 68(7), 1062-1068.
- No, H. K., Meyers, S. P., Prinyawiwatkul, W., & Xu, Z. (2007). Applications of chitosan for improvement of quality and shelf life of foods: a review. *Journal of food science*, 72(5), R87-R100.
- Nuruzzaman, M., Rahman, M. M., Liu, Y., & Naidu, R. (2016). Nanoencapsulation, nano-guard for pesticides: a new window for safe application. *Journal of agricultural and food chemistry*, 64(7), 1447-1483.
- Pardo-Lopez, L., Soberon, M., & Bravo, A. (2013). *Bacillus thuringiensis* insecticidal three-domain Cry toxins: mode of action, insect resistance and consequences for crop protection. *FEMS microbiology reviews*, 37(1), 3-22.
- Patel, K. D., Bhanshali, F., & Ingle, S. S. (2011). Diversity and characterization of *Bacillus thuringiensis* isolates from alluvial soils of Mahi river basin, India. *J Adv Dev Res*, 2(1), 14-20.
- Photchanachai, S., Singkaew, J., & Thamthong, J. (2006). Effects of chitosan seed treatment on colletotrichum sp. and seedling growth of chili cv. 'jinda'. In *IV International Conference on Managing Quality in Chains-The Integrated View on Fruits and Vegetables Quality 712* ( 585-590).

- Pigott, C. R., & Ellar, D. J. (2007). Role of receptors in *Bacillus thuringiensis* crystal toxin activity. *Microbiology and Molecular Biology Reviews*, 71(2), 255-281.
- Prado, A. G., Moura, A. O., & Nunes, A. R. (2011). Nanosized silica modified with carboxylic acid as support for controlled release of herbicides. *Journal of agricultural and food chemistry*, 59(16), 8847-8852.
- Promdonkoy, B., & Ellar, D. J. (2003). Investigation of the pore-forming mechanism of a cytolytic  $\delta$ -endotoxin from *Bacillus thuringiensis*. *Biochemical Journal*, 374(1), 255-259.
- Qaim, M., & Zilberman, D. (2003). Yield effects of genetically modified crops in developing countries. *Science*, 299(5608), 900-902.
- Rabea, E. I., Badawy, M. E. T., Stevens, C. V., Smagghe, G., & Steurbaut, W. (2003). Chitosan as antimicrobial agent: applications and mode of action. *Biomacromolecules*, 4(6), 1457-1465.
- Rahman, M. M., Roberts, H. L., Sarjan, M., Asgari, S., & Schmidt, O. (2004). Induction and transmission of *Bacillus thuringiensis* tolerance in the flour moth *Ephesia kuehniella*. *Proceedings of the National academy of Sciences of the United States of America*, 101(9), 2696-2699.
- Rajamohan, F., Alcantara, E., Lee, M. K., Chen, X. J., Curtiss, A., & Dean, D. H. (1995). Single amino acid changes in domain II of *Bacillus thuringiensis* CryIAb delta-endotoxin affect irreversible binding to *Manduca sexta* midgut membrane vesicles. *Journal of bacteriology*, 177(9), 2276-2282.
- Rodda & Ubertini (2004) The Basis of Civilisation-Water Science? International Association of Hydrological Science .ISBN 1-901502-57-0
- Rodríguez, A. T., Ramírez, M. A., Cárdenas, R. M., Hernández, A. N., Velázquez, M. G., & Bautista, S. (2007). Induction of defense response of *Oryza sativa* L. against

- Pyricularia grisea (Cooke) Sacc. by treating seeds with chitosan and hydrolyzed chitosan. *Pesticide Biochemistry and Physiology*, 89(3), 206-215.
- S. Boonlertnirun, E.D. Sarobol, S. Meechoui, I. Sooksathan, Drought recovery and grain yield potential of rice after chitosan application, *Kasetsart J. Nat.Sci.* 41 (2007) 1–6.
- Sansinenea, E. (2012). *Bacillus thuringiensis biotechnology* (pp. 3-18). Dordrecht; Heidelberg; London; New York, NY: Springer.
- Sansinenea, E. (2012). Discovery and description of *Bacillus thuringiensis*. *Bacillus thuringiensis Biotechnology*, Springer: 3-18.
- Schnepf, E., Crickmore, N. V., Van Rie, J., Lereclus, D., Baum, J., Feitelson, J., ... & Dean, D. H. (1998). *Bacillus thuringiensis* and its pesticidal crystal proteins. *Microbiology and molecular biology reviews*, 62(3), 775-806.
- Shah, P. A., & Pell, J. K. (2003). Entomopathogenic fungi as biological control agents. *Applied microbiology and biotechnology*, 61(5), 413-423.
- Shelton, A. M., Zhao, J. Z., & Roush, R. T. (2002). Economic, ecological, food safety, and social consequences of the deployment of Bt transgenic plants. *Annual review of entomology*, 47(1), 845-881.
- Singh, V. P., & Yadava, R. N. (Eds.). (2003). Water Resources System Operation: Proceedings of the International Conference on Water and Environment (WE-2003), December 15-18, 2003, Bhopal, India (Vol. 5). Allied Publishers.
- Skovmand, O., & Sanogo, E. (1999). Experimental formulations of *Bacillus sphaericus* and *B. thuringiensis israelensis* against *Culex quinquefasciatus* and *Anopheles gambiae* (Diptera: Culicidae) in Burkina Faso. *Journal of medical entomology*, 36(1), 62-67.
- Smith, R. A., & Couche, G. A. (1991). The phylloplane as a source of *Bacillus thuringiensis* variants. *Applied and Environmental Microbiology*, 57(1), 311-315.

- Stein, J. (1998). *Running steel, running America: Race, economic policy and the decline of liberalism*. Univ of North Carolina Press.
- Swiecicka I, Fiedoruk K, Bednarz G (2002) The occurrence and properties of *Bacillus thuringiensis* isolated from free-living animals. *Lett Appl Microbiol* 34: 194–198.
- Szczzech, Magdalena, and Robert Maciorowski. "Microencapsulation Technique with Organic Additives for Biocontrol Agents." *Journal of Horticultural Research* 24, no. 1 (2016): 111-122.
- T.L. Couch, C.M. and Ignoffo H.D. Burges (1981) (Ed.), Formulation of insect pathogens, in microbial control of pests and plant diseases, Academic Press, London, pp. 621–634.
- Tabashnik, B. E., Sisterson, M. S., Ellsworth, P. C., Dennehy, T. J., Antilla, L., Liesner, L., Yelich, A. J. (2010). Suppressing resistance to Bt cotton with sterile insect releases. *Nature biotechnology*, 28(12), 1304-1307.
- Tamez-Guerra, P., Garcia-Gutierrez, C., Medrano-Roldán, H., Galán-Wong, L. J., & Sandoval-Coronado, C. F. (1999). Spray-dried microencapsulated *Bacillus thuringiensis* formulations for the control of *Epilachna varivestis* Mulsant. *Southwest. Entomol*, 24, 37-48.
- Tamez-Guerra, P., McGuire, M. R., Behle, R. W., Shasha, B. S., & Galí Wong, L. J. (2000). Assessment of microencapsulated formulations for improved residual activity of *Bacillus thuringiensis*. *Journal of economic entomology*, 93(2), 219-225.
- Tang, J. D., Gilboa, S., Roush, R. T., & Shelton, A. M. (1997). Inheritance, stability, and lack-of-fitness costs of field-selected resistance to *Bacillus thuringiensis* in diamondback moth (Lepidoptera: Plutellidae) from Florida. *Journal of Economic Entomology*, 90(3), 732-741.

- Toledo, J., Liedo, P., Williams, T., & Ibarra, J. (1999). Toxicity of *Bacillus thuringiensis*  $\beta$ -exotoxin to three species of fruit flies (Diptera: Tephritidae). *Journal of economic entomology*, 92(5), 1052-1056.
- Valaitis, A. P., Jenkins, J. L., Lee, M. K., Dean, D. H., & Garner, K. J. (2001). Isolation and partial characterization of gypsy moth BTR-270, an anionic brush border membrane glycoconjugate that binds *Bacillus thuringiensis* Cry1A toxins with high affinity. *Archives of insect biochemistry and physiology*, 46(4), 186-200.
- Wan P, Huang Y, Wu H, Huang M, Cong S et al. (2012) Increased frequency of pink bollworm resistance to Bt toxin Cry1Ac in China. PLOS ONE 7: e29975.
- Wang, L., Li, X., Zhang, G., Dong, J., & Eastoe, J. (2007). Oil-in-water nanoemulsions for pesticide formulations. *Journal of colloid and interface science*, 314(1), 230-235.
- Wei, J., Guo, Y., Liang, G., Wu, K., Zhang, J., Tabashnik, B. E., & Li, X. (2015). Cross-resistance and interactions between Bt toxins Cry1Ac and Cry2Ab against the cotton bollworm. *Scientific reports*, 5, 7714.
- Wisseman S. U. & Williams, W. S. (1994) Ancient Technologies and Archaeological Materials, Routledge ISBN 2-88124-632-
- Wojdyła, A. T. (2003). Chitosan (biochikol 020 PC) in the control of some ornamental foliage diseases. *Communications in agricultural and applied biological sciences*, 69(4), 705-715.
- Wolfersberger, M. G., Chen, X. J., & Dean, D. H. (1996). Site-directed mutations in the third domain of *Bacillus thuringiensis* delta-endotoxin CryIAa affect its ability to increase the permeability of *Bombyx mori* midgut brush border membrane vesicles. *Applied and environmental microbiology*, 62(1), 279-282.
- Wood, B. (1996). The Origins and Spread of Agriculture and Pastoralism in Eurasia, by D. Harris. *Nature*, 382(6586), 36-37.

- Yang, F. L., Li, X. G., Zhu, F., & Lei, C. L. (2009). Structural characterization of nanoparticles loaded with garlic essential oil and their insecticidal activity against *Tribolium castaneum* (Herbst)(Coleoptera: Tenebrionidae). *Journal of agricultural and food chemistry*, 57(21), 10156-10162.
- Yin, Yan-hua, Qing-ming GUO, Han Yun, Ling-jing WANG, and Shu-qing WAN. "Preparation, characterization and nematicidal activity of lansiumamide B nano-capsules." *Journal of Integrative Agriculture* 11, no. 7 (2012): 1151-1158.