

List of Figures

Sr No.	Table no.	Title	Page no.
1	Figure 1.1	Chemical Reaction for DNSA Method	20
2	Figure 1.2	Life cycle of <i>S. oryzae</i> : a) egg, b) &c) larvae, d) pupa and e) adults	24
3	Figure 1.3a	Female Identification Characters	24
4	Figure 1.3b	Male Identification Characters	25
5	Figure 1.4 a	A. Aedeagus of male genitalia B. Y-sclerite of female genitalia c. Antennae of <i>S. oryzae</i>	25
6	Figure 1.4b	Epipharyngeal rod of larva. B. Apical sensory organs on the labial palp of larva	25
7	Figure 1.5	Infestation of <i>S. oryzae</i> in all four selected host grain	27
8	Figure 1.6	Incubation time taken by <i>S. oryzae</i> in different grains	29
9	Figure 1.7	No. of Adult <i>S. oryzae</i> emerge in different grains	29
10	Figure 1.8	Amount of grain weight loss by <i>S. oryzae</i>	30
11	Figure 1.9	Body length of male and female in different grains	30
12	Figure 1.10	Effect of <i>S. oryzae</i> infestation on protein content of the different grains	32
13	Figure 1.11	Effect of <i>S. oryzae</i> infestation on Carbohydrate content of the different grains	32
14	Figure 1.12	Effect of different grains on fecundity rate of <i>S. oryzae</i>	34
15	Figure 1.13	Effect of different grains on longevity of <i>S. oryzae</i>	34
16	Figure 1.14	Male : Female ratio in newly emerge <i>S. oryzae</i> in different grains	35
17	Figure 2.1	Shows the cultured cell density with Grace's and TNF-FH media till 96 hrs of duration	60
18	Figure 2.2	Cell density in different incubation media at 24 hrs	61
19	Figure 2.3	Cell density in different incubation media at 48 hrs	63
20	Figure 2.4	Cell density in different incubation media at 72 hrs	65
21	Figure 2.5	Cell density in different incubation media at 96 hrs	67
22	Figure 2.6	A Growth curve of SoMG cell line in Grace's medium containing 25% fetal bovine serum + 10pmol ecdysone+ vitamin mixture	69
23	Figure 2.7	<i>S.oryzae</i> midgut cultured cell growth in Grace's media having 25% FBS +10pmol Ecdysone . A) at 24 hrs B) at 48 hrs C) at 72 hrs D) at 96 hrs (100X magnification)	71
24	Figure 2.8	Gel image of COI, lane 1 DNA ladder, lane 2 tissue, Lane 3. SoMG cell line	73
25	Figure 2.9	Gel image of 16s rRNA. lane 1 DNA ladder, lane 2	74

		tissue, Lane 3. SoMG cell line	
26	Figure 2.10	A. Semi-quantitative RT-PCR to compare gene expression patterns in SoMG cell line and tissues. B. Quantitative qPCR amplification for 3 surface marker.	75
27	Figure 2.11	Karyotypes of SoMG cell line. (Magnification 10 x 100, 5 μ M)	76
28	Figure 3.1	SoMG cell mortality against different concentration of Fenitrothion	97
29	Figure 3.2	SoMG cell mortality against different concentration of Malathion	98
30	Figure 3.3	Sf9 cell mortality against different concentration of Fenitrothion	98
31	Figure 3.4	Sf9 cell mortality against different concentration of Malathion	99
32	Figure 3.5	Results of Acridine orange/Ethidium bromide assay for (A) SoMG and (B) Sf9 cells after 24 hrs of treatment with sublethal concentrations (LD, MD and HD) of Fenitrothion	100
33	Figure 3.6	Results of Acridine orange/Ethidium bromide assay for (A) SoMG and (B) Sf9 cells after 48 hrs of treatment with sublethal concentrations (LD, MD and HD) of Fenitrothion	101
34	Figure 3.7	Results of Acridine orange/Ethidium bromide assay for (A) SoMG and (B) Sf9 cells after 72 hrs of treatment with sublethal concentrations (LD, MD and HD) of Fenitrothion	102
35	Figure 3.8	Results of Acridine orange/Ethidium bromide assay for (A) SoMG and (B) Sf9 cells after 96 hrs of treatment with sublethal concentrations (LD, MD and HD) of Fenitrothion	103
36	Figure 3.9	: Results of Acridine orange/Ethidium bromide assay for (A) SoMG and (B) Sf9 cells after 24 hrs of treatment with sublethal concentrations (LD, MD and HD) of Malathion	104
37	Figure 3.10	Results of Acridine orange/Ethidium bromide assay for (A) SoMG and (B) Sf9 cells after 48 hrs of treatment with sublethal concentrations (LD, MD and HD) of Malathion	105
38	Figure 3.11	Results of Acridine orange/Ethidium bromide assay for (A) SoMG and (B) Sf9 cells after 72 hrs of treatment with sublethal concentrations(LD, MD and HD) of Malathion	106
39	Figure 3.12	Results of Acridine orange/Ethidium bromide assay for (A) SoMG and (B) Sf9 cells after 96 hrs of treatment with sublethal concentrations(LD, MD and	107

		HD) of Malathion	
40	Figure 3.13	Depicts the fluorescence in SoMG and Sf9 cells treated with sub-lethal doses of fenitrothion at 24 hrs and stain with AO/EB	110
41	Figure 3.14	Depicts the fluorescence in SoMG and Sf9 cells treated with sub-lethal doses of fenitrothion at 48 hrs and stain with AO/EB	110
42	Figure 3.15	Depicts the fluorescence in SoMG and Sf9 cells treated with sub-lethal doses of fenitrothion at 72 hrs and stain with AO/EB	111
43	Figure 3.16	Depicts the fluorescence in SoMG and Sf9 cells treated with sub-lethal doses of fenitrothion at 96 hrs and stain with AO/EB	111
44	Figure 3.17	Depicts the fluorescence in SoMG and Sf9 cells treated with sub-lethal doses of malathion at 24 hrs and stain with AO/EB	112
45	Figure 3.18	Depicts the fluorescence in SoMG and Sf9 cells treated with sub-lethal doses of malathion at 48 hrs and stain with AO/EB	112
46	Figure 3.19	Depicts the fluorescence in SoMG and Sf9 cells treated with sub-lethal doses of malathion at 72 hrs and stain with AO/EB	113
47	Figure 3.20	Depicts the fluorescence in SoMG and Sf9 cells treated with sub-lethal doses of malathion at 96hrs and stain with AO/EB	113
48	Figure 3.21	Results of DAPI staining for SoMG cell line after (A) 24 hrs and (B) 48 hrs of treatment with sublethal concentrations(LD, MD and HD) of Fenitrothion	116
49	Figure 3.22	Results of DAPI staining for SoMG cell line after (A) 72 hrs and (B) 96 hrs of treatment with sublethal concentrations(LD, MD and HD) of Fenitrothion	117
50	Figure 3.23	Depicts the fluorescence in SoMG cells treated with sub-lethal doses of fenitrothion for 24, 48, 72 and 96 hrs and stain with DAPI	119
51	Figure 3.24	Results of DCF-DH staining for ROS Generation in SoMG cell line after (A) 24 hrs and (B) 48 hrs of treatment with sublethal concentrations(LD, MD and HD) of Fenitrothion	121
52	Figure 3.25	Results of DCF-DH staining for ROS Generation in SoMG cell line after (A) 72 hrs and (B) 96 hrs of treatment with sublethal concentrations(LD, MD and HD) of Fenitrothion	122
53	Figure 3.26	Depicts the fluorescence in SoMG cells treated with sub-lethal doses of fenitrothion for 24, 48, 72 and 96 hrs and stain with DCFH-DA	123
54	Figure 3.27	Depicts the level of SOD (in folds) in SoMG cells	124

		treated with sub-lethal doses of fenitrothion for 24, 48, 72 and 96 hrs	
55	Figure 3.28	Depicts the level of GST (in folds) in SoMG cells treated with sub-lethal doses of fenitrothion for 24, 48, 72 and 96 hrs	125
56	Figure 3.29	Depicts the level of CAT (in folds) in SoMG cells treated with sub-lethal doses of fenitrothion for 24, 48, 72 and 96 hrs	125
57	Figure 3.30	Flow cytometry dot plots with double Annexin V-FITC/PI staining for a) control cells, b) cells exposed to LD of fenitrothion c) cells exposed to MD of fenitrothion and d) cells exposed to HD of fenitrothion at 24 hrs	127
58	Figure 3.31	Flow cytometry dot plots with double Annexin V-FITC/PI staining for a) control cells, b) cells exposed to LD of fenitrothion c) cells exposed to MD of fenitrothion and d) cells exposed to HD of fenitrothion at 48 hrs	128
59	Figure 3.32	Flow cytometry dot plots with double Annexin V-FITC/PI staining for a) control cells, b) cells exposed to LD of fenitrothion c) cells exposed to MD of fenitrothion and d) cells exposed to HD of fenitrothion at 72 hrs	129
60	Figure 3.33	Flow cytometry dot plots with double Annexin V-FITC/PI staining for a) control cells, b) cells exposed to LD of fenitrothion c) cells exposed to MD of fenitrothion and d) cells exposed to HD of fenitrothion at 96 hrs	130
61	Figure 3.34	Depicts the summary data of FAC _S analysis of apoptosis where SoMG cells were treated with fenitrothion at 24 hrs	131
62	Figure 3.35	Depicts the summary data of FACS analysis of apoptosis where SoMG cells were treated with fenitrothion at 48 hrs	131
63	Figure 3.36	Depicts the summary data of FACS analysis of apoptosis where SoMG cells were treated with fenitrothion at 72 hrs	132
64	Figure 3.37	Depicts the summary data of FACS analysis of apoptosis where SoMG cells were treated with fenitrothion at 24 hrs	132
65	Figure I	The diagram depicting the host preference of <i>S. oryzae</i> . Maize grains was found to be most preferred with lowest incubation period giving the highest number of progeny, body length, fecundity, longevity, and female: male ratio compared to rice, chickpea, and millet .	158

66	Figure II	Illustration showing the isolation of midgut cells of <i>S. oryzae</i> cultured in two insect media. Grace's media with supplements resulted in an appreciable growth of the cells. Molecular characterization of newly developed SoMG cells was confirmed by COI and 16sRNA sequencing. Karyotyping and its ectodermal origin were authenticated by expression of cup1, cup4, and hmg176.	159
67	Figure III	Summarizing the results of the toxic potential of organophosphate insecticides - Fenitrothione and Malathion on SoMG primary cell-line. Cytotoxic assay confirmed the Fenitrothion to be highly toxic with increase apoptosis in AO/EB staining. Furthermore, DNA damage was also confirmed by DAPI staining and ROS generation by DCFDA stains. Therefore, activation of antioxidants SOD, CAT and GSH were confirmed by validating the gene expression followed by quantification of apoptosis and live cells by FACs analysis.	160