

REFERENCES

- AACC International Method 02-52.01. *Hydrogen-Ion Activity (pH)—Electrometric Method*. doi.org/10.1094/AACCIntMethod-02-52.01.
- AACC International Method 28-41.03. *Acid Hydrolysis Method for Extracting Insect Fragments and Rodent Hairs—Light Filth in White Flour*. doi.org/10.1094/AACCIntMethod-28-41.03
- AACC International Method 28-44.01. *Iodine Method, for Insect Eggs in Flour*. doi.org/10.1094/AACCIntMethod-28-44.01.
- AACC International Method, 30-10.01. *Crude Fat in Flour, Bread, and Baked Cereal Products Not Containing Fruit*.
<https://doi.org/10.1094/aaccintmethod-30-10.01>.
doi.org/10.1094/AACCIntMethod-30-10.01.
- AACC International Method 44-01.01. *Percent Moisture*.
doi.org/10.1094/AACCIntMethod-44-01.01.
- Abbas, M., Sheikh, A. D., Sabir, H. M., & Nighat, S. (2005). Factors Responsible for Low Wheat Productivity in Central Punjab. *Pakistan Journal of Agricultural Sciences*, 42(3–4), 79–82.
- Abbasipour, H.; Mahmoudvand, M.; Rastegar, F.; Hosseinpour, M. H. (2011). Bioactivities of jimsonweed extract, *Datura stramonium* L.(Solanaceae), against *Tribolium castaneum* (Coleoptera: Tenebrionidae). *Turkish Journal of Agriculture and Forestry*, 35(6), 623–629.
<https://doi.org/10.3906/tar-1004-874>
- Abbaszadeh, S., Sharifzadeh, A., Shokri, H., Khosravi, A. R., & Abbaszadeh, A. (2014). Antifungal efficacy of thymol, carvacrol, eugenol and menthol as alternative agents to control the growth of food-relevant fungi. *Journal de Mycologie Medicale*, 24(2), e51–e56.
<https://doi.org/10.1016/j.mycmed.2014.01.063>
- Abdelgaleil, S. A. M., Abbassy, M. A., Belal, A. S. H., & Abdel Rasoul, M. A. A. (2008). Bioactivity of two major constituents isolated from the

- essential oil of *Artemisia judaica* L. *Bioresource Technology*, 99(13), 5947–5950. <https://doi.org/10.1016/j.biortech.2007.10.043>
- Ahluwalia, M. S. (1978). The Journal of Development Rural poverty and agricultural performance in India. *The Journal of Development Studies*, 14(3), 298–323. <https://doi.org/10.1080/00220387808421677>
- Ahmed, H. (1983). Losses incurred in stored foos grains by insect pets- A review. *PPakistan Journal of Agriculture Research*, 4(3), 198–207.
- Ajayi, F. A., & Rahman, S. A. (2006). Susceptibility of some staple processed meals to red flour beetle, *Tribolium castaneum* (Herbst)(Coleoptera: Tenebrionidae). *Pakistan Journal of Biological Sciences*, 9(9), 1744–1748.
- Ali, M., & Gupta, S. (2012). Carrying capacity of Indian agriculture: Pulse crops. *Current Science*, 102(6), 874–881.
- Anbar, A. D.; Yung, Y. L.; Chavez, F. P. (1996). Methyl bromide: Ocean sources, ocean sinks, and climate sensitivity. *Global Biogeochemical Cycles*, 10(1), 175–190.
- Anita, S., Sujatha, P., & Prabhudas, P. (2012). Efficacy of pulverised leaves of *Annona squamosa* (L .), *Moringa oleifera* (Lam .) and *Eucalyptus globulus* (Labill .) against the stored grain pest , *Tribolium castaneum* (Herbst .). *Recent Research in Science and Technology*, 4(2), 19–23.
- Arnaud, L., Haubruge, E., & Gage, M. J. G. (2005). The malathion-specific resistance gene confers a sperm competition advantage in *Tribolium castaneum*. *Functional Ecology*, 19(6), 1032–1039. <https://doi.org/10.1111/j.1365-2435.2005.01055.x>
- Aulakh, J. and Regmi, A. (2013). Post-harvest food losses estimation- development of consistent methodology. In *Rome: FAO*. <https://doi.org/10.1016/j.jns.2003.09.014>
- Babarinde, S. A., Oyegoke, O. O., & Adekunle, A. E. (2011). Larvicidal and insecticidal properties of *Ricinus communis* seed extracts obtained by different methods against *Tribolium castaneum* Herbst (Coleoptera:

- Tenebrionidae). *Archives of Phytopathology and Plant Protection*, 44(5), 451–459. <https://doi.org/10.1080/03235400903093220>
- Bachrouch, O., Jemâa, J. M. Ben, Talou, T., Marzouk, B., & Abderraba, M. (2010). Fumigant toxicity of *Pistacia lentiscus* essential oil against *Tribolium castaneum* and *Lasioderma serricorne*. *Bulletin of Insectology*, 63(1), 129–135.
- Bakkali, F., Averbeck, S., Averbeck, D., & Idaomar, M. (2008). Biological effects of essential oils - A review. *Food and Chemical Toxicology*, 46(2), 446–475. <https://doi.org/10.1016/j.fct.2007.09.106>
- Bakula, L., Bakula, T., Baranowski, M., & Czarnewicz, A. (2011). The carcinogenic effects of benzoquinones produced by the flour beetle. *Polish Journal of Veterinary Sciences*, 14(1), 159–164. <https://doi.org/10.2478/v10181-011-0025-8>
- Bamaiyi, L. J., Onu, I., Amatobi, C. I., & Dike, M. C. (2006). Effect of *Callosobruchus maculatus* infestation on nutritional loss on stored cowpea grains. *Archives of Phytopathology and Plant Protection*, 39(2), 119–127. <https://doi.org/10.1080/03235400500180743>
- Bell, C. H. (2000). Fumigation in the 21st century. *Crop Protection*, 19(8–10), 563–569. [https://doi.org/10.1016/S0261-2194\(00\)00073-9](https://doi.org/10.1016/S0261-2194(00)00073-9)
- Benhalima, H., Chaudhry, M. Q., Mills, K. A., & Price, N. R. (2004). Phosphine resistance in stored-product insects collected from various grain storage facilities in Morocco. *Journal of Stored Products Research*, 40(3), 241–249. [https://doi.org/10.1016/S0022-474X\(03\)00012-2](https://doi.org/10.1016/S0022-474X(03)00012-2)
- Benzi, V., Stefanazzi, N., Murray, A. P., Werdin González, J. O., & Ferrero, A. (2014). Composition, Repellent, and Insecticidal Activities of Two South American Plants against the Stored Grain Pests *Tribolium castaneum* and *Tribolium confusum* (Coleoptera: Tenebrionidae). *ISRN Entomology*, 2014, 1–5. <https://doi.org/10.1155/2014/175827>
- Berg, A. D. (1968). Malnutrition and national development. *Journal of Tropical Pediatrics*, 14(3), 116–123.

<https://doi.org/10.1093/oxfordjournals.tropej.a057292>

- Bergerson, O., & Wool, D. (1988). The process of adaptation of flour beetles to new environments. *Genetica*, 77(1), 3–13.
<https://doi.org/10.1007/BF00058546>
- Bett, P. K., Deng, A. L., Ogendo, J. O., Kariuki, S. T., Kamatenesi-Mugisha, M., Mihale, J. M., & Torto, B. (2016). Chemical composition of *Cupressus lusitanica* and *Eucalyptus saligna* leaf essential oils and bioactivity against major insect pests of stored food grains. *Industrial Crops and Products*, 82, 51–62.
<https://doi.org/10.1016/j.indcrop.2015.12.009>
- Bhaskar Mi, B., & Tripathi, S. P. (2011). Repellent Activity of Plant Derived Essential Oils against *Sitophilous oryzae* (Linnaeus) and *Tribolium castaneum* (Herbst). *Singapore Journal of Scientific Research*, Vol. 1, pp. 173–178. <https://doi.org/10.3923/sjsres.2011.173.178>
- Bloomquist, J. R. (2001). GABA and glutamate receptors as biochemical sites for insecticide action. In *Biochemical sites of insecticide action and resistance* (pp. 17–41). Berlin, Heidelberg: Springer.
- Bora, K. S., & Sharma, A. (2011). The genus *Artemisia*: A comprehensive review. *Pharmaceutical Biology*, 49(1), 101–109.
<https://doi.org/10.3109/13880209.2010.497815>
- Boyer, S., Zhang, H., & Lempérière, G. (2012). A review of control methods and resistance mechanisms in stored-product insects. *Bulletin of Entomological Research*, 102(2), 213–229.
<https://doi.org/10.1017/S0007485311000654>
- Brari, J., & Thakur, D. R. (2018). Larvicidal effects of eight essential oils against *Plodia interpunctella* and *Tribolium castaneum*, serious pests of stored products worldwide. *Journal of Entomological and Acarological Research*, 50(1), 738–742. <https://doi.org/10.4081/jear.2018.7469>
- Buege, J. A., & Aust, S. D. (1978). [30] Microsomal lipid peroxidation. In *Methods in Enzymology* (pp. 302–310). Cambridge, Massachusetts,

United States: Academic Press, Elsevier.

- Bui, D. T., Nicolas, J., Maksimenko, A., Desmaële, D., & Couvreur, P. (2014). Multifunctional squalene-based prodrug nanoparticles for targeted cancer therapy. *Chemical Communications*, 50(40), 5336–5338. <https://doi.org/10.1039/c3cc47427e>
- Butt, M. S., Nasir, M., Akhtar, S., & Sharif, K. (2004). Effect OF Moisture and Packaging on the Shelf Life of Wheat Flour. *Internet Journal of Food Safety*, 4, 1–6.
- Caballero-Gallardo, K., Olivero-Verbel, J., & Stashenko, E. E. (2012). Repellency and toxicity of essential oils from *Cymbopogon martinii*, *Cymbopogon flexuosus* and *Lippia origanoides* cultivated in Colombia against *Tribolium castaneum*. *Journal of Stored Products Research*, 50, 62–65. <https://doi.org/10.1016/j.jspr.2012.05.002>
- Cao, J. Q., Guo, S. S., Wang, Y., Pang, X., Geng, Z. F., & Du, S. S. (2018). Contact toxicity and repellency of the essential oils of *Evodia lenticellata* Huang and *Evodia rutaecarpa* (Juss.) Benth. Leaves against three stored product insects. *Journal of Oleo Science*, 67(8), 1027–1034. <https://doi.org/10.5650/jos.ess17251>
- Cardiet, G., Fuzeau, B., Barreau, C., & Fleurat-Lessard, F. (2011). Contact and fumigant toxicity of some essential oil constituents against a grain insect pest *Sitophilus oryzae* and two fungi, *Aspergillus westerdijkiae* and *Fusarium graminearum*. *Journal of Pest Science*, 85(3), 351–358. <https://doi.org/10.1007/s10340-011-0400-3>
- Chakravarti, A. K. (1974). Regional preference for food : Some aspects of food habit patterns in India. *Canadian Geographer/Le Géographe Canadien*, 18(4), 395–410.
- Champ, B. R., & Campbell-Brown, M. J. (1970). Insecticide resistance in Australian *Tribolium castaneum* (Herbst)-I. A test method for detecting insecticide resistance. *Journal of Stored Products Research*, 6(1), 53–70. [https://doi.org/10.1016/0022-474X\(70\)90027-5](https://doi.org/10.1016/0022-474X(70)90027-5)

- Chaubey, M. K. (2007). Insecticidal activity of *Trachyspermum ammi* (Umbelliferae), *Anethum graveolens* (Umbelliferae) and *Nigella sativa* (Ranunculaceae) essential oils against stored-product beetle *Tribolium castaneum* Herbst (Coleoptera: Tenebrionidae). *African Journal of Agricultural Research*, 2(11), 596–600.
- Chauhan, N., Kumar, P., Mishra, S., Verma, S., Malik, A., & Sharma, S. (2015). Insecticidal activity of *Jatropha curcas* extracts against housefly, *Musca domestica*. *Environmental Science and Pollution Research*, 22(19), 14793–14800. <https://doi.org/10.1007/s11356-015-4686-1>
- Clevenger, J. F. (1928). Apparatus for the determination of volatile oil. *The Journal of the American Pharmaceutical Association* 1912, 17(4), 345–349. <https://doi.org/10.1002/jps.3080170407>
- Cohen, E. (1986). Glutathione-S-transferase activity and its induction in several strains of *Tribolium castaneum*. *Entomologia Experimentalis et Applicata*, 41(1), 39–44. <https://doi.org/10.1111/j.1570-7458.1986.tb02169.x>
- Conti, B., Canale, A., Cioni, P. L., Flamini, G., & Rifici, A. (2011). *Hyptis suaveolens* and *Hyptis spicigera* (Lamiaceae) essential oils: Qualitative analysis, contact toxicity and repellent activity against *Sitophilus granarius* (L.) (Coleoptera: Dryophthoridae). *Journal of Pest Science*, 84(2), 219–228. <https://doi.org/10.1007/s10340-010-0343-0>
- Cosimi, S., Rossi, E., Cioni, P. L., & Canale, A. (2009). Bioactivity and qualitative analysis of some essential oils from Mediterranean plants against stored-product pests: Evaluation of repellency against *Sitophilus zeamais* Motschulsky, *Cryptolestes ferrugineus* (Stephens) and *Tenebrio molitor* (L.). *Journal of Stored Products Research*, 45(2), 125–132. <https://doi.org/10.1016/j.jspr.2008.10.002>
- Daft, J. L. (1991). Fumigants and related chemicals in foods: Review of residue findings, contamination sources, and analytical methods. *Science of the Total Environment*, The, 100, 501–518. [https://doi.org/10.1016/0048-9697\(91\)90390-Z](https://doi.org/10.1016/0048-9697(91)90390-Z)

- Dahham, S. S., Tabana, Y. M., Iqbal, M. A., Ahamed, M. B. K., Ezzat, M. O., Majid, A. S. A., & Majid, A. M. S. A. (2015). The anticancer, antioxidant and antimicrobial properties of the sesquiterpene β -caryophyllene from the essential oil of *Aquilaria crassna*. *Molecules*, *20*(7), 11808–11829. <https://doi.org/10.3390/molecules200711808>
- Dangash, A. (2017). *Amendment of artemisinin content in Artemisia annua L. through in vivo and in vitro approach* (The Maharaja Sayajirao University of Baroda). <https://doi.org/10.1192/bjp.112.483.211-a>
- Das, S. K. (2014). Recent Development and future of Botanical Pesticides in India. *Popular Kheti*, *2*(2), 93–99.
- Devi, M. B., & Devi, N. V. (2015). Biology of rust-red flour beetle, *Tribolium castaneum* (Herbst) (Tenebrionidae: Coleoptera). *Indian Journal of Entomology*, *77*(1), 81. <https://doi.org/10.5958/0974-8172.2015.00015.2>
- Dorman, H. J. D., & Deans, S. G. (2000). Antimicrobial agents from plants: Antibacterial activity of plant volatile oils. *Journal of Applied Microbiology*, *88*(2), 308–316. <https://doi.org/10.1046/j.1365-2672.2000.00969.x>
- Driscoll, R., Longstaff, B. C., & Beckett, S. (2000). Prediction of insect populations in grain storage. *Journal of Stored Products Research*, *36*(2), 131–151. [https://doi.org/10.1016/S0022-474X\(99\)00032-6](https://doi.org/10.1016/S0022-474X(99)00032-6)
- Duke, S. O.; Paul, R. N. (1993). Development and Fine Structure of the Glandular Trichomes of *Artemisia annua* L. *International Journal of Plant Sciences*, *154*(1), 107–118. [doi/10.1086/297096](https://doi.org/10.1086/297096).
- Durden, K., Sellars, S., Cowell, B., Brown, J. J., & Pszczolkowski, M. A. (2011). *Artemisia annua* extracts, artemisinin and 1,8-cineole, prevent fruit infestation by a major, cosmopolitan pest of apples. *Pharmaceutical Biology*, *49*(6), 563–568. <https://doi.org/10.3109/13880209.2010.528433>
- Eastman, R. T., & Fidock, D. A. (2009). Artemisinin-based combination therapies: A vital tool in efforts to eliminate malaria. *Nature Reviews Microbiology*, *7*(12), 864–874. <https://doi.org/10.1038/nrmicro2239>

- El-Mofty, M. M.; Sakr, S. A.; Osman, S. I.; Toulan, B. A. (1989).
Carcinogenic effect of biscuits made of flour infested with *Tribolium castaneum* in *Bufo regularis*. *Oncology*, 46(1), 63–65.
- Ellman, G. L.; Courtney, K. D.; Andres Jr, V.; Featherstone, R. M. (1961). A new and rapid colorimetric determination of acetylcholinesterase activity. *Biochemical Pharmacology*, 7(2), 88-95.
- Enan, E. E. (2005). Molecular and pharmacological analysis of an octopamine receptor from American cockroach and fruit fly in response to plant essential oils. *Archives of Insect Biochemistry and Physiology*, 59(3), 161–171. <https://doi.org/10.1002/arch.20076>
- Evans, P. D. (1984). Studies on the mode of action of octopamine, 5-hydroxytryptamine and proctolin on a myogenic rhythm in the locust. *Journal of Experimental Biology*, 110(1), 231–251.
- Falowo, A. B., Mukumbo, F. E., & Muchenje, V. (2019). Phytochemical Constituents and Antioxidant Activity of *Artemisia Afra* and *Bidens Pilosa* Essential Oil in Ground Pork. *Journal of Essential Oil-Bearing Plants*, 22(1), 176–186. <https://doi.org/10.1080/0972060X.2019.1574212>
- FAO. (2013). *The State of Food Insecurity in the World The multiple dimensions of food security 2013*.
- FAO. (2014). *The State of Food and Agriculture*.
- Farrar, R. R.; Barbour, J. D.; Kennedy, G. G. (1989). Quantifying Food Consumption and Growth in Insects. *Annals of the Entomological Society of America*, 82(5), 593–598. <https://doi.org/10.1093/aesa/82.5.593>
- Fedina, T. Y., & Lewis, S. M. (2008). An integrative view of sexual selection in *Tribolium* flour beetles. *Biological Reviews*, 83(2), 151–171. <https://doi.org/10.1111/j.1469-185X.2008.00037.x>
- Fields, P. G.; White, N. D. (2002). Alternatives to methyl bromide treatments for stored-product and quarantine insect. *Annual Review of Entomology*, 47(1), 331–359. <https://doi.org/10.1227/01.NEU.0000103446.26057.78>
- Gawande, S. B., & Patil, D. I. (2015). A Review on Causes for Damaged

- Sorghum and Corn Grains. *PRATIBHA: International Journal Of Science, Spirituality, Business And Technology (IJSSBT)*, 3(2), 5–9.
<https://doi.org/10.1017/CBO9781107415324.004>
- Gonzalez-Coloma, A., Bailen, M., Diaz, C. E., Fraga, B. M., Martínez-Díaz, R., Zuñiga, G. E., ... Burillo, J. (2012). Major components of Spanish cultivated *Artemisia absinthium* populations: Antifeedant, antiparasitic, and antioxidant effects. *Industrial Crops and Products*, 37(1), 401–407.
<https://doi.org/10.1016/j.indcrop.2011.12.025>
- Good, N. E. (1933). Biology of the flour beetles, *Tribolium confusum* Duv. and *T. ferrugineum* Fab. *Journal of Agricultural Research*, 46(4), 327–334.
- Gorham, J. R. (1979). Human health of insects in food. *Annual Review of Entomology*, 24(1), 209–224.
- Guo, S., Zhang, W., Liang, J., You, C., Geng, Z., Wang, C., & Du, S. (2016). Contact and Repellent Activities of the Essential Oil from *Juniperus formosana* against Two Stored Product Insects. *Molecules*, 21(4).
<https://doi.org/10.3390/molecules21040504>
- Gupta, P. N. Sen. (1980). *Food consumption and nutrition of regional tribes of India Regional tribes of India †*. 0244(May).
<https://doi.org/10.1080/03670244.1980.9990587>
- Habig, W. H.; Pabst, M. J.; Jakoby, W. B. (1974). Glutathione S-transferases the first enzymatic step in mercapturic acid formation. *Journal of Biological Chemistry*, 249(22), 7130–7139.
- Hagstrum, D. (2017). Infestation records. In *Atlas of stored-product insects and mites* (pp. 474–483). Minnesota: Elsevier.
- Haider, S. Z., Mohan, M., Pandey, A. K., & Singh, P. (2017). Use of *Tanacetum tomentosum* and *Ta. dolichophyllum* essential oils as botanical repellents and insecticidal agents against storage pest *Tribolium castaneum* (Coleoptera: Tenebrionidae). *Entomological Research*, 47(5), 318–327. <https://doi.org/10.1111/1748-5967>

- Haider, S. Z., Mohan, M., Pandey, A. K., & Singh, P. (2015). Repellent and fumigant activities of *Tanacetum nubigenum* Wallich. ex DC essential oils against *Tribolium castaneum* (Herbst) (Coleoptera: Tenebrionidae). *Journal of Oleo Science*, 64(8), 895–903.
<https://doi.org/10.5650/jos.ess15094>
- Haridasan, P., Gokuldas, M., & Ajaykumar, A. P. (2017). Antifeedant Effects of Vitex Negundo L. Leaf Extracts on the Stored Product Pest, *Tribolium castaneum* H. (Coleoptera: Tenebrionidae). *International Journal of Pharmacy and Pharmaceutical Sciences*, 9(3), 17.
<https://doi.org/10.22159/ijpps.2017v9i3.15600>
- Hashem, A. S., Awadalla, S. S., Zayed, G. M., Maggi, F., & Benelli, G. (2018). *Pimpinella anisum* essential oil nanoemulsions against *Tribolium castaneum*—insecticidal activity and mode of action. *Environmental Science and Pollution Research*, 25(19), 18802–18812.
<https://doi.org/10.1007/s11356-018-2068-1>
- Hasspieler, B. M., Arnason, J. T., & Downe, A. E. R. (1990). Modes of action of the plant-derived phototoxin α -terthienyl in mosquito larvae. *Pesticide Biochemistry and Physiology*, 38(1), 41–47.
[https://doi.org/10.1016/0048-3575\(90\)90146-S](https://doi.org/10.1016/0048-3575(90)90146-S)
- Hernandez-Lambraño, R., Pajaro-Castro, N., Caballero-Gallardo, K., Stashenko, E., & Olivero-Verbel, J. (2015). Essential oils from plants of the genus *Cymbopogon* as natural insecticides to control stored product pests. *Journal of Stored Products Research*, 62, 81–83.
<https://doi.org/10.1016/j.jspr.2015.04.004>
- Hertog, M. L. A. T. M., Uysal, I., McCarthy, U., Verlinden, B. M., & Nicolai, B. M. (2014). Shelf life modelling for first-expired-first-out warehouse management. *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 372(2017).
<https://doi.org/10.1098/rsta.2013.0306>
- Hodges, Rick; Robinson, R' Hall, D. (1995). Quinone Contamination of Dehusked Rice by *Tribolium castaneum* (Herbst) (Coleoptera :

- Tenebrionidae). *Journal of Stored Products Research*, 32(1), 31–37.
[https://doi.org/10.1016/0022-474X\(95\)00036-7](https://doi.org/10.1016/0022-474X(95)00036-7)
- Hu, J., Wang, W., Dai, J., & Zhu, L. (2019). Chemical composition and biological activity against *Tribolium castaneum* (Coleoptera: Tenebrionidae) of *Artemisia brachyloba* essential oil. *Industrial Crops and Products*, 128, 29–37. <https://doi.org/10.1016/j.indcrop.2018.10.076>
- Huang, Y., & Ho, S. H. (1998). Toxicity and antifeedant activities of cinnamaldehyde against the grain storage insects, *Tribolium castaneum* (Herbst) and *Sitophilus zeamais* Motsch. *Journal of Stored Products Research*, 34(1), 11–17. [https://doi.org/10.1016/S0022-474X\(97\)00038-6](https://doi.org/10.1016/S0022-474X(97)00038-6)
- Huang, Y., Tan, J. M. W. L., Kini, R. M., & Ho, S. H. (1997). Toxic and antifeedant action of nutmeg oil against *Tribolium castaneum* (Herbst) and *Sitophilus zeamais* Motsch. *Journal of Stored Products Research*, 33(4), 289–298. [https://doi.org/10.1016/S0022-474X\(97\)00009-X](https://doi.org/10.1016/S0022-474X(97)00009-X)
- Huang, Yan, Ho, S. H., Lee, H. C., & Yap, Y. L. (2002). Insecticidal properties of eugenol, isoeugenol and methyleugenol and their effects on nutrition of *Sitophilus zeamais* Motsch. (Coleoptera: Curculionidae) and *Tribolium castaneum* (Herbst) (Coleoptera: Tenebrionidae). *Journal of Stored Products Research*, 38(5), 403–412.
[https://doi.org/10.1016/S0022-474X\(01\)00042-X](https://doi.org/10.1016/S0022-474X(01)00042-X)
- Huang, Z., Shi, P., Dai, J., & Du, J. (2004). Protein metabolism in *Spodoptera litura* (F.) is influenced by the botanical insecticide azadirachtin. *Pesticide Biochemistry and Physiology*, 80(2), 85–93.
<https://doi.org/10.1016/j.pestbp.2004.07.001>
- Iram, N., Arshad, M., & Akhter, N. (2013). Evaluation of Botanical and Synthetic Insecticide for the Control of *Tribolium castaneum* (Herbst) (Coleoptera: Tenebrionidae). *Bioassay*, 8(3), 1–10. Retrieved from www.bioassay.org.br
- Isman B., M. (2000). Plant essential Oils for Pest and disease management. *Crop Protection*, 19, 603–608. Retrieved from <http://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.456.3876&rep>

=rep1&type=pdf

- Jagadeesan, R., Collins, P. J., Daglish, G. J., Ebert, P. R., & Schlipalius, D. I. (2012). Phosphine resistance in the rust red flour beetle, *Tribolium castaneum* (Coleoptera: Tenebrionidae): Inheritance, gene interactions and fitness costs. *PLoS ONE*, 7(2).
<https://doi.org/10.1371/journal.pone.0031582>
- Janssen, A. M., Vendsen, A. B. S., & Laboratories, G. (1987). Antimicrobial activities of essential oils. *Pharmaceutisch Weekblad*, 9(4), 193–197.
- Jaya, Singh, P., Prakash, B., & Dubey, N. K. (2014). Insecticidal activity of *Ageratum conyzoides* L., *Coleus aromaticus* benth. And *Hyptis suaveolens* (L.) poit essential oils as fumigant against storage grain insect *Tribolium castaneum* herbst. *Journal of Food Science and Technology*, 51(9), 2210–2215. <https://doi.org/10.1007/s13197-012-0698-8>
- Jayakumar, M.; Arivoli, S.; Raveen, R.; Tennyson, S. (2017). Repellent activity and fumigant toxicity of a few plant oils against the adult rice weevil *Sitophilus oryzae* Linnaeus 1763 (Coleoptera: Curculionidae). *Journal of Entomology and Zoology Studies*, 5(2), 324–335.
- Jeyasankar, A., Chennaiyan, V., & Chinnamani, T. (2016). Evaluation of five essential plant oils as a source of Repellent and larvicidal activities against larvae of *Tribolium castaneum* (Herbst) (Coleoptera: Tenebrionidae). *Journal of Entomology*, 13(3), 98–103.
<https://doi.org/10.3923/je.2016.98.103>
- Jollow, D. J.; Mitchell, J. R.; Potter, W. Z.; Davis, D. C.; Gillette, J. R.; Brodie, B. B. (1973). Acetaminophen-induced hepatic necrosis. II. Role of covalent binding in vivo. *Journal of Pharmacology and Experimental Therapeutics*, 187, 195–202.
- Jood, S., & Kapoor, A. C. (1993). Protein and uric acid contents of cereal grains as affected by insect infestation. *Food Chemistry*, 46(2), 143–146.
[https://doi.org/10.1016/0308-8146\(93\)90027-D](https://doi.org/10.1016/0308-8146(93)90027-D)
- Jood, S., Kapoor, A. C., & Singh, R. (1993). Available carbohydrates of cereal

- grains as affected by storage and insect infestation. *Plant Foods for Human Nutrition*, 43(1), 45–54. <https://doi.org/10.1007/BF01088095>
- Kanda, D., Kaur, S., & Koul, O. (2017). A comparative study of monoterpenoids and phenylpropanoids from essential oils against stored grain insects: acute toxins or feeding deterrents. *Journal of Pest Science*, 90(2), 531–545. <https://doi.org/10.1007/s10340-016-0800-5>
- Karunakaran, C., Jayas, D. S., & White, N. D. G. (2004). Identification of wheat kernels damaged by the red flour beetle using X-ray images. *Biosystems Engineering*, 87(3), 267–274. <https://doi.org/10.1016/j.biosystemseng.2003.12.002>
- Khani, A., & Heydarian, M. (2014). Fumigant and repellent properties of sesquiterpene-rich essential oil from *Teucrium polium* subsp. capitatum (L.). *Asian Pacific Journal of Tropical Medicine*, 7(12), 956–961. [https://doi.org/10.1016/S1995-7645\(14\)60169-3](https://doi.org/10.1016/S1995-7645(14)60169-3)
- Kim, S. Il, & Lee, D. W. (2014). Toxicity of basil and orange essential oils and their components against two coleopteran stored products insect pests. *Journal of Asia-Pacific Entomology*, 17(1), 13–17. <https://doi.org/10.1016/j.aspen.2013.09.002>
- Kim, S. Il, Park, C., Ohh, M. H., Cho, H. C., & Ahn, Y. J. (2002). Contact and fumigant activities of aromatic plant extracts and essential oils against *Lasioderma serricorne* (Coleoptera: Anobiidae). *Journal of Stored Products Research*, 39(1), 11–19. [https://doi.org/10.1016/S0022-474X\(02\)00013-9](https://doi.org/10.1016/S0022-474X(02)00013-9)
- Klayman, D. L. (1985). Qinghaosu (artemisinin): An antimalarial drug from China. *Science*, 228(4703), 1049–1055. <https://doi.org/10.1126/science.3887571>
- Koodalingam, A., Mullainadhan, P., & Arumugam, M. (2011). Effects of extract of soapnut *Sapindus emarginatus* on esterases and phosphatases of the vector mosquito, *Aedes aegypti* (Diptera: Culicidae). *Acta Tropica*, 118(1), 27–36. <https://doi.org/10.1016/j.actatropica.2011.01.003>

- Kostyukovsky, M., Rafaeli, A., Gileadi, C., Demchenko, N., & Shaaya, E. (2002). Activation of octopaminergic receptors by essential oil constituents isolated from aromatic plants: Possible mode of action against insect pests. *Pest Management Science*, *58*(11), 1101–1106. <https://doi.org/10.1002/ps.548>
- Kumar, R., Kumar, A., Prasa, C., Dubey, N., & Samant, R. (2008). Insecticidal activity *Aegle marmelos* (L.) Correa essential oil against four stored grain insect pests. *Internation Journal of Food Safety*, *10*, 39–49. Retrieved from <http://www.internetjfs.org/articles/ijfsv10-7.pdf>
- KUMAR, R., SRIVASTAVA, M., & DUBEY, N. K. (2006). Evaluation of *Cymbopogon martinii* Oil Extract for Control of Postharvest Insect Deterioration in Cereals and Legumes. *Journal of Food Protection*, *70*(1), 172–178. <https://doi.org/10.4315/0362-028x-70.1.172>
- Ladisch, R. K.; Ladisch, S. K.; Howe, P. M. (1967). Quinoid secretions in grain and flour beetles. *In Proceedings of the Pennsylvania Academy of Science*, 213–219. Pennsylvania: Pennsylvania Academy of Science.
- Lale, N. E. S. (1992). A laboratory study of the comparative toxicity of products from three spices to the maize weevil. *Postharvest Biology and Technology*, *2*(1), 61–64.
- Langeveld, W. T., Veldhuizen, E. J. A., & Burt, S. A. (2014). Synergy between essential oil components and antibiotics: A review. *Critical Reviews in Microbiology*, *40*(1), 76–94. <https://doi.org/10.3109/1040841X.2013.763219>
- Lee, B. H., Annis, P. C., Tumaalii, F., & Choi, W. S. (2004). Fumigant toxicity of essential oils from the Myrtaceae family and 1,8-cineole against 3 major stored-grain insects. *Journal of Stored Products Research*, *40*(5), 553–564. <https://doi.org/10.1016/j.jspr.2003.09.001>
- Lee, B. H., Choi, W. S., Lee, S. E., & Park, B. S. (2001). Fumigant toxicity of essential oils and their constituent compounds towards the rice weevil, *Sitophilus oryzae* (L.). *Crop Protection*, *20*(4), 317–320. [https://doi.org/10.1016/S0261-2194\(00\)00158-7](https://doi.org/10.1016/S0261-2194(00)00158-7)

- Lee, H. K., & Lee, H. S. (2016). Toxicities of active constituent isolated from *Thymus vulgaris* flowers and its structural derivatives against *Tribolium castaneum* (Herbst). *Applied Biological Chemistry*, 59(6), 821–826.
<https://doi.org/10.1007/s13765-016-0230-3>
- Leelaja, B. C., Rajashekar, Y., & Rajendran, S. (2007). Detection of eggs of stored-product insects in flour with staining techniques. *Journal of Stored Products Research*, 43(3), 206–210.
<https://doi.org/10.1016/j.jspr.2006.05.003>
- Leonard LeCato, G. (1975). Red Flour Beetle: Population Growth on Diets of Corn, Wheat, Rice or Shelled Peanuts Supplemented with Eggs or Adults of the Indian Meal Moth. *Journal of Economic Entomology*, 68(6), 763–765. <https://doi.org/10.1093/jee/68.6.763>
- Lhaloui, S., Hagstrum, D. W., Keith, D. L., Holtzer, T. O., & Ball, H. J. (1988). Combined Influence of Temperature and Moisture on Red Flour Beetle (Coleoptera: Tenebrionidae) Reproduction on Whole Grain Wheat. *Journal of Economic Entomology*, 81(2), 488–489.
<https://doi.org/10.1093/jee/81.2.488>
- Li, W. Q.; Jiang, C. H.; Chu, S. S.; Zuo, M. X.; Liu, Z. L. (2010). Chemical composition and toxicity against *Sitophilus zeamais* and *Tribolium castaneum* of the essential oil of *Murraya exotica* aerial parts. *Molecules*, 15(8), 5831–5839. <https://doi.org/10.3390/molecules15085831>
- Li, Y. (2012). Qinghaosu (artemisinin): Chemistry and pharmacology. *Acta Pharmacologica Sinica*, 33(9), 1141–1146.
<https://doi.org/10.1038/aps.2012.104>
- Liang, J. Y., Guo, S. S., Zhang, W. J., Geng, Z. F., Deng, Z. W., Du, S. S., & Zhang, J. (2018). Fumigant and repellent activities of essential oil extracted from *Artemisia dubia* and its main compounds against two stored product pests. *Natural Product Research*, 32(10), 1234–1238.
<https://doi.org/10.1080/14786419.2017.1331227>
- Licciardello, F., Muratore, G., Suma, P., Russo, A., & Nerín, C. (2013). Effectiveness of a novel insect-repellent food packaging incorporating

- essential oils against the red flour beetle (*Tribolium castaneum*). *Innovative Food Science and Emerging Technologies*, 19, 173–180.
<https://doi.org/10.1016/j.ifset.2013.05.002>
- Liu, C. H., Mishra, A. K., Tan, R. X., Tang, C., Yang, H., & Shen, Y. F. (2006). Repellent and insecticidal activities of essential oils from *Artemisia princeps* and *Cinnamomum camphora* and their effect on seed germination of wheat and broad bean. *Bioresource Technology*, 97(15), 1969–1973. <https://doi.org/10.1016/j.biortech.2005.09.002>
- Liu, Z. L., & Ho, S. H. (1999). Bioactivity of the essential oil extracted from *Evodia rutaecarpa* Hook f. et Thomas against the grain storage insects, *Sitophilus zeamais* Motsch. and *Tribolium castaneum* (Herbst). *Journal of Stored Products Research*, 35(4), 317–328.
[https://doi.org/10.1016/S0022-474X\(99\)00015-6](https://doi.org/10.1016/S0022-474X(99)00015-6)
- Liu, Zhi Long, Chu, S. S., & Jiang, G. H. (2011a). Insecticidal activity and composition of essential oil of *Ostericum sieboldii* (Apiaceae) against *Sitophilus zeamais* and *Tribolium castaneum*. *Records of Natural Products*, 5(2), 74–81.
- Liu, Zhi Long, Chu, S. S., & Jiang, G. H. (2011b). Toxicity of *Schizonpeta multifida* essential oil and its constituent compounds towards two grain storage insects. *Journal of the Science of Food and Agriculture*, 91(5), 905–909. <https://doi.org/10.1002/jsfa.4263>
- López, M. D., Jordán, M. J., & Pascual-Villalobos, M. J. (2008). Toxic compounds in essential oils of coriander, caraway and basil active against stored rice pests. *Journal of Stored Products Research*, 44(3), 273–278.
<https://doi.org/10.1016/j.jspr.2008.02.005>
- Macedo, M. R., & Freire, M. D. G. M. (2011). Insect digestive enzymes as a target for pest control. *Invertebrate Survival Journal*, 8(2), 190–198.
<https://doi.org/10.1016/j.jlumin.2013.04.047>
- Mamedov, N. (2012). Medicinal Plants Studies: History, Challenges and Prospective. *Medicinal & Aromatic Plants*, 01(08), e133.
<https://doi.org/10.4172/2167-0412.1000e133>

- Manuwoto, S.; Mark Scriber, J. (1982). Consumption and Utilization of Three Maize Genotypes by the Southern Armyworm. *Journal of Economic Entomology*, 75(2), 163–167. <https://doi.org/10.1093/jee/75.2.163>
- Masotti, V., Juteau, F., Bessière, J. M., & Viano, J. (2003). Seasonal and Phenological Variations of the Essential Oil from the Narrow Endemic Species *Artemisia molinieri* and Its Biological Activities. *Journal of Agricultural and Food Chemistry*, 51(24), 7115–7121. <https://doi.org/10.1021/jf034621y>
- Matsuda, K.; Buckingham, S. D.; Kleier, D.; Rauh, J. J.; Grauso, M.; Sattelle, D. B. (2001). Neonicotinoids: insecticides acting on insect nicotinic acetylcholine receptors. *Trends in Pharmacological Sciences*, 22(11), 573–580.
- Mezzomo, N., Mileo, B. R., Friedrich, M. T., Martínez, J., & Ferreira, S. R. S. (2010). Supercritical fluid extraction of peach (*Prunus persica*) almond oil: Process yield and extract composition. *Bioresource Technology*, 101(14), 5622–5632. <https://doi.org/10.1016/j.biortech.2010.02.020>
- Micu, L. M., & Petanec, D. I. (2011). Research regarding the changes in the content of micro-elements (Cu, Zn, Mn) in stored wheat after infestation by *Rhizopertha dominica* F. *Rom. Agri. Res*, 28, 2067–5720.
- Mills, C., Cleary, B. V., Walsh, J. J., & Gilmer, J. F. (2004). Inhibition of acetylcholinesterase by Tea Tree oil. *Journal of Pharmacy and Pharmacology*, 56(3), 375–379. <https://doi.org/10.1211/0022357022773>
- Mishra, A. K., & Dubey, N. K. (1994). Evaluation of Some Essential Oils for Their Toxicity against Fungi Causing Deterioration of Stored Food Commodities. *Applied and Environmental Microbiology*, 60(4), 1101–1105.
- Mishra, B. B., Tripathi, S. P., & Tripathi, C. P. M. (2012). Response of *Tribolium castaneum* (Coleoptera: Tenebrionidae) and *Sitophilus oryzae* (Coleoptera: Curculionidae) to potential insecticide derived from essential oil of *Mentha arvensis* leaves. *Biological Agriculture and Horticulture*, 28(1), 34–40.

<https://doi.org/10.1080/01448765.2012.662792>

Mishra, B. B., Tripathi, S. P., & Tripathi, C. P. M. (2016). *Impact of Syzygium aromaticum (L.) essential oil as fumigant against Tribolium castaneum (Coleoptera: Tenebrionidae)*. *Journal of Entomology and Zoological Studies*, 4(6), 811–816.

Modgil, R., & Mehta, U. (1996). Effect of *Callosobruchus chinensis* (L.) (Coleoptera: Bruchidae) on carbohydrate content of chickpea, green gram and pigeon pea. *Nahrung - Food*, 40(1), 41–43.
<https://doi.org/10.1002/food.19960400111>

Moleyar, V.; Narasimham, P. (1986). Antifungal activity of some essential oil components. *Food Microbiology*, 3(4), 331–336.

Mossa, A. T. H. (2016). Green Pesticides: Essential oils as biopesticides in insect-pest management. *Journal of Environmental Science and Technology*, 9(5), 354–378. <https://doi.org/10.3923/jest.2016.354.378>

Müller, O., & Krawinkel, M. (2005). Malnutrition and health in developing countries. *Cmaj*, 173(3), 279–286.
<https://doi.org/10.1177/019394599101300605>

Nathanson, M. (1975). The effect of resource limitation on competing populations of flour beetles, *Tribolium* spp. (Coleoptera, Tenebrionidae). *Bulletin of Entomological Research*, 65(1), 1–12.
<https://doi.org/10.1017/S0007485300005691>

Nattudurai, G., Irudayaraj, S. S., Paulraj, M. G., Baskar, K., & Ignacimuthu, S. (2015). Insecticidal and Repellent Activities of *Toddalia asiatica* (L.) Lam. Extracts against Three Major Stored Product Pests. *Entomology, Ornithology and Herpetology: Current Research*, 04(02), 1–5.
<https://doi.org/10.4172/2161-0983.1000148>

Negahban, M., Moharramipour, S., & Sefidkon, F. (2007). Fumigant toxicity of essential oil from *Artemisia sieberi* Besser against three stored-product insects. *Journal of Stored Products Research*, 43(2), 123–128.
<https://doi.org/10.1016/j.jspr.2006.02.002>

- Nenaah, G. E. (2014). Chemical composition, toxicity and growth inhibitory activities of essential oils of three *Achillea* species and their nano-emulsions against *Tribolium castaneum* (Herbst). *Industrial Crops and Products*, 53(February 2014), 252–260.
<https://doi.org/10.1016/j.indcrop.2013.12.042>
- Ni, L., Acharya, K., Hao, X., & Li, S. (2012). Isolation and identification of an anti-algal compound from *Artemisia annua* and mechanisms of inhibitory effect on algae. *Chemosphere*, 88(9), 1051–1057.
<https://doi.org/10.1016/j.chemosphere.2012.05.009>
- NITI Aayog. (2015). Raising Agricultural Productivity and Making Farming Remunerative for Farmers. *An Occasional Paper, GoI*, (December).
- Obeng-Ofori, D., & Reichmuth, C. (1997). Bioactivity of eugenol, a major component of essential oil of *Ocimum suave* (wild.) against four species of stored-product coleoptera. *International Journal of Pest Management*, 43(1), 89–94. <https://doi.org/10.1080/096708797229040>
- Obeng-Ofori, D., Reichmuth, C. H., Bekele, A. J., & Hassanali, A. (1998). Toxicity and protectant potential of camphor, a major component of essential oil of *Ocimum kilimandscharicum*, against four stored product beetles. *International Journal of Pest Management*, 44(4), 203–209.
<https://doi.org/10.1080/096708798228112>
- Ogendo, J. O., Kostyukovsky, M., Ravid, U., Matasyoh, J. C., Deng, A. L., Omolo, E. O., ... Shaaya, E. (2008). Bioactivity of *Ocimum gratissimum* L. oil and two of its constituents against five insect pests attacking stored food products. *Journal of Stored Products Research*, 44(4), 328–334.
<https://doi.org/10.1016/j.jspr.2008.02.009>
- Okonkwo, E. U., & Okoye, W. I. (1996). The efficacy of four seed powders and the essential oils as protectants of cowpea and maize grains against infestation by *Callosobruchus maculatus* (Fabricus)(Coleoptera: Bruchidae) and *Sitophilus zeamais* (Motschulsky)(Coleoptera: Curculionidae) in Nigeria. *International Journal of Pest Management*, 42(3), 143–146. <https://doi.org/10.1080/09670879609371985>

- Okwute, S. K. (2012). Plants as potential sources of pesticidal agents: a review. In *Pesticides—Advances in chemical and botanical pesticides* (pp. 207–232). <https://doi.org/http://dx.doi.org/10.5772/46225>
- Olivero-Verbel, Jesús, Nerio, L. S., & Stashenko, E. E. (2010). Bioactivity against *Tribolium castaneum* Herbst (Coleoptera: Tenebrionidae) of *Cymbopogon citratus* and *Eucalyptus citriodora* essential oils grown in Colombia. *Pest Management Science*, 66(6), 664–668. <https://doi.org/10.1002/ps.1927>
- Olivero-Verbel, Jesus, Tirado-Ballestas, I., Caballero-Gallardo, K., & Stashenko, E. E. (2013). Essential oils applied to the food act as repellents toward *Tribolium castaneum*. *Journal of Stored Products Research*, 55, 145–147. <https://doi.org/10.1016/j.jspr.2013.09.003>
- Olmedo, R., Herrera, J. M., Lucini, E. I., Zunino, M. P., Pizzolitto, R. P., Dambolena, J. S., & Zygadlo, J. A. (2015). Essential oil of *Tagetes filifolia* against the flour beetle *Tribolium castaneum* and its relation to acetylcholinesterase activity and lipid peroxidation. *Agriscientia*, 32(2), 113–121.
- Opit, G. P., Phillips, T. W., Aikins, M. J., & Hasan, M. M. (2012). Phosphine resistance in *Tribolium castaneum* and *Rhyzopertha dominica* from stored wheat in Oklahoma. *Journal of Economic Entomology*, 105(4), 1107–1114. <https://doi.org/10.1603/ec12064>
- Owusu, E. O. (2001). Effect of some Ghanaian plant components on control of two stored-product insect pests of cereals. *Journal of Stored Products Research*, 37(1), 85–91.
- Padín, S., Dal Bello, G., & Fabrizio, M. (2002). Grain loss caused by *Tribolium castaneum*, *Sitophilus oryzae* and *Acanthoscelides obtectus* in stored durum wheat and beans treated with *Beauveria bassiana*. *Journal of Stored Products Research*, 38(1), 69–74. [https://doi.org/10.1016/S0022-474X\(00\)00046-1](https://doi.org/10.1016/S0022-474X(00)00046-1)
- Pandey, A. K.; Kumar, P.; Singh, P.; Tripathi, N. N.; Bajpai, V. K. (2017). Essential oils: sources of antimicrobials and food preservatives. *Frontiers*

- in Microbiology*, 7(2161), 1–14.
<https://doi.org/10.3390/medicines4030068>
- Pandey, M. (2009). Indian Agriculture – an Introduction. *Apcaem*, (February), 1–39. Retrieved from <http://un-csam.org/Activities Files/A0902/in-p.pdf>
- Pant, M., Dubey, S., Patanjali, P. K., Naik, S. N., & Sharma, S. (2014). Insecticidal activity of eucalyptus oil nanoemulsion with karanja and jatropha aqueous filtrates. *International Biodeterioration and Biodegradation*, 91, 119–127. <https://doi.org/10.1016/j.ibiod.2013.11.019>
- Pingale, S. V., Rao, M. N., & Swaminathan, M. (1954). Effect of insect infestation on stored grain. I.—studies on soft wheat. *Journal of the Science of Food and Agriculture*, 5(1), 51–54.
<https://doi.org/10.1002/jsfa.2740050108>
- Priestley, C. M., Williamson, E. M., Wafford, K. A., & Sattelle, D. B. (2003). Thymol, a constituent of thyme essential oil, is a positive allosteric modulator of human GABA A receptors and a homo-oligomeric GABA receptor from *Drosophila melanogaster*. *British Journal of Pharmacology*, 140(8), 1363–1372.
<https://doi.org/10.1038/sj.bjp.0705542>
- Pugazhvendan, S., Elumalai, K., Ross, P. R., & Soundararajan, M. (2009). Repellent activity of chosen plant species against *Tribolium castaneum*. *World Journal of Zoology*, 4(3), 188–190. Retrieved from [http://www.idosi.org/wjz/wjz4\(3\)2009/7.pdf](http://www.idosi.org/wjz/wjz4(3)2009/7.pdf)
- Rajashekar, Y., Raghavendra, A., & Bakthavatsalam, N. (2014). Acetylcholinesterase inhibition by biofumigant (Coumaran) from leaves of *Lantana camara* in stored grain and household insect pests. *BioMed Research International*, 2014, 1–6. <https://doi.org/10.1155/2014/187019>
- Rattan, R. S. (2010). Mechanism of action of insecticidal secondary metabolites of plant origin. *Crop Protection*, 29(9), 913–920.
<https://doi.org/10.1016/j.cropro.2010.05.008>
- Reichling, J., Schnitzler, P., Suschke, U., & Saller, R. (2009). Essential oils of

aromatic plants with antibacterial, antifungal, antiviral, and cytotoxic properties - An overview. *Forschende Komplementarmedizin*, 16(2), 79–90. <https://doi.org/10.1159/000207196>

Rimi, S. A., Hossain, S., Islam, S., Islam, Z., Chhabi, S. B., & Islam, N. (2017). Bioactive Potentials of *Cleome viscosa* L. Extracts: Dose-mortality, Insect Repellency and Brine Shrimp Lethality. *Journal of Scientific Research*, 9(4), 375–382. <https://doi.org/10.3329/jsr.v9i4.32592>

Sakuma, M.; Fukumi, H. (1985). The linear track olfactometer: an assay device for taxes of the German cockroach, *Blattella germanica* (L.)(Dictyoptera: Blattellidae) toward their aggregation pheromone. *Applied Entomology and Zoology*, 20(4), 387–402.

Salem, N., Bachrouch, O., Sriti, J., Msaada, K., Khammassi, S., Hammami, M., ... Mediouni Ben Jemaa, J. (2017). Fumigant and repellent potentials of *Ricinus communis* and *Mentha pulegium* essential oils against *Tribolium castaneum* and *Lasioderma serricorne*. *International Journal of Food Properties*, 20, S2899–S2913. <https://doi.org/10.1080/10942912.2017.1382508>

Salimi Khorshidi, A., Storsley, J., Malunga, L. N., Thandapilly, S. J., & Ames, N. (2018). Advancing the science of wheat quality evaluation using nuclear magnetic resonance (NMR) and ultrasound-based techniques. *Cereal Chemistry*, 95(3), 347–364. <https://doi.org/10.1002/cche.10040>

Sánchez-MariñEz, R. I., Cortez-Rocha, M. O., Ortega-Dorame, F., Morales-Valdes, M., & Silveira, M. I. (1997). End-use quality of flour from *Rhyzopertha dominica* infested wheat. *Cereal Chemistry*, 74(4), 481–483. <https://doi.org/10.1094/CCHEM.1997.74.4.481>

Sarada, V., Swamy, S. V. S. G., Madhumathi, T., & Varma, P. K. (2018). Preference and progeny development of pulse beetle in response to black gram treated with botanicals and inert materials. *Journal of Entomology and Biology Studies*, 6(3), 1812–1815.

Sarwar, M., & Salman, M. (2015). Toxicity of oils formulation as a new useful tool in crop protection for insect pests control. *International Journal of*

Chemical and Biomolecular Science, 1(4), 297–302. Retrieved from <http://files.aiscience.org/journal/article/pdf/70420059.pdf>

- Sasidharan, S.; Chen, Y.; Saravanan, D.; Sundram, K. M.; Latha, L. Y. (2011). Extraction, isolation and characterization of bioactive compounds from plants' extracts. *African Journal of Traditional, Complementary and Alternative Medicines.*, 8(1), 1–10. https://doi.org/10.1007/978-3-642-56936-4_2
- Shaaya, E., Kostjukovski, M., Eilberg, J., & Sukprakarn, C. (1997). Plant oils as fumigants and contact insecticides for the control of stored-product insects. *Journal of Stored Products Research*, 33(1), 7–15. [https://doi.org/10.1016/S0022-474X\(96\)00032-X](https://doi.org/10.1016/S0022-474X(96)00032-X)
- Shafique, M., Ahmad, M., & Chaudry, M. A. (2006). Feeding Preference and Development of *Tribolium castaneum* (Herbst.) in Wheat Products. *Pakistan Journal of Zoology*, 38(1), 27–31.
- Sharon, M., Abirami, C. V. K., & Alagusundaram, K. (2014). Grain storage management in India. *Journal of Postharvest Technology*, 02(01), 012–014. Retrieved from www.jpht.info
- Singh, P., & Santosh Satya, and S. N. N. (2013). Effect of Insect Infestation on Quality Parameters of Wheat. In *International Conference on Food Security and Nutrition IPCBEE*, 67, 35–39. <https://doi.org/10.7763/IPCBEE>
- Singh, V., Yadav, K. S., Tripathi, A. K., Tandon, S., & Yadav, N. P. (2016). Exploration of various essential oils as fumigant to protect stored grains from insect damage. *Annals of Phytomedicine : An International Journal*, 5(2), 87–90. <https://doi.org/10.21276/ap.2016.5.2.10>
- Sinha, R. N. (1994). Stored grain ecosystem. In *The stored grain ecosystem* (pp. 1–31). United States: CRC Press, Taylor & Francis.
- Smirle, M. J., Lowery, D. T., & Zurowski, C. L. (1996). Influence of neem oil on detoxication enzyme activity in the obliquebanded leafroller, *Choristoneura rosaceana*. *Pesticide Biochemistry and Physiology*, 56(3),

220–230. <https://doi.org/10.1006/pest.1996.0075>

- Stefanazzi, N., Stadler, T., & Ferrero, A. (2011). Composition and toxic, repellent and feeding deterrent activity of essential oils against the stored-grain pests *Tribolium castaneum* (Coleoptera: Tenebrionidae) and *Sitophilus oryzae* (Coleoptera: Curculionidae). *Pest Management Science*, 67(6), 639–646. <https://doi.org/10.1002/ps.2102>
- Tak, J. H., Jovel, E., & Isman, M. B. (2015). Comparative and synergistic activity of *Rosmarinus officinalis* L. essential oil constituents against the larvae and an ovarian cell line of the cabbage looper, *Trichoplusia ni* (Lepidoptera: Noctuidae). *Pest Management Science*, 72(3), 474–480. <https://doi.org/10.1002/ps.4010>
- Tapondjou, L. A., Adler, C., Bouda, H., & Fontem, D. A. (2002). Efficacy of powder and essential oil from *Chenopodium ambrosioides* leaves as post-harvest grain protectants against six-stored product beetles. *Journal of Stored Products Research*, 38(4), 395–402. [https://doi.org/10.1016/S0022-474X\(01\)00044-3](https://doi.org/10.1016/S0022-474X(01)00044-3)
- Thakore, Y. (2006). The biopesticide market for global agricultural use. *Industrial Biotechnology*, 2(3), 194–208. <https://doi.org/10.1192/bjp.112.483.211-a>
- Tripathi, A. K.; Prajapati, V.; Verma, N.; Bahl, J. R.; Bansal, R. P.; Khanuja, S. P. S.; Kumar, S. (2002). Bioactivities of the leaf essential oil of *Curcuma longa* (var. ch-66) on three species of stored-product beetles (Coleoptera). *Journal of Economic Entomology*, 95(1), 183–189. <https://doi.org/10.1603/0022-0493-95.1.183>
- Tripathi, A. K., Prajapati, V., Aggarwal, K. K., Khanuja, S. P. S., & Kumar, S. (2000). Repellency and Toxicity of Oil from *Artemisia annua* to Certain Stored-Product Beetles. *Journal of Economic Entomology*, 93(1), 43–47. <https://doi.org/10.1603/0022-0493-93.1.43>
- Tripathi, Arun K., Prajapati, V., Aggarwal, K. K., & Kumar, S. (2001). Toxicity, Feeding Deterrence, and Effect of Activity of 1,8-Cineole from *Artemisia annua* on Progeny Production of *Tribolium castaneum*

- (Coleoptera: Tenebrionidae). *Journal of Economic Entomology*, 94(4), 979–983. <https://doi.org/10.1603/0022-0493-94.4.979>
- Upadhyay, N., Dwivedy, A. K., Kumar, M., Prakash, B., & Dubey, N. K. (2018). Essential Oils as Eco-friendly Alternatives to Synthetic Pesticides for the Control of *Tribolium castaneum* (Herbst) (Coleoptera: Tenebrionidae). *Journal of Essential Oil-Bearing Plants*, 21(2), 282–297. <https://doi.org/10.1080/0972060X.2018.1459875>
- Upadhyay, R. K., & Jaiswal, G. (2007). Evaluation of biological activities of *Piper nigrum* oil against *Tribolium castaneum*. *Bulletin of Insectology*, 60(1), 57–61.
- USDA. (2019). World agricultural production. In *Foreign Agricultural Service*. <https://doi.org/10.32317/2221-1055.201907059>
- Varma, J., & Dubey, N. K. (2001). Efficacy of essential oils of *Caesulia axillaris* and *Mentha arvensis* against some storage pests causing biodeterioration of food commodities. *International Journal of Food Microbiology*, 68(3), 207–210. [https://doi.org/10.1016/S0168-1605\(01\)00506-2](https://doi.org/10.1016/S0168-1605(01)00506-2)
- Wang, Y., You, C. X., Yang, K., Wu, Y., Chen, R., Zhang, W. J., ... Han, J. (2015). Bioactivity of Essential Oil of *Zingiber purpureum* Rhizomes and Its Main Compounds against Two Stored Product Insects. *Journal of Economic Entomology*, 108(3), 925–932. <https://doi.org/10.1093/jee/tov030>
- Weaver, L. T. (1994). Feeding the weanling in the developing world: Problems and solutions. *International Journal of Food Sciences and Nutrition*, 45(2), 127–134. <https://doi.org/10.3109/09637489409166151>
- Xie, Y. S.; Bodnaryk, R. P.; Fields, P. G. (1996). A rapid and simple flour-disk bioassay for testing substances active against stored-product insects. *The Canadian Entomologist*, 128(5), 865–875.
- Yang, R. Z., & Tang, C. S. (1988). Plants used for pest control in China: A literature review. *Economic Botany*, 42(3), 376–406.

<https://doi.org/10.1007/BF02860162>

You, C. X., Zhang, W. J., Guo, S. S., Wang, C. F., Yang, K., Liang, J. Y., ...
Deng, Z. W. (2015). Chemical composition of essential oils extracted
from six *Murraya* species and their repellent activity against *Tribolium
castaneum*. *Industrial Crops and Products*, 76, 681–687.
<https://doi.org/10.1016/j.indcrop.2015.07.044>

Zapata, N., & Smagghe, G. (2010). Repellency and toxicity of essential oils
from the leaves and bark of *Laurelia sempervirens* and *Drimys winteri*
against *Tribolium castaneum*. *Industrial Crops and Products*, 32(3), 405–
410. <https://doi.org/10.1016/j.indcrop.2010.06.005>

Zhang, Z., Guo, S., Zhang, W., Geng, Z., & Liang, J. (2017). Essential oil and
polyacetylenes from *Artemisia ordosica* and their bioactivities against
Tribolium castaneum Herbst (Coleoptera : Tenebrionidae). *Industrial
Crops & Products*, 100, 132–137.
<https://doi.org/10.1016/j.indcrop.2017.02.020>