References

1. Acosta, A., Sammarco, P. W. and Duarte, L. F. 2005. New fission processes in the zoanthid *Palythoa caribaeorum*: description and quantitative aspects. Bulletin of Marine Sciences. 76:1-26.

2. Alcock, A. and Anderson, A. R. S. 1894. Natural history notes from RIMSS 'Investigator'. Ser. 2, No. 17. List of shore and shallow water Brachyura collected during the season 1893-1896. Journal of Asiatic Society of Bengal. 65: 88-106.

3. Alcock, A. and Anderson, A. R. S. 1899. Natural history notes from H. M. Royal Indian Marine Survey Ship, 'Investigator', Commander T. H., Heming, R. N., commanding. Series 3, No. 2. An account of the Deep-Sea Crustacea dredged during the surveying season of 1897-1898. The Annals and Magazine of Natural History. Series. 7, 3: 1-27.

4. Alcock, A.W. 1898. Materials for a carcinological fauna of India. No. 3. The BrachyuraCyclometopa. Part 1. The Family Xanthidae. Journal of the Asiatic Society of Bengal. 67: 67-233.

5. Aller, R. C. 2001. Transport and reactions in the bioirrigated zone, in The Benthic Boundary Layer. Transport Processes and Biogeochemistry, edited by B. P. Boudreau and B. B. Jorgensen. Oxford University Press, Oxford, 269-301.

6. Anger, K. 1995. Starvation resistance in larvae of a semiterrestrial crab, *Sesarma curacaoense* (Decapoda: Grapsidae). Journal of Experimental Marine Biology and Ecology. 187, 2: 161-174.

7. Ansari, Z. A., Ingole, B. S., Banerjee, G., and Parulekar, A. H. 1986. Spatial and temporal changes in benthic macrofauna from Mandovi and Zuari estuaries of Goa, West coast of India. Indian Journal of Marine Sciences. 15: 223-229.

8. Apte, D. 1998. Book of Indian shells. Bombay Natural History Society, Mumbai, India. 115pp.

9. Attrill, M. J. and Rundle, S. D. 2002. Ecotone or ecocline: Ecological boundaries in estuaries. Estuarine, Coastal and Shelf Science. 55: 929-936.

10. Bacci, G. and Sella, G. 1970. Correlation between characters and environmental condition in Patella of the coerulea group. Pubbl. Staz. Zool. Napoli. 38: 1-17.

11. Bandekar, P. D., Neelkantan, K. and Kakati, V. S. 2011. Biodiversity of crabs in Karwar mangrove environment west coast of India. Recent Research in Science and Technology. 3, 4: 1-5.

12. Barros, F. 2001. Ghost crabs as a tool for rapid assessment of human impacts on exposed sandy shores. Biological Conservation. 97: 399-404.

13. Bertness, M. D. and Miller, T. 1984. The distribution and dynamics of *Uca pugnax* (Smith) burrowsin a New England salt marsh. Journal of Experimental Marine Biology and Ecology. 83: 211-237.

14. Bhattji, N. S., Shah, D. G., Desai, N. D. and Mankodi, P. C. 2010. A new record of genus Palythoa, a Zoanthid from the Gulf of Kachchh, Gujarat, India: A great concern for Reef Ecosystem, Seshaiyana. 1: 18.

15. Bolam, S. G. 2003. Spatial patterns of estuarine macrobenthic assemblages: relationship with hydrodynamic regime. Estuarine Process Research Project, CEFAS, March 2003 Report.

16. Boschi, E. E. 2000. Biodiversity of marine decapod brachyurans of the Americas. Journal of Crustacean Biology. 2: 337-342.

17. Botto F., Iribarne, O. and Martinez, M. M. 1998. The Effect of Migratory Shorebirds on the benthic species of three southwestern Atlantic Argentinean estuaries. Estuaries. 21, 4 (B): 700-709.

18. Boxshall, G. A., Mees, J., Costello, M. J., Hernandez, F., et. al., (above 50 authors). 2015. World Register of Marine Species. Available from http://www.marinespecies.org at VLIZ.

19. Braithwaite, C. J. R. and Talbot, M. R. 1972. Crustacean burrows in the Seychelles, Indian Ocean. Palaeogeography, Palaeoclimatology, Paleoecology. 11: 265–285.

20. Brown, A.C., and McLachlan, A. 1990.*Ecology of sandy shores*. Amsterdam: Elsevier Publishers.

21. Brown, A. C. 1983. The eco physiology of sandy beach animals-a partial review. In 'Sandy Beaches as Ecosystems'. (Eds A. McLachlan and T. Erasmus.) pp. 575-605.

22. Burnett, W. J., Benzie, J. A. H., Beardmore, J. A. and Ryland, J. S. 1997. Zoanthids (Anthozoa, Hexacorallia) from the Great Barrier Reef and Torres Strait, Australia: systematics, evolution and a key to species. Coral Reefs. 16: 55-68.

23. Castro, P., Ng, P.K .L. and Ahyong, S. T. 2004. Phylogeny and systematics of the Trapeziidae Miers, 1886 (Crustacea: Brachyura), with the description of a new family. Zootaxa. 643: 1-70.

24. Chace F. A. Jr. 1951. The number of species of decapoda and stomatopod Crustacea. Journal of the Washington Academy of Sciences. 41, 11: 369-372.

25. Chakrabarti, A., Chakrabarti, R. and Hertweck, G. 2006. Surface traces and bioturbate textures from bubbler crabs: an indicator of subtropical to tropical tidal flat environments. Senckenbergiana maritime. 36, 1:19-2706.

26. Chakrabarti, A. 1981. Burrow pattern of *Ocypode ceratophthalma* (Pallas) and their environmental significance. Journal of Paleontology. 187: 113-130.

27. Chakraborthy, S. K., Poddar, T. K. and Choudhury, A. 1992. Species diversity of macrozoobenthos of Sagar Island, Sunderbans, India. Proceeding of Zoological Society, Calcutta. 45 (A): 435-444.

28. Chan, K. K. B., Chan, K. K. Y. and Leung, P. C. M. 2006. Burrow architecture of the ghost crab *Ocypode ceratophthalma*on a sandy shore in Hong Kong. Hydrobiologia. 560: 43-49.

29. Chandy, M. 1973. New records of brachyuran decapods from the Gulf of Kutch. Journal of Bombay Natural History Society. 70 (2): 401-402.

30. Chandy, M. 1969. Thalamita poissonii (Audouin and Savignyi) De Man (Crustacea: Brachyura), a new record to Indian coasts. Journal of Bombay Natural History Society. 66(3): 635-637, text-figs 1-2.

31. Chavanich, S. and Wilson, K. A. 2000. Rocky intertidal zonation of gammaridean amphipods in Long Island Sound, Connecticut. Crustaceana. 73: 835-846.

32. Chhapgar, B. F. 1955. On two new species and a variety of crabs (Decapoda: Brachyura) from Bombay state. Records of the Indian Museum. 53: 251-257.

33. Chhapgar, B. F. 1957a. On the marine crabs (Decapoda: Brachyura) of Bombay state, part 1. Journal of Bombay Natural History Society. 54(2): 399-439, pls 1-11.

34. Chhapgar, B. F. 1957b. On the marine crabs (Decapoda: Brachyura) of Bombay state, part 2. Journal of Bombay Natural History Society. 54(3): 503-549, pls B and 12-16.

35. Chhapgar, B. F., Desai, B. G. and Patel, S. J. 2004. On two interesting marine crabs (Decapoda: Brachyura) from Mandvi, Kutch. Journal of Bombay Natural History Society. 101 (1): 184-186.

36. Chhapgar, B. F. and Mundkur, T. 1995. Occurrence of the crab *Euxanthus exsculptus* (Herbst) in Gujarat. Journal of Bombay Natural History Society. 92 (2):286.

37. Chhapgar, B. F. 1958. Addition to the crab fauna of Bombay State.Journal of Bombay Natural History Society. 55 (3): 582-585.

38. Chhapgar, B. F. 1969. More additions to the crab fauna of Bombay State.Journal of Bombay Natural History Society. 65 (3): 608-617.

39. Cooke, W. J. 1976. Reproduction, growth, and some tolerance of *Zoanthus pacificus* and *Palythoa vestitus* in Kaneohe Bay, Hawaii. In GO Mackie (ed) Coelenterate ecology and behavior. Plenum Press, New York. 281-288pp.

40. Christy, J. H. 1982. Burrow structure and use in the sand fiddlercrab, *Uca pugilator* (Bosc). Animal Behaviour. 31: 687–694.

41. Dahl, E. 1952. Some aspects ofthe ecology and zonation of the fauna on sandy beaches. Oikos. 4: 1-27.

42. Davenport, J. and Davenport, J. L. 2005. Effects of shore height, wave exposure and geographical distance on thermal niche width of intertidal fauna. Marine Ecology Progress Series. 292: 41-50.

43. Davie, P.J. F. 1982. A preliminary checklist of Brachyura (Crustacea: Decapoda) associated with Australian mangrove forests. Operculum. 5: 204-207.

44. Denny, M. W. 1988. Biology and the mechanics of the wave-swept environment.Princeton University Press, Princeton.des Sciences. Paris. 259: 1576-9.

45. Deshmukhe, G., Ramamoorthy, K. and Gupta, S. R. 2000. On the coral reefs of the Gulf of Kachchh. Current Science. 79, 2: 160-162.

46. Dev Roy, M. K. and Nandi, N. C. 2012. Brachyuran crabs (Crustacea). Fauna of Andaman and Nicobar Islands, State Fauna Series. 19, 1: 185-236. Zoological Survey of India, Kolkata.

47. Dev Roy, M. K. 2013. Diversity and distribution of marine brachyuran crab communities inhabiting west coast of India: in Ecology and Conservation of Tropical Marine Faunal Communities Part 1. (edt. K. Venkatraman, C. Sivaperuman and C. Raghunathan)Springer Berlin Heidelberg. pp 147-169.

48. Dev Roy, M.K. and Das, A.K. 2000. Taxonomy, ecobiology and distribution pattern of Brachyuran Crabs of mangrove ecosystem in Andaman Islands. Records of Zoological Survey of India. 185: 1-21.

49. Dexter, D. M. 1984. Temporal and spatial variability in the community structure of the fauna of four sandy beaches in south-eastern New South Wales. Australian Journal of Marine and Freshwater Research. 35: 663-72.

50. Diaz, R. J. and Cutter, G. R. 2001. In situ measurement of organism–sediment interaction: rates of burrow formation/abandonment and sediment oxidation/reduction, in Organism–Sediment Interactions, edited by J. Y. Aller, S. A. Woodin, and R. C. Aller, University of South Carolina Press, Columbia. 19-32pp.

51. Diesel, R. and Horst, D. 1995. Breeding in a snail shell: ecology and biology of the Jamaican montane crab *Sesarma jarvisi* (Decapoda: Grapsidae). Journal of Crustacean Biology. 15: 179-195.

52. Doty, M. S. 1957. Rocky Intertidal Surfaces. In: Treatise on Marine Ecology and Paleoecology, Joel W. Hedgepeth, ed. The Geological Society of America, Memoiries. 67.

53. Dubey, S. K., Chakraborty, D. P., Chakraborty, S. and Choudhury, A. 2013. Burrow architecture of red ghost crab Ocypode macrocera (H. Milne-Edwards, 1852): a case study in Indian Sundarbans. Exploratory Animal and Medical Research. 3, 2: 136-144.

54. Edgar, G.J. and Aoki, M. 1993. Resource limitation and fish predation: their importance to mobile epifauna associated with Japanese *Sargassum*. Oecologia. 95: 122-133.

55. Elliott, A. N. 1993. Global Marine Biological Diversity: A Strategy for Building Conservation into Decision Making. Island Press, Washington, DC.

56. Ellis, D. V. 2003. Rocky shore intertidal zonation as a means of monitoring and assessing shoreline biodiversity recovery. Marine Pollution Bulletin. 46: 305-307.

57. Emson, R.H. and Faller-Fritsch, R. J. 1976. An experimental investigation into the effect of crevice availability on abundance and size structure in a population of *Littorina rudis* (Maton). Gastropoda; Prosobranchia. Journal of Experimental Marine Biology and Ecolology. 23: 285-297.

58. Esselink, P. and Zwarts, L. 1989. Seasonal trend in burrow depth and tidal variation in feeding activityof *Nereis diversicolor*. Marine EcologyProgress Seies. 56: 243-254.

59. Eston, V.R. and Bussab, W.O. 1990. An experimental analysis of ecological dominance in a rocky subtidal macroalgal community. Journal of Experimental Marine Biology and Ecology. 136: 170-195.

60. Fabricius, J. C. 1775. Systema Entomologiae, sistens insectorum Classes, Ordines, Genera, Species, Adjectis Synonymies, Locis, Descriptionibus,

Observationibus, pp 1-832. Libraria Kortii: Flensburgi, Lipsiae endogenee. Compte rendu hebdomadaire des Seances de I'Academie.

61. Fabricius, J. C. 1793. Entomologia Systematica Emendata et Aucta, Secundum, Classes, Ordines, Genera, Species, Adjectis Synonimas, Locis, Observationibus, Descriptionibus. Hafniae. 2 (8): 1-519.

62. Farrow, G. E. 1971. Back reef and lagoonal environment of Aldabra Atoll, distinguished by their crustacean burrows. In Stoddart, D. R. & C. M. Yonge (eds), Regional Variation in Indian Ocean Coral Reefs. Proceedings of the Symposium ofthe Zoological Society of London. 455-500 pp.

63. Fautin, D. G. 2010. Hexacorallians of the world.

64. Fenical, W. 1996. Marine Biodiversity and the Medicine Cabinet: The Status of New Drugs from Marine Organisms. Oceanography. 9(1): 23-24.

65. Fischer, P. (1966). Disposition gregaire chez um Cerithium Indo- Pacifique. Journal. Conchyl, Paris. 105: 49-52.

66. Folk, R. L. and Ward, W. C. 1957. Brazos River Bar: a study in the significance of grain sizeparameters: Journal of Sedimentary Petrology. 27: 3-26.

67. Fossa, S. A. and Nilsen, A. J. 1998. The modern coral reef aquarium. Vol. 2. Bornheim, Sweden: Birgit Schmettkamp Verlag. 479 pp.

68. Fransozo, A. and Negreiros-Fransozo, M.L. 1996. Brazilian coastal CrustaceaDecapoda. In: de Bicudo C.E. and Menezes M.N.A. (eds) Biodiversity in Brazil: A First Approach. Proceedings of the Workshop Methods for the assessment of biodiversity in plants and animals, Campos do Jorda˜ o, Sa˜o Paulo, Brazil, 26-30 May 1996, CNPq – ConselhoNacional de DesenvolvimentoCientı´fico e Tecnolo´ gico, Sa˜o Paulo, Brazil. pp. 275-287.

69. Fransozo, A., Negreiros-Fransozo, M. L. and Mantelatto, F. L. 1992. Composicao e distribuicao dos Brachyura (Crustacea, Decapoda) do sublitoralnãoconsolidadonaEnseada da Fortaleza, Ubatuba (SP). Revista Brasileira de Zoologi. 52(4): 667-675.

70. Giese, A. C. 1959. Comparative physiology: Annual reproductive cycles of marine invertebrates. Annual Review of Physiology. 21: 547-576.

71. Gohil, B. and Kundu, R. 2011. Ecological status of *Rhinoclavis sinensis* at Dwarka Coast, Gujarat (India). The Ecoscan. 5: 131-134.

72. Golley, F., Odum, H.T. and Wilson, R.L. 1962. The structure and metabolism of Puertrican red mangrove forest in May. Ecology. 43: 9-18.

73. Gravely, F. H. 1927. Orders Decapoda (except Paguridea) and Stomatopoda. In: The Littoral Fauna of Krusadai Island in Gulf of Mannar Bulletin of Madras Government Museum. 1(1): 135-155, pls 20-26.

74. Griffis, R. B. and Suchanek, T. H. 1991. A model of burrow architecture and trophic modes inthalassinidean shrimp (Decapoda: Thalassinidea). Marine Ecology Progress Series. 79: 171-183.

75. Hebling, N. J., Mantelatto, F. L. and Negreiros-Fransozo, M. L. 1994. Levantamentoe distribuicao de braquiúros e Anomuros (Crustacea, Decapoda) dos sediment ssublitorais da região da IlhaAnchieta, Ubatuba (SP). Bolm. Inst. Pesca. 21: 1-9.

76. Henderson, J. R. 1893. A contribution to Indian carcinology. Transactions of the Linnaean Society of London, Zoology, Series 2. 5: 325-458.

77. Hill, G. W. and Hunter, R. E. 1973. Burrows of the ghost crab *Ocypode quadrata* (Facricius) on the Barrier Islands, southcentral Texas coast. Journal of Sedimentary Petrology. 43:24–30.

78. Hiyodo, C. M. and Fransozo,A. 1994. Biologia populacional de *Acanthonyx petiverii* H. Milne Edwards, 1834 (Crustacea, Decapoda, Majidae). pp. 373. In: 2 Congresso de Ecologia do Brasil. Universidade Estadual de Londrina, Londrina, Brasil.

79. Hornell, J. 1916. Report to the Government of Baroda on the marine zoology of Okhamandal in Kattiawar, Williams and Norgate.

80. Houbrick, R. S. 1992. Monograph of the genus Cerithium Bruguière in the Indo-Pacific (Cerithiidae: Prosobranchia). Smithsonian Contributions to Zoology. 510: 1-211.

81. Hughes, R. N. 1980. *Predation and community structure in the shore Environment* Vol 2. Ecosystems (eds. D. E C, Irvine and W F Farnham) (Systematics Association Special Volume No 17B) Academic Press London. Pp. 699-728.

82. ICMAM. 2002. Geographical Information System for Gulf of Kachchh, Integrated Coastal and Marine Area Management Report.

83. Irei, Y., Nozawa, Y.and Reimer, J. D. 2011. Distribution patterns of five zoanthid species at Okinawa Island, Japan. Zoological Studies. 50(4):426-433.

84. Jaramillo, E. and McLachlan, A. 1993. Community and population responses of the macroinfauna to physical factors over a range of exposed sandy beaches in South-central Chile. Estuarine, Coastal and Shelf Science. 37, 615-624.

85. Jaramillo, E., McLachlan, A. and Coetzee, P. 1993. Intertidal zonation patterns of macrofauna over a range of exposed sandy beaches in South-Central Chile. Marine Ecology Progress Series. 101: 105-118.

86. Jardine, J. W. 1985.Ecology of Rocky Coasts. (eds. Moor, P.G. and Seed, R.) Hodder and Stoughton, London, Sydney, Auckland, Toronto, pp. 117-135.

87. Jeyabaskaran, R. S. and Wafar, M. 2002. CD on Brachyuran Crabs of West Coast, India. National Institute of Oceanography, Dona Paula, Goa, India.

88. Jeyabaskaran, R., Khan, A. S. and Ramaiyan, V. 2000. Biodiversity Project on Gulf of Mannar Biosphere Reserve. Centre of Advanced Study in Marine Biology, Annamalai University, Parangipettai, India.

89. Jirapunpipat, K., Aungtenya, C. and Watanbe, S. 2008. Morphological study and application of multivariate analysis for the mud crab Scylla in Klongangao mangrove, Rangong province. Phuket Marine Biology Centre Research Bulletin. 69: 7-24.

90. Joel, D. R. and Raj, P. J. S. 1980. Taxonomic remarks on two species of the genus Scylla de Haan (Portunidae: Brachyura) from Pulicat Lake. Journal of Inland Fishries Society of India. 12(2): 39-50.

91. Joel, D. R., Raj, P. J. S. and Raghavan, R. 1985. Distribution and zonation of shore crabs in the Pulicat Lake. Proceedingsof Indian Academy of Animal Sciences. 95: 437-445.

92. Johannesson, L. T., Stevens, R. L. and Alexanderson, J. H. 2000. Sediment character in a micro-tidal, harbor-estuary environment, Goteborg, Sweden, Estuaries. 23: 400-410.

93. Jones, D. A. 1984. Crabs of the mangal ecosystem. In: Por, F.D. and I. Dor. (eds.), Hydrobiology of the Mangal. Dr. W. Junk, Publisher, The Hague, Netherlands. 89-109.

94. Karlson, R. H. 1983. Disturbance and monopolization of a spatial resource by *Zoanthus sociatus* (Coelenterata, Anthozoa). Bulletin of Marine Sciences. 33: 118-131.

95. Kasinathan, R. and Shanmugam, A. 1986. Benthic macrofauna of the Pichavaram mangroves, South India. J. Annamalai University science. Part-B, 34: 109-119.

96. Keenan, C. P., Davie, P. J. F. and Mann, D. L. 1998. A revision of the genus Scylla de Haan, 1833 (Crustacea: Decapoda: Brachyura: Portunidae). Raffles Bulletin of Zoology. 46(1):217-245.

97. Kemp, S. 1917. Crustacea Decapoda in the Indian Museum. 10. Hymenosomatidae. Records of Indian Museum. 13(4): 243-279.

98. Kemp, S. 1919a. Notes on Crustacea Decapoda in the Indian Museum. 12. Scopimera. Records of Indian Museum. 16(5): 305-348, pls 12, 13.

99. Khan, A. S., Raffi, S. M. and Lyla, P. S. 2005. Brachyuran crab diversity in natural (Pitchavaram) and artificially developed mangroves (Vellar estuary). Current Science. 88 (8): 1316-1324.

100. Kumar, A. A. J. and Wesley, S. G. 2010. Spatio- temporal distribution of assemblages of brachyuran crabs at Lammu atoll, Maldives. Atoll Research Bulletin. 588: 1-16.

101. Kumaralingam, S., Sivaperuman, C. and Raghunathan, C. 2012. Diversity and Distribution of Brachyuran Crabs from Ritchieo's Archipelago. International Journal of Oceanography and Marine Ecological System. 1: 60-66.

102. Leber, K. M. 1982. Seasonality of macro invertebrates on a temperate, high wave energy sandy beach. Bulletin of Marine Science. 32: 86-98.

103. Lee, S. Y. 1998. Ecological role of grapsid crabs in mangrove ecosystems: a review. Marine and Freshwater Research. 49: 335-343.

104. Lenihan, H. S and Micheli, F. 2001. Soft-sediment communities. In: Bertness MD, Gaines SD, Hay ME (eds) Marine Community Ecology. Sinauer Associates Inc, Sunderland, Massachusetts. pp 253-287.

105. Levinton, J. S. 2001. Marine Biology: Function, Biodiversity, Ecology. Oxford University Press.

106. Lim, S. S. L. 2006. Fiddler crab burrow morphology: How do burrow dimensions and bio turbative activities compare in sympatric populations of *Uca vocans* (Linnaeus, 1758) and *U. annulipes* (H. Milne Edwards, 1837). Crustaceana. 79: 525-540.

107. Linsley, R. M., 1978a. Locomotion rates and shell form in gastropoda. Malacologi. 17(2): 193-206.

108. Lohrer, A. M., Fukui,Y., Wada,K. andWhitlatch, R. B. 2000. Structural complexity and vertical zonation of intertidal crabs, with focus on habitat requirements of the invasive Asian shore crab, *Hemigrapsus sanguineus* (de Haan).Journal of Experimental Marine Biology and Ecology. 244: 203-217.

109. Lohrer, A. M., Fukui,Y., Wada, K., Whitlatch, R. B. 2000. Structural complexity and vertical zonation of intertidal crabs, with focus on habitat requirements of the invasive Asian shore crab, Hemigrapsus sanguineus (de Haan). Journal of Experimental Marine Biology and Ecology. 244: 203-217.

110. Loosanoff, V. L. and Devis, H. C. 1952. Repeated semiannual spawning of northern oysters, Science. 115: 675-676.

111. Lucrezi, S., Schlacher, T. A. and Simon, W. 2009. Monitoring human impacts on sandy shore ecosystems: a test of ghost crabs (Ocypode spp.) as

biological indicators on an urban beach. Environment Monitoring and Assessment. 152: 413-424.

112. MacArthur, R. H. 1972. Geographical ecology: patterns in distribution of species. New York: Harper and Row.

113. MacArthur, R. H. 1972. Geographical ecology: patterns in distribution of species. New York: Harper and Row.Mantelatto, F. L., Fabíola C. R. F., Biagi, Rand Melo. G. A. S. 2004. Majoid Crabs Community (Crustacea: Decapoda) from Infralittoral Rocky/sandy Bottom of Anchieta Island, Ubatuba. Brazilian Archives of Biology and Technology. 47 (2): 273-279.

114. Malli, P. C. 1983. Ecological studies on two turbinid gastropod *Turbo cornatus* (Gmelin) and *Turbo intercoastalis* (Manke) from Saurashtra, West coast of India. Ph.D. Thesis. Saurashtra University, Rajkot, India.

115. Mantelatto, F. L., Fabiola, C. R. F., Biagi, R. and Melo, G. A. S. 2004.Majoid Crabs Community (Crustacea: Decapoda) from Infralittoral Rocky/sandy Bottom of Anchieta Island, Ubatuba. Brazilian Archives of Biology and Technology. 47 (2): 273-279.

116. McAllister, D. 1995. Status of the World Ocean and Its Biodiversity. Sea Wind. 9(4): 14.

117. Melo, G.A.S. 1996. Manual de identificac¸a˜o dos Brachyura (caranguejos siris) do litoralbrasileiro.Pleˆiade/FAPESP Ed., Sa˜o Paulo, Brazil. 604 pp.

118. Mendonca-Neto, J. P., Ferreira, C. E. L., Chaves, L. C. T and Pereira, R. C. 2008. Influence of Palythoa caribaeorum (Anthozoa, Cnidaria) zonation on site-attached reef fishes. An. Acad. Bras. Ciênc. 80: 495-513.

119. Milne Edwards, H. 1852. Observations sur les affinities Zoologiques et la classification naturelle des Crustacés. Annales des Sciences Naturelles, Zoologie, Paris, Series, 3. 18: 109-166, pls 3-4.

120. Misra, S. 2004. Studies on the biodiversity and the impact of anthropogenic pressure on the ecology and biology of certain intertidal macrofauna. Ph.D.Thesis, Saurashtra University, Rajkot.

121. Misra, S. and Kundu, R. 2005. Seasonal variations in population dynamics of key intertidal molluscs at two contrasting locations. Aquatic Ecology. 39: 315-324.

122. Montouchet, P.C.G. 1979. Sur la communauté des ani- maux vagiles associés à *Sargassum cymosum* C. Agardh, à Ubatuba, Etat de São Paulo, Brésil. Studies on Neotropical Fauna and Environment. 14:33-64.

123. Morton, B. & J. Morton, 1983. The Seashore Ecology of HongKong. Hong Kong University Press, Hong Kong.

124. Moser, S., Macintosh, D., Laoprasert, S. and Tongdee, N. 2005. Population ecology of the mud crab *Scylla olivacea*: a study in the Ranong mangrove ecosystem, Thailand, with emphasis on juvenile recruitment and mortality. Fisheries Research. 71: 27-41.

125. Myers, A. A. and Mcgrath, D. 1993. Settlement and growth of the intertidal gastropod *Melarhaphe neritoides* on the south coast of Ireland. Journal of the Marine Biological Association of the United Kingdom. 73: 313-319.

126. Nayar, N.K. and Appukuttan, K. K. 1983.*Trochus* and *Turbo* resources. CMFRI, Bulletin. 34: 81-84.

127. Neves, F. M. and Bemvenuti, C. E. 2006. The ghost crab *Ocypode quadrata* (Fabricius, 1787) as a potential indicator of anthropogenic impact along the Rio Grande do Sul coast, Brazil. Biological Conservation. 133: 431-435.

128. Newell, R. C. 1979. Biology of intertidal animals. Marine Ecological Surveys, Ltd., Kent.

129. Ng, P. K. L. and Davie, P. J. F. 2007. On identity of *Atergatis floridus* (Linnaeus, 1767) and recognition of *Atergatis ocyroe* (Herbst, 1906) as a valid species from Indian Ocean (Crustacea: Brachyuran: Xanthidae). The Raffles Bulletin of Zoology. 16: 169-175.

130. Ng, P. K. L., Guinot, D. and Davie, P. J. F. 2008. Syestema brachyuroram part 1. An annotated checklist of extant brachyuran crabs of the world. The Raffles Bulletin of Zoology. 17: 1-286.

131. Nickell, L. A. and Atkinson, R. J. A. 1995. Functional morphology of burrows and trophic modes of three thalassinidean shrimp species, and a new approach to the classification of thalassinidean burrow morphology. Marine Ecology Progress Series. 128: 181-197.

132. NIO, 2002. Status of Flora and Fauna of Gulf of Kachchh, India. National Institute of Oceanography, Goa, India.

133. Nobili, G. 1903. Crostacei di Pondichéry, Mahé, Bombay etc. Boll Musei Zool Ant Comp R Univ. 18:1-24.

134. Ono, S., Reimer,J. D. andSukahara, J. T. 2003. Long-term changes of Zoanthus spp. in the infra-littoral zone at Taisho Lava Field, Sakurajima, Kagoshima, Japan. Galaxea. 5: 21-31.

135. Osman, R. W., and Whitlatch, R. B. 1978. Patterns of species diversity: fact or artefact. Paleobiology. 4:41-54.

136. Pandya, K. M. and Mankodi, P. C. 2013. Brachycnemic Zooxanthellate Zoanthids (Cnidaria: Zoantharia) of Saurashtra Coast: A Preliminary Survey. Research Journal of Marine Sciences. 1(1): 10-13.

137. Pandya, K. M., Dave, C. S. and Mankodi, P. C. 2012. Interactions between scleratinians and zoanthids in the rocky intertidal area of Saurashtra coast, Gujarat, India. Proceedings of Biodiversity and Conservation of Coastal and Marine Ecosystems of India, Mumbai, India. 67-69.

138. Pandya, P. J. 2011. Benthic community structure of Mahi River estuary with special reference to animalsediment relationship. PhD Thesis, TheMaharaja Sayajirao University of Baroda, Vadodara, Gujarat, India.

139. Pandya, P. J. and Vachhrajani, K. D. 2010. Spatial distribution and substratum preference of the brachyuran crab *Macrophthalmus depressus* (Decopoda, Ocypodidae) along the lower estuarine mudflat of MahiRiver, Gujarat, India. Crustaceana. 83, 1055-1067.

140. Pandya, P. J. andVachharajani, K. D. 2011. Life under ecological stress: An estuarine case study. In: Animal Diversity, Natural History and Conservation (Vol. 1) (Gupta, V. K., Verma, A, K, eds). 427-436, Daya Publication House,New Delhi, India.

141. Patel, A. N. 1984. Ecological studies on two cerithiid gastropod, *Cerrithium caerulum* (sowerby) and *Clypeomorus monoliferus* (Miener) from Saurashtra Coasts, India, Ph D. Thesis, Saurashtra University, India.

142. Patel, A. N. 1984. Ecological studies on two cerithiid gastropod, *Cerrithium caerulum* (sowerby) and *Clypeomorus monoliferus* (Miener) from Saurashtra Coasts, India, Ph D. Thesis, Saurashtra University, India.

143. Paula, E.J. and Oliveira Filho, E. C. 1980. Phenology of two populations of *Sargassum cymosum* (Phaeophyta Fucales) of São Paulo State coast, Brazil. Boletim de Botânica da Universidade de São Paulo. 8: 21-39.

144. Pemberton, G. S., Risk, M. J. and Buckley, D. E. 1976. Supershrimp: Deep-bioturbation in the straitof Canso, Nova Scotia. Science.192: 790-791.

145. Perez, C. D., Vila-Nova, D. A. and Santos, A. M. 2005. Associated community with the zoanthid Palythoa caribaeorum (Duchassaing and Michelotti, 1860) (Cnidaria, Anthozoa) from littoral of Pernambuco, Brazil. Hydrobiologia. 548: 207-215.

146. Peter, M. J., Middelburg,J. J. and Heip, H. R. 2001. Benthic community structure and sediment processes on an intertidal flat: results from the ECOFLAT project. Continental Shelf Research. 21(18-19):2055-2071.

147. Pillai, N. K. 1951. Decapoda (Brachyura) from Travancore. Bulletin of Central Research Institute, University of Travancore, series C, 2(1): 1-46.

148. Place, S.P., O'Donnell, M. J. and Hofmann, G. E. 2008. Gene expression in the intertidal mussel *Mytilus californianus*: physiological response to environmental factors on a biogeographic scale. Marine Ecology Progress Series. 356, 1-14.

149. Pohle, G., Iken, K. and Clarke, K. R. 2011. Aspects of benthic decapode diversity and distribution from rocky near shore habitat of geographically widely dispersed sites PLoS ONE. 6(4): 1-11.

150. Portner, H. O., Bennett, A.F., Bozinovic, F., Clarke, A., Lardies, M.A., Lucassen, M., Pelster, B., Schiemer, F. and Stillman, J. H. 2006. Trade-offs in thermal adaptation: the need for a molecular to ecological integration. Physiological and Biochemical Zoology. 79, 295-313.

151. Prasad, M. N. 1984. Ecological studies on two limpets *Cellana radiata* (Born) and *Siphonaria siphonaria* (Sowerby) from Saurashtra coast, India. Ph.D. Thesis, Saurashtra University, Rajkot, India.

152. Prasad, M.N. and Mansuri, A. P. 1982. Population density of the limpet, *Cellana radiata* (Born) in polluted waters at Porbandar, west coast of India. Indian Journal of Marine Sciences.11: 180-181.

153. Quayle, D. B. 1951. The rate of growth of *Venerupis pullastra* (Montagu) at Millport,Scotland, Proc. R. Soc. E. Dinb, B. 64, 384, 406.

154. Raffaelli, D., Karakassis, I. and Galloway, A. 1991. Zonation schemes on sandy shores: a multivariate approach. Journal of Experimental Marine Biology and Ecology. 148, 241-53.

155. Raghunathan, C., Sen Gupta, R., Wangikar, U. and Lakhmapurkar, J.2004. A record of live corals along the Saurashtra coast of Gujarat, Arabian Sea. Current Science. 87(8): 1131-1138.

156. Rao, B.M. and Ganpati. 1971. Ecological studies on a tropical limpet *Cellana radiata*. Structural variations in the shell inrelation to distribution. Marine Biology.10:236-243.

157. Ravichandran S., Soundarapandian, P. and Kannupandi, T. 2001. Zonation and distribution of crabs in Pichavaram mangrove swamp, southeast coast of India. Indian Journal Fisheries. 48 (2): 221-226.

158. Ravichandran, S., Anbuchezhian, R., Sivasubramaniyan, K. and Rameshkumar, G. 2010. Expansion of occurrence of two bat crab (Crustacea: Decapoda: Brachyura: Parthenopidae) from the West to Southeast Coast of India. Biotemas.23 (4): 163-167.

159. Reimer, J. D. and Miyake, H. 2009. Recent research on ignored taxa: diversity of zoanthids (Zoantharia: Hexacorallia). Kaiyo Month. 41: 302-312. (in Japanese).

160. Reimer, J. D. and Hickman, C. P. 2009. Preliminary survey of zooxanthellate zoanthids (cnidaria: hexacorallia) of the galapagos, and associated symbiotic dinoflagellates (*Symbiodinium* spp.) Galapagos Research. 66:14-19.

161. Reimer, J. D., Takishita, K., Ono,S. and Maruyama, T. 2007. Diversity and evolution in the zoanthid genus Palythoa (Cnidaria: Hexacorallia) based on nuclear ITS-rDNA. Coral Reefs. 26: 399-410.

162. Reimer, J. D., Ono,S., Iwama,A., Takishita,K., Tsukahara,J. and Maruyama, T. 2006. Morphological and molecular revision of Zoanthus (Anthozoa: Hexacorallia) from southwestern Japan, with descriptions of two new species. Zoological Science. 23: 261-275.

163. Rhoads, D. C. and Boyer, L. F. 1982. The effects of marine benthos on physical properties of sediments. A successional perspective, in Animal-Sediment Relations, edited by P. L. McCall andM. J. S. Tevesz, 3-52, Plenum Publishing Company, New York.

164. Samantray, K. C. 1979. Edible limpet *Cellana radiata (Born)* (Mollusca : Patellidae at Veraval coast of Saurashtra. M.Sc. Dessertation, Saurashtra University, Rajkot, India.

165. Sankarnkutty, C. 1969. On the new species of *Xenophthalmus* White (Crustacea: Brachyura: Pinnotheridae) from Cochin. Journal of Bombay Natural History Society. 66 (1): 92-98.

166. Sankarnkutty, C. 1975. On the new species of *Hexapus* De Haan (Decapoda: Goneplacidae) from Cochin. Crustaceana. 28 (1): 1-6.

167. Santos, C. G. and Correia, M.D. 1995. Fauna associada ao fital *Halimeda opuntia* (Linnaeus) Lamouroux (Chlorophyta) do recife da Ponta Verde, Maceió, Alagoas, Brasil. Revista Brasileira de Zoologia da Universidade de São Paulo. 12: 263-271.

168. Sarvaiya, R. T. 1977. Studies on Mollusca of Saurashtra Coasts-l. Fisheries Technology. 1: 27-32.

169. Sasekumar, A. 1974. Distribution of macrofauna on a Malayan mangrove shore. Journal of Animal Ecology. 43: 51-69.

170. Schlacher, T. A., Jager, R. D. and Nielsen, T. 2011. Vegetation and ghost crabs in coastal dunes as indicators of putative stressors from tourism. Ecological Indicators. 11 (2): 284-294.

171. Sebens, K. S. 1982. Intertidal distribution of zoanthids on the Caribbean coast of Panama: effects of predation and desiccation. Bulletin of Marine Science. 32: 316-335.

172. Seike, K. and Nara, M. 2008. Burrow morphologies of the ghost crabs *Ocypode ceratophthalma* and *O. Sinensis* in foreshore, backshore, and dune subenvironments of a sandy beach in Japan. Journal of Geological Society of Japan. 114(11):591-596.

173. Serene, R. 1972. On the brachyuran fauna of the Indo-Pacific coral reefs. In: Mukundan, C., Gopindha Pillai, C.S. (Eds.), Proceedings of the First International Symposium on Corals and Coral Reefs. Marine Biological Association of India, Mandapam Camp. pp. 419-424.

174. Sethuramalingam, S. and Khan, A. S. 1991. Brachyuran Crabs of Parangipettai Centre of Advance Study in Marine Biology, Annamalai University, India.

175. Shelton, C. R., and Robertson, P. B. 1981. Community structure of intertidal macrofauna on two surf-exposed Texas sandy beaches. Bulletin of Marine Science. 31, 833-42.

176. Shih, H. T., Mok, H. K. and Chang, H. W. 2005. Chimney Building by Male *Ucaformosensis*Rathbun, 1921 (Crustacea: Decapoda: Ocypodidae) after Pairing: A New Hypothesis for Chimney Function. Zoological Studies. 44(2): 242-251.

177. Short, A.D. 1997. Overview of sandy beach morphodynamics. Revista Chilena de Historia Natural. 69: 589-604.

178. Shuchman, E. and Warburg, M. R. 1978. Dispersal, population structure and burrow shape of *Ocypode cursor*. Marine Biology. 49: 255-263.

179. Shukla, M.L., Patel, B. K., Trivedi, J.N. and Vachhrajani, K.D. 2013. Brachyuran crabs diversity of Mahi and Dhadhar estuaries, Gujarat, India. Journal of Marine Sciences. 1(2): 8-11.

180. Silas, E. G. and Sankarankutty, C. 1965. Field investigations on the shore crabs of the Gulf of Mannar and Palk bay, with special reference to the

ecology and behaviour of the pellet crab *Scopimera proxima*. Proceedings of the Symposium on Crustacea. Part 2: 1008-1025.

181. Smith, J. R., Fong, P. and Ambrose, R. F. 2006. Dramatic declines in mussel bed community diversity: response to climate change. Ecology. 87, 1153-1161.

182. Snively, G. 1978. Exploring the Seashore in British Columbia, Washington and Oregon. Gordon Soules Book Publishers, Inc.

183. Somero, G. N. 2002. Thermal physiology and vertical zonation of intertidal animals: optima, limits, and costs of living. Integrative and Comparative Biology. 42, 780-789.

184. Stephenson, T. A. and Stephenson, A. 1949. The universal features of zonation between tide-marks on rocky coast. Journal of Ecology. 37: 289-305.

185. Stephenson, T. A. and Stephenson, A. 1972. Life between tidemarks on rocky shores. W.H.Freeman and Company, San Francisco.

186. Stillman, J. H. and Somero, G. N. 2000. A comparative analysis of the upper thermal tolerance limits of eastern Pacific porcelain crabs, genus *Petrolisthes*: influences of latitude, vertical zonation, acclimation, and phylogeny. Physiological and Biochemical Zoology. 73, 200-208.

187. Stimpson, W. 1907. Report on the crustacean (Brachyura and Anomura) collected by the North Pacific Exploring Expedition, 1853-1856. (Smithsonian Institution: Washington, DC). 240 pp.

188. Strachan, P. H., Smith, R. C., Hamilton, D. A. B., Taylor, A. C. and Atkinson, R. J. A. 1999. Studies on the ecology and behavior of the ghost crab *Ocypode cursor* (L.) in Northern Cyprus. Scientia Marina. 63 (1): 51-60.

189. Subba Rao, N. V. 1991.Molluscain Animal Resources of India. Zoological Survey of India, Calcutta. 125-147.

190. Suchanek, T. H. and Green, D. J. 1981. Interspecific competition between *Palythoa caribaeorum* and other sessile invertebrates on St. Croix reefs, U.S.

Virgin Islands. Proceedings of 4th International Coral Reef Symposium. 2: 679-684.

191. Sukumaran, S. and Krishnaswamy, S. 1962. Reactions of *Cellana radiata (Born)* to salinity changes. Proceedings of the Indian Academy of Science. 54: 122-129.

192. Suseelan, C. 1972. *Ixa inremis* Leach (Crustacea: Decapoda: Brachyura) from Indian waters. Journal of Marine Biological Association of India. 22 (1-2): 159-161.

193. Takahashi, S. 1932. On the burrows of *Ocypode ceratophthalma* Fabricius. Kwagaku. 2: 329–335 (in Japanese).

194. Talley, T. S. and Dayton, P. K. 2000. Tidal flat macrofaunal communities and their associated environments in estuaries of Southern California and Northern Baja California, Mexico. Estuaries. 23(1):97-114.

195. Tan, C.G.S. and Ng, P.K.L. 1994. An annotated checklist of mangrove braychyuran crabsfrom Malaysia and Singapore. Hydrobiologia. 285: 75-84.

196. Tikader, B. K. 1965. Marine fauna of Deogadh coast (Ratnagiri district) Maharashtra, Part 2. Brachyura (Crabs). Journal of University of Poona Science and Technology Section. 30: 31-32.

197. Trivedi, J. N., Arya, S. and Vachhrajani, K. D. 2013. Gastropod shell utilization preferences of hermit crab *Clibanarius zebra* (Dana, 1852) (Diogenidae: Anomura). Taprobanica. 5(1): 12-18.

198. Trivedi, J. N., Gadhavi, M. K. and Vachhrajani, K. D. 2011. Diversity and habitat preference of brachyuran crabs in Gulf of Kutch, Gujarat, India. Arthropods. 1(1):13-23.

199. Ummerkutty, A. N. P. and Deb, M. 1972. Studies on the crustacean fauna of Maysore coast. 1. Decapoda: Brachyura. Records of Zoological Survey of India. 66 (1-4): 191-196.

200. Underwood, A. J. 2000. Experimental ecology of rocky intertidal habitats: what are we learning? Journal of Experimental Marine Biology and Ecology, 250: 51-76.

201. Underwood, A.J. and Denley, E. J. 1984. Ecological Communities, Conceptual Issues and the Evidence (eds. Strong, D.R., Simberloff, D., Abele, L.G. and Thistle, A.B.) Princeton University Press, Princeton, NH, pp. 151-180.

202. Vaghela, A. 2010. Spatial and temporal variations in population dynamics of few key rocky intertidal macrofauna at anthropogenically influenced intertidal shoreline. Ph. D. Thesis, Saurashtra University, Gujarat, India.

203. Venkataraman, K. and Wafar, M. 2005. Coastal and marine biodiversity of India, Indian Journal of Marine Sciences. Vol. 34(1) pp.57-75.

204. Vogel, F. 1983. A redescription of *Dotilla malabarica* Nobili, 1903 (Decapoda: Brachyura: Ocypodidae). Crustaceana. 45(2): 157-163.

205. Wethey, W. S. 1983. Eographic limits and local zonation: the barnacles *Semibalanus* (*Balanus*) and *Chthamalus* in New England. The Biological Bulletin. 165: 330-341.

206. Williams, E. E. 1964. The growth and distribution of *Littorina littorea* (L.) on a rocky shore in wales. Journal of Animal Ecology. 33: 413-432.

207. Wolfrath, B. 1992. Burrowing of the fiddler crab, *Uca tangeri*, in the Ria Formosa in Portugal and its influence on sediment structure. MarineEcology Progress Series. 85: 237-243.

208. Xin, P., Jin, G., Li, L. and Barry, D. A. 2009. Effects of crab burrows on pore water flows in salt marshes, Advances in Water Resources. Advances in Water Resource. 32 (3): 439-449.

209. Xu, L., Sun, H., Wu, G. and Feng, Q. 1992. A preliminary ecological survey on intertidal benthic animals of a sandy beach along the coast of Chengmaijiao, Hainan Island. Tropical Oceanology. 11(1): 15.21.

210. Young, A. M. 1980. Temperature-Salinity tolerances of three hermit crab species. *Clibanarius vittatus* (Bsc.) *Pagurus longicarous* Say and, *P. Pollicaris* say (Crustacea: Decapoda: Anomura) Ophelia. 19:27-35.