LIST OF FIGURES

Figures	Title	Page
Figure 1.1	Log scale showing size of nanoparticles (NPs) compared to biological component	3
Figure 1.2	Types of nanomaterials by size of their structural elements	4
Figure 1.3	Various shapes of nanoparticles	5
Figure 1.4	Aggregation/Agglomeration state of nanoparticles	6
Figure 1.5	Sources of natural and engineered nanoparticles	8
Figure 1.6	Types of engineered nanoparticles	10
Figure 1.7	Number of products containing commercial NMs	11
Figure 1.8	Possible ways through which nanoparticles enter into environment and humans	17
Figure 1.9	Physicochemical properties of engineered nanomaterials leading to nanotoxicology	18
Figure 1.10	Possible mechanism of TiO_2 NPs induced cellular toxicity	29

Figure 2.1	The mean hydrodynamic diameter of metal oxide nanoparticles	45
Figure 2.2	Percent hemolysis of chicken RBCs incubated with different metal oxide nanoparticles	46
Figure 2.3	The cytotoxic potential of metal oxide nanoparticles on chicken embryonic fibroblasts	47
Figure 2.4	LDH release in chicken embryonic fibroblasts	48
Figure 2.5 (A)	Phase contrast images depicting the changes in the morphology of chicken embryonic fibroblasts	48
Figure 2.5 (B)	Fluorescent images of chromosome condensation by DAPI staining in chicken embryonic fibroblasts	48
Figure 2.6	Florescence photomicrographs of DCF-DA and RHO-123 stained fibroblast cells	49
Figure 2.7	Semi-quantitative RT-PCR analysis of apoptotic genes in chicken embryonic fibroblasts	50
Figure 2.8	Possible mechanism of ZnO NPs induced cellular toxicity	54
Figure 3.1	Reaction of CAM to metal oxide nanoparticles exposure	64
Figure 3.2	Chicken embryonic development after 19 days of incubation in metal oxide nanoparticles treated groups	65-66
Figure 3.3 (A, B)	Percentage viability and morphometry of TiO_2 NPs-treated chicken embryos	67

Figure 3.4 (A, B)	Percentage viability and morphometry of Fe_2O_3 NPs-treated chicken embryos	69
Figure 3.5 (A, B)	Percentage viability and morphometry of ZnO NPs-treated chicken embryos	71
Figure 3.6 (A, B)	Percentage viability and morphometry of MgO NPs-treated chicken embryos	73
Figure 3.6 (A, B)	Percentage viability and morphometry of Al_2O_3 NPs-treated chicken embryos	75
Figure 4.1	Schematic drawing of nanoparticles induced cytotoxicity	83
Figure 4.2	Borosil Mansingh Survismeter	84
Figure 4.3	Schematic diagram of Borosil Mansingh Survismeter	86
Figure 4.4 (A-F)	Density, Viscosity, Surface tension, Friccohesity, Activation energy and Molecular radii (A-F) of Fe_2O_3 NPs in egg albumen	96-97
Figure 4.5 (A-F)	Density, Viscosity, Surface tension, Friccohesity, Activation energy and Molecular radii (A-F) of TiO_2 NPs in egg albumen	100-101
Figure 5.1	FTIR spectra of native egg albumen and TiO_2 NPs + Albumen	118-120

Figure 5.2	Concentration of Ti in embryos of control and TiO_2 NPs-treated group (4 day old) by ICP-OES.	101
Figure 5.3	Embryonic development in TiO_2 NPs-treated chickens	121
Figure 5.4	Photographs showing endoskeleton (bone and cartilage) of 19-day-old chick embryos	121
Figure 5.5	Expression of Wnt signaling pathway-related genes in TiO_2 NPs-treated chicken embryos	122-123
Figure 5.6	Expression of limb development gene HOXD13 in TiO_2 NPs-treated chicken embryos	124
Figure 5.7	Somite numbers in control and TiO_2 NPs-treated embryos 24 h after treatment in shell-less culture (HH-23)	124
Figure 5.8	The possible mechanism of TiO2 NPs induced omphalocele and flexed limbs in chicken embryo	130
Figure 6.1	Comparative FTIR spectra of native egg albumen and Fe_2O_3 NPs + Albumen	140
Figure 6.2 (A,B)	Gaussian curve fitting of secondary derivative of Native egg Albumen (A) and Fe_2O_3 NPs + egg Albumen (B)	141
Figure 6.3 (A)	Digitally enlarged images of healthy neuron and degenerating neuron	142
Figure 6.3 (B)	Histopathological evaluation of brain sections (5 µm) stained with HXE of Fe_2O_3 NPs-treated chicken embryos	143
Figure 6.3 (C)	Percentage neuronal loss in the brain of Fe_2O_3 NPs-treated chicken embryos	144

Figure 6.4 (A,B)	Percentage viability and morphometry of Fe_2O_3 NPs-immersed chicken embryos	145
Figure 6.5 (A)	Histopathological evaluation of brain sections (5 µm) stained with HXE of Fe_2O_3 NPs-immersed chicken embryos	147
Figure 6.5 (B)	Percentage neuronal loss in the brain of Fe_2O_3 NPs-immersed chicken embryos	148
Figure 7.1	Summary of screening of metal oxide nanoparticles for possible toxicity *in ovo* and in *in vitro*	155
Figure 7.2	Possible mechanism of TiO_2 and Fe_2O_3 NPs induced toxicity to chicken embryo	158