

LIST OF FIGURE

Figure no.	Caption	Page no.
FIGURE 2.1	Anatomy of skin	13
FIGURE 2.2	Illustration of different types of wound	15
FIGURE 2.3	Phases of wound healing process	16
FIGURE 2.4	<i>Calendula officinalis</i> L (Pot Marigold): (a) plant and (b) flower	45
FIGURE 3.1	DSC thermogram of <i>Calendula</i> flower extract	83
FIGURE 3.2	FT-IR spectra of <i>Calendula</i> flower extract	84
FIGURE 3.3	Standard calibration plot of Gallic acid solution	89
FIGURE 3.4	Standard curve of quercetin dehydrate solution	92
FIGURE 3.5	Standard calibration curve of %Inhibition vs. Concentration of quercetin solution	95
FIGURE 4.1	Preparation of CFE loaded PVA hydrogel sheet by γ -rays irradiation	100
FIGURE 4.2	Measurement of mechanical properties of CFE-Hydrogel sheet by CT3 Texture analyzer	103
FIGURE 4.3	Response surface (3D) plots showing combined effect of (a) PVA content and carrageenan content, (b) PVA content and radiation dose, (c) CFE content and PVA content (d) radiation dose and carrageenan content, (e) CFE content and carrageenan content and (f) CFE content and radiation dose on % fluid absorptivity of hydrogel sheet	115
FIGURE 4.4	Response surface (3D) plot showing combined effect of (a) PVA content and carrageenan content; (b) PVA content and radiation dose; (c) CFE content and PVA content; (d) Radiation dose and carrageenan content; (e) CFE content and carrageenan content; and (f) CFE	120

LIST OF FIGURE

Figure no.	Caption	Page no.
	content and radiation dose on gel fraction of hydrogel sheet	
FIGURE 4.5	Response surface (3D) plot showing combined effect of (a) PVA content and carrageenan content; (b) PVA content and radiation dose; (c) CFE content and PVA content; (d) radiation dose and carrageenan content; (e) CFE content and carrageenan content; and (f) CFE content and radiation dose on hardness of hydrogel sheet	124
FIGURE 4.6	Software generated overlaid plot of design space	126
FIGURE 4.7	Physical appearance of calendula flower extract loaded hydrogel sheet	127
FIGURE 4.8	Result of hardness study of CFE loaded hydrogel sheet by Texture analyzer	128
FIGURE 4.9	Load vs. distance plot generated by texture analyzer	129
FIGURE 4.10	FT IR spectra of Calendula flower extract	131
FIGURE 4.11	FT IR spectra of polyvinyl alcohol	132
FIGURE 4.12	FT-IR spectra of carrageenan	132
FIGURE 4.13	FT IR spectra of physical mixture of PVA, carrageenan, CFE	133
FIGURE 4.14	FT IR spectra of CFE loaded hydrogel sheet	133
FIGURE 4.15	DSC thermogram of CFE	134
FIGURE 4.16	DSC thermogram of PVA	135
FIGURE 4.17	DSC thermogram of carrageenan	135
FIGURE 4.18	DSC thermogram of physical mixture of CFE, PVA, carrageenan	136

LIST OF FIGURE

Figure no.	Caption	Page no.
FIGURE 4.19	DSC thermogram of CFE loaded hydrogel sheet	136
FIGURE 5.1	Flow diagram for the preparation of CFE-Collagen film	145
FIGURE 5.2	Gel Electrophoresis: preparation of separating gel and stacking gel	151
FIGURE 5.3	CFE-collagen film	152
FIGURE 5.4	Standard calibration plot of Gallic acid solution	156
FIGURE 5.5	Standard curve of quercetin dehydrate solution	157
FIGURE 5.6	Standard calibration curve of %Inhibition vs. Concentration of quercetin solution	158
FIGURE 5.7	SDS-PAGE profile of (1) collagen, (2) collagen film and (3) marker protein	160
FIGURE 5.8	Optical microscopy of CFE-collagen film (magnification 20x)	162
FIGURE 5.9	Atomic force microscope topography images of collagen film (i) 2D images, (ii) 3D images and (iii) Height-profile taken of the dashes line part shown in figure 5.9 (i) 2D image	163
FIGURE 5.10	DSC thermogram of CFE	164
FIGURE 5.11	DSC thermogram of Collagen	165
FIGURE 5.12	DSC thermogram of EDTA	166
FIGURE 5.13	DSC thermogram of CFE-collagen film	167
FIGURE 5.14	FT-IR spectra of CFE	168

LIST OF FIGURE

Figure no.	Caption	Page no.
FIGURE 5.15	FT-IR spectra of Collagen	169
FIGURE 5.16	FT-IR spectra of CFE-Collagen film	170
FIGURE 6.1	<i>In-vitro</i> cumulative total polyphenol compound release profile from CFE loaded hydrogel sheet	178
FIGURE 6.2	Hixon-crowell cube root release kinetics profile of total polyphenol release from CFE loaded hydrogel sheet	180
FIGURE 6.3	First order release kinetics profile of total polyphenol release from CFE loaded hydrogel sheet	180
FIGURE 6.4	Korsmeyer peppas release kinetics profile of total polyphenol release from CFE loaded hydrogel sheet	181
FIGURE 6.5	Higuchi release kinetics profile of total polyphenol release from CFE loaded hydrogel sheet	182
FIGURE 6.6	Zero order release kinetics profile of total polyphenol release from CFE loaded hydrogel sheet	182
FIGURE 7.1	Swiss 3T3 albino fibroblasts	186
FIGURE 7.2	Result of <i>In-vitro</i> cell viability study with CFE and CFE loaded hydrogel sheet	190
FIGURE 7.3	Result of <i>in-vitro</i> cell viability study with CFE loaded collagen film	191
FIGURE 7.4	Single representative area of cell well after wounding (a) Only media (control), (b) CFE, (c) extract of collagen film without CFE, (d) extract of CFE loaded collagen film, (e) extract of hydrogel sheet with CFE and (f) extract of CFE loaded collagen film	193
FIGURE 8.1	Animal study certificate from IAEC for the cutaneous excision wound healing model	198

LIST OF FIGURE

Figure no.	Caption	Page no.
FIGURE 8.2	Animal study certificate from IAEC for the burn wound healing model	199
FIGURE 8.3	Rat with shaved back side	200
FIGURE 8.4	Rat having skin incision (2cm x 2 cm)	201
FIGURE 8.5	Rat's wound covered with secondary dressing	201
FIGURE 8.6	Macroscopic changes of wound over different post-excision days of Group I (non treated animal), Group II (collagen film without CFE treatment), Group III (CFE loaded collagen film treatment) and Group IV (market product treatment)	205
FIGURE 8.7	% Wound contraction in rat during 21 days <i>in-vivo</i> animal study: I (non treated animal), Group II (Collagen film without CFE treatment), Group III (CFE loaded collagen film treatment) and Group IV (market product treatment)	206
FIGURE 8.8	Photomicrograph of histopathological wound tissue section: (a) Non treated group, (b) Placebo collagen film treated group, (c) Market product treated group and (d) collagen film containing CFE treated group	207
FIGURE 8.9	Rat having shaved back side	209
FIGURE 8.10	Rat having burn wound (Day 0)	209
FIGURE 8.11	Macroscopic changes of wound over different post-excision days of Group I(non treated animal), Group II (Hydrogel sheet without CFE treatment), Group III (CFE loaded hydrogel sheet treatment) and Group IV (market product treatment)	213
FIGURE 8.12	% Wound contraction in rat during 21 days <i>in-vivo</i> animal study: I (non treated animal), Group II (Hydrogel sheet without CFE treatment), Group III (CFE loaded	215

LIST OF FIGURE

Figure no.	Caption	Page no.
	hydrogel sheet treatment) and Group IV (market product treatment)	
FIGURE 8.13	Histopathology study of skin after 21 days (a) Non treated group (b) Placebo hydrogel sheet treated group (c) Market product treated group (d) Hydrogel sheet containing CFE treated group	216