

List of Tables		Page No.
Table 1.1	Rational Design of an “Ideal” Nanocarrier for Cancer Therapy Taking into Account all the Biological Barriers and Requirements	3
Table 2.1	Molecular characteristics of the block copolymer	58
Table 3.1	Radius of Gyration (R_g), Hydrodynamic Radii (R_H), and Polydispersity Index	72
Table 3.2	Effect of initial concentration of copolymer in DMSO and final concentration of copolymer in nanoprecipitation	76
Table 4.1	Loading content of PolyDOX at different feed weight ratios of DOX/block copolymer	96
Table 4.2	Stability study of PolyDOX at RT and 4°C	103
Table 4.3	<i>In Vitro</i> half maximal inhibitory concentration (IC_{50}) values measured for free DOX and PolyDOX on C6 cells at different incubation time	106
Table 5.1	<i>In Vitro</i> Cytotoxicity of PolyDOX and Free DOX (Concentration in μM). IC_{50}	129
Table 6.1	<i>In vitro</i> stability of ^{99m}Tc -labeled compound in presence of DTPA	150
Table 6.2	<i>In vitro</i> stability of ^{99m}Tc -labeled compounds in PBS and serum	151
Table 6.3	<i>In vitro</i> stability after 100 times dilution of ^{99m}Tc -labeled compounds	151
Table 6.4	<i>In vivo</i> stability of ^{99m}Tc -labeled compounds	151
Table 6.5	Pharmacokinetic parameters of ^{99m}Tc -Labeled compounds in rabbits	154
Table 6.6	Body weight changes in treated mice at different doses of DOX, PolyDOX and Blank-POLY after 25 days from first injection	160
Table 6.7	Blood parameters in healthy mice on day 7 after last intravenous doses of PolyDOX injections (5mg/kg/day), (4×5mg/kg/day), (3×5mg/kg/day), free DOX injection (3×5mg/kg/day), (5mg/kg/day) and Polymersomes (20mg/kg/day)	161
Table 6.8	Blood parameters in healthy mice on day 21 after last intravenous doses of PolyDOX injections (5mg/kg/day), (4×5mg/kg/day), (3×5mg/kg/day), free DOX injection, (5mg/kg/day) and Polymersomes (20mg/kg/day)	162

Table 6.9	Serum biochemistry in healthy BalB/c mice on day 7 and 21 after last intravenous doses of PolyDOX injections (5mg/kg/day), (4×5mg/kg/day), (3×5mg/kg/day), free DOX injections (3×5mg/kg/day), (5mg/kg/day) and Polymersomes (20mg/kg/day)	163
Table 7.1	Drug loading content and efficacy of PolyDOC	186
Table 7.2	DLS result of PolyDOC formulations before and after probe sonication (10 minute)	187
Table 7.3	<i>In Vitro</i> Cytotoxicity of PolyDOC and Free DOC (Concentration in nM). IC ₅₀	192
Table 7.4	<i>In vitro</i> stability of ^{99m} Tc-labelled compound in presence of DTPA	194
Table 7.5	<i>In vitro</i> stability of ^{99m} Tc-labelled compound in PBS and serum	194
Table 7.6	<i>In vitro</i> stability after 100 times dilution of ^{99m} Tc-labeled compounds	194
Table 7.7	<i>In vivo</i> stability of ^{99m} Tc-labeled compounds	195
Table 7.8	Pharmacokinetic parameters of ^{99m} Tc-Labeled compounds in rabbits	196