



LIST OF TABLES

Table No.	Title	Page No.
1.1	Important medicinal plants used in liver disorders	5
2.1	Chemical constituents isolated from various parts of <i>F. limonia</i>	13-15
2.2	Chemical constituents isolated from various parts of <i>T. undulata</i>	29-31
4.1	Characteristic macroscopic features of leaves stem bark and root bark of <i>F. limonia</i>	73
4.2	Quantitative microscopic features of <i>Feronia limonia</i> leaves	74
4.3	Leaf constants of <i>Feronia limonia</i>	75
4.4	Physico-chemicals constants of leaves, stem bark and root bark of <i>Feronia limonia</i>	79
4.5	Elemental analysis of leaves, stem bark and root bark of <i>Feronia limonia</i>	80
4.6	Successive extractive values of the leaves, stem bark and root bark of <i>Feronia limonia</i>	82
4.7	Preliminary phytochemical investigation of <i>Feronia limonia</i> leaves, stem bark and root bark	83
4.8	Quantitative evaluation of total phenolic, flavanoid and flavanol content of <i>Feronia limonia</i> leaves	84
4.9	Method validation parameter for quantification of MR-1 using proposed HPTLC densitometric method	101
4.10	Method validation parameters for quantification of MR-1 using proposed HPTLC densitometric method	107
4.11	Chemical shifts and multiplicity of MR-2 ¹³ C NMR spectra	113
4.12	Effect of FLs and sylimarin on CCL ₄ induced hepatotoxicity	118

4.13	Effect of FSBs and sylimarin on CCL ₄ induced hepatotoxicity	119
4.14	Effect of FRBs and sylimarin on CCL ₄ induced hepatotoxicity	120
4.15	Effect of FL-7, FL-9, MR-2 and sylimarin on hepatic antioxidants level and lipid peroxidation during CCL ₄ induced hepatotoxicity	127
4.16	Effect of FSB-7 and sylimarin on hepatic antioxidants and lipid peroxidation during CCL ₄ induced hepatotoxicity	128
4.17	Effects of FRB-7 and MR-1 on hepatic antioxidants and lipid peroxidation during CCL ₄ induced hepatotoxicity	129
4.18	Effects of FL-7, FL-9, MR-2 and sylimarin on plasma levels of hepatic injury markers during CCL ₄ induced hepatotoxicity	130
4.19	Effects of FSB-7 and sylimarin on plasma levels of hepatic injury markers during CCL ₄ induced hepatotoxicity	131
4.20	Effects of FRB-7 and MR-1 on plasma levels of hepatic injury markers during CCL ₄ induced hepatotoxicity	132
5.1	Characteristic macroscopic features of <i>Tecomella undulata</i> leaves and stem bark	137
5.2	Physicochemical analysis of <i>Tecomella undulata</i> leaves and stem bark	142
5.3	Elemental analysis values of <i>Tecomella undulata</i> leaves and stem bark	143
5.4	Quantitative evaluation of total phenolic, flavanoid and flavanol content of leaves and stem bark of <i>Tecomella undulata</i>	143
5.5	Successive extractive values of <i>Tecomella undulata</i> leaves	144
5.6	Preliminary phytochemical investigations of <i>Tecomella undulata</i> leaves and stem bark	145
5.7	TLC studies on successive extracts of leaves <i>Tecomella</i>	146

	<i>undulata</i>	
5.8	TLC studies on successive extracts of stem bark	147
	<i>Tecomella undulata</i>	
5.9	Method validation parameter for quantification of MS-1 using proposed HPTLC Densitometric method	158
5.10	Effect of <i>Tecomella undulata</i> leaf extracts, fractions and sylimarin on CCl ₄ induced hepatotoxicity	166
5.11	Effect of <i>Tecomella undulata</i> stem bark extracts, fractions, MS-2 and sylimarin on CCl ₄ induced hepatotoxicity	167
5.12	Effect of TL-2 and sylimarin on hepatic antioxidant level and lipid peroxidation during CCl ₄ induced hepatotoxicity	172
5.13	Effect of TSB-7, TSB-9, MS-2 and sylimarin on hepatic antioxidants and lipid peroxidation during CCl ₄ induced hepatotoxicity	173
5.14	Effects of TL-2 and sylimarin on plasma levels of hepatic injury markers during CCl ₄ induced hepatotoxicity	174
5.15	Effects of TSB-7, TSB-9, MS-2 and sylimarin on plasma levels of hepatic injury markers during CCl ₄ induced hepatotoxicity	174