

LIST OF FIGURES

Fig. No.	Title	Page No.
1.1	Application of chemical markers in research area	3
2.1	Taxonomical classification and photograph of <i>Feronia limonia</i>	8
2.2	Vernacular names and photograph of <i>Feronia limonia</i>	8
2.3	Graphical outline containing traditional uses, phytochemistry and pharmacology of <i>Feronia limonia</i>	9
2.4	Chemical structures of isolated constituents from <i>Feronia limonia</i>	18
2.5	Taxonomical classification, common names and photographs of leaves, Flower and plant of <i>Tecomella undulata</i>	25
2.6	Graphical outline containing traditional uses, phytochemistry and pharmacology of <i>Tecomella undulata</i>	27
2.7	Chemical structures of isolated constituents from <i>Tecomella undulata</i>	31
2.8	Diagram depicting anterior surface and posterior surface of Liver	35
2.9	Figure depicting various mechanisms of Injury to hepatocytes	37
2.10	Flow diagram of various liver function tests	39
2.11	Mechanisms of drug-induced Liver Injury	48
3.1	Schematic diagram of <i>in vitro</i> cytotoxicity assay	66
3.2	Schematic protocol of <i>in vivo</i> hepatotoxicity studies	68
4.1	Photographs of stem bark, leaves and root bark of <i>Feronia limonia</i>	72
4.2	Transverse sections showing various characteristic features of <i>Feronia limonia</i> leaves	75
4.3	Transverse section showing stomata of <i>Feronia limonia</i> leaf	75
4.4	Transverse sections showing various characteristic features of <i>F. limonia</i> stem bark	76
4.5	Transverse sections showing various characteristic features of <i>Feronia limonia</i> root bark	77
4.6	Various powder microscopic features of <i>Feronia limonia</i> leaves	78

4.7	Various powder microscopic features of <i>F. limonia</i> stem bark	78
4.8	Various powder microscopic features of <i>Feronia limonia</i> root bark	78
4.9	Fluorescence analysis of the methanolic extract of <i>Feronia limonia</i> leaves	81
4.10	Fluorescence analysis of the methanolic extract of stem bark	81
4.11	Fluorescence analysis of the methanolic extract of root bark	81
4.12	TLC of the extracts obtained from leaves in successive extraction at 254 nm	85
4.13	TLC of the extracts obtained from leaves in successive extraction at 366 nm	85
4.14	TLC of the extracts obtained from leaves in successive extraction at 540 nm	85
4.15	TLC of the extracts obtained from stem bark in successive extraction at 254 nm	86
4.16	TLC of the extracts obtained From stem bark in successive extraction at 366 nm	86
4.17	TLC of the extracts obtained from stem bark in successive extraction at 540 nm	86
4.18	TLC of the extracts obtained from root bark in successive extraction at 254 nm	87
4.19	TLC of the extracts obtained from root bark in successive extraction at 366 nm	87
4.20	TLC of the extracts obtained from root bark in successive extraction at 540 nm	87
4.21	HPLC chromatogram of <i>Feronia limonia</i> leaves (methanolic extract) at 254 nm	88
4.22	HPLC chromatogram of <i>Feronia limonia</i> leaves (methanolic extract) at 366 nm	88
4.23	HPLC chromatogram of <i>Feronia limonia</i> stem bark (methanolic extract) at 254 nm	89
4.24	HPLC chromatogram of <i>Feronia limonia</i> stem bark (methanolic extract) at 366 nm	89

4.25	HPLC chromatogram of <i>Feronia limonia</i> root bark (methanolic extract) at 254 nm	90
4.26	HPLC chromatogram of <i>Feronia limonia</i> root bark (methanolic extract) at 366 nm	90
4.27	Graphical representation of MR-2 isolation from <i>Feronia limonia</i> leaves	91
4.28	Graphical representations of MR-1 and MR-2 isolation from <i>Feronia limonia</i> stem bark	92
4.29	Graphical representation of MR-1 isolation from <i>Feronia limonia</i> root bark	93
4.30	TLC fingerprinting and three-dimensional overlaid chromatogram of extracts, fractions and isolated compound from <i>Feronia limonia</i> leaf at UV 366 nm.	
4.31	TLC fingerprinting and three-dimensional overlaid chromatogram of extracts, fractions and isolated compound from <i>Feronia limonia</i> stem bark at UV 366 nm.	95
4.32	TLC fingerprinting and three-dimensional overlaid chromatogram of extracts, fractions and isolated compound from <i>Feronia limonia</i> root bark at UV 366 nm	95
4.33	HPLC chromatogram of isolated compound MR-1	96
4.34	HPLC chromatogram of isolated compound MR-2	96
4.35	TLC fingerprinting and chromatogram for detection of total flavanoids in extracts and fractions of <i>Feronia limonia</i> leaves at UV 366 nm	97
4.36	TLC fingerprinting and chromatogram for detection of total flavanoids in extract of <i>Feronia limonia</i> stem bark at UV 366 nm	98
4.37	TLC fingerprinting and chromatogram for detection of total flavanoids in extract of <i>Feronia limonia</i> root bark at UV 366 nm	98
4.38	TLC chromatogram of stem bark methanolic extract of <i>Feronia Limonia</i>	102
4.39	TLC chromatogram, for a standard MR-1 in methanol	102
4.40	Calibration curve and HPLC chromatogram of isolated compound MR-1	104

4.41	HPLC chromatogram of FL-7 at UV 280 nm	104
4.42	HPLC chromatogram of FSB-7 at UV 280 nm	105
4.43	HPLC chromatogram of FRB-7 at UV 280 nm	105
4.44	Chemical structure of MR-1	108
4.45	IR spectrum of MR-1	109
4.46	¹³ C NMR of MR-1	110
4.47	Mass spectrum of MR-1	110
4.48	CHN analysis of MR-1	111
4.49	Oxygen analysis of MR-1	111
4.50	Chemical structure of MR-2	112
4.51	IR spectrum of MR-2	112
4.52	¹³ C NMR of MR-2	113
4.53	Mass spectrum of MR-2	114
4.54	Cytotoxicity evaluation of extracts, fractions and isolated compound from <i>Feronia limonia</i> leaves	116
4.55	Cytotoxicity evaluation of extracts, fractions and isolated compound from <i>Feronia limonia</i> stem bark	116
4.56	Cytotoxicity evaluation of extracts, fractions and isolated compound from <i>Feronia limonia</i> rootbark	116
4.57	Phase contrast and AO/EB stained photomicrographs of HepG2 cells exposed to 1% CCL ₄ alone or in presence of FLs or sylimarin	123
4.58	Phase contrast and AO/EB stained photomicrographs of HepG2 cells exposed to 1% CCL ₄ alone or in presence of FSB7 or sylimarin	124
4.59	Phase contrast and AO/EB stained photomicrographs of HepG2 cells exposed to 1% CCL ₄ alone or in presence of FRB7, MR-1 or sylimarin.	125
4.60	Morphological, haematoxyline-eosin and picro-sirus stained photomicrographs of rat liver treated with CCL ₄ alone or in presence of FSB-7 or sylimarin	133
4.61	Morphological, haematoxyline-eosin and picro-sirus stained photomicrographs of rat liver treated with CCL ₄ alone or in	134

	presence of FSB-7 or sylimarin	
4.62	Effect of FRB-7, MR and sylimarin on hepatic collagen deposition (picro-sirus red staining) during CCL ₄ induced hepatotoxicity	135
4.63	Effect of FRB-7, MR and sylimarin on hepatic histopathological alterations during CCL ₄ induced hepatotoxicity	135
5.1	Photographs of leaves and stem bark of <i>Tecomella undulata</i>	136
5.2	Charestiristic features of Transverse section of <i>Tecomella undulata</i> leaves	139
5.3	Charestiristic features of powder microscopy of <i>Tecomella undulata</i> leaves	141
5.4	Charecteristic features of transverse section of <i>Tecomella undulata stem bark</i>	140
5.5	Characteristic features of powder microscopy of <i>Tecomella undulata stem bark</i>	141
5.6	HPLC chromatogram of <i>Tecomella undulata</i> leaves methanolic extract at 254 nm	148
5.7	HPLC chromatogram of <i>Tecomella undulata</i> leaves methanolic extract at 366 nm	148
5.8	HPLC chromatogram of <i>Tecomella undulata</i> stem bark methanolic extract at 254 nm	149
5.9	chromatogram of <i>Tecomella undulata</i> stem bark methanolic extract at 366 nm	149
5.10	HPLC chromatogram of <i>T. undulata</i> leaves methanolic extract at 205 nm	151
5.11	HPLC chromatogram of betulinic acid at 205 nm	152
5.12	HPLC chromatogram of <i>T. undulata</i> leaves extract, betulinic acid and Leaves sample after adding betulinic acid at 205 nm	152
5.13	HPLC chromatogram of <i>T. undulata</i> stem bark (TSB-7), betulinic acid and TSB-7 after adding betulinic acid at 205 nm	154
5.14	HPLC chromatogram of <i>T. undulata</i> ethyl acetate fraction (TSB-9) at 205 nm	154
5.15	Representing chromatograms and calibration curve for isolated	156

	marker (MS-1; 50-250µg/ml)	
5.16	HPLC chromatogram showing separation of MS-1 in TSB-7 at UV 262 nm	156
5.17	Chemical structure of MS-1	159
5.18	IR spectrum of isolated compound (MS-1)	160
5.19	Overlaid IR spectrum of isolated compound (MS-1) and standard lapachol	160
5.20	Chemical structure of MS-2	161
5.21	IR spectrum of MS-2	161
5.22	Overlaid IR spectra of MS-2 along with standard betulinic acid	162
5.23	Overlaid UV spectrum of MS-2 along with standard betulinic acid	162
5.24	Cytotoxicity assessment of <i>Tecomella undulata</i> TLs	164
5.25	Cytotoxicity assessment of <i>Tecomella undulata</i> TSBs	164
5.26	Phase contrast and AO/ EB stained photomicrographs of HepG2 cells exposed to 1% CCL ₄ alone / in presence of TL-2 or sylimarin	169
5.27	Phase contrast and AO/ EB stained photomicrographs of HepG2 cells exposed to 1% CCL ₄ alone / in presence of TL-2 or sylimarin	170
5.28	Morphological, haematoxyline-eosin photomicrographs of rat liver treated with CCL ₄ alone / in presence of TL-2 or sylimarin	176
5.29	Morphological, haematoxyline-eosin photomicrographs of rat liver treated with CCL ₄ alone / in presence of TSB-7, TSB-9, MS-2 or sylimarin	177