

List of Figures

Figure 1.1	Graphical representation of Spherulites assembly	5
Figure 2.1	A complete, schematic view of the human respiratory system with their parts and functions.	10
Figure 2.2	Schematic representation of normal cell cycle.	14
Figure 2.3	Cellular death due to damage or alteration in normal cell cycle	15
Figure 2.4	Cancer cell division.	15
Figure 2.5	Types of cancer.	17
Figure 2.6	Share of population with cancer in 2016.	19
Figure 2.7	Share of population suffering from cancer types, World, 2016.	19
Figure 2.8	Types of Lung cancer.	27
Figure 2.9	Graphical representation of novel drug delivery systems; A) Polymeric nanoparticles, B) Polymeric micelles, C) Dendrimers, D) Liposomes, E) Viral-based nanoparticles, F) Carbon nanotubes.	33
Figure 2.10	Advantages of PEGylation of a novel drug delivery system.	39
Figure 3.1	Overlay plot of Gemcitabine Hydrochloride in distilled water at 266 nm.	70
Figure 3.2	Calibration plot of Gemcitabine Hydrochloride in distilled water.	70
Figure 3.3	Overlay plot of Gemcitabine Hydrochloride in PBS pH 7.4 at 269 nm.	72
Figure 3.4	Calibration plot of Gemcitabine Hydrochloride in PBS pH 7.4.	72
Figure 3.5	Chromatograms of Gemcitabine Hydrochloride calibration plot showing retention time of 2.89 min (A- Blank, B- 20 ng/mL, C- 100 ng/mL, D- 250 ng/mL, E- 500 ng/mL, F- 1000 ng/mL, G- 2000 ng/mL).	76
Figure 3.6	Standard Calibration plot of Gemcitabine Hydrochloride in rat plasma by LCMS MS method.	77

Figure 3.7	Standard Calibration plot of Gemcitabine Hydrochloride extracted from rat heart homogenate by LCMS-MS method.	78
Figure 3.8	Standard Calibration plot of Gemcitabine Hydrochloride extracted from rat lungs homogenate by LCMS-MS method.	80
Figure 3.9	Standard Calibration plot of Gemcitabine Hydrochloride extracted from rat liver homogenate by LCMS-MS method.	81
Figure 3.10	Standard Calibration plot of Gemcitabine Hydrochloride extracted from rat kidney homogenate by LCMS-MS method.	83
Figure 3.11	Standard Calibration plot of Gemcitabine Hydrochloride extracted from rat spleen homogenate by LCMS-MS method.	84
Figure 3.12	Overlay plot of Vinorelbine Tartrate in distilled water at 271 nm.	86
Figure 3.13	Calibration plot of Vinorelbine Tartrate in distilled water.	86
Figure 3.14	Overlay plot of Vinorelbine Tartrate in PBS pH 7.4 at 273 nm.	88
Figure 3.15	Calibration plot of Vinorelbine Tartrate in PBS pH 7.4.	88
Figure 3.16	Chromatograms of Vinorelbine tartrate calibration plot showing retention time of 2.72 min (A- Blank, B- 5 ng/mL, C- 10 ng/mL, D- 50 ng/mL, E- 100 ng/mL, F- 200 ng/mL, G- 500 ng/mL).	92
Figure 3.17	Standard Calibration plot of Vinorelbine tartrate in rat plasma by LCMS-MS method.	93
Figure 3.18	Standard Calibration plot of Vinorelbine tartrate extracted from rat heart homogenate by LCMS-MS method.	94
Figure 3.19	Standard Calibration plot of Vinorelbine tartrate extracted from rat lungs homogenate by LCMS-MS method.	96
Figure 3.20	Standard Calibration plot of Vinorelbine tartrate extracted from rat liver homogenate by LCMS-MS method.	97
Figure 3.21	Standard Calibration plot of Vinorelbine tartrate extracted from rat kidney homogenate by LCMS-MS method.	99
Figure 3.22	Standard Calibration plot of Vinorelbine tartrate extracted from rat spleen homogenate by LCMS-MS method.	100
Figure 4.1	Various types of Preformulation studies.	103

Figure 4.2	FTIR Spectra of GCH.	106
Figure 4.3	DSC thermogram of GCH.	107
Figure 4.4	FTIR Spectra of physical mixture of GCH and excipients.	108
Figure 4.5	DSC thermogram of physical mixture of GCH and excipients.	109
Figure 4.6	FTIR Spectra of VLB.	110
Figure 4.7	DSC thermogram of VLB.	111
Figure 4.8	FTIR Spectra of physical mixture of VLB and excipients.	111
Figure 4.9	DSC thermogram of physical mixture of VLB and excipients.	112
Figure 4.10	Microscopic images of preliminary optimization batches of spherulites prepared by varying concentration of SPC; A) 30-46%, B) 48-62%.	114
Figure 4.11	Homogenizer tube; A) with smooth inner surface, B) with serrated inner surface and C) Teflon probe.	118
Figure 5.1	Ishikawa diagram showing the developmental process of GCH loaded Spherulites.	125
Figure 5.2	Flow chart depicting the preparation of GCH loaded Spherulites.	126
Figure 5.3	Response surface plot depicting the effect of independent variables on GCH loaded spherulites size.	130
Figure 5.4	Bubble plot depicting the effect of independent variables on GCH loaded spherulites size.	130
Figure 5.5	Response surface plot depicting the effect of independent variables on GCH loaded spherulites %EE.	132
Figure 5.6	Bubble plot depicting the effect of independent variables on GCH loaded spherulites %EE.	133
Figure 5.7	Desirability Plot for Optimization of GCH loaded spherulites.	134
Figure 5.8	Ishikawa diagram showing the developmental process of VLB loaded Spherulites.	136
Figure 5.9	Flow chart depicting the preparation of VLB loaded Spherulites.	137

Figure 5.10	Response surface plot depicting the effect of independent variables on VLB loaded spherulites size.	140
Figure 5.11	Bubble plot depicting the effect of independent variables on VLB loaded spherulites size.	141
Figure 5.12	Response surface plot depicting the effect of independent variables on VLB loaded spherulites %EE.	142
Figure 5.13	Bubble plot depicting the effect of independent variables on VLB loaded spherulites %EE.	142
Figure 5.14	Desirability Plot for Optimization of VLB loaded spherulites.	144
Figure 6.1	GCH loaded Non-PEGylated Spherulites size analysis.	152
Figure 6.2	GCH loaded PEGylated Spherulites size analysis.	152
Figure 6.3	GCH loaded Non-PEGylated Spherulites zeta potential analysis.	153
Figure 6.4	GCH loaded PEGylated Spherulites zeta potential analysis.	153
Figure 6.5	SEM image of GCH loaded Non-PEGylated Spherulites.	155
Figure 6.6	SEM image of GCH loaded PEGylated Spherulites.	155
Figure 6.7	TEM image of GCH loaded Non-PEGylated Spherulites.	156
Figure 6.8	TEM image of GCH loaded PEGylated Spherulites.	157
Figure 6.9	In Vitro drug release study of Standard GCH solution, GCH loaded Non-PEGylated Spherulites and GCH loaded PEGylated Spherulites. Data represents mean of three experimental values (n=3; mean±SD).	158
Figure 6.10	Graphical representation of Electrolyte-induced flocculation study of GCH loaded Non-PEGylated Spherulites and GCH loaded PEGylated Spherulites. Data represents mean of three experimental values (n=3; mean±SD).	160
Figure 6.11	VLB loaded Non-PEGylated Spherulites size analysis.	161
Figure 6.12	VLB loaded PEGylated Spherulites size analysis.	162
Figure 6.13	VLB loaded Non-PEGylated Spherulites zeta potential analysis.	162

Figure 6.14	VLB loaded PEGylated Spherulites zeta potential analysis.	163
Figure 6.15	SEM image of VLB loaded Non-PEGylated Spherulites.	164
Figure 6.16	SEM image of VLB loaded PEGylated Spherulites.	165
Figure 6.17	TEM image of VLB loaded Non-PEGylated Spherulites.	166
Figure 6.18	TEM image of VLB loaded PEGylated Spherulites.	166
Figure 6.19	In vitro drug release study of Standard VLB solution, VLB loaded Non-PEGylated Spherulites and VLB loaded PEGylated Spherulites. Data represents mean of three experimental values (n=3; mean±SD).	168
Figure 6.20	Graphical representation of Electrolyte-induced flocculation study of VLB loaded Non-PEGylated Spherulites and VLB loaded PEGylated Spherulites. Data represents mean of three experimental values (n=3; mean±SD).	169
Figure 7.1	Schematic representation of steps involved in cell cycle and its analysis by using DNA intercalating fluorescence probe in flow cytometry.	173
Figure 7.2	Results of cell cytotoxicity by MTT assay A) at 8 hours, B) at 16 hours and C) at 24 hours (where, Standard GCH Solution Vs. GCH loaded Spherulites; *p<0.05, *** p<0.001). Experiment was performed in triplicate (n=3). Data represents mean±SD.	176-177
Figure 7.3	IC50 values with respect to time of standard GCH drug and GCH loaded spherulites in A549 cell line.	179
Figure 7.4	Results of cell cytotoxicity by MTT assay A) at 8 hours, B) at 16 hours and C) at 24 hours (where, Standard VLB Solution Vs. VLB loaded Spherulites; **p<0.01, *** p<0.001). Experiment was performed in triplicate (n=3). Data represents mean±SD.	180-181
Figure 7.5	IC50 values with respect to time of standard VLB drug and VLB loaded spherulites in A549 cell line.	183

Figure 7.6	Graphical representation of Cell Cycle Analysis of A549 cell line with respect to time A) After 12 hours and B) After 24 hours by PI staining using FACS technique (where, Standard GCH Solution Vs. GCH loaded Spherulites; * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$) Experiment was performed in triplicate (n=3). Data represents mean \pm SD.	185
Figure 7.7	Cell cycle analysis of A549 cells; where A) Untreated cells, B) Cells treated with Std GCH, C) Cells treated with GCH loaded spherulites.	186
Figure 7.8	Graphical representation of Cell Cycle Analysis of A549 cell line with respect to time A) After 12 hours and B) After 24 hours by PI staining using FACS technique (where, Standard VLB Solution Vs. VLB loaded Spherulites; *** $p < 0.001$). Experiment was performed in triplicate (n=3). Data represents mean \pm SD.	188
Figure 7.9	Cell cycle analysis of A549 cells; where A) Untreated cells, B) Cells treated with Standard VLB, C) Cells treated with VLB loaded spherulites.	189
Figure 8.1	Microscopic images of RBCs for understanding the compatibility with drug solution and formulation; A) Normal saline, B) 1%v/v Triton x-100, C) Standard GCH solution, D) Standard VLB solution, E) GCH loaded Spherulites and F) VLB loaded Spherulites.	201
Figure 8.2	Plasma concentration Vs. Time profile of Standard GCH solution, GCH loaded non-PEGylated Spherulites and GCH loaded PEGylated Spherulites administered intravenously via tail vein in Female SD rats (n=6). (Data represents mean \pm SD).	203
Figure 8.3	Plasma concentration Vs. Time profile of Standard VLB solution, VLB loaded non-PEGylated Spherulites and VLB loaded PEGylated Spherulites administered intravenously via tail vein in Female SD rats (n=6). (Data represents mean \pm SD).	205

Figure 8.4	Biodistribution of Standard GCH solution in rats in highly perfused organs as function of time. Data represents % dose injected of total dose estimated from organs. (n=3; mean±SD).	207
Figure 8.5	Biodistribution of GCH loaded non-PEGylated Spherulites in rats in highly perfused organs as function of time. Data represents % dose injected of total dose estimated from organs. (n=3; mean±SD).	208
Figure 8.6	Biodistribution of GCH loaded PEGylated Spherulites in rats in highly perfused organs as function of time. Data represents % dose injected of total dose estimated from organs. (n=3; mean±SD).	209
Figure 8.7	Comparison of lung targeting potential of Standard GCH Solution, GCH loaded Non PEGylated spherulites and GCH loaded PEGylated spherulites. Data represents % dose injected of total dose estimated from Lungs. (n=3; mean±SD) (Statistically analyzed by Two way ANOVA; p<0.0001).	210
Figure 8.8	Biodistribution of Standard VLB solution in rats in highly perfused organs as function of time. Data represents % dose injected of total dose estimated from organs. (n=3; mean±SD).	212
Figure 8.9	Biodistribution of VLB loaded non-PEGylated Spherulites in rats in highly perfused organs as function of time. Data represents % dose injected of total dose estimated from organs. (n=3; mean±SD).	213
Figure 8.10	Biodistribution of VLB loaded PEGylated Spherulites in rats in highly perfused organs as function of time. Data represents % dose injected of total dose estimated from organs. (n=3; mean±SD).	214
Figure 8.11	Comparison of lung targeting potential of Standard VLB Solution, VLB loaded Non PEGylated spherulites and VLB loaded PEGylated spherulites. Data represents % dose	215

- injected of total dose estimated from Lungs. (n=3; mean±SD) (Statistically analyzed by Two way ANOVA; p<0.0001).
- Figure 8.12 ^{99m}Tc-GCH complex DTPA challenge study (n=3; mean±SD). 217
- Figure 8.13 Gamma scintigraphy images at different time points viz. A1) 0.25 hr, B1) 0.5 hr, C1) 1 hr, D1) 2 hrs, E1) 3 hrs, F1) 4 hrs, of rat administered with ^{99m}Tc labelled GCH plain drug solution showing non specific distribution of plain drug evidenced by scattered gamma quanta and A2) 0.25 hr, B2) 0.5 hr, C2) 1 hr, D2) 2 hrs, E2) 3 hrs, F2) 4 hrs, of rat administered with ^{99m}Tc labelled GCH loaded Non-PEGylated Spherulites depicting distribution of the formulation. 219
- Figure 8.14 Gamma scintigraphy images at different time points viz. A) 0.25 hr, B) 0.5 hr, C) 1 hr, D) 2 hrs, E) 3 hrs, F) 4 hrs, of rat administered with ^{99m}Tc labelled GCH loaded PEGylated Spherulites showing targeting as well as retention of the formulation at the desired site i.e. lungs (region of interest marked by arrows). 220
- Figure 8.15 Biodistribution of (A) ^{99m}Tc labelled GCH plain drug solution (B) ^{99m}Tc labelled GCH loaded Non-PEGylated Spherulites and (C) ^{99m}Tc labelled GCH loaded PEGylated Spherulites in rats in different organs as function of time. 221
- Figure 8.16 Comparison of lung targeting potential of GCH plain drug, GCH loaded Non-PEGylated spherulites and GCH loaded PEGylated spherulites, where: A) Left Lung, B) Right Lung. (Where, For Plain drug Solution Vs. Non-PEGylated Spherulites; *** p<0.0001, ** p<0.01 and * p< 0.05. For Plain drug Solution Vs. PEGylated Spherulites; ### p<0.0001. For Non-PEGylated Spherulites Vs. PEGylated Spherulites; \$\$\$ p<0.0001). 222
- Figure 8.17 ^{99m}Tc-VLB complex DTPA challenge study (n=3; mean±SD). 224

- Figure 8.18 In vivo biodistribution of ^{99m}Tc labelled VLB plain drug solution injected intravenously in Sprague-Dawley rats, visualized by Gamma scintigraphy at different time points viz. A) 0.25 hr, B) 0.5 hr, C) 1 hr, D) 2 hrs, E) 3 hrs, F) 4 hrs. 226
- Figure 8.19 % dose injected intravenously, of ^{99m}Tc labelled VLB plain drug solution in Sprague-Dawley rats (n=3), where data depicts concentration found in highly perfused organs with respect to time. Data represents mean \pm SD. 227
- Figure 8.20 In vivo biodistribution of ^{99m}Tc labelled VLB loaded Non-PEGylated Spherulites injected intravenously in Sprague-Dawley rats, visualized by Gamma scintigraphy at different time points viz. A) 0.25 hr, B) 0.5 hr, C) 1 hr, D) 2 hrs, E) 3 hrs, F) 4 hrs. 227
- Figure 8.21 % dose injected intravenously, of ^{99m}Tc labelled VLB loaded Non-PEGylated Spherulites in Sprague-Dawley rats (n=3), where data depicts concentration found in highly perfused organs with respect to time. Data represents mean \pm SD. 228
- Figure 8.22 In vivo biodistribution of ^{99m}Tc labelled VLB loaded PEGylated Spherulites injected intravenously in Sprague-Dawley rats, visualized by Gamma scintigraphy at different time points viz. A) 0.25 hr, B) 0.5 hr, C) 1 hr, D) 2 hrs, E) 3 hrs, F) 4 hrs. 228
- Figure 8.23 % dose injected intravenously, of ^{99m}Tc labelled VLB loaded PEGylated Spherulites in Sprague-Dawley rats (n=3), where data depicts concentration found in highly perfused organs with respect to time. Data represents mean \pm SD. 229
- Figure 8.24 Lung targeting potential of VLB plain drug, VLB loaded Non-PEGylated spherulites and VLB loaded PEGylated spherulites at different time points, where: A) Left Lung, B) Right Lung. (Where, For Plain drug Solution Vs. Non-PEGylated Spherulites; *** p<0.001. For Plain drug Solution Vs. PEGylated Spherulites; ### p<0.001. For Non-PEGylated Spherulites Vs. PEGylated Spherulites; \$\$\$ p<0.001) 229

statisically analyzed by repeated measure of Two way ANOVA followed by Bonferroni Post-tests.