

Table index

Table 1. 1 Genes involved in depression	9
Table 3. 1 Purity and concentration of p11 cDNA using NanoDrop instrument.....	52
Table 3. 2 Standard deviation between obtained and actual concentrations of cDNA solutions.....	53
Table 3. 3 Accuracy measurements of cDNA concentrations	53
Table 3. 4 Inter-day and intraday precision of the method.....	54
Table 3. 5 Peak area and % RSD for actual and observed concentrations of cDNA	56
Table 3. 6 Absorbance of BSA in linearity range from 1 to 150 $\mu\text{g}/\mu\text{L}$	58
Table 4(B). 1 Nomenclature of synthesised lipids for further reference	86
Table 4(B). 2 Conjugation efficiency of protected amino acids to DOPE	94
Table 4(B). 3 Concentration of DCM present in samples	98
Table 5. 1 Effects of various process parameters on lipids film and subsequently on liposome formation.....	105
Table 5. 2 Variables and response parameters for design optimization of liposome formulation.....	107
Table 5. 3 Actual and coded values for optimization parameters.....	107
Table 5. 4 Constrains applied to design for optimization	107
Table 5. 5 Primer characteristics for mice p11 cDNA and GAPDH gene.....	123
Table 5. 6 Components for RT-PCR reaction.....	124
Table 5. 7 RT-PCR temperature cycle.....	125
Table 5. 8 Liposome optimization with the help of design matrix	127
Table 5. 9 ANOVA results summary for particle size.....	128
Table 5. 10 ANOVA results for special cubic model	129
Table 5. 11 Summary of ANOVA result for special cubic model.....	129
Table 5. 12 ANOVA results summary for PDI	140
Table 5. 13 ANOVA results for quadratic model.....	141
Table 5. 14 Summary of ANOVA result for special cubic model.....	141
Table 5. 15 Constraints applied for selection of optimised batch.....	149
Table 5. 16 Optimized batch parameters based on desirability criteria.....	151
Table 5. 17 Size and PDI of predicted batches	152
Table 5. 18 Complexation efficiencies of liposomes at optimised L/P ratio and zeta potential of formed lipoplex	155
Table 5. 19 Effects of sodium sulphate concentration on modified liposomes	160

Table 5. 20 effect of heparin concentration on complexation efficiency	161
Table 5. 21 cDNA retention efficiency of lipoplex in presence of serum	162
Table 5. 22 Haemolytic potential of modified lipoplex.....	164
Table 5. 23 % Gene expression in SHSY5Y cells after treatment with interferon, naked DNA and lipoplex formulations	171
Table 6. 1 Composition of nasal spray formulation.....	178
Table 6. 2 Test tube contents for quantification of sulfhydryl group	187
Table 6. 3 Acrylamide percentage required for the separation of proteins.....	189
Table 6. 4 Resolving gel and stacking gel compositions.....	190
Table 6. 5 Compositions and conditions for staining and destaining solutions.....	191
Table 6. 6 Observation of sign for toxicity after intravenous injection.....	195
Table 6. 7 Results of nasal spray characterization.....	207
Table 6. 8 Ex-vivo nasal permeation study sheep mucosa	208
Table 6. 9 Absorbance and concentration of sulfhydryl group present in antibody containing samples.....	211
Table 6. 10 Sighting study for dose selection in swiss albino mice	218
Table 6. 11 Mortality study in Swiss albino mice at varied doses.....	219
Table 6. 12 Average change in weight prior and during study (gram).....	219
Table 6. 13 Feed consumption by group per cage (gram)	220
Table 6. 14 Results of CBC study after intravenous injection of formulation	220