

BIBLIOGRAPHY


BIBLIOGRAPHY

- Abdel-Aleem, H., Ahmed, A., Sabra, A.M., Zakari, A.M., Soliman, M. and Hamed, H. (1996): Serum alpha-L fucosidase enzyme activity in ovarian and other female genital tract tumours. *Int Jr. Gynecol. Obstet.* **55**: 273-279.
- Adan, A. and Farber, E. R. (1982): Immunochemical and electrophoretic study of glycoproteins in cancer patients. *Clin. Biochem.* **15**: 22-23.
- Akamatsu, S., Yazawa, S., Tachikawa, T., Furuta, T., Okaichi, Y., Nakamura, T., Asao, T. and Nagamachi, Y. (1996): Alpha 2-3 sialyltransferase associated with the synthesis of CA19-9 in colorectal tumours. *Cancer* **77**: 1694-1700.
- Alhadeff, J.A. (1989): Malignant cell glycoproteins and glycolipids. *CRC Crit. Rev. Oncol Hematol.* **9**: 37-107.
- Alhadeff, J.A. and Holzinger, R.T. (1982): Sialyl transferase, sialic acid and sialoglycoconjugates in metastatic tumor and human liver tissue. *Int. Jr. Biochem.* **14**: 119-126.
- Altara, S. (1993): Cancer associated serum globulin determination in patients with cervical, endometrial and ovarian cancer. *Acta. Med. Crotica.* **47**: 19-22.
- Amano, N., Moriyama, M., Hibi, N., Tasukada, Y. (1992): Elevation of serum fucosyl transferase activities in malignant disease -A sensitive tumour marker. *Rinso. Byari.* **40**: 182-188.
- Aoyagi, Y., Suzuki, Y., Isemura, M., et al. (1988): The fucosylation index of Alpha fetoprotein and its usefulness in early diagnosis of hepatocellular carcinoma. *Cancer* **61**: 769-774.

- Aoyagi, Y., Sazuki, Y., Igarashi, K., Yokoto, T., Mori, S., Nomoto, M., Isemura, M. and Asakura, H. (1991): The usefulness of simultaneous determination of glycosaminylation and fucosylation indices of alpha fetoprotein in the differential diagnosis of neoplastic disease of the liver. *Cancer* **67**: 2390-2394.
- Aoyagi, Y., Sazuki, Y., Igarashi, K., Yokoto, T., Mori, S., Suda, T., Naitoh, A., Isemura, M. and Asakura, H. (1993): Highly enhanced fucosylation of alpha feto protein in patients with germ cell tumour. *Cancer* **72**: 615-618
- Arivazhagan, S., Kavitha, K. and Nagini, S. (1998): Glycoconjugate profile in plasma and erythrocytes of gastric cancer patients. *Indian Jr. Physiol. Pharmacol.* **42**: 123-126.
- Asano, K., Sakamoto, H., Sasaki, H., Ochiya, T., Yoshida, T., Ohishi, Y and Machida T.(1995): Tumorigenicity and gene amplification potential of cyclin-D Over expressing NIH3T3 cells. *Biochem.Biophys.Res. Commun.* **217**: 1169-1176.
- Asao, T., Yazawa, S., Nagamachi, Y., Abbas, S.A. and Matta, K.L. (1989): Serum α -(1-3)-L-fucosyltransferase carcinoembryonic antigen and sialyl Lewis X- antigen levels in lung cancer. *Cancer* **64**: 2541-2545.
- Atkinson, M.A. and Bramwell, M.E. (1981): Studies on the surface properties of hybrid cells III A membrane glycoprotein found on the surface of a wide range of malignant cells. *J. Cell Sci.* **48**: 147-170.
- Aziz, S., Lodi, T.Z. and Hassan, T. (1988): Serum protein electrophoresis in healthy subjects. *Jama. J. pak. Assoc.* **38**: 18-20.
- Baker, M.A., Kanani, A., Brockhausen, I., Schachter, H., Hindenburg, A. and Raub, R.N. (1987): Presence of cytidine 5'-monophospho-N-acetyl neuraminic acid: Gal β -1-3 Gal NAc-R α (2-3)-sialyltransferase in normal human leukocytes and increased activity of the enzyme in granulocytes from chronic myelogenous leukemia patients. *Cancer Res.* **47**: 2763 - 2766.

- Balaram, P., Pillai, M.R. and Abraham, T.C. (1987): Immunology of pre-malignant and malignant conditions of oral cavity – II Circulating immune complexes. *J. Oral Pathol.* **16**: 389-391.
- Barlow, J.J., DiCioccio, R.A., Dillard, P.H., Blumenson, L.E. and Matta, K.L. (1981): Frequency of an allele for low activity of α -L-fucosidase in sera. Possible increase in epithelial ovarian cancer patients. *J. Natl. Cancer Inst.* **87**: 1005-1009.
- Baselar, M.W., Maxim, P.E. and Veltri, R.W. (1987): Circulating IgA immune complexes in head and neck cancer, nasopharyngeal carcinoma, lung cancer and colon cancer. *Cancer* **59**: 1727-1731.
- Basu, S.S., Basu, M., Li, Z., Basu, S.S. (1996): Characterization of two glycolipid: Alpha 2-3 sialyl transferase, SAT-3 (CMP-NeuAc: nLcose4Cer Alpha 2-3 sialyltransferase) and SAT-4 (CMP-NeuAc : GgOse4Cer alpha 2-3 sialyltransferase) from human colon carcinoma (colo 205) cell lines. *Biochemistry* **35**: 5166-5174.
- Bates, S.E. (1991): Clinical application of serum tumour markers. *Ann.Int. Med.* **115**: 623-638.
- Bauer, C.H., Reutter, W.G., Erhart, K.K., Kottgen, E. and Gerok, W. (1978): Decrease of human serum fucosyltransferase as an indicator of successful tumour therapy. *Science* **201**: 1232-1233.
- Baxi, B.R., Patel, P.S., Adhvaryu, S.G. and Dayal, P.K. (1991): Usefulness of serum glycoconjugates in precancerous and cancerous disease of the oral cavity. *Cancer* **67**: 135-140.
- Bellahcene, A., Menard, S., Bufalino, R., Moreau, L., Casronovo, V. (1996): Expression of bone sialoprotein in primary human breast cancer is associated with poor survival. *Int. J. Cancer.* **69**: 350-353.
- Bernacki R J. Niedbala M J.Karytny K W.(1985) Glycosidase in cancer and invasion. *Cancer Metast. Rev.* **4**: 81-101.

- Bhavanandan, V.P. and Davidson, E.A. (1982): Cell surface glycoprotein markers for neoplasia. *Methods. Cancer Res.* **19**: 53-58
- Bhuvaramurthy, V., Balasubramanian, N., Vijaykumar, S. and Govindasamy, S. (1995): Effect of radiation on serum glycoprotein and glycosidase in patients with cervical carcinoma. *Int.J.Gynecol.Obstet.* **48**: 49-54.
- Black, P.,H. (1980): Shedding from the cell surface of normal and cancer cells. *Adv. Cancer. Res:* **32**: 75-119.
- Bloscher, J.G., Bruyneel, E.A., Van Rooy, H., Schallier, D.C., Mareel, N.M. and Smets, L.A. (1989): Decreased fucose incorporation in cell-surface carbohydrates is associated with inhibition of invasion. *Clin. Exp. Metast.* **7**: 557-569,
- Bolmer, S.D. and Davidson, E.A (1981): Preparation and properties of a glycoprotein associated with malignancy. *Biochemistry* **20**: 1047-1054.
- Bradley, W.P., Blasco, A.P., Weiss, J.P., Alexander, Jr.J.C., Silverman, N.A. and Chretien, P.B. (1977): Correlations among serum protein bound carbohydrates serum glycoproteins lymphocyte reactivity and tumour burden in cancer patients.*Cancer* **40**: 2264-2272.
- Bresalier, R.S., Rockwell, R.W., Dahiya, R., Duh, Q.Y. and Kim, Y.S. (1990): Cell-surface sialoprotein alterations in metastatic murine colon cancer cell lines selected in an animal model for colon-cancer metastasis. *Cancer Res.* **50**: 1299-1307.
- Bruyneel, E.A., Debray, H., De Mets, M., Marcel, M.M. and Montrcuil, J. (1990): Altered glycosylation of madin-Dabry canine kidney (MDCK) cells after transformation by murine sarcoma virus. *Clin. Exp. Metasts.* **8**: 241-253.
- Chandrasekaran, E.V. and Davidson, E.A. (1979): Sialoglycoproteins of human mammary cells: Partial characterization of sialo glycopeptides. *Biochemistry* **18**: 5615-5620.



Chandrasekaran, E.V., Jain, R.K. and Matta, K.L. (1992): Ovarian Cancer α -1-3-L-fucosyl transferase. Differentiation of distinct catalytic species with unique substrate, 3'sulfo N-acetyl lactosamine in conjunction with other synthetic acceptors. *J. Biol. Chem.* **267**: 23806-23814.

Chatterjee, S.K., Kim, U. and Bielat, K. (1976): Plasma membrane bound enzyme of mammary tumours as the biochemical indication of the metastatic capacity. Analysis of enriched plasma membrane preparation. *Br. J. Cancer* **33**: 15-29.

Chatterjee, S.K., Bhattacharya, M., Barlow, J.J. (1979): Glycosyltransferase and glycosidase activities in ovarian cancer patients. *Cancer Res.* **39**: 1943-1951.

Chen, Y.H. and Magalhaes, M.C. (1990): Hypoalbuminemia in patients with multiple myeloma. *Arch. Int. Med.* **150**: 605-610.

Chester, S.J., Maimonis, P., Meitner, P. A. and Vezeridis, M.P. (1990): An analysis of immunocomplexes for the detection of early stages of colon cancer. *Cancer* **65**: 1338-1344.

Chorvath, B. Dara, J., Babusikova, O., Gyarfás, J. (1983): Cell surface proteins and glycoprotein electrophoretic pattern in some human hematopoietic malignancies. *Neoplasma* **30**: 715-724.

Chuman, Y., Sumizawa, T., Takebayashi, Y., Niwa, K., Yamada, K., Haraguchi, M., Furukawa, T., Akiyama, S., Aikou, T. (1996): Expression of multidrug resistance associated protein (MRP) gene in human colorectal, gastric associated and non small cell lung carcinomas. *Int. J. Cancer* **66**: 274-279.

Collard, J.G., Schijven, J.F., Bikker, A., LaRiviere, G., Bolscher, J.G. and Roose, E. (1986): Cell surface sialic acid and the invasive and metastatic potential of T-cell hybridomas. *Cancer Res.* **46**: 3521-3527.

- Consenses Report. (1987): Proceeding of a workshop on the biochemical and immunological diagnosis of cancer. *Tumour Biol.* **8**: 65-84.
- Cox, A.R. (1972): Regression analysis model and life table. *Jr. Stat. Soc. Br.* **34**: 187- 198.
- Dall'olio, F. and Trere, D. (1993): Expression of alpha 2,6 sialylated sugar chain in normal and neoplastic colon tissues. Detection by digoxigenin conjugated sambucus nigra agglutinin. *Eur. Jr. Histochem.* **37** : 257-265.
- Dao, T.L., Ip, C. and Patel, J. (1980): Serum sialyltransferase and 5'nucleotidase as reliable biomarkers in women with Breast cancer. *Jr.Natl .Canecr Inst.* **65**: 529-534.
- De, A.K. and Hardy, R.E. (1990): Elucidation of sialyltransferase as a tumour marker. *Ind. Jr. Biochem. & Biophys.* **27**: 452-455.
- Dennis, J. and Kerbal, R. (1981): Charecterization of a deficiency in fucose metabolism in lectin resistant varient of amurine tumour showing altred tumorigenic and metastatic capacities in vivo. *Cancer Research* **41**: 98-104.
- Dennis, J.W. (1992): Changes in glycosylation assosiated with malignant transformation and tumour progression. In: Fukuda, M. (Ed.): Cell surface carbohydrate and cell development. Boca Raton F L, CRC Press. pp 161-194.
- Dermer, G.B., Silverman, L.M., Gendler, S.J., and Tokes, Z.A. (1980): Incidence of asplit alpha 2-glycoprotein band in electrophoretis pattern from serum of adeno carcinoma patientsb. *Clin.Chem* **26**: 392-395.
- Deugnier, Y., David, V., Brissot, p., Mabou, P., Delamair, d., Massner, M., Bourel, M., Legall, J.Y. (1984): Serum alpha-L-fucosidase: A new marker for diagnosis of primary hepatic carcinoma? *Hepatology* **5**: 889-892.

- Dische, Z. and Shettles, L.B. (1948): A specific colour reaction of methyl pentoses and a spectrophotometric micromethod for their determination. *Jr. Biol. Chem.* **175**: 595-603.
- Dische, Z. and Shettles, L.B. (1963): Reciprocal reaction between fucose and sialic acid and mammalian glycoprotein. *Ann. N Y Acad. Sci.* **106**: 259-262.
- Domino, S.E., Hiraiwa, N. and Lowe, J.B. (1997): Molecular cloning chromosome assignment and tissue specific expression of a murine alpha (1-2) fucosyltransferase expressed in thymic and epidymal epithelial cells. *Biochem. J.* **327**: 105-115.
- Doyle, L.A., Ross, D.D., Sridhara, R., Fojo, A.T., Kaufmann, S.H., Lee, E.J. and Schiffer C.A. (1995): Expression of a 95 Kda Membrane protein is associated with low Daunorubicin accumulation in leukemic blast cells. *Br. J. Cancer* **71**: 52-58.
- Durnald, P., Gatti, R. and Borrone, C. (1982): Fucosidosis. In: Durnand, P., and O'Brien, J.S. (Eds), Genetic errors of glycoprotein metabolism, Ermes, Milano, Italy, pp 49-87.
- Evans, A.S., Dolan, M.F., Sobocinski, P.Z. and Quinn, F.A. (1974): Utility of serum protein bound neutral hexoses and l-fucose for estimations of malignant tumour extent and evaluation of efficacy of therapy. *Cancer Res.* **34**:538-542.
- Fairbanks, G., Steck, T.K. and Wallach D.F. (1971): Electrophoretic analysis of the major polypeptides of the human erythrocyte membrane. *Biochemistry* **10**: 2606-2617.
- Feijoo, C., Paez-de-la, Cadena, M. and Rodriguezberrocat, F.J. (1997): Sialic acid level in serum and tissue from colorectal cancer patients. *Cancer Lett.* **112**: 155-160.

- Feinstein, A.R. (1985): Clinical epidemiology. In: The architecture of clinical research. W.B. Saunders, Philadelphia. pp 601-609.
- Feizi, T. (1981): Carbohydrate differentiation. *TIBS* **6**: 333-335.
- Feizi, T. and Childs, R.A. (1987): Carbohydrates as antigenic determinants of glycoproteins, *Biochem. J.* **245**: 1-11.
- Fernandez-Rodriguez, J., Paez-de-la-cadena, M., Martinez-zorzano, V.S. (1997): Fucose level in serum and in tumor of colorectal adenocarcinoma patients. *Cancer Lett.* **121**: 147-153.
- Filipitis, M., Suchomel, A., Lecher, K. and Priker, R. (1997): Immunocytochemical detection of the multidrug resistance associated protein and P-glycoprotein in acute myeloid leukemia: Impact of antibodies sample source and disease status. *Leukemia* **11**: 1073-1077.
- Finne .J., Tao, T.W. and Burger, M.M. (1980): Carbohydrate changes in glycoproteins of a poorly metastasizing wheat germ agglutinin resistant melanoma clone. *Cancer research* **40**: 2580-2587
- Flowers, H.M. (1981): Chemistry and biochemistry of D- and L-fucose. *Adv. Carbohydr. Chem. Biochem* **39**: 279-345.
- Fukuda, M. (1994): In Molecular Glycobiology: (M. Fukuda and O. Hindsgual, eds.) Frontiers in molecular biology series Oxford Univ. Press, Oxford, U.K. pp. 1-52.
- Fukushima, K. Hirota, M., Terasaki, P.I., Wakisaka, A., Togashi, H., Chia, D. Suyama, N., Fukushi, Y., Nudelman, E. and Wakomari, S. (1984): Characterization of sialosylated Lewix X as a new tumour associated antigen. *Cancer Res.* **44**: 5279-5285.
- Ganzinger, U. and Deutsch, E. (1980): Serum sialyltransferase level as a parameter in the diagnosis and follow-up of gastrointestinal tumors. *Cancer Res.* **40**: 1300-1304.

- Geolz, S.E., Hession, C., Goff, D., Griffiths, B., Tizard, R., Newman, B., Chi-Rosso, G. and Lobb, R. (1990): ELFT: a gene that directs the expression of an ELAM-1 ligand. *Cell* **63**: 1349-1356.
- Gessner, P., Riedl, S., Quentmaier, A. and Kemmno, W. (1993): Enhanced activity of CMP-NeuAc:Gal beta 1-4GlcNAc:alpha 2-6Sialyl transferase in metastasizing tumour tissue and serum of tumour patients. *Cancer Lett.***75**:143-149.
- Gil-Martin, E., Gil-seijo. S., Nieto-Novoa, C. and Fernandez-Briera, A. (1996): Elevation of acid glycosidase activities in thyroid and gastric tumours. *Int J. Biol. Chem. Cell Biol.* **28**:651-657.
- Glick, M.C.(1978): Fucosylation A role in cell function. In : Walbo E.F. (Ed.) *Glycoproteins and glycolipids in disease processes* Washington D.C. American Chemical Society. Pp 405-411.
- Gonzalez, A., Vizoso, F., Allen-De, M.T. Sanchez, M.T., Balibrea, T.L. (1996): pre operative CEA and TAG-72 serum levels as pronostic indicators in resectable gastric carcinoma. *Int. J. Biol. Mark.* **11**: 165-171.
- Gosh, M. and Nayak, B.R. (1991): Serum sialic acid, fucose, sialic acid/fucose ratio as tumour marker in oral cancer. *Annals of Density* **50**: 33-35.
- Grant, C.W. and Peters, M.W. (1984): Lectin membrane interactions. Information from model systems. *Biochim. Biophys. Acta* **779**: 403-422.
- Griffiths, J. and Reynold, S. (1982): Plasma sialyltransferase total and isoenzyme activity in the diagnosis of cancer of colon. *Clin. Biolchem.***15**: 46-48.
- Gupta, S.C., Singh P.A., Sukla, H.S., Mehrotra T.N. (1980): Serum immuoglobulin profile in patients with cancer cervix. *Ind. Jr. Med. Res.***71**: 893-896.

- Hada, T. (1997): Increased plasma α -1-3 fucosyltransferase activities in patient with hepatocellular carcinoma. *Glycoconj. J.* **12**: 627-631.
- Hakomari, S. (1981): Glycosphingolipids in cellular interactions differentiation and oncogenesis. *Annu. Rev. Biochem.* **50**: 733-764.
- Hakomori S.I. and Mirakami W.T. (1968): Glycolipid of hamster fibroblast and derived malignant transformed cell lines. *Proc.Natl.Acad.Sci.* **59**: 254-261.
- Hakomori, S. (1989): Aberrant glycosylation in tumour and tumour-associated carbohydrate antigens. *Adv.Cancer Res.* **52**: 257-331.
- Harmenberg, U., Frodin, J.E., Ljungdahl-Stahl, E., Mellstedt, H., Wahren, B. and Stigbrand, T. (1989): Significance of alkaline phosphatase isoenzymes in the monitoring of patients with colorectal carcinoma. *Tumor Biol.* **10**: 225-231.
- Harvey, B.E., Toth, C.A., Wagner, H.E., Steele, J.G.D. and Thomas, P. (1992): Sialyltransferase activity and hepatic tumour growth in nude mouse model of colorectal cancer metastases. *Cancer Res.* **52**: 1775-1779.
- Hasilik, A., Klein, U., Waheed, A., Strecker, G. and Von-figure, K. (1980): Phosphorylated oligosaccharides in lysosomal enzymes. Identification of alpha-N-acetyl glucosamine (1) phospho (6) mannose diester groups. *Proc .Natl. Acad.Sci.* **77**: 7074-7078.
- Holmes, E., Osterander, G. and Hakomori, S. (1986): Biosynthesis of the sialyl -LeX determinant carried out by type-2 chain glycosphingolipids (IV 3 NeuAcIII3 FucnLc4, VI3NeuAcV3FucnLc6 and VI3 NeuAcIII3 V3 Fuc2 nLc6)in human lung carcinoma cell lines. *J. Biol. Chem.* **261**: 3737-3743.
- Huglund, C., Ylatupa, S., Mertaniemi, P., Partanen, P. (1997): Cellular fibronectin concentration in the plasma of patients with malignant and benign diseases: a comparison with CA19-9 and CEA. *Br. J. Cancer* **76**: 777-783.

- Hutchinson, W.L., Johnson, P.J., Du, M.Q. and Williams, R. (1991): Serum and tissue α -L-fucosidase activity in the pre-clinical and clinical stage of hepatocellular carcinoma. *Clin. Sci.* **81**: 117-182.
- Inoue, M., Nakayama, M. and Tanizawa, O. (1990): Altered expression of Lewis blood group and related antigens in fetal, normal adult and malignant tissue of the uterine endometrium. *Virchows. Arch. A. Pathol. Anat.* **416**: 221-228.
- Jaernfelt, J., Finne, J., Krusius, T. and Rauvala, H. (1978): Protein bound oligosaccharides of cell membranes. *TIBS* **3**: 110-114.
- Kaplan, F. and Moskowitz, Y. (1963): Enzymes and isoenzymes of carbohydrate metabolism. *Clin. Chim. Acta* **48**: 259-267.
- Kaplan, E.L. and Meier, P. (1958): Nonparametric estimation from incomplete observations. *J. Am. Stat. Assoc.* **53**: 457-481.
- Kelm, S. and Schauer, R. (1997): Sialic acids in molecular and cellular interaction. *Int. Rev. Cytol.* **175**: 137-240.
- Kessel, D. and Allen, J. (1975): Elevated plasma sialyltransferase in the cancer patients. *Cancer Res.* **35**: 670-672.
- Kim, Y.S., Kim, Y.W., Siddiqui, B. and Tsao, D. (1982): Membrane associated fucose containing glycoproteins and glycolipids of cultured epithelial cells from human colonic adenocarcinoma and fetal intestine. *Eur. J. Cancer Clin. Oncol.* **18**: 1329-1336.
- King, I.A., Tabiwo, A. and Williams, R.H. (1980): Incorporation of 1-(3H) fucose and d-(3H) glucosamine into cell surface associated glycoconjugates in epidermis of cultured pig skin slices. *Biochem. J.* **190**: 65-77.
- Kobata, A. and Takasaki, S. (1993): Structural characterization of oligosaccharides from glycoproteins. In: M. Fukuda and A. Kobata (eds) *Glycobiology: A practical approach*. Oxford University Press, New York. pp 165-185.

- Koj, A. (1974): Acute-phase reactants: Their synthesis turnover and biological significance. In structure and functions of plasma proteins. Allison, A.C. (ed.) Vol. 1, Plenum Press, London, pp. 73-125
- Kondo, A., Hosokawa, Y., Kiso, M., Hasigawa, A. and kato I. (1994): Analysis of oligosaccharides of human IgG from serum of leukemia patients. *Biochem. Mol. Biol. Int.* **32**: 897-902.
- Kornfeld, R. and Kornfeld, S. (1980): Structure of glycoproteins and their oligosaccharide units. In: W.J. Lennarz (ed.) The biochemistry of glycoprotein and proteoglycans. Plenum Publishing Corp., New York, pp 1-34.
- Laemmli, U.K.(1970): Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* **227**: 680-685.
- Lagana, A., Pardo Martinez, B., Marino, A., Fago, G. and Bizzarri, M. (1998): Determination of serum total, lipid and free N-acetyl neuraminic acid in genitourinary malignancies by fluorimetric high performance liquid chromatography. Relevance of free N-acetyl neuraminic acid as tumor marker. *Clin. Chim. Acta.* **243**: 165-179.
- Lileng, R., Tved, K., Nesland J M., Reed, W., Erikstein, B .K., Funderud, S. (1993): CDW75 antigen expression in breast cancer. *Pathol Res Pract* **189**: 394-398.
- Lis, H. and Sharon, N. (1986): Lectins as molecules and as tools. *Annu. Rev. Biochem.* **55**: 35-67.
- Listinsky, J.J., Siegal, G.P. and Listinsky, C.M. (1998): Alpha-L-fucose: A potentially critical molecule in pathologic processes including neoplasia. *Am. J. Clin. Pathol.* **110**: 425-440.

- Lowe, J.B., Kudowska-Latallo, J.F., Nair, R.P., Larsen, R.D., Markes, R.M., Macher, B.A., Rely, R.J. and Ernst, L.K. (1991): Molecular cloning of a human fucosyltransferase gene that determines expression of the Lewis X and VIM-2 epitope but not ELAM-1 dependent cell adhesion. *J. Biol. Chem.* **266**: 17467-17477.
- Lowry, O.H., Rosenbrough, N.J., Farr, A.L., Randell, R.J. (1951): Protein measurement with folin phenol reagent. *J. Biol. Chem.* **193**: 265-275.
- Macbeth, R.A.L. and Bekesi, J.G. (1962): Plasma glycoproteins in various disease states including carcinoma. *Cancer Res.* **22**: 1170-1176.
- Macher, B.A., Holmes, E.H., Swiedler, S.J., Stults, C.L.M. and Srnka, C.A. (1991): Human alpha 1-3 fucosyltransferase. *Glycobiology.* **1**: 577-584.
- Maisin, J., Couvreur, P., Ochrymowicz, I.P., Van-Duyse, E. (1972): Electrophoretic pattern of sera of normal and cancer patients. *Eur. J. Cancer* **8**: 217-219.
- Majuri, M.L., Pinola, M., Niemela, R., Tiisala, S., Nutunen, J., Renkonen, O. and Renkonen, R. (1994): α -1,3 fucosyltransferase-dependent synthesis of sialyl Lewis X an essential digosaccharide present on L-selectin counter receptor in culture endothelial cells. *Eur. J. Immunol.* **24**: 3205-3210.
- Marth E. , Flaschka G. , Stiegler S. and Mose J. (1988): Sialic acid as a marker for differentiation between benign and malignant intracranial tumours. *Clinica. Chimica. Acta.* **176**: 251-258.
- Martin, S., Jansen, F., Bokelmann, J., Kolb, H. (1997): Soluble CD44 splice variants in metastasizing human breast cancer. *Int. J. Cancer* **74**: 443-445.

- Mas, E., Pas, E., Caillol, N., El-Battari, A., Crotte, C., Lombardo, D., Sacloulet, M.O. (1998): Fucosyl transferase activities in human pancreatic tissue. Comparative study between cancer tissues and established tumoral cell lines. *Glycobiology* **6**: 605-615.
- Matei, L.C. (1997): Plasma proteins glycosylation and its alteration in disease. *Rom. J. Int. Med.* **35**: 3-11.
- Miliotes, G., Lyman, G.H., Cruse, C.W., Puleo, C., Albertini, P.A., Rapaport, D., Glass, F., Fenske, N., Soriano, T., Cuny, C., Van-Voorhis, N. and Reintgen, D. (1996): Evaluation of new putative tumor markers for melanoma. *Annal. Surg. Oncol.* **3**: 558-563.
- Miller, A.B., Hogstaten, B., Staquet, M. and Winkler, A.(1981): Reporting results of cancer treatment. *Cancer* **47**:207-214.
- Miyagi (1994)Metastatic potential of transformed rat 3Y1 cell lines is inversely correlated with lysosomal type sialidase activity. *FEBS Lett.* **349**: 255-259.
- Muller J. (1838) In : *CANCER, The biology of malignant disease*. Graham Currie and Angela Currie eds. Edward arnold Publishers. pp 79-96.
- Muryama,T., Zuber, C., Seelentag W.K., Li,W.P., Kemmner, W., Heiz P.U. and Roth, J. (1997): Colon carcinoma glycoprotein carrying alpha2,6-linked sialic acid reactive with sambucus nigra agglutinin are not constitutively expressed in normal human colon mucosa and are distinct from sialyl Tn antigen. *Int. J. Cancer* **70**: 575-581.
- Mustafapour, M.K. and GoldsteinI, J. (1993): Cultured Eherlich ascitis tumour cell show increased N-linked alpha 2,6 sialyltransferase activity. *Arch. Bio. Chem .Biophys.* **303**: 255-259.
- Nakata, B., Chung, K.H., Maguruma, K., Yamashita, Y., Inoue, T., Matsuoka, T., Onoda, N., Kato, Y., Sakurai, M. and Sowa, M.(1998): Changes in tumor marker levels as a predictor of chemo-therapeutic effect in patients with gastric carcinoma. *Cancer* **83**: 19-24.

- Neuberger, A., Marshel, R..D. and Gottschalk, A. (1966): Some aspects of chemistry of the component sugars of glycoproteins, In: Gottschalk, A. (Ed) Glycoproteins their composition structure and function. American Elsevier Publishing Co. Inc. New York. pp 158
- Ng, I.O., Lam, K.Y., Ng, M., Kwong, D.L., Sham, J.S. (1998): Expression of P-glycoprotein a multidrug resistance gene product, It induced by radiotherapy in patients with oral squamous cell carcinoma. *Cancer* **83**: 851-857.
- Nicolson, G.L, (1976): Trans membrane control of receptors on normal and tumor cells. II Surface changes associated with transformation and malignancy. *Biochim. Biophys. Acta* **458**: 1-72.
- Nigam, B., Joshi, S.P., Nagar, S., Chavan, R. and Rendharker, D. (1996): Some biochemical markers in various neoplasms: A multivariate analysis. *Ind. Jr. Clin.Biochem.* **11**: 124-128.
- Nigam, V.N. and Cantero, A. (1973): Polysaccharides in cancer. Glycoproteins and glycolipids. *Adv. Cancer. Res.* **17**: 1-80.
- Noursis, M.J.(1996): SPSS for windows .SPSS Advanced stastic release 7.5 SPSS Inc. Chicago.
- Nuddleman, E., Fukushi, Y., Leverly, S.B., Higuchi, T. and Mori, S. (1986): Novel fucolipids of human adenocarcinoma: Disialosyl LeA antigen (III4FucIII6NeuAcIV3NeuAcLc4) of haman colonic adenocarcinoma and the monoclonal antibody (FH7) defining this structure. *J. Biol. Chem.* **261**: 5487-5495.
- O'Brien, M.I.R., Souberbielle, B.E., Cowan, M.E., Allen, C.A., Luesley, D.M., Mould, J.J., Blackledge, G.R.R., and Skinner, G.R.B. (1991): Glycoprotein pattern in normal and malignant cervical tissue. *Can. Lett.* **58**: 247-254.

- Ogoshi, K., Miyaji, M., Nakamura, K., Kondoh, Y., Makuuchi, H. and Tajima, T. (1997): Immunotherapy and combined assay of serum level of CEA and acute phase reactants. *Cancer Immunol. Immunotherapy* **46**: 14-20.
- Olden, K., Parent, J.B. and White, S.L. (1982): Carbohydrate moieties of glycoproteins. A re-evaluation of their function. *Biochim. Biophys. Acta.* **650**: 209-232.
- Painbeni, T., Gamelin, E., Cailleux, A., Le-Bouil, A., Boisdron-Celle, M., Daver, A., Larra, F., Allain, P. (1997): Plasma sialic acid as a marker of the effect of the treatment on metastatic colorectal cancer. *Eur. J. Cancer* **33**: 2216-2220.
- Passaniti, A. and Hart, G. W. (1988): Cell surface sialylation and tumour cell metastatic potential of B16 melanoma variants correlates with their relative number of specific penultimate with oligosaccharide structure. *J. Biol. Chem.* **263**: 7591-7603.
- Paszkowaska, A., Berbec, H., Semczuk, A. and Cybulski, M. (1998): Sialic acid concentration in serum and tissue of endometrial cancer patients. *Eur. J. Obstet. Gynecol. Reprod. Biol.* **76**: 211-215.
- Patel, P.S., Adhvaryu, S.G. and Balar, D.B. (1994): Serum lactate dehydrogenase and its isoenzymes in leukemia patients. Possible role in diagnosis and treatment monitoring. *Neoplasma* **41**: 55-59.
- Patel, P.S., Raval, G.N., Patel, M.M., Balar, D.B. and Patel, D.D. (1995): Electrophoretic pattern of serum glycoproteins on polyacrylamide disc gel in patients with breast cancer. *Anticancer Res.* **16**: 2089-2094.
- Patel, P.S., Rawal, R.M., Raval, G.N., Patel, Y.V., Shah, S.G., Balar, D.B. and Patel, D.D. (1997): Glycoprotein electrophoretic pattern in sera of patients with oral precancerous conditions and upper aerodigestive tract cancer. *J. Exp. Clin. Cancer Res.* **16**: 65-70.

- Paulson, J.C. (1989): Glycoproteins: what are the sugar chains for? *TIBS*. **14**: 272-276.
- Paulson, J.C. and Colley, K.I. (1989): Glycosyl transferase structure, localization and control of cell-type specific glycosylation. *J. Biol. Chem.* **264**: 17615-17618.
- Periera, M.E. and Kabat, E.A. (1974): Blood group specificity of the lectin from lotus tetragonolobus. *Ann. N. Y. Acad. Sci.* **234**: 301-305.
- Petryniak, J. and Goldstein, I.J. (1986): Immunochemical studies on the interaction between synthetic glycoconjugates and α -L-fucosyl binding lectins. *Biochemistry* **25**: 2829-2838.
- Pettijon, D.E., Pfenninger, O., Brown, J., Duke, R. and Olsson, L. (1988): Tumorigenic human squamous lung-cancer cells have defined cell-surface carbohydrates that are absent from non-tumorigenic cells. *Proc. Natl. Acad. Sci.* **85**: 802-806.
- Philips, M.L., Nudelman, E., Gaeta, F.C.A., Perez, M., Singhal, A.K., Hakomori, S. and Paulson, J.C. (1990): ELAM-1 mediates cell adhesion by recognition of a carbohydrate ligand, sialyl-LeX. *Science* **250**: 1132-1135.
- Population based and hospital based registry (1994): Annual report, The Gujarat Cancer and Research Institute, Ahmedabad, INDIA
- Pos. O. Van der. Stelt M.E. Woltink G.J. Nijsten M.W.N Van der temple G.I. Van .dijk W.(1990) Chnges in the serum concentration and glycosylation of human alpha-1 acid glycoprotein and alpha-1 protease inhibitor in severly burned persons relations to interleukin-6 levels. *Clin. Exp. Immunol.* **82**: 579-582.
- Potier, M., Mameli, L., Belisle, L., .Dollaire, L. and Melancon, S.B. (1979): Flouorimetric assay of neuraminidase with a sodium (4-Methylumbelliferyl alpha-D-N-Acetyl neuraminate substrate. *Anal. Biochem.* **94**: 287-296.

- Putnam, F.W. and Udin, B. (1953): Proteins in multiple myeloma. I. Physiochemical study of serum proteins. *J. Biol. Chem.* **202**: 727-743.
- Rademacher, T.W., Parekh, R.B., Dwek, R.A., (1988): Glycobiology. *Annu. Rev. Biol. Chem.* **57**: 785-838.
- Reutter, W. and Bauer, C. (1978): In morris Hepatomas: Mechanism of regulation, W. Criss and H.P. Morris (eds.), Plenum, New York, pp 405-437.
- Romppanen, J., Eskelinen, M., Tikanojas, Mononcn, I. (1997): Total and lipid bound serum sialic acid in benign and malignant breast diseases. *Anti Cancer Res.* **17**: 1249-1253.
- Ronquist, G. and Nou, E. (1983): Serum sialyltransferase and fucosyl transferase activities in patients with bronchial carcinoma. *Cancer* **52**: 1679-1683.
- Rosato, F.E. and Radvin R.G. (1968): Serum protein bound hexoses in patients with breast cancer. *Cancer Res.* **28**: 44-45.
- Rothenberg, R.E., Laraja, R.D., Imran-ul-hag, saber, A.A., Nardkarni, M. (1996): Correlation between low neuraminidase blood levels and a predisposition towards family-related breast cancer. *The American Surgon* **62**: 949-951.
- Rothenberg, R.E., LaRaja, R.D., Muller, O.T. and Pryce, E.H. (1994): Sialic acid and neuraminidase levels in benign and malignant breast disease: A preliminary study. *Breast Dis.* **3**: 197-202.
- Ruoslahti, E., Yaveri, A., Kuusela, P. and Linder, E. (1973): Fibroblast surface antigen. A new serum protein. *Biochim. Biophys. Acta.* **322**: 350-358.
- Rye P.D. and Walker, R.A. (1989): Analysis of glycoproteins released from benign and malignant human breast: Change in size and fucosylation with malignancy. *Eur. J. Can. Clin. Oncol.* **25**: 65-72.

- Sakai, T., Yamamoto, K., Hokozaiki-Usui, K., Hino, F. and Kato, I. (1990): Rapid, simple enzymatic assay of free L-fucose in serum and urine, and its use as a marker for cancer, cirrhosis and gastric. *Clin. Chem.* **36**: 474-476.
- Salasvov, A., Sanchez-Hidalgo, V.M., Robago-Velsco, M., Sanchez-Leon, P., Vergas, C., Salas-valdea, R. (1980): Polyacrylamide gel electrophoresis in human serum proteins in normal individual and cancer patients. *Arch. Invest. Med. Mex.* **11**: 31-47.
- Santer, U. and Glick, M.C. (1983): Presence of fucosyl residue on oligosaccharide antennae of membrane glycopeptides of human neuroblastoma cells. *Cancer Res.* **43**: 4159-4166.
- Sashikantha, M.C. and Rao, B.B. (1994): Study of serum fucose and serum sialic acid levels in oral squamous cell carcinoma. *Indian J. Dent. Res.* **5**: 119-124.
- Sata, T., Roth, J., Zuber, C., Stamm, B. and Heitz, P.U. (1991): Expression of α -2-6 linked sialic acid residues in neoplastic but not in normal colonic mucosa. *Amer. Histochem. Cytochem.* **37**: 1577-1588.
- Scambia, G., Benedetti-panici, P., Perrore, L., Sonsini, C., Gjannelli, S., Gallow, A., Natali, P.G. and Mancoso, S. (1990): Serum levels of tumour associated glycoprotein (TAG-72) in patients with gynecological malignancy. *Br. J. Cancer* **62**: 147-151.
- Schachter, H. (1986): Biosynthetic controls that determines the branching and microheterogeneity of protein-bound oligosaccharides. *Adv. Exp. Med. Sci.* **205**: 53-85.
- Schauer, R. and Corfield, A.P. (1982): Occurrence of sialic acids - chemistry, metabolism and function. In cell biology monographs (Schauer, R. ed.), Springer Verlag, New York. Vol. 10: pp 5-50.

- Schauer, R., Kelm, S., Reuter, G and Shaw, L. (1995): New insights in to the moce and role of enzymatic sialic acid modifications. *Glycoconj. J.* **10**: 327
- Schauer, R., Kelm, S., Reuter, G., Roggentin, P. and Shaw, L. (1995): Biochemistry and role of sialic acid. In : Biology of sialic acids. (Abraham Rosenberg Ed.)Plenum press. New York. pp7-67.
- Schawartz, R., Schirrmacher, V. and Muhlradt, P.F. (1984): Glycoconjugates of murine tumour cell lines with different metastatic capacities. I: Difference in fucose utilization and in glycoprotein patterns. *Int.J.Cancer* **33**: 503-509.
- Schumacher, U.J., Mukhtar, D., Stehling, P., Reutter, W. (1996): Is the lectin binding pattern of human breast and colon cancer cells influenced by modulation of sialic acid metabolism. *Histochem.Cell.Biol.* **106**: 599-564.
- Sell, S. (1993): Detection of cancer by tumour markers in the blood: A review to the future. *Crit.Rev.Oncogenesis.* **4**: 419- 433.
- Sen, U., Guha, S. and Roychowdhury, J. (1984): Serum fucose levels as an index of progreession and regression of transplanted tumours. *IRCS Med. Sci.***12**:789-790.
- Sen, U., Guha, S., and Chowdhury, J.R. (1983): Serum fucosyl transferase activity and serum focose levels as diagnostic tools in malignancy. *Acta Med. Okayama* **37**: 457-462.
- Shahangian, S., Fristsche, H.A., Hughes, J.I., Katopodis, N. and Foemmel, R.S. (1991): Plasma protein bound sialic acid in patients with colorectal cancer. *Clin. Chem.* **37**: 200-204.
- Shamberger, R.J. (1984): Serum sialic acid in normal and in cancer patients. *Jr. Clin. Chem. Clin. Biochem.* **22**: 647-651.
- Shetlar, M.R. (1961): Serum glycoprotein their origin and significance. *Ann.Ny.Acd.Sci* **94**:44-54.

- Shimada, I. Shoji, M., Futu-suya. R., Katoh, T., Kominato, Y., Sakamoto, T., Fujikura, T. (1995): Elevation of ratio of urinary N-acetyl Neuraminlactose to free sialic acids in some advanced cancer patients. *J. Gastroentrol.* **30**: 21-27.
- Silverstri, B., Galderazzo, F., Coppola, V., Rosato, A., Icobelli, S., Natoli, C., Ullrivh, A., Sures, I., Azam, M., Brackebush, C., Chieco-Bianchi, L., and Amaderi, A. (1998): Differential effect on TCR: CD3 stimulation of a 90 Kda glycoprotein (gp90/Mac230) a member of scavenger receptor cystein domine protein family. *Clin. Expt. Immunol.* **113**: 394-400.
- Singh, N.K., Gupta, R.M., Patra, H., Sharma, M. and Sanyal, B. (1984): Multiple myeloma: A clinopathological and roentgenographic study of 44 cases. *Indian J. Cancer* **21**: 184-187.
- Skipski, V.P., Barclay, M., Archibald, F.M., Lynch, T.P. and Stock, C.C. (1971) : A new proteio lipid apparently Associated with cancer. *Proc. Soc. Exp. Biol. Med.* **136**: 1261-1264.
- Skoza, L. and Mohos, S. (1976): Stable thiobarbituric acid chromophore with dimethyl sulfoxide. Application to sialic acid assay in analytical De-O-acetylation. *Biochem. J.* **159**: 457-462.
- Smets, L.A. and Van-Beek, W.P (1984): Carbohydrates of tumour cell surface. *Biochim. Biophys. Acta* **738**: 237-249.
- Smets, L.A., Van-Beek, W.P. and Van-Nie, R. (1977): Membrane glycoprotein changes in primary mammary tumours associated with autonomous growth. *Cancer lett.* **3**: 133-138.
- Stefanelli, N., Klotes, H., Engel, A. and Bauer, P. (1985): Serum sialic acid in normal and in cancer patients. *J. Cancer. Res. Oncol.* **109**: 55-59.

- Suaraz-Nieto, G., Cuesta-Garcia, A. and Fernandez-Bustillo, F. (1986): Serum glycoprotein and prognosis in cancer of head and neck. *Clin. Otolaryngol.* **11**:41-45.
- Suer, S., Sonmez, H., Karaaslan, I., Baloglu, H., Kokoglu, E. (1996): Tissue sialic acid and fibronectin levels in human prostatic cancer. *Cancer Lett.* **99**: 135-137.
- Sattar, N., Scott, H.R., McMillan, D.C., Talwar, D., O'Reilly, D.S. and Fell, G.S. (1997): Acute-phase reactants and plasma trace element concentrations in non-small cell lung cancer. *Nutr. Cancer* **28**: 308-312.
- Takahasi, H., Salbara, T., Iwamura, S., Tomiba, A., Maeda, T., Onishi, S., Yamamoto, Y. and Enzan, H. (1994): Serum α -L fucosidase activity and tumour size in Hepatocellular carcinoma. *Hepatology* **19**: 1414-1417.
- Tatsumura, T., Sato, H., Mori, A. Komori, Y., Yamamoto, K., Fukatani, G. and Kuno, I. (1977): Clinical significance of fucose level in glycoprotein fraction of serum in patients with malignant tumours. *Cancer Res.* **37**: 4101-4103.
- Thaw, P.A. and Albutt, E.C. (1980): A critical evaluation of a serum seromuroid assay and its replacement by a serum alpha -1 Acid glycoprotein assay. *Ann. Clin. Biochem.* **17**: 142-143.
- Thompson, S. and Turner, G.A. (1987): Elevated levels of abnormally fucosylated heptoglobins in cancer sera. *Br. Jr. Cancer* **56**: 605-610.
- Thompson, S., Cantwell, B.M.J., Matta, K.L. and Turner, G.A. (1992): Parallel changes in blood levels of abnormally fucosylated heptoglobin and α -1-3 fucosyl transferase in relationship to tumour burden: More evidence for a disturbance of fucose metabolism in cancer. *Cancer Lett.* **65**: 115-121.

- Thompson, S., Gurthrie, D., Turner, G.A. (1988): Fucosylated form of α 1-antitrypsin that predict unresponsiveness to chemotherapy in ovarian cancer. *Br. J. Can.* **58**: 589-593.
- Tiemeyer, M., Swiedler, S.J., Ishihara, M., Moreland, M., Schweigruber, H., Hirtzer, P and Brandly, B.K. (1991): Carbohydrate ligand for endothelial-leukocyte adhesion molecule. *Proc. Natl. Acad. Sci.* **88**: 1138-1142.
- Turner, G. A., Goodarzi, M.T. and Thompson, S. (1995): Glycosylation of alpha-1-protease inhibitor and heptoglobin in ovarian cancer. Evidence for two different mechanism. *Glycoconj. J.* **12**: 211-215.
- Turner, G.A. (1992): N-Glycosylation of serum proteins in disease and its investigation using Lectins. *Clin. Chim. Acta.* **208**: 149-171.
- Turner, G.A., Skillen, A.W., Buamah, P., Guthrie, D., Welsh, J., Harrison, J. and Kowalski, A. (1985): Relation between raised concentrations of fucose sialic acids and acute phase proteins in serum from patients with cancer: choosing suitable serum glycoprotein markers. *J. Clin. Pathol.* **38**: 588-592.
- UICC TNM classification of malignant tumours (1980): UICC Technical Report series :Seller, A. H. (ed): 1st edition 51:7.
- Van Dijk, W., Pos, O., Van-der, Stelt, E. and Eap, C.B. (1991): Inflammation induced changes in excretion and glycosylation of genetic variant of α 1- acid glycoprotein studies with human sera. Primary culture of human hepatocytes in transgenic mice. *BioChem. J.* **276**: 343-347.
- Van Dijk, W., Turner, G.A. and Mackiewicz, A. (1994): Changes in glycosylation of acute-phase proteins in health and disease: occurrence, regulation and function. *Glycosylation and disease* **1**: 5-14.

- Vangsted, A., Drivsholm, L., Andersen, I., Pallesen, T., Zeuthen, J., Wallin H. (1994): New serum markers for small cell lung cancer. The ganglioside fucosyl GM3. *Cancer. Detect. Preven* **18**: 578-81.
- Varki, A. (1993): Biological roles of oligosaccharides: All of the theories are correct. *Glycobiology* **3**: 97-130.
- Varly, H., Gownlock, H. and Bell. M. (1984): The plasma proteins. In: Practical clinical biochemistry. Ed-5 The white Friars Press, Tonbridge, pp535-595.
- Verazin, G., Riley, W.M., Gregory, J., Tautu, C., Prorok J.J. and Alhadeff, J.A. (1990): Serum sialic acid and carcinoembryonic antigen levels in the detection and monitoring of colorectal cancer. *Dis.Colon. Rectum* **33**: 139-142.
- Vierbuchen, M.J., Fruectnicht, W., Brackrock, S., Krause, K.T., and Zienkiewicz, T.J. (1995): Quantitative lectin histochemical and immunochemical studies on the occurrence of alpha 2,6-linked sialic acid residues in colorectal carcinomas. Relation to clinicopathological features. *Cancer* **76**: 727- 735.
- Waalkes, J.E., Dinsmore, S.R., Tormey, D.C. and Waalkes, T.P. (1978): Serum protein bound carbohydrates for following the course of disease in patients with metastatic breast carcinoma. *J. Natl. Cancer Inst.* **61**: 703-707.
- Walker, C. and Gray, B.N. (1983): Acute phase reactant proteins and carcinoembryonic antigen in cancer of the colon and rectum. *Cancer* **52**: 150-154.
- Walz, G., Aruffo, A., Kolanus, W., Bevilacqua, M. and Seed, B. (1990): Recognition by ELAM-1 of the sialy-Lex determinant on myeloid and tumour cells. *Science* **250**: 1132-1135.

- Wang, J.W., Ambros, R.A., Weber, P.B. and Rosano, T.G. (1995): Fucosyl transferase and α -L fucosidase and fucose level in normal and malignant endometrial tissue. *Cancer Res.* **55**: 3654-3658.
- Warren, L., Buck, C.A. and Tuszynski, G.P. (1978): Glycopeptid changes and malignant transformation: A possible role for carbohydrate in malignant behaviour. *Biochim. Biophys. Acta.* **516**: 97-127.
- Wiederschain, G.Y., Rosenfeld, E.L., Brusilovsky, A.L. and Kolibaba, L.G. (1971): α -L-fucosidase and other glycosidase in human placenta, fetus liver and amniotic fluid at various stages of gestation. *Clin. Chim. Acta* **35**: 99-107.
- Winzler R.J. (1955): Determination of serum glycoproteins. In: Methods of biochemical analysis. Vol-II. Interscience publishers. New York, pp 279-311.
- Yamamoto, H., Kaneko, Y., Vandermulan, D., Kersey, D., Mkrdichian, E., Leestma, J. and Moskal, J.R. (1995): The expression of CMP-NeUAc: Gal beta 1,4 Glc Nac, alpha 2,6 sialyltransferase (E.C 2.4.99.1} the glycoprotein bearing alpha 2,6-linked sialic acids in human brain tumours. *Glycoconj. J.* **12**: 848-856.
- Yazawa, S., Asao, T., Nagamachi, Y., Abbas, S.A. and Matta, K.L. (1989): Tumor-related elevation of serum (α -1-3)-L-fucosyltransferase activity in gastric cancer. *J. Cancer Res. Clin. Oncol.* **115**: 451-455.
- Yazawa, S., Madiyalakan, R., Izawa, H., Asano, T., Furukawa, K. and Matta, K.L. (1988): Cancer associated elevation of α (1-3) fucosyl transferase activity in human serum. *Cancer* **62**: 516-520.
- Yogeeswaran, G. (1981): Metastatic potential is positively correlated with cell surface sialylation of cultured murine cell lines. *Science* **212**: 1514-1516.

- Yogeeswaran, G. (1983): Cell surface glycolipids and glycoproteins in malignant transformation. *Adv. Cancer Res.* **38**: 289-350
- Yoshimura, S., Tamaoki, N., Ueyama, Y. and Hata, J.I. (1978): Plasma protein production by human tumors xenotransplanted in nude mice. *Can. Res.* **38**: 3474-3478.
- Zieske, J.D. and Bernstein, I.A. (1982): Modification of cell surface glycoprotein: Addition of fucosyl residues during epidermal differentiation. *J.Cell.Biol.* **95**: 626-631.
- Zweig, M.M. and Campbell, G. (1993): Receiver-operating characteristic (ROC) plots. A fundamental evaluation tool in clinical medicine. *Clin.Chem.* **39**: 561-577.