

LIST OF FIGURES

Biochemical and Molecular Approaches for Developing Phosphate Solubilizing Plant Growth Promoting Rhizobacteria (PGPR)		
Figure No.	TITLE	Page No.
Chapter 1 Review of Literature and Introduction		
Fig. 1.1	PGPR as Phosphate solubilizer	12
Fig. 1.2	Direct oxidation pathway and its role in P solubilization	16
Fig. 1.3	Global distribution of phosphorus in soil	28
Chapter 2 Biochemical Characterization of Variation in the Nature of Organic Acid Secretion and Rock Phosphate Solubilization by Rhizobacteria, <i>Citrobacter</i> sp. DHRSS on Various Carbon Sources		
Fig. 2.1	Acidification on RP minimal medium containing methyl red by <i>Citrobacter</i> sp. DHRSS	34
Fig. 2.2	PCR amplification and cloning of rDNA of <i>Citrobacter</i> sp. DHRSS	34
Fig. 2.3 (a)	Partial 16S rDNA sequence of <i>Citrobacter</i> sp. DHRSS	35
Fig. 2.3 (b)	Homology search result of partial 16S rDNA of <i>Citrobacter</i> sp. DHRSS using RDP II	35
Fig. 2.3 (c)	Genbank accession result of partial 16S rDNA sequence of <i>Citrobacter</i> sp. DHRSS	36
Fig. 2.4 (a) and (b)	Growth and acidification pattern of <i>Citrobacter</i> Sp. DHRSS on buffered RP medium with different C sources	37
Fig. 2.5	Substrate specificity of glucose dehydrogenase activity of <i>Citrobacter</i> Sp. DHRSS using different sugars as substrates	40

Figure No.	TITLE	Page No.
Chapter 3 Repression of Mineral Phosphate Solubilizing (mps) Phenotype in the Presence of Weak Organic Acids in Plant Growth Promoting Fluorescent <i>Pseudomonads</i>		
Fig. 3.1	HPLC Profile of <i>P. aeruginosa</i> M3 and SP1 on 100mM Tris Cl (pH 8.0) and 100 mM glucose RP minimal medium	49
Fig. 3.2	Fluorescence of <i>P. aeruginosa</i> M3 and SP1 on Pseudomonas agar plate	50
Fig. 3.3 a	Partial 16S rDNA sequence of <i>P. aeruginosa</i> M3 and SP1	50
Fig. 3.3 b	Homology search result of partial 16S rDNA of <i>P. aeruginosa</i> M3 and SP1 using RDP II	51
Fig. 3.3 c	GenBank accession result of partial 16S rDNA sequence of <i>P. aeruginosa</i> M3 and SP1	52
Fig. 3.4:	Acidification by <i>P. aeruginosa</i> M3 and SP1 on methyl red containing minimal media agar plates with RP and 50 mM Tris Cl (pH 8.0)	54
Fig. 3.5	Acidification by <i>P. aeruginosa</i> (a) M3 and (b) SP1 in the presence of organic acids in RP containing minimal medium	56
Fig. 3.6	GDH activity of <i>E. asburiae</i> PSI3 in the presence of glucose with malate	57
Fig. 3.7	GDH and GAD enzyme activities of <i>P. aeruginosa</i> (1) M3 and (2) SP1 grown in the presence of glucose with malate	58
Fig. 3.8	GDH and GAD enzyme activities of <i>P. aeruginosa</i> (1) M3 and (2) SP1 grown in the presence of glucose with succinate	59
Fig. 3.9	GDH and GAD enzyme activities of <i>P. aeruginosa</i> (1) M3 and (2) SP1 grown in the presence of glucose with malate and succinate	60

Figure No.	TITLE	Page No.
Chapter 4 Screening of <i>Arabidopsis thaliana</i> cDNA Library for Mineral Phosphate Solubilizing (mps) Ability		
Fig 4.1	Restriction map of pYES	67
Fig 4.2	Site specific recombination between lox sites to release pYES	68
Fig. 4.3	Determination of the titre of <i>A. thaliana</i> λYES cDNA library	74
Fig. 4.4	Confirmation of conversion of λYES cDNA library of <i>A. thaliana</i> into pYES	77
Fig. 4.5	<i>Eco</i> RI digestion of pYES from cDNA transformants obtained on different carbon sources	78
Fig. 4.6	P solubilization by <i>E. coli</i> cDNA transformants on 70mM Tris Cl buffered RP minimal medium	79
Fig. 4.7 (a)	HPLC profile of organic acids secreted by <i>E. coli</i> cDNA transformants on xylose as C source	80
Fig. 4.7 (b)	HPLC profile of organic acids secreted by <i>E. coli</i> cDNA transformants on arabinose as C source	80
Fig. 4.8	P solubilization by <i>P. fluorescens</i> cDNA transformants on RP minimal medium with glucose	82
Fig. 4.9 a and (b)	Acidification and P releasing ability of <i>P. fluorescens</i> cDNA transformants on glucose with 100 mM Tris Cl (pH 8.0)	83
Fig. 4.10	Acidification by <i>P. fluorescens</i> cDNA transformants on glucose with 10mM KH ₂ PO ₄ as soluble Pi source with 100 mM Tris Cl (pH 8.0) buffered media	83
Fig. 4.11	Characterization of nature and amount of organic acid secreted by <i>P. fluorescens</i> cDNA transformants on glucose with 100 mM Tris Cl (pH 8.0)	84

Figure No.	TITLE	Page No.
Fig. 4.12	P solubilization by <i>P. fluorescens</i> cDNA transformants on 100mM xylose and 50 mM Tris Cl pH (8.0) buffered RP minimal medium	86
Fig. 4.13	PCR amplification of the cDNA from randomly obtained pYES clones from <i>E. coli</i>	86
Fig. 4.14	Schematic representation of integration of pYES in the genome of <i>P. fluorescens</i> and its effect on PCR amplification of the cDNA	89