

LIST OF FIGURES

CHAPTER-2

- Figure 2.1. MRI images of white matter lesions and structural abnormalities associated with FXTAS.
Figure 2.2. Classification of FMRI allele based on number of expanded CGG repeats at 5UTR of FMR1
Figure 2.3. Correlation of CGG repeat length, FMR1 RNA levels, FMRP expression and clinical outcomes.
Figure 2.4. Two basic proposed mechanisms of CGG-repeat induced toxicity in PM carriers.
Figure 2.5. Structural and functional domains of FMRP.
Figure 2.6. Role of Phosphorylated FMRP (P-FMRP) in regulation of translation.
Figure 2.7. Schematic representation of mitochondrial dysfunction involved in premutation condition.

CHAPTER-3

- Figure 3.1. Schematic representation of possible molecular mechanisms for mitochondrial dysfunctions in premutation condition.

CHAPTER-5

- Figure 5.1. Plasmids used to induce expression of 99XCGG repeats mimicking in vitro premutation condition.
Figure 5.2. Expression of plasmids containing expanded 99XCGG repeats affects cellular metabolism and causes decreased cell viability.
Figure 5.3. Expression of expanded CGG repeats causes decreased cell survival and increased apoptosis.
Figure 5.4. Expression of expanded CGG repeats activates caspases and causes apoptotic cell death.

CHAPTER-6

- Figure 6.1. Translation of CGG repeats forms FMRpolyG nuclear inclusions.
Figure 6.2. FMRpolyG forms small cytosolic inclusions.
Figure 6.3. Cytosolic inclusions of FMRpolyG are in the vicinity of the mitochondria in HEK293 cells.
Figure 6.4. FMRpolyG forms cytosolic small aggregates and show association with mitochondria.
Figure 6.5. Mitochondrial functions altered by expression of premutation allele.
Figure 6.6. FMRpolyG formed due to expanded CGG repeats disrupts mitochondrial respiratory chain supercomplexes (SCs) assembly and decreases activity of individual complexes.
Figure 6.7. Quantification of the area for band intensities of mitochondrial supercomplexes and individual complexes.
Figure 6.8. FMRpolyG alters transcripts encoded by mitochondrial genome.

CHAPTER-7

- Figure 7.1. Expression of expanded CGG repeats causes differential expression of cellular and mitochondrial miRNAs.
Figure 7.2. FMRpolyG translated due to CGG repeats causes altered pattern of mitochondrial associated miRNAs.
Figure 7.3. Bioinformatics analysis for targets of altered associated miRNAs with mitochondria in expanded CGG repeats transfected conditions.
Figure 7.4. Validation of candidate miRNAs by qPCR and localization of miR-320a in mitoplast under premutation condition.
Figure 7.5. Validation of endogenous controls and mitoplasmic localization of miR-320a.
Figure 7.6. Putative seeding sequences (7-8mers) present in miR-320a (red) which can interact with 3' region of mitotranscripts (green) identified by BLASTn tool.
Figure 7.7. miR-320a regulates levels of mitochondrial transcripts by modulating RISC assembly with Ago2 and mitotranscripts.
Figure 7.8. Interaction of miR-320a with Ago2 gets altered under premutation condition.
Figure 7.9. Transfection of miR-320a rescues mitochondrial transcripts and mitochondrial functions.
Figure 7.10. miR-320a causes decreased cellular ROS levels and increased survival in premutation condition.
Figure 7.11. Schematic representation of different pathways by which miRNA present in mitochondrial matrix can regulate mitochondrial RNA processing and ultimately mito-transcript levels.