

List of Figures

No.	TITLE	Page no.
1	Electron micrograph of a <i>Magnetospirillum gryphiswaldense</i> Magnetosome particles isolated from <i>M. gryphiswaldense</i>	2
2	Magnetic fluid being influenced by the presence of external magnet	2
3	Isolation of nucleic acid binding molecules	14
4	Magnetocapture protein interaction assays	17
5	Drug targeting using MTCs	30
6	PNP calibration curve	64
7	Standard curve of bovine serum albumin (BSA) by Bradford dye binding method	69
8	Standard curve of ALP by Bradford dye binding method	69
9	<i>In vitro</i> closed circulatory model system	75
10	Transmission electron micrograph of magnetic nanoparticles	82
11	Infrared spectrum of magnetite nanoparticles	83
12	(A) Agarose gel electrophoresis of genomic DNA isolated from blood (B) Agarose gel electrophoresis of isolated genomic DNA from PBMCs	93
13	(A) Agarose gel electrophoresis of genomic DNA isolated from buffy-coat (B) Agarose gel electrophoresis of genomic DNA isolated from cultured cells (HCT116).	94
14	Agarose gel electrophoresis of genomic DNA isolated using magnetite particles from different mammalian cell types	97
15	Comparison of the isolated genomic DNA from blood cells using conventional, silica magnetic beads and naked magnetite	100
16	DNA extracted from whole blood. Magnetic Vs Qiagen	104
17	DNA extracted from cultured cell lines: Magnetic Vs Qiagen	104

18	Agarose gel electrophoresis of genomic DNA isolated from <i>Streptomyces flaviscleroticus</i>	109
19	(A) Agarose gel electrophoresis of genomic DNA isolated from <i>E coli</i> (B) Agarose gel electrophoresis of genomic DNA isolated from <i>S typhi</i>	112
20	(A) Agarose gel electrophoresis of plasmid DNA (pBR322) isolated from <i>E coli</i> using magnetite as solid-phase adsorbent (B) Restriction endonuclease digestion of extracted plasmid DNA	115
21	(A) DNA isolated from <i>S cerevisiae</i> (B) DNA isolated from <i>D discoideum</i>	119
22	(A) Agarose gel electrophoresis of DNA isolated from plant tissue (B) Restriction digestion analysis of isolated genomic DNA from plant tissue	126
23	(A) Agarose gel electrophoresis of genomic DNA isolated from human blood cells using magnetic nanoparticles (B) Agarose gel electrophoresis of DNA extracted from agarose gel	129
24	(A) DNA extracted from agarose gel using magnetite as solid-phase support (B) Restriction analysis of DNA	131
25	Restriction endonuclease digestion of extracted DNA from blood and cultured cells	134
26	Restriction endonuclease digestion of extracted DNA	134
27	Restriction digest of DNA bound and unbound to magnetite particles	136
28	Restriction digestion of DNA extracted from mammalian tissues	137
29	(A) Restriction digests of <i>Streptomyces</i> genomic DNA (B) Restriction digests of <i>E coli</i> genomic DNA	138
30	Restriction digests of <i>Salmonella</i> genomic DNA	138
31	Restriction digests of genomic DNA extracted from yeast and <i>Dictyostelium</i>	140
32	Agarose gel electrophoresis of PCR amplicon	143

33	Agarose gel electrophoresis of 226-bp amplicon of GAPDH gene	143
34	Mechanism of direct linking of protein on naked magnetite	157
35	FTIR spectra of ALP alone and ALP bound to magnetic particles	167
36	Storage stability of free and immobilized ALP covalently linked to magnetite with the shaking method	169
37	Storage stability of free and immobilized ALP covalently linked to magnetite with the sonication method	170
38	Restriction digestion of pcDNA 3	174
39	Agarose gel electrophoresis of plasmid DNA (pcDNA 3 – 5 kb) following ligation reaction	174
40	Streptavidin (SA) binding to magnetic particles	179
41	Biotin-ALP activity of magnetic streptavidin nanoparticles immobilized at different pH	179
42	Comparison of biotin-binding capacity of magnetic streptavidin nanoparticles immobilized with shaking and sonication method, respectively	184
43	FTIR spectra of pure streptavidin and streptavidin bound to magnetic nanoparticles	185
44	Storage stability of magnetic streptavidin nanoparticles using shaking and sonication method, respectively	186
45	Colour and texture of MLs and magnetite	192
46	Magnetic property of MLs	192
47	TEM of ML	192
48	Spectral shift of DOX	194
49	Dilution induced drug leaching from liposomal/magnetoliposomal preparation	197
50	Serum induced DOX release from liposomes	200
51	Serum induced DOX release from magnetoliposomes	200

52	<i>In vitro</i> circulation stability of liposome encapsulated DOX in normal saline	203
53	<i>In vitro</i> circulation stability of magnetoliposomes encapsulated DOX in normal saline	203
54	<i>In vitro</i> circulation stability of liposomes encapsulated DOX in 5% dextran	204
55	<i>In vitro</i> circulation stability of magnetoliposomes encapsulated DOX in 5% dextran	204
56	Biodistribution of DOX after intravenous administration in rat	206
57	Mice tumor model	209
58	ML-DOX group animal with subcutaneously implanted magnet	209
59	% Reduction in the size of the tumor	210