

## Table of Tables

<b>Chapter 1</b>		
<b>Table 1.1</b>	Drugs previously used as therapeutic for obesity (Rodgers et al., 2012). DA: Dopamine; NA: Noradrenaline; 5-HT: 5-hydroxytryptamine	22
<b>Table 1.2</b>	Proposed health benefit of probiotics	27
<b>Table 1.3</b>	Commercially available probiotic in market	30
<b>Table 1.4</b>	Probiotic effect, mechanisms and fitness factors of EcN (Schultz et al., 2011)	32
<b>Table 1.5</b>	List of natural probiotics used for management of obesity	40
<b>Table 1.6</b>	List of natural and recombinant probiotic strains used in infection, allergy and inflammation model	47
<b>Table 1.7</b>	EcN in treatment of IBD (Malchow et al., 1997; Kruis et al., 1997; Rembacken et al., 1996; Kruis et al., 2004; Henker et al., 2008)	49
<b>Table 1.8</b>	Different prebiotics (Macfarlane et al., 2006).	62
<b>Chapter 2</b>		67
<b>Table 2.1</b>	Bacterial strains and plasmid	72
<b>Table 2.2</b>	Fecal PQQ concentration of all rat groups at the end of the experiment	78
<b>Table 2.3</b>	Serum SGOT and SGPT levels. Values are expressed as Mean $\pm$ SEM (n=6). ***P $\leq$ 0.001, ** P $\leq$ 0.01 in comparison to control rat group. ## P $\leq$ 0.01 in comparison to Vitamin C group.	79
<b>Table 2.4</b>	Effect of orally fed PQQ, different constructs of <i>E. coli</i> CFR 16 probiotics and Vitamin-C on SOD (Superoxide dismutase), CAT (Catalase) and GPx (Glutathione peroxidase) activity in control and experimental rats. Values are expressed as Mean $\pm$ SEM (n=6 each group). a P $\leq$ 0.05, a* P $\leq$ 0.001 compared with DMH group. b P $\leq$ 0.05, b* P $\leq$ 0.001 compared with 16:: <i>vgb-gfp</i> ( <i>pqq</i> ) group.	84
<b>Chapter 3</b>		92

<b>Table 3.1</b>	Bacterial strains and plasmids	106
<b>Table 3.2</b>	Effect of modified EcN and Ec16 on body weight gain, food intake, fasting glucose, serum insulin and serum uric levels of rats. Values are expressed as mean $\pm$ SEM (n = 6 each group). *P $\leq$ 0.05, ** P $\leq$ 0.01 and *** P $\leq$ 0.001 compared with control group. # # P $\leq$ 0.01 compared with fructose control group.	107
<b>Table 3.3</b>	PQQ concentration in liver tissue and fecal sample of rats fed with modified strains of EcN and Ec16 at the end of experiment: Values are expressed as mean $\pm$ SEM (n = 6 each group). ** P $\leq$ 0.01 and ***P $\leq$ 0.001 compared with control.	108
<b>Table 3.4</b>	Hepatic lipid profile. Values are expressed as mean $\pm$ SEM (n = 6 each group). *P $\leq$ 0.05, ** P $\leq$ 0.01 and ***P $\leq$ 0.001 compared with control. ##P $\leq$ 0.01 compared with fructose control. Hepatic values are expressed as mg/g Tissue.	108
<b>Table 3.5</b>	Colonic short chain fatty acid (SCFA). Values are expressed as mean $\pm$ SEM (n = 6 each group). *P $\leq$ 0.05 and** P $\leq$ 0.01 compared with control group. # P $\leq$ 0.05 and ##P $\leq$ 0.001 compared with Fructose control group. Concentration ( $\mu$ Moles/g Fecal matter)	113
<b>Chapter 4</b>		118
<b>Table 4.1</b>	Plasmid constructs and strains	123
<b>Table 4.2</b>	Effect of genetically engineered probiotic <i>E. coli</i> Nissle 1917 on liver lipid profile of rats. Values are expressed as mean $\pm$ SEM (n=6 each group). ** P $\leq$ 0.01 compared with control. ##P $\leq$ 0.01 compared with Sucrose control. Values are expressed as mg/g Tissue.	130
<b>Table 4.3</b>	Effect of genetically modified probiotic <i>E. coli</i> Nissle 1917 on Colonic SCFA. Values are expressed as mean $\pm$ SEM (n=6 each group). *P $\leq$ 0.05 and** P $\leq$ 0.01 compared with control group. Concentration ( $\mu$ Moles/ gm Fecal matter).	133
<b>Table 4.4</b>	Liver lipid profile of rats. Values are expressed as mean $\pm$ SEM (n=6 each group). * P $\leq$ 0.05, ** P $\leq$ 0.01 compared with	139

	control. #P ≤ 0.05, ##P ≤ 0.01 compared with Sucrose control. Values are expressed as mg/g Tissue.	
--	---	--