REFERENCES

CHAPTER 7

7. References

Aathira, R., and Jain, V. (2014). Advances in management of type 1 diabetes mellitus. World J Diabetes *5*, 689-696.

Abdi, R., Fiorina, P., Adra, C.N., Atkinson, M., and Sayegh, M.H. (2008). Immunomodulation by mesenchymal stem cells. Diabetes *57*, 1759-1767.

Abraham, E.J., Leech, C.A., Lin, J.C., Zulewski, H., and Habener, J.F. (2002). Insulinotropic hormone glucagon-like peptide-1 differentiation of human pancreatic islet-derived progenitor cells into insulin-producing cells. Endocrinology *143*, 3152-3161.

Afelik, S., and Rovira, M. (2017). Pancreatic beta-cell regeneration: Facultative or dedicated progenitors? Molecular and cellular endocrinology *445*, 85-94.

Ahlgren, U., Jonsson, J., Jonsson, L., Simu, K., and Edlund, H. (1998). β-Cell-specific inactivation of the mouseIpf1/Pdx1 gene results in loss of the β-cell phenotype and maturity onset diabetes. Genes & development *12*, 1763-1768.

Akiyama, T., Takasawa, S., Nata, K., Kobayashi, S., Abe, M., Shervani, N.J., Ikeda, T., Nakagawa, K., Unno, M., and Matsuno, S. (2001a). Activation of Reg gene, a gene for insulin-producing β-cell regeneration: poly (ADP-ribose) polymerase binds Reg promoter and regulates the transcription by autopoly (ADP-ribosyl) ation. Proceedings of the National Academy of Sciences *98*, 48-53.

Al-Hasani, K., Pfeifer, A., Courtney, M., Ben-Othman, N., Gjernes, E., Vieira, A., Druelle, N., Avolio, F., Ravassard, P., and Leuckx, G. (2013). Adult duct-lining cells can reprogram into β-like cells able to counter repeated cycles of toxin-induced diabetes. Developmental cell *26*, 86-100.

Altmeyer, M., Messner, S., Hassa, P.O., Fey, M., and Hottiger, M.O. (2009). Molecular mechanism of poly (ADP-ribosyl) ation by PARP1 and identification of lysine residues as ADP-ribose acceptor sites. Nucleic acids research *37*, 3723-3738.

Ambrose, H.E., Willimott, S., Beswick, R.W., Dantzer, F., De Murcia, J.M., Yelamos, J., and Wagner, S.D. (2009). Poly (ADP-ribose) polymerase-1 (Parp-1)-deficient mice demonstrate abnormal antibody responses. Immunology *127*, 178-186.

Amé, J.C., Spenlehauer, C., and de Murcia, G. (2004). The PARP superfamily. Bioessays *26*, 882-893.

Andrali, S.S., Sampley, M.L., Vanderford, N.L., and Özcan, S. (2008). Glucose regulation of insulin gene expression in pancreatic β-cells. Biochemical Journal *415*, 1-10.

Andreone, T., Meares, G.P., Hughes, K.J., Hansen, P.A., and Corbett, J.A. (2012a). Cytokine-mediated beta-cell damage in PARP-1-deficient islets. American journal of physiology Endocrinology and metabolism *303*, E172-179.

Andreone, T., Meares, G.P., Hughes, K.J., Hansen, P.A., and Corbett, J.A. (2012b). Cytokine-mediated β-cell damage in PARP-1-deficient islets. American Journal of Physiology-Endocrinology and Metabolism *303*, E172-E179.

Apelqvist, Å., Ahlgren, U., and Edlund, H. (1997). Sonic hedgehog directs specialised mesoderm differentiation in the intestine and pancreas. Current Biology *7*, 801-804.

Artner, I., Blanchi, B., Raum, J.C., Guo, M., Kaneko, T., Cordes, S., Sieweke, M., and Stein, R. (2007). MafB is required for islet beta cell maturation. Proceedings of the National Academy of Sciences of the United States of America *104*, 3853-3858.

Association, A.D. (2000). Nutrition recommendations and principles for people with diabetes mellitus. Diabetes care *23*, S43.

Association, A.D. (2003). Report of the expert committee on the diagnosis and classification of diabetes mellitus. Diabetes care *26*, s5-s20.

Atkinson, M.A., and Eisenbarth, G.S. (2001). Type 1 diabetes: new perspectives on disease pathogenesis and treatment. The Lancet *358*, 221-229.

Baeyens, L., De Breuck, S., Lardon, J., Mfopou, J., Rooman, I., and Bouwens, L. (2005). In vitro generation of insulin-producing beta cells from adult exocrine pancreatic cells. Diabetologia *48*, 49-57.

Baeyens, L., Lemper, M., Leuckx, G., De Groef, S., Bonfanti, P., Stangé, G., Shemer, R., Nord, C., Scheel, D.W., and Pan, F.C. (2014). Transient cytokine treatment induces acinar cell reprogramming and regenerates functional beta cell mass in diabetic mice. Nature biotechnology *32*, 76-83.

Bai, P., Houten, S.M., Huber, A., Schreiber, V., Watanabe, M., Kiss, B., de Murcia, G., Auwerx, J., and Ménissier-de Murcia, J. (2007). Poly (ADP-ribose) polymerase-2 [corrected] controls adipocyte differentiation and adipose tissue function through the regulation of the activity of the retinoid X receptor/peroxisome proliferator-activated receptor-gamma [corrected] heterodimer. The Journal of biological chemistry *282*, 37738-37746.

Banerjee, M., and Bhonde, R.R. (2003). Islet generation from intra islet precursor cells of diabetic pancreas: in vitro studies depicting in vivo differentiation. Jop *4*, 137-145.

Banerjee, M., Kanitkar, M., and Bhonde, R.R. (2005). Approaches towards endogenous pancreatic regeneration. Rev Diabet Stud *2*, 165-176.

Beger, H.G., Buchler, M., Kozarek, R., Lerch, M., Neoptolemos, J.P., Warshaw, A., Whitcomb, D., and Shiratori, K. (2009). The pancreas: an integrated textbook of basic science, medicine, and surgery (John Wiley & Sons).

Bell, G.I., Broughton, H.C., Levac, K.D., Allan, D.A., Xenocostas, A., and Hess, D.A. (2011). Transplanted human bone marrow progenitor subtypes stimulate endogenous islet regeneration and revascularization. Stem cells and development *21*, 97-109.

Bensley, R.R., and dquo (1911). Studies on the pancreas of the guinea pig. American Journal of Anatomy *12*, 297-388.

Bertolino, P., Holmberg, R., Reissmann, E., Andersson, O., Berggren, P.O., and Ibanez, C.F. (2008). Activin B receptor ALK7 is a negative regulator of pancreatic beta-cell function. Proceedings of the National Academy of Sciences of the United States of America *105*, 7246-7251.

Boj, S.F., Hwang, C.-I., Baker, L.A., Chio, I.I.C., Engle, D.D., Corbo, V., Jager, M., Ponz-Sarvise, M., Tiriac, H., and Spector, M.S. (2015). Organoid models of human and mouse ductal pancreatic cancer. Cell *160*, 324-338.

Bonner-Weir, S., Baxter, L.A., Schuppin, G.T., and Smith, F.E. (1993). A second pathway for regeneration of adult exocrine and endocrine pancreas: a possible recapitulation of embryonic development. Diabetes *42*, 1715-1720.

Bonner-Weir, S., Taneja, M., Weir, G.C., Tatarkiewicz, K., Song, K.-H., Sharma, A., and O'Neil, J.J. (2000). In vitro cultivation of human islets from expanded ductal tissue. Proceedings of the National Academy of Sciences *97*, 7999-8004.

Bonner-Weir, S., Trent, D., and Weir, G. (1983). Partial pancreatectomy in the rat and subsequent defect in glucose-induced insulin release. Journal of Clinical Investigation *71*, 1544.

Borch-Johnsen, K., Neil, A., Balkau, B., and Larsen, S. (1999). Glucose tolerance and mortality: comparison of WHO and American Diabetes Association diagnostic criteria. The Lancet *354*, 617.

Bretzel, R.G., Eckhard, M., and Brendel, M.D. (2004). Pancreatic islet and stem cell transplantation: new strategies in cell therapy of diabetes mellitus. Panminerva Med *46*, 25-42.

Brockenbrough, J.S., Weir, G.C., and Bonner-Weir, S. (1988). Discordance of exocrine and endocrine growth after 90% pancreatectomy in rats. Diabetes *37*, 232-236.

Brown, M.L., and Schneyer, A.L. (2010). Emerging roles for the TGFβ family in pancreatic β-cell homeostasis. Trends in Endocrinology & Metabolism *21*, 441-448.

Brownlee, M. (2001). Biochemistry and molecular cell biology of diabetic complications. Nature *414*, 813-820.

Brun, T., Franklin, I., St-Onge, L., Biason-Lauber, A., Schoenle, E.J., Wollheim, C.B., and Gauthier, B.R. (2004). The diabetes-linked transcription factor PAX4 promotes β-cell proliferation and survival in rat and human islets. J Cell Biol *167*, 1123-1135.

Bruni, A., Gala-Lopez, B., Pepper, A.R., Abualhassan, N.S., and Shapiro, A. (2014). Islet cell transplantation for the treatment of type 1 diabetes: recent advances and future challenges. Diabetes Metab Syndr Obes *7*, 211-223.

Burkart, V., Wang, Z.Q., Radons, J., Heller, B., Herceg, Z., Stingl, L., Wagner, E.F., and Kolb, H. (1999). Mice lacking the poly(ADP-ribose) polymerase gene are resistant to pancreatic beta-cell destruction and diabetes development induced by streptozocin. Nature medicine *5*, 314-319.

Caiafa, P., Guastafierro, T., and Zampieri, M. (2009). Epigenetics: poly (ADP-ribosyl) ation of PARP-1 regulates genomic methylation patterns. The FASEB Journal *23*, 672-678.

Caiafa, P., and Zlatanova, J. (2009). CCCTC-binding factor meets poly (ADP-ribose) polymerase-1. Journal of cellular physiology *219*, 265-270.

Cerf, M.E. (2006). Transcription factors regulating β-cell function. European Journal of Endocrinology *155*, 671-679.

Chamberlain, G., Fox, J., Ashton, B., and Middleton, J. (2007). Concise review: mesenchymal stem cells: their phenotype, differentiation capacity, immunological features, and potential for homing. Stem cells *25*, 2739-2749.

Chandra, V., G, S., Phadnis, S., Nair, P.D., and Bhonde, R.R. (2009). Generation of pancreatic hormone-expressing islet-like cell aggregates from murine adipose tissue-derived stem cells. Stem Cells *27*, 1941-1953.

Chao, E.C. (2014). SGLT-2 Inhibitors: A New Mechanism for Glycemic Control. Clinical diabetes : a publication of the American Diabetes Association 32, 4-11.

Chao, K.C., Chao, K.F., Fu, Y.S., and Liu, S.H. (2008). Islet-like clusters derived from mesenchymal stem cells in Wharton's Jelly of the human umbilical cord for transplantation to control type 1 diabetes. PloS one *3*, e1451.

Chen, L.B., Jiang, X.B., and Yang, L. (2004). Differentiation of rat marrow mesenchymal stem cells into pancreatic islet beta-cells. World J Gastroenterol *10*, 3016-3020.

Chen, S., Borowiak, M., Fox, J.L., Maehr, R., Osafune, K., Davidow, L., Lam, K., Peng, L.F., Schreiber, S.L., and Rubin, L.L. (2009). A small molecule that directs differentiation of human ESCs into the pancreatic lineage. Nature chemical biology *5*, 258-265.

Chera, S., Baronnier, D., Ghila, L., Cigliola, V., Jensen, J.N., Gu, G., Furuyama, K., Thorel, F., Gribble, F.M., and Reimann, F. (2014). Diabetes recovery by age-dependent conversion of pancreatic [dgr]-cells into insulin producers. Nature *514*, 503-507.

Chopp, M., Zhang, X.H., Li, Y., Wang, L., Chen, J., Lu, D., Lu, M., and Rosenblum, M. (2000). Spinal cord injury in rat: treatment with bone marrow stromal cell transplantation. Neuroreport *11*, 3001-3005.

Collombat, P., and Mansouri, A. (2009). [Pax4 transdifferentiates glucagon-secreting alpha cells to insulin-secreting beta endocrine pancreatic cells]. Medecine sciences : M/S *25*, 763-765.

Collombat, P., Mansouri, A., Hecksher-Sørensen, J., Serup, P., Krull, J., Gradwohl, G., and Gruss, P. (2003). Opposing actions of Arx and Pax4 in endocrine pancreas development. Genes & development *17*, 2591-2603.

Collombat, P., Xu, X., Ravassard, P., Sosa-Pineda, B., Dussaud, S., Billestrup, N., Madsen, O.D., Serup, P., Heimberg, H., and Mansouri, A. (2009). The ectopic expression of Pax4 in the mouse pancreas converts progenitor cells into α and subsequently β cells. Cell *138*, 449-462.

Consultation, W. (1999). Definition, diagnosis and classification of diabetes mellitus and its complications (Part).

Courtney, M., Pfeifer, A., Al-Hasani, K., Gjernes, E., Vieira, A., Ben-Othman, N., and Collombat, P. (2011). In vivo conversion of adult α-cells into β-like cells: a new research avenue in the context of type 1 diabetes. Diabetes, Obesity and Metabolism *13*, 47-52.

Creusot, R.J., Battaglia, M., Roncarolo, M.G., and Fathman, C.G. (2016). Concise Review: Cell-Based Therapies and Other Non-Traditional Approaches for Type 1 Diabetes. STEM CELLS.

D'Amour, K.A., Bang, A.G., Eliazer, S., Kelly, O.G., Agulnick, A.D., Smart, N.G., Moorman, M.A., Kroon, E., Carpenter, M.K., and Baetge, E.E. (2006). Production of pancreatic hormone–expressing endocrine cells from human embryonic stem cells. Nature biotechnology *24*, 1392-1401.

D'amours, D., Desnoyers, S., D'silva, I., and Poirier, G.G. (1999). Poly (ADP-ribosyl) ation reactions in the regulation of nuclear functions. Biochemical Journal *342*, 249-268.

Dadheech, N., Soni, S., Srivastava, A., Dadheech, S., Gupta, S., Gopurappilly, R., Bhonde, R.R., and Gupta, S. (2013). A small molecule swertisin from Enicostemma littorale differentiates NIH3T3 cells into Islet-Like clusters and restores normoglycemia upon transplantation in diabetic balb/c mice. Evidence-Based Complementary and Alternative Medicine *2013*.

Dadheech, N., Srivastava, A., Paranjape, N., Gupta, S., Dave, A., Shah, G.M., Bhonde, R.R., and Gupta, S. (2015). Swertisin an Anti-Diabetic Compound Facilitate Islet Neogenesis from Pancreatic Stem/Progenitor Cells via p-38 MAP Kinase-SMAD Pathway: An In-Vitro and In-Vivo Study. PloS one *10*, e0128244.

Dahl, M., Maturi, V., Lönn, P., Papoutsoglou, P., Zieba, A., Vanlandewijck, M., van der Heide, L.P., Watanabe, Y., Söderberg, O., and Hottiger, M.O. (2014). Fine-tuning of Smad protein function by poly (ADP-ribose) polymerases and poly (ADP-ribose) glycohydrolase during transforming growth factor β signaling. PloS one *9*, e103651.

Dalvi, M.P., Umrani, M.R., Joglekar, M.V., and Hardikar, A.A. (2009). Human pancreatic islet progenitor cells demonstrate phenotypic plasticity in vitro. J Biosci *34*, 523-528.

De Murcia, J.M., Niedergang, C., Trucco, C., Ricoul, M., Dutrillaux, B., Mark, M., Oliver, F.J., Masson, M., Dierich, A., and LeMeur, M. (1997). Requirement of poly (ADP-ribose) polymerase in recovery from DNA damage in mice and in cells. Proceedings of the National Academy of Sciences *94*, 7303-7307.

De Murcia, J.M., Ricoul, M., Tartier, L., Niedergang, C., Huber, A., Dantzer, F., Schreiber, V., Amé, J.C., Dierich, A., and LeMeur, M. (2003). Functional interaction between PARP-1 and PARP-2 in chromosome stability and embryonic development in mouse. The EMBO journal *22*, 2255-2263.

Deans, R.J., and Moseley, A.B. (2000). Mesenchymal stem cells: biology and potential clinical uses. Experimental hematology *28*, 875-884.

Desai, B.M., Oliver-Krasinski, J., De Leon, D.D., Farzad, C., Hong, N., Leach, S.D., and Stoffers, D.A. (2007). Preexisting pancreatic acinar cells contribute to acinar cell, but not islet β cell, regeneration. The Journal of clinical investigation *117*, 971-977.

Dieckmann, A., Kriebel, M., Andriambeloson, E., Ziegler, D., and Elmlinger, M. (2012). Treatment with Actovegin(R) improves sensory nerve function and pathology in streptozotocin-diabetic rats via mechanisms involving inhibition of PARP activation.

Experimental and clinical endocrinology & diabetes : official journal, German Society of Endocrinology [and] German Diabetes Association *120*, 132-138.

Doege, C.A., Inoue, K., Yamashita, T., Rhee, D.B., Travis, S., Fujita, R., Guarnieri, P., Bhagat, G., Vanti, W.B., and Shih, A. (2012). Early-stage epigenetic modification during somatic cell reprogramming by Parp1 and Tet2. Nature *488*, 652-655.

Dor, Y., Brown, J., Martinez, O.I., and Melton, D.A. (2004). Adult pancreatic β-cells are formed by self-duplication rather than stem-cell differentiation. Nature *429*, 41-46.

Du, X., Matsumura, T., Edelstein, D., Rossetti, L., Zsengeller, Z., Szabo, C., and Brownlee, M. (2003). Inhibition of GAPDH activity by poly(ADP-ribose) polymerase activates three major pathways of hyperglycemic damage in endothelial cells. The Journal of clinical investigation *112*, 1049-1057.

Edlund, H. (2001). Developmental biology of the pancreas. Diabetes *50 Suppl 1*, S5-9.

El-Gohary, Y., Tulachan, S., Guo, P., Welsh, C., Wiersch, J., Prasadan, K., Paredes, J., Shiota, C., Xiao, X., and Wada, Y. (2013). Smad signaling pathways regulate pancreatic endocrine development. Developmental biology *378*, 83-93.

El-Gohary, Y., Wiersch, J., Tulachan, S., Xiao, X., Guo, P., Rymer, C., Fischbach, S., Prasadan, K., Shiota, C., and Gaffar, I. (2016). Intraislet pancreatic ducts can give rise to insulin-positive cells. Endocrinology *157*, 166-175.

Eng, J., Kleinman, W.A., Singh, L., Singh, G., and Raufman, J.P. (1992). Isolation and characterization of exendin-4, an exendin-3 analogue, from Heloderma suspectum venom. Further evidence for an exendin receptor on dispersed acini from guinea pig pancreas. J Biol Chem *267*, 7402-7405.

Erener, S., Pétrilli, V., Kassner, I., Minotti, R., Castillo, R., Santoro, R., Hassa, P.O., Tschopp, J., and Hottiger, M.O. (2012). Inflammasome-activated caspase 7 cleaves PARP1 to enhance the expression of a subset of NF-κB target genes. Molecular cell *46*, 200-211.

Ezquer, F.E., Ezquer, M.E., Parrau, D.B., Carpio, D., Yañez, A.J., and Conget, P.A. (2008). Systemic administration of multipotent mesenchymal stromal cells reverts hyperglycemia and prevents nephropathy in type 1 diabetic mice. Biology of Blood and Marrow Transplantation *14*, 631-640.

Fagerberg, L., Hallström, B.M., Oksvold, P., Kampf, C., Djureinovic, D., Odeberg, J., Habuka, M., Tahmasebpoor, S., Danielsson, A., and Edlund, K. (2014a). Analysis of the human tissue-specific expression by genome-wide integration of transcriptomics and antibody-based proteomics. Molecular & Cellular Proteomics *13*, 397-406.

Falin, L. (1967). The development and cytodifferentiation of the islets of Langerhans in human embryos and foetuses. Cells Tissues Organs *68*, 147-168.

Feng, N., Hao, G., Yang, F., Qu, F., Zheng, H., Liang, S., and Jin, Y. (2016). Transplantation of mesenchymal stem cells promotes the functional recovery of the central nervous system following cerebral ischemia by inhibiting myelin-associated inhibitor expression and neural apoptosis. Experimental and Therapeutic Medicine *11*, 1595-1600.

Fernandes, A., King, L., Guz, Y., Stein, R., Wright, C., and Teitelman, G. (1997). Differentiation of New Insulin-Producing Cells Is Induced by Injury in Adult Pancreatic Islets 1. Endocrinology *138*, 1750-1762.

Figliuzzi, M., Cornolti, R., Perico, N., Rota, C., Morigi, M., Remuzzi, G., Remuzzi, A., and Benigni, A. (2009). Bone marrow–derived mesenchymal stem cells improve islet graft function in diabetic rats. Paper presented at: Transplantation proceedings (Elsevier).

Fleming, A., and Rosenberg, L. (2007). Prospects and challenges for islet regeneration as a treatment for diabetes: a review of islet neogenesis associated protein. Journal of diabetes science and technology *1*, 231-244.

Florio, P., Luisi, S., Marchetti, P., Lupi, R., Cobellis, L., Falaschi, C., Sugino, H., Navalesi, R., Genazzani, A., and Petraglia, F. (2000). Activin A stimulates insulin secretion in cultured human pancreatic islets. Journal of endocrinological investigation *23*, 231-234.

Froud, T., Ricordi, C., Baidal, D.A., Hafiz, M.M., Ponte, G., Cure, P., Pileggi, A., Poggioli, R., Ichii, H., and Khan, A. (2005). Islet transplantation in type 1 diabetes mellitus using cultured islets and steroid-free immunosuppression: Miami experience. American Journal of Transplantation *5*, 2037-2046.

Fu, Z., Gilbert, E.R., and Liu, D. (2013). Regulation of insulin synthesis and secretion and pancreatic Beta-cell dysfunction in diabetes. Current diabetes reviews *9*, 25-53.

Gabr, M.M., Sobh, M.M., Zakaria, M.M., Refaie, A.F., and Ghoneim, M.A. (2008). Transplantation of insulin-producing clusters derived from adult bone marrow stem cells to treat diabetes in rats. Exp Clin Transplant *6*, 236-243.

Gagné, J.-P., Hendzel, M.J., Droit, A., and Poirier, G.G. (2006). The expanding role of poly (ADP-ribose) metabolism: current challenges and new perspectives. Current opinion in cell biology *18*, 145-151.

Gagné, J.-P., Moreel, X., Gagné, P., Labelle, Y., Droit, A., Chevalier-Paré, M., Bourassa, S., Mcdonald, D., Hendzel, M.J., and Prigent, C. (2008). Proteomic investigation of phosphorylation sites in poly (ADP-ribose) polymerase-1 and poly (ADP-ribose) glycohydrolase. Journal of proteome research *8*, 1014-1029.

Gao, F., Kwon, S.W., Zhao, Y., and Jin, Y. (2009). PARP1 poly (ADP-ribosyl) ates Sox2 to control Sox2 protein levels and FGF4 expression during embryonic stem cell differentiation. Journal of Biological Chemistry *284*, 22263-22273.

Gao, F., Wu, D.-q., Hu, Y.-h., and Jin, G.-x. (2008). Extracellular matrix gel is necessary for in vitro cultivation of insulin producing cells from human umbilical cord blood derived mesenchymal stem cells. Chinese medical journal *121*, 811-818.

Garcia Soriano, F., Virag, L., Jagtap, P., Szabo, E., Mabley, J.G., Liaudet, L., Marton, A., Hoyt, D.G., Murthy, K.G., Salzman, A.L., *et al.* (2001). Diabetic endothelial dysfunction: the role of poly(ADP-ribose) polymerase activation. Nature medicine *7*, 108-113.

Gerich, J.E. (1989). Oral hypoglycemic agents. New England Journal of Medicine *321*, 1231-1245.

Gershengorn, M.C., Hardikar, A.A., Wei, C., Geras-Raaka, E., Marcus-Samuels, B., and Raaka, B.M. (2004). Epithelial-to-mesenchymal transition generates proliferative human islet precursor cells. Science *306*, 2261-2264.

Gibson, B.A., and Kraus, W.L. (2012). New insights into the molecular and cellular functions of poly (ADP-ribose) and PARPs. Nature reviews Molecular cell biology *13*, 411-424.

Gittes, G.K. (2009). Developmental biology of the pancreas: a comprehensive review. Developmental biology *326*, 4-35.

Gonzalez-Flores, A., Aguilar-Quesada, R., Siles, E., Pozo, S., Rodriguez-Lara, M., López–Jiménez, L., Lopez-Rodriguez, M., Peralta-Leal, A., Villar, D., and Martin-Oliva, D. (2014). Interaction between PARP-1 and HIF-2α in the hypoxic response. Oncogene *33*, 891-898.

Goodall, T.A., and Halford, W.K. (1991). Self-management of diabetes mellitus: a critical review. Health Psychology *10*, 1.

Gopurappilly, R., Bhat, V., and Bhonde, R. (2013). Pancreatic tissue resident mesenchymal stromal cell (MSC)-like cells as a source of in vitro islet neogenesis. Journal of cellular biochemistry *114*, 2240-2247.

Gorasia, D.G., Dudek, N.L., Veith, P.D., Shankar, R., Safavi-Hemami, H., Williamson, N.A., Reynolds, E.C., Hubbard, M.J., and Purcell, A.W. (2015). Pancreatic beta cells are

highly susceptible to oxidative and ER stresses during the development of diabetes. Journal of proteome research *14*, 688-699.

Govindasamy, V., Ronald, V., Abdullah, A., Nathan, K.G., Ab. Aziz, Z., Abdullah, M., Musa, S., Kasim, N.A., and Bhonde, R. (2011). Differentiation of dental pulp stem cells into islet-like aggregates. Journal of dental research *90*, 646-652.

Gradwohl, G., Dierich, A., LeMeur, M., and Guillemot, F. (2000). neurogenin3 is required for the development of the four endocrine cell lineages of the pancreas. Proceedings of the National Academy of Sciences *97*, 1607-1611.

Guariguata, L., Whiting, D.R., Hambleton, I., Beagley, J., Linnenkamp, U., and Shaw, J.E. (2014). Global estimates of diabetes prevalence for 2013 and projections for 2035. Diabetes research and clinical practice *103*, 137-149.

Guastafierro, T., Cecchinelli, B., Zampieri, M., Reale, A., Riggio, G., Sthandier, O., Zupi, G., Calabrese, L., and Caiafa, P. (2008). CCCTC-binding factor activates PARP-1 affecting DNA methylation machinery. Journal of Biological Chemistry *283*, 21873-21880.

Gupta S, Dadheech N, Singh A, and, S.S., and RR, B. (2010). Enicostemma littorale: A new therapeutic for islet neogenesis. International Journal of Integrative biology *9*, 49-53.

Guz, Y., Nasir, I., and Teitelman, G. (2001). Regeneration of pancreatic beta cells from intra-islet precursor cells in an experimental model of diabetes. Endocrinology *142*, 4956-4968.

Hakmé, A., Wong, H.K., Dantzer, F., and Schreiber, V. (2008). The expanding field of poly (ADP-ribosyl) ation reactions. EMBO reports *9*, 1094-1100.

Hanai, S., Uchida, M., Kobayashi, S., Miwa, M., and Uchida, K. (1998). Genomic organization of Drosophila poly (ADP-ribose) polymerase and distribution of its mRNA during development. Journal of Biological Chemistry *273*, 11881-11886.

Hassa, P.O., Buerki, C., Lombardi, C., Imhof, R., and Hottiger, M.O. (2003). Transcriptional coactivation of nuclear factor-κB-dependent gene expression by p300 is regulated by poly (ADP)-ribose polymerase-1. Journal of Biological Chemistry *278*, 45145-45153.

Hassa, P.O., Covic, M., Bedford, M.T., and Hottiger, M.O. (2008). Protein arginine methyltransferase 1 coactivates NF-κB-dependent gene expression synergistically with CARM1 and PARP1. Journal of molecular biology *377*, 668-678.

Hassa, P.O., Covic, M., Hasan, S., Imhof, R., and Hottiger, M.O. (2001). The enzymatic and DNA binding activity of PARP-1 are not required for NF-κB coactivator function. Journal of Biological Chemistry *276*, 45588-45597.

Hassa, P.O., Haenni, S.S., Buerki, C., Meier, N.I., Lane, W.S., Owen, H., Gersbach, M., Imhof, R., and Hottiger, M.O. (2005). Acetylation of poly (ADP-ribose) polymerase-1 by p300/CREB-binding protein regulates coactivation of NF-κB-dependent transcription. Journal of biological chemistry *280*, 40450-40464.

Hassa, P.O., Haenni, S.S., Elser, M., and Hottiger, M.O. (2006). Nuclear ADP-ribosylation reactions in mammalian cells: where are we today and where are we going? Microbiology and Molecular Biology Reviews *70*, 789-829.

Heller, B., Wang, Z.Q., Wagner, E.F., Radons, J., Burkle, A., Fehsel, K., Burkart, V., and Kolb, H. (1995). Inactivation of the poly(ADP-ribose) polymerase gene affects oxygen radical and nitric oxide toxicity in islet cells. The Journal of biological chemistry *270*, 11176-11180.

Hematti, P., Kim, J., Stein, A.P., and Kaufman, D. (2013). Potential role of mesenchymal stromal cells in pancreatic islet transplantation. Transplantation Reviews *27*, 21-29.

Henschler, R., Deak, E., and Seifried, E. (2008). Homing of mesenchymal stem cells. Transfusion Medicine and Hemotherapy *35*, 306-312.

Herbst, R.S. (2004). Review of epidermal growth factor receptor biology. Int J Radiat Oncol Biol Phys *59*, 21-26.

Hess, D., Li, L., Martin, M., Sakano, S., Hill, D., Strutt, B., Thyssen, S., Gray, D.A., and Bhatia, M. (2003). Bone marrow–derived stem cells initiate pancreatic regeneration. Nature biotechnology *21*, 763-770.

Holland, A.M., Elefanty, A.G., and Stanley, E.G. (2009). Pancreatic differentiation from pluripotent stem cells: tweaking the system. Cell Res *19*, 395-396.

Hottiger, M.O., Hassa, P.O., Lüscher, B., Schüler, H., and Koch-Nolte, F. (2010). Toward a unified nomenclature for mammalian ADP-ribosyltransferases. Trends in biochemical sciences *35*, 208-219.

Hsu, F.L., Huang, C.F., Chen, Y.W., Yen, Y.P., Wu, C.T., Uang, B.J., Yang, R.S., and Liu, S.H. (2013). Antidiabetic effects of pterosin A, a small-molecular-weight natural product, on diabetic mouse models. Diabetes *62*, 628-638.

Huang, D., Wang, Y., Wang, L., Zhang, F., Deng, S., Wang, R., Zhang, Y., and Huang, K. (2011). Poly (ADP-ribose) polymerase 1 is indispensable for transforming growth factor-β Induced Smad3 activation in vascular smooth muscle cell. PLoS One *6*, e27123.

Huang, H.-P., Liu, M., El-Hodiri, H.M., Chu, K., Jamrich, M., and Tsai, M.-J. (2000). Regulation of the Pancreatic Islet-Specific GeneBETA2 (neuroD) by Neurogenin 3. Molecular and cellular biology *20*, 3292-3307.

Huch, M., Bonfanti, P., Boj, S.F., Sato, T., Loomans, C.J., Van De Wetering, M., Sojoodi, M., Li, V.S., Schuijers, J., and Gracanin, A. (2013). Unlimited in vitro expansion of adult bi-potent pancreas progenitors through the Lgr5/R-spondin axis. The EMBO journal *32*, 2708-2721.

Hui, H., Wright, C., and Perfetti, R. (2001). Glucagon-like peptide 1 induces differentiation of islet duodenal homeobox-1-positive pancreatic ductal cells into insulin-secreting cells. Diabetes *50*, 785-796.

Immervoll, H., Hoem, D., Steffensen, O.J., Miletic, H., and Molven, A. (2011). Visualization of CD44 and CD133 in Normal Pancreas and Pancreatic Ductal Adenocarcinomas Non-overlapping Membrane Expression in Cell Populations Positive for Both Markers. Journal of Histochemistry & Cytochemistry *59*, 441-455.

Ito, T., Itakura, S., Todorov, I., Rawson, J., Asari, S., Shintaku, J., Nair, I., Ferreri, K., Kandeel, F., and Mullen, Y. (2010). Mesenchymal stem cell and islet co-transplantation promotes graft revascularization and function. Transplantation *89*, 1438-1445.

Jiang, F.X., and Morahan, G. (2014). Pancreatic stem cells remain unresolved. Stem cells and development *23*, 2803-2812.

Jiang, J., Au, M., Lu, K., Eshpeter, A., Korbutt, G., Fisk, G., and Majumdar, A.S. (2007). Generation of insulin-producing islet-like clusters from human embryonic stem cells. Stem cells *25*, 1940-1953.

Jiang, W., Sui, X., Zhang, D., Liu, M., Ding, M., Shi, Y., and Deng, H. (2011). CD24: a novel surface marker for PDX1-positive pancreatic progenitors derived from human embryonic stem cells. Stem cells *29*, 609-617.

Jin, L., Feng, T., Shih, H.P., Zerda, R., Luo, A., Hsu, J., Mahdavi, A., Sander, M., Tirrell, D.A., and Riggs, A.D. (2013). Colony-forming cells in the adult mouse pancreas are expandable in Matrigel and form endocrine/acinar colonies in laminin hydrogel. Proceedings of the National Academy of Sciences *110*, 3907-3912.

Jinnin, M., Ihn, H., and Tamaki, K. (2006). Characterization of SIS3, a novel specific inhibitor of Smad3, and its effect on transforming growth factor-β1-induced extracellular matrix expression. Molecular pharmacology *69*, 597-607.

Joglekar, M.V., Parekh, V.S., and Hardikar, A.A. (2007). New pancreas from old: microregulators of pancreas regeneration. Trends in Endocrinology & Metabolism *18*, 393-400.

Johnson, J.D., Ahmed, N.T., Luciani, D.S., Han, Z., Tran, H., Fujita, J., Misler, S., Edlund, H., and Polonsky, K.S. (2003). Increased islet apoptosis in Pdx1+/–mice. The Journal of clinical investigation *111*, 1147-1160.

Jonsson, J., Ahlgren, U., Edlund, T., and Edlund, H. (2003). IPF1, a homeodomain protein with a dual function in pancreas development. International Journal of Developmental Biology *39*, 789-798.

Joslin, E.P., and Kahn, C.R. (2005). Joslin's Diabetes Mellitus: Edited by C. Ronald Kahn...[et Al.] (Lippincott Williams & Wilkins).

Ju, B.-G., Solum, D., Song, E.J., Lee, K.-J., Rose, D.W., Glass, C.K., and Rosenfeld, M.G. (2004). Activating the PARP-1 sensor component of the groucho/TLE1 corepressor complex mediates a CaMKinase IIδ-dependent neurogenic gene activation pathway. Cell *119*, 815-829.

Jubin, T., Kadam, A., Gani, A.R., Singh, M., Dwivedi, M., and Begum, R. (2017). Poly ADP-ribose polymerase-1: Beyond transcription and towards differentiation. Paper presented at: Seminars in cell & developmental biology (Elsevier).

Kalra, S., Baruah, M.P., Sahay, R.K., Unnikrishnan, A.G., Uppal, S., and Adetunji, O. (2016). Glucagon-like peptide-1 receptor agonists in the treatment of type 2 diabetes: Past, present, and future. Indian journal of endocrinology and metabolism 20, 254-267.

Karamanou, M., Protogerou, A., Tsoucalas, G., Androutsos, G., and Poulakou-Rebelakou, E. (2016). Milestones in the history of diabetes mellitus: The main contributors. World journal of diabetes *7*, 1-7.

Karaoz, E., Ayhan, S., Okçu, A., Aksoy, A., Bayazıt, G., Osman Gürol, A., and Duruksu, G. (2011). Bone marrow-derived mesenchymal stem cells co-cultured with pancreatic islets display β cell plasticity. Journal of tissue engineering and regenerative medicine *5*, 491-500.

Karp, J.M., and Teo, G.S.L. (2009). Mesenchymal stem cell homing: the devil is in the details. Cell stem cell *4*, 206-216.

Katdare, M.R., Bhonde, R.R., and Parab, P.B. (2004). Analysis of morphological and functional maturation of neoislets generated in vitro from pancreatic ductal cells and their suitability for islet banking and transplantation. J Endocrinol *182*, 105-112.

Kauppinen, T.M., Chan, W.Y., Suh, S.W., Wiggins, A.K., Huang, E.J., and Swanson, R.A. (2006). Direct phosphorylation and regulation of poly (ADP-ribose) polymerase-1 by extracellular signal-regulated kinases 1/2. Proceedings of the National Academy of Sciences *103*, 7136-7141.

Khorsandi, L., Saremy, S., Khodadadi, A., and Dehbashi, F. (2016). Effects of Exendine-4 on The Differentiation of Insulin Producing Cells from Rat Adipose-Derived Mesenchymal Stem Cells. Cell Journal (Yakhteh) *17*, 720.

Kim, H.S., Hong, S.H., Oh, S.H., Kim, J.H., Lee, M.S., and Lee, M.K. (2013). Activin A, exendin-4, and glucose stimulate differentiation of human pancreatic ductal cells. The Journal of endocrinology *217*, 241-252.

Kim, M.Y., Mauro, S., Gévry, N., Lis, J.T., and Kraus, W.L. (2004). NAD+-dependent modulation of chromatin structure and transcription by nucleosome binding properties of PARP-1. Cell *119*, 803-814.

Kim, M.Y., Zhang, T., and Kraus, W.L. (2005). Poly (ADP-ribosyl) ation by PARP-1:PAR-laying'NAD+ into a nuclear signal. Genes & development *19*, 1951-1967.

Kim, S.K., and Hebrok, M. (2001). Intercellular signals regulating pancreas development and function. Genes & development *15*, 111-127.

Kim, S.K., Hebrok, M., Li, E., Oh, S.P., Schrewe, H., Harmon, E.B., Lee, J.S., and Melton, D.A. (2000). Activin receptor patterning of foregut organogenesis. Genes & development *14*, 1866-1871.

Kim, S.Y., Lee, S., Hong, S.W., Min, B.H., Lee, K.U., Bendayan, M., and Park, I.S. (2010). Nestin action during insulin-secreting cell differentiation. The journal of histochemistry and cytochemistry : official journal of the Histochemistry Society *58*, 567-576.

Kishi, Y., Fujihara, H., Kawaguchi, K., Yamada, H., Nakayama, R., Yamamoto, N., Fujihara, Y., Hamada, Y., Satomura, K., and Masutani, M. (2015). PARP inhibitor PJ34 suppresses osteogenic differentiation in mouse mesenchymal stem cells by modulating BMP-2 signaling pathway. International journal of molecular sciences *16*, 24820-24838.

Kiss, L., and Szabo, C. (2005). The pathogenesis of diabetic complications: the role of DNA injury and poly(ADP-ribose) polymerase activation in peroxynitrite-mediated cytotoxicity. Memorias do Instituto Oswaldo Cruz *100 Suppl 1*, 29-37.

Kobayashi, S., Akiyama, T., Nata, K., Abe, M., Tajima, M., Shervani, N.J., Unno, M., Matsuno, S., Sasaki, H., Takasawa, S., *et al.* (2000). Identification of a receptor for reg (regenerating gene) protein, a pancreatic beta-cell regeneration factor. The Journal of biological chemistry *275*, 10723-10726.

Kojima, I., and Umezawa, K. (2006). Conophylline: a novel differentiation inducer for pancreatic beta cells. The international journal of biochemistry & cell biology *38*, 923-930.

Kondo, S., Muneta, T., Nakagawa, Y., Koga, H., Watanabe, T., Tsuji, K., Sotome, S., Okawa, A., Kiuchi, S., and Ono, H. (2016). Transplantation of autologous synovial mesenchymal stem cells promotes meniscus regeneration in aged primates. Journal of Orthopaedic Research.

Kopinke, D., Brailsford, M., Shea, J.E., Leavitt, R., Scaife, C.L., and Murtaugh, L.C. (2011). Lineage tracing reveals the dynamic contribution of Hes1+ cells to the developing and adult pancreas. Development *138*, 431-441.

Kopinke, D., and Murtaugh, L.C. (2010). Exocrine-to-endocrine differentiation is detectable only prior to birth in the uninjured mouse pancreas. BMC developmental biology *10*, 38.

Kopp, J.L., Dubois, C.L., Hao, E., Thorel, F., Herrera, P.L., and Sander, M. (2011a). Progenitor cell domains in the developing and adult pancreas. Cell cycle *10*, 1921-1927.

Kopp, J.L., Dubois, C.L., Schaffer, A.E., Hao, E., Shih, H.P., Seymour, P.A., Ma, J., and Sander, M. (2011b). Sox9+ ductal cells are multipotent progenitors throughout development but do not produce new endocrine cells in the normal or injured adult pancreas. Development *138*, 653-665.

Kotz, J. (2012). PARP target practice. SciBX: Science-Business eXchange *5*.

Krakowski, M.L., Kritzik, M.R., Jones, E.M., Krahl, T., Lee, J., Arnush, M., Gu, D., and Sarvetnick, N. (1999). Pancreatic expression of keratinocyte growth factor leads to differentiation of islet hepatocytes and proliferation of duct cells. Am J Pathol *154*, 683-691.

Kraus, W.L. (2008). Transcriptional control by PARP-1: chromatin modulation, enhancer-binding, coregulation, and insulation. Current opinion in cell biology *20*, 294-302.

Kraus, W.L., and Hottiger, M.O. (2013). PARP-1 and gene regulation: progress and puzzles. Molecular aspects of medicine *34*, 1109-1123.

Kraus, W.L., and Lis, J.T. (2003). PARP goes transcription. Cell *113*, 677-683.

Krishnakumar, R., and Kraus, W.L. (2010a). PARP-1 regulates chromatin structure and transcription through a KDM5B-dependent pathway. Molecular cell *39*, 736-749.

Krishnakumar, R., and Kraus, W.L. (2010b). The PARP side of the nucleus: molecular actions, physiological outcomes, and clinical targets. Molecular cell *39*, 8-24.

Kunisada, Y., Tsubooka-Yamazoe, N., Shoji, M., and Hosoya, M. (2012). Small molecules induce efficient differentiation into insulin-producing cells from human induced pluripotent stem cells. Stem cell research *8*, 274-284.

Lee, C.S., Sund, N.J., Behr, R., Herrera, P.L., and Kaestner, K.H. (2005). Foxa2 is required for the differentiation of pancreatic α-cells. Developmental biology *278*, 484-495.

Lee, C.S., Sund, N.J., Vatamaniuk, M.Z., Matschinsky, F.M., Stoffers, D.A., and Kaestner, K.H. (2002). Foxa2 controls Pdx1 gene expression in pancreatic β-cells in vivo. Diabetes *51*, 2546-2551.

Lee, J., Sugiyama, T., Liu, Y., Wang, J., Gu, X., Lei, J., Markmann, J.F., Miyazaki, S., Miyazaki, J.-i., and Szot, G.L. (2013). Expansion and conversion of human pancreatic ductal cells into insulin-secreting endocrine cells. Elife *2*, e00940.

Lee, R.H., Seo, M.J., Reger, R.L., Spees, J.L., Pulin, A.A., Olson, S.D., and Prockop, D.J. (2006). Multipotent stromal cells from human marrow home to and promote repair of pancreatic islets and renal glomeruli in diabetic NOD/scid mice. Proceedings of the National Academy of Sciences *103*, 17438-17443.

Lehmann, M., Pirinen, E., Mirsaidi, A., Kunze, F.A., Richards, P.J., Auwerx, J., and Hottiger, M.O. (2015). ARTD1-induced poly-ADP-ribose formation enhances PPARγ ligand binding and co-factor exchange. Nucleic acids research *43*, 129-142.

Lenzen, S. (2008). The mechanisms of alloxan-and streptozotocin-induced diabetes. Diabetologia *51*, 216-226.

Levetan, C. (2010). Distinctions between islet neogenesis and beta-cell replication: implications for reversal of Type 1 and 2 diabetes. Journal of diabetes *2*, 76-84.

Li, F., Drel, V.R., Szabo, C., Stevens, M.J., and Obrosova, I.G. (2005a). Low-dose poly(ADP-ribose) polymerase inhibitor-containing combination therapies reverse early peripheral diabetic neuropathy. Diabetes *54*, 1514-1522.

Li, F., Szabo, C., Pacher, P., Southan, G.J., Abatan, O.I., Charniauskaya, T., Stevens, M.J., and Obrosova, I.G. (2004). Evaluation of orally active poly(ADP-ribose) polymerase inhibitor in streptozotocin-diabetic rat model of early peripheral neuropathy. Diabetologia *47*, 710-717.

Li, L., Seno, M., Yamada, H., and Kojima, I. (2001). Promotion of beta-cell regeneration by betacellulin in ninety percent-pancreatectomized rats. Endocrinology *142*, 5379-5385.

Li, L., Seno, M., Yamada, H., and Kojima, I. (2003). Betacellulin improves glucose metabolism by promoting conversion of intraislet precursor cells to β-cells in streptozotocin-treated mice. American Journal of Physiology-Endocrinology And Metabolism *285*, E577-E583.

Li, W.-C., Rukstalis, J.M., Nishimura, W., Tchipashvili, V., Habener, J.F., Sharma, A., and Bonner-Weir, S. (2010). Activation of pancreatic-duct-derived progenitor cells during pancreas regeneration in adult rats. J Cell Sci *123*, 2792-2802.

Li, Y., Cao, X., Li, L.-X., Brubaker, P.L., Edlund, H., and Drucker, D.J. (2005b). β-Cell Pdx1 expression is essential for the glucoregulatory, proliferative, and cytoprotective actions of glucagon-like peptide-1. Diabetes *54*, 482-491.

Lin, P., Chen, L., Yang, N., Sun, Y., and Xu, Y. (2009). Evaluation of stem cell differentiation in diabetic rats transplanted with bone marrow mesenchymal stem cells. Paper presented at: Transplantation proceedings (Elsevier).

Lina Sui, J.Q., Fei Yi and Guang-Hui Liu (2013). Stem Cell Therapy for Diabetes: A Call for Efficient Differentiation of Pancreatic Progenitors. J Regen Med *2*.

Lönn, P., van der Heide, L.P., Dahl, M., Hellman, U., Heldin, C.-H., and Moustakas, A. (2010). PARP-1 attenuates Smad-mediated transcription. Molecular cell *40*, 521-532.

Luo, X., and Kraus, W.L. (2012). On PAR with PARP: cellular stress signaling through poly (ADP-ribose) and PARP-1. Genes & development *26*, 417-432.

Mabley, J.G., and Soriano, F.G. (2005). Role of nitrosative stress and poly(ADP-ribose) polymerase activation in diabetic vascular dysfunction. Current vascular pharmacology *3*, 247-252.

Mak, J., Jablonski, C., Leonard, C., Dunn, J., Raharjo, E., Matyas, J., Biernaskie, J., and Krawetz, R. (2016). Intra-articular injection of synovial mesenchymal stem cells improves cartilage repair in a mouse injury model. Scientific Reports *6*.

Mamin, A., and Philippe, J. (2007). Activin A decreases glucagon and arx gene expression in alpha-cell lines. Molecular endocrinology *21*, 259-273.

Marshall, S. (2003). Oral hypoglycaemic agents. Practical Diabetes International *20*, S1-S4.

Mashima, H., Shibata, H., Mine, T., and Kojima, I. (1996). Formation of insulin-producing cells from pancreatic acinar AR42J cells by hepatocyte growth factor. Endocrinology *137*, 3969-3976.

Masutani, M., Suzuki, H., Kamada, N., Watanabe, M., Ueda, O., Nozaki, T., Jishage, K., Watanabe, T., Sugimoto, T., Nakagama, H., *et al.* (1999). Poly(ADP-ribose) polymerase gene disruption conferred mice resistant to streptozotocin-induced diabetes. Proceedings of the National Academy of Sciences of the United States of America *96*, 2301-2304.

Mato, E., Lucas, M., Petriz, J., Gomis, R., and Novials, A. (2009). Identification of a pancreatic stellate cell population with properties of progenitor cells: new role for stellate cells in the pancreas. Biochemical Journal *421*, 181-191.

Matsuoka, T.-a., Kaneto, H., Stein, R., Miyatsuka, T., Kawamori, D., Henderson, E., Kojima, I., Matsuhisa, M., Hori, M., and Yamasaki, Y. (2007). MafA regulates expression of genes important to islet β-cell function. Molecular endocrinology *21*, 2764-2774.

Menge, B.A., Tannapfel, A., Belyaev, O., Drescher, R., Müller, C., Uhl, W., Schmidt, W.E., and Meier, J.J. (2008). Partial pancreatectomy in adult humans does not provoke β-cell regeneration. Diabetes *57*, 142-149.

Messner, S., Altmeyer, M., Zhao, H., Pozivil, A., Roschitzki, B., Gehrig, P., Rutishauser, D., Huang, D., Caflisch, A., and Hottiger, M.O. (2010). PARP1 ADP-ribosylates lysine residues of the core histone tails. Nucleic acids research, gkq463.

Min, W., and Wang, Z.-Q. (2008). Poly (ADP-ribose) glycohydrolase (PARG) and its therapeutic potential. Frontiers in bioscience (Landmark edition) *14*, 1619-1626.

Minami, K., Okuno, M., Miyawaki, K., Okumachi, A., Ishizaki, K., Oyama, K., Kawaguchi, M., Ishizuka, N., Iwanaga, T., and Seino, S. (2005). Lineage tracing and characterization of insulin-secreting cells generated from adult pancreatic acinar cells. Proceedings of the National Academy of Sciences of the United States of America *102*, 15116-15121.

Minami, K., and Seino, S. (2008). Pancreatic acinar-to-beta cell transdifferentiation in vitro. Frontiers in bioscience : a journal and virtual library *13*, 5824-5837.

Minchenko, A.G., Stevens, M.J., White, L., Abatan, O.I., Komjati, K., Pacher, P., Szabo, C., and Obrosova, I.G. (2003). Diabetes-induced overexpression of endothelin-1 and endothelin receptors in the rat renal cortex is mediated via poly(ADP-ribose) polymerase activation. FASEB journal : official publication of the Federation of American Societies for Experimental Biology *17*, 1514-1516.

Morrison, H. (1937). Contributions to the Microscopic Anatomy of the Pancreas, by PAUL LANGERHANS (Berlin 1869). With an English Translation and an Introductory Essay. Bulletin of the History of Medicine *5*, 259.

Mu, D., Zhang, X.-L., Xie, J., Yuan, H.-H., Wang, K., Huang, W., Li, G.-N., Lu, J.-R., Mao, L.-J., and Wang, L. (2016). Intracoronary Transplantation of Mesenchymal Stem Cells with Overexpressed Integrin-Linked Kinase Improves Cardiac Function in Porcine Myocardial Infarction. Scientific reports *6*.

Murtaugh, L.C. (2007). Pancreas and beta-cell development: from the actual to the possible. Development *134*, 427-438.

Nagaya, M., Katsuta, H., Kaneto, H., Bonner-Weir, S., and Weir, G.C. (2009). Adult mouse intrahepatic biliary epithelial cells induced in vitro to become insulin-producing cells. Journal of Endocrinology *201*, 37-47.

Nakagawa, Y., Muneta, T., Otabe, K., Ozeki, N., Mizuno, M., Udo, M., Saito, R., Yanagisawa, K., Ichinose, S., and Koga, H. (2016). Cartilage Derived from Bone Marrow Mesenchymal Stem Cells Expresses Lubricin In Vitro and In Vivo. PloS one *11*, e0148777.

Naor, D., and Assayag-Asherie, N. (2013). CD44 expression on insulin-secreting {beta} cells enhances the development of type 1 diabetes (P5109). The Journal of Immunology *190*, 129.119.

Naya, F.J., Huang, H.-P., Qiu, Y., Mutoh, H., DeMayo, F.J., Leiter, A.B., and Tsai, M.-J. (1997). Diabetes, defective pancreatic morphogenesis, and abnormal enteroendocrine differentiation in BETA2/neuroD-deficient mice. Genes & development *11*, 2323-2334.

Naya, F.J., Stellrecht, C., and Tsai, M.-J. (1995). Tissue-specific regulation of the insulin gene by a novel basic helix-loop-helix transcription factor. Genes & development *9*, 1009-1019.

Negi, G., Kumar, A., Kaundal, R.K., Gulati, A., and Sharma, S.S. (2010a). Functional and biochemical evidence indicating beneficial effect of Melatonin and Nicotinamide alone and in combination in experimental diabetic neuropathy. Neuropharmacology *58*, 585-592.

Negi, G., Kumar, A., and Sharma, S.S. (2010b). Concurrent targeting of nitrosative stress-PARP pathway corrects functional, behavioral and biochemical deficits in experimental diabetic neuropathy. Biochemical and biophysical research communications *391*, 102-106.

Nishimura, W., Kondo, T., Salameh, T., El Khattabi, I., Dodge, R., Bonner-Weir, S., and Sharma, A. (2006). A switch from MafB to MafA expression accompanies differentiation to pancreatic β-cells. Developmental biology *293*, 526-539.

Noguchi, H. (2010). Pancreatic stem/progenitor cells for the treatment of diabetes. Rev Diabet Stud *7*, 105-111.

Nostro, M.C., Sarangi, F., Ogawa, S., Holtzinger, A., Corneo, B., Li, X., Micallef, S.J., Park, I.-H., Basford, C., and Wheeler, M.B. (2011). Stage-specific signaling through TGFβ family members and WNT regulates patterning and pancreatic specification of human pluripotent stem cells. Development *138*, 861-871.

Obrosova, I.G., Li, F., Abatan, O.I., Forsell, M.A., Komjati, K., Pacher, P., Szabo, C., and Stevens, M.J. (2004a). Role of poly(ADP-ribose) polymerase activation in diabetic neuropathy. Diabetes *53*, 711-720.

Obrosova, I.G., Minchenko, A.G., Frank, R.N., Seigel, G.M., Zsengeller, Z., Pacher, P., Stevens, M.J., and Szabo, C. (2004b). Poly(ADP-ribose) polymerase inhibitors counteract diabetes- and hypoxia-induced retinal vascular endothelial growth factor overexpression. International journal of molecular medicine *14*, 55-64.

Ogino, H., Nozaki, T., Gunji, A., Maeda, M., Suzuki, H., Ohta, T., Murakami, Y., Nakagama, H., Sugimura, T., and Masutani, M. (2007). Loss of Parp-1 affects gene expression profile in a genome-wide manner in ES cells and liver cells. BMC genomics *8*, 41.

Ohlsson, H., Karlsson, K., and Edlund, T. (1993). IPF1, a homeodomain-containing transactivator of the insulin gene. The EMBO journal *12*, 4251.

Oka, S., Kato, J., and Moss, J. (2006). Identification and characterization of a mammalian 39-kDa poly (ADP-ribose) glycohydrolase. Journal of Biological Chemistry *281*, 705-713.

Oliver, F.J., Menissier-de Murcia, J., Nacci, C., Decker, P., Andriantsitohaina, R., Muller, S., de la Rubia, G., Stoclet, J.C., and de Murcia, G. (1999). Resistance to endotoxic shock as a consequence of defective NF-kappaB activation in poly (ADP-ribose) polymerase-1 deficient mice. The EMBO journal *18*, 4446-4454.

Otonkoski, T., Beattie, G.M., Mally, M.I., Ricordi, C., and Hayek, A. (1993). Nicotinamide is a potent inducer of endocrine differentiation in cultured human fetal pancreatic cells. J Clin Invest *92*, 1459-1466.

Pacher, P., Liaudet, L., Soriano, F.G., Mabley, J.G., Szabo, E., and Szabo, C. (2002). The role of poly(ADP-ribose) polymerase activation in the development of myocardial and endothelial dysfunction in diabetes. Diabetes *51*, 514-521.

Pan, F.C., Bankaitis, E.D., Boyer, D., Xu, X., Van de Casteele, M., Magnuson, M.A., Heimberg, H., and Wright, C.V. (2013). Spatiotemporal patterns of multipotentiality in Ptf1a-expressing cells during pancreas organogenesis and injury-induced facultative restoration. Development *140*, 751-764.

Pang, K., Mukonoweshuro, C., and Wong, G.G. (1994). Beta cells arise from glucose transporter type 2 (Glut2)-expressing epithelial cells of the developing rat pancreas. Proceedings of the National Academy of Sciences of the United States of America *91*, 9559-9563.

Pangas, S.A., and Woodruff, T.K. (2000). Activin signal transduction pathways. Trends in Endocrinology & Metabolism *11*, 309-314.

Pardo, M., Lang, B., Yu, L., Prosser, H., Bradley, A., Babu, M.M., and Choudhary, J. (2010). An expanded Oct4 interaction network: implications for stem cell biology, development, and disease. Cell stem cell *6*, 382-395.

Park, K.-S., Kim, Y.-S., Kim, J.-H., Choi, B., Kim, S.-H., Tan, A.H.-K., Lee, M.-S., Lee, M.-K., Kwon, C.-H., and Joh, J.-W. (2010). Trophic molecules derived from human mesenchymal stem cells enhance survival, function, and angiogenesis of isolated islets after transplantation. Transplantation *89*, 509-517.

Pasquali, L., Fan, Y., Trucco, M., and Ringquist, S. (2006). Rehabilitation of adaptive immunity and regeneration of beta cells. Trends in biotechnology *24*, 516-522.

Pasquali, L., Giannoukakis, N., and Trucco, M. (2008). Induction of immune tolerance to facilitate β cell regeneration in type 1 diabetes. Advanced drug delivery reviews *60*, 106-113.

Pavri, R., Lewis, B., Kim, T.-K., Dilworth, F.J., Erdjument-Bromage, H., Tempst, P., De Murcia, G., Evans, R., Chambon, P., and Reinberg, D. (2005). PARP-1 determines specificity in a retinoid signaling pathway via direct modulation of mediator. Molecular cell *18*, 83-96.

Peters, K., Panienka, R., Li, J., Klöppel, G., and Wang, R. (2005). Expression of stem cell markers and transcription factors during the remodeling of the rat pancreas after duct ligation. Virchows Archiv *446*, 56-63.

Petesch, S.J., and Lis, J.T. (2012). Activator-induced spread of poly (ADP-ribose) polymerase promotes nucleosome loss at Hsp70. Molecular cell *45*, 64-74.

Petrou, P., Gothelf, Y., Argov, Z., Gotkine, M., Levy, Y.S., Kassis, I., Vaknin-Dembinsky, A., Ben-Hur, T., Offen, D., and Abramsky, O. (2016). Safety and Clinical Effects of Mesenchymal Stem Cells Secreting Neurotrophic Factor Transplantation in Patients With Amyotrophic Lateral Sclerosis: Results of Phase 1/2 and 2a Clinical Trials. JAMA neurology, 1-8.

Pieper, A.A., Brat, D.J., Krug, D.K., Watkins, C.C., Gupta, A., Blackshaw, S., Verma, A., Wang, Z.Q., and Snyder, S.H. (1999). Poly(ADP-ribose) polymerase-deficient mice are protected from streptozotocin-induced diabetes. Proceedings of the National Academy of Sciences of the United States of America *96*, 3059-3064.

Pileggi, A. (2012). Mesenchymal stem cells for the treatment of diabetes. Diabetes *61*, 1355-1356.

Pittenger, M.F., Mackay, A.M., Beck, S.C., Jaiswal, R.K., Douglas, R., Mosca, J.D., Moorman, M.A., Simonetti, D.W., Craig, S., and Marshak, D.R. (1999). Multilineage potential of adult human mesenchymal stem cells. science *284*, 143-147.

Poirier, G.G., De Murcia, G., Jongstra-Bilen, J., Niedergang, C., and Mandel, P. (1982). Poly (ADP-ribosyl) ation of polynucleosomes causes relaxation of chromatin structure. Proceedings of the National Academy of Sciences *79*, 3423-3427.

Qian, H., Xu, J., Lalioti, M.D., Gulle, K., and Sakkas, D. (2010). Oocyte numbers in the mouse increase after treatment with 5-aminoisoquinolinone: a potent inhibitor of poly(ADP-ribosyl)ation. Biology of reproduction *82*, 1000-1007.

Rabinowe, S.L., and Eisenbarth, G.S. (1984). Type I diabetes mellitus: a chronic autoimmune disease? Pediatric Clinics of North America *31*, 531-543.

Rackham, C.L., Dhadda, P.K., Chagastelles, P.C., Simpson, S.J.S., Dattani, A.A., Bowe, J.E., Jones, P.M., and King, A.J.F. (2013). Pre-culturing islets with mesenchymal stromal cells using a direct contact configuration is beneficial for transplantation outcome in diabetic mice. Cytotherapy *15*, 449-459.

Rafaeloff, R., Pittenger, G.L., Barlow, S.W., Qin, X.F., Yan, B., Rosenberg, L., Duguid, W.P., and Vinik, A.I. (1997). Cloning and sequencing of the pancreatic islet neogenesis associated protein (INGAP) gene and its expression in islet neogenesis in hamsters. The Journal of clinical investigation *99*, 2100.

Rajawat, J., Mir, H., and Begum, R. (2011). Differential role of poly (ADP-ribose) polymerase in D. discoideum growth and development. BMC developmental biology *11*, 14.

Rakieten, N., Gordon, B.S., Beaty, A., Cooney, D.A., Davis, R.D., and Schein, P.S. (1971). Pancreatic islet cell tumors produced by the combined action of streptozotocin and nicotinamide. Proceedings of the Society for Experimental Biology and Medicine Society for Experimental Biology and Medicine *137*, 280-283.

Ramiya, V.K., Maraist, M., Arfors, K.E., Schatz, D.A., Peck, A.B., and Cornelius, J.G. (2000). Reversal of insulin-dependent diabetes using islets generated in vitro from pancreatic stem cells. Nature medicine *6*, 278-282.

Reale, A., De Matteis, G., Galleazzi, G., Zampieri, M., and Caiafa, P. (2005). Modulation of DNMT1 activity by ADP-ribose polymers. Oncogene *24*, 13-19.

Rekittke, N.E., Ang, M., Rawat, D., Khatri, R., and Linn, T. (2016). Regenerative Therapy of Type 1 Diabetes Mellitus: From Pancreatic Islet Transplantation to Mesenchymal Stem Cells. Stem Cells International *2016*.

Ringel, J., Jesnowski, R., Schmidt, C., Ringel, J., Köhler, H.J., Rychly, J., Batra, S.K., and Löhr, M. (2001). CD44 in normal human pancreas and pancreatic carcinoma cell lines. Teratogenesis Carcinogenesis and Mutagenesis *21*, 97-106.

Robertson, R.P. (2004). Islet transplantation as a treatment for diabetes—a work in progress. New England Journal of Medicine *350*, 694-705.

Robertson, R.P., Davis, C., Larsen, J., Stratta, R., and Sutherland, D.E. (2000). Pancreas and islet transplantation for patients with diabetes. DIABETES CARE-ALEXANDRIA VA-*23*, 112-116.

Rosenberg, L., Brown, R.A., and Duguid, W.P. (1983). A new approach to the induction of duct epithelial hyperplasia and nesidioblastosis by cellophane wrapping of the hamster pancreas. Journal of Surgical Research *35*, 63-72.

Rosenberg, L., and Vinik, A. (1989). Induction of endocrine cell differentiation: a new approach to management of diabetes. The Journal of laboratory and clinical medicine *114*, 75-83.

Rother, K.I., and Harlan, D.M. (2004). Challenges facing islet transplantation for the treatment of type 1 diabetes mellitus. The Journal of clinical investigation *114*, 877-883.

Rovira, M., Scott, S.-G., Liss, A.S., Jensen, J., Thayer, S.P., and Leach, S.D. (2010). Isolation and characterization of centroacinar/terminal ductal progenitor cells in adult mouse pancreas. Proceedings of the National Academy of Sciences *107*, 75-80.

Rowan, J.A., Hague, W.M., Gao, W., Battin, M.R., and Moore, M.P. (2008). Metformin versus insulin for the treatment of gestational diabetes. New England Journal of Medicine *358*, 2003-2015.

Rukstalis, J.M., and Habener, J.F. (2009). Neurogenin3: a master regulator of pancreatic islet differentiation and regeneration. Islets *1*, 177-184.

Sakamaki, J.-i., Daitoku, H., Yoshimochi, K., Miwa, M., and Fukamizu, A. (2009). Regulation of FOXO1-mediated transcription and cell proliferation by PARP-1. Biochemical and biophysical research communications *382*, 497-502.

Sakata, N., Goto, M., Yoshimatsu, G., Egawa, S., and Unno, M. (2011). Utility of co-transplanting mesenchymal stem cells in islet transplantation. World journal of gastroenterology *17*, 5150-5155.

Sala, A., La Rocca, G., Burgio, G., Kotova, E., Di Gesù, D., Collesano, M., Ingrassia, A.M., Tulin, A.V., and Davide, F. (2008). The nucleosome-remodeling ATPase ISWI is regulated by poly-ADP-ribosylation. PLoS Biol *6*, e252.

Samson, S.L., and Chan, L. (2006). Gene therapy for diabetes: reinventing the islet. Trends in Endocrinology & Metabolism *17*, 92-100.

Sander, M., Sussel, L., Conners, J., Scheel, D., Kalamaras, J., Cruz, F.D., Schwitzgebel, V., Hayes-Jordan, A., and German, M. (2000). Homeobox gene Nkx6. 1 lies downstream of Nkx2. 2 in the major pathway of beta-cell formation in the pancreas. Development *127*, 5533-5540.

Sasada, R., Ono, Y., Taniyama, Y., Shing, Y., Folkman, J., and Igarashi, K. (1993). Cloning and expression of cDNA encoding human betacellulin, a new member of the EGF family. Biochem Biophys Res Commun *190*, 1173-1179.

Schein, P.S., Cooney, D.A., and Vernon, M.L. (1967). The use of nicotinamide to modify the toxicity of streptozotocin diabetes without loss of antitumor activity. Cancer research *27*, 2324-2332.

Schnedl, W.J., Ferber, S., Johnson, J.H., and Newgard, C.B. (1994). STZ transport and cytotoxicity: specific enhancement in GLUT2-expressing cells. Diabetes *43*, 1326-1333.

Schreiber, V., Amé, J.-C., Dollé, P., Schultz, I., Rinaldi, B., Fraulob, V., Ménissier-de Murcia, J., and de Murcia, G. (2002). Poly (ADP-ribose) polymerase-2 (PARP-2) is required for efficient base excision DNA repair in association with PARP-1 and XRCC1. Journal of Biological Chemistry *277*, 23028-23036.

Schreiber, V., Dantzer, F., Ame, J.-C., and De Murcia, G. (2006). Poly (ADP-ribose): novel functions for an old molecule. Nature reviews Molecular cell biology *7*, 517-528.

Scobie, I.N. (2006). Atlas of diabetes mellitus (CRC Press).

Seaberg, R.M., Smukler, S.R., Kieffer, T.J., Enikolopov, G., Asghar, Z., Wheeler, M.B., Korbutt, G., and van der Kooy, D. (2004). Clonal identification of multipotent precursors from adult mouse pancreas that generate neural and pancreatic lineages. Nature biotechnology *22*, 1115-1124.

Segev, H., Fishman, B., Schulman, R., and Itskovitz-Eldor, J. (2012). The expression of the class 1 glucose transporter isoforms in human embryonic stem cells, and the potential use of GLUT2 as a marker for pancreatic progenitor enrichment. Stem cells and development *21*, 1653-1661.

Segev, H., Fishman, B., Ziskind, A., Shulman, M., and Itskovitz-Eldor, J. (2004a). Differentiation of human embryonic stem cells into insulin-producing clusters. Stem Cells *22*, 265-274.

Segev, H., Fishman, B., Ziskind, A., Shulman, M., and Itskovitz-Eldor, J. (2004b). Differentiation of human embryonic stem cells into insulin-producing clusters. Stem cells *22*, 265-274.

Serafimidis, I., Rakatzi, I., Episkopou, V., Gouti, M., and Gavalas, A. (2008). Novel Effectors of Directed and Ngn3-Mediated Differentiation of Mouse Embryonic Stem Cells into Endocrine Pancreas Progenitors. Stem cells *26*, 3-16.

Seymour, P.A. (2014). Sox9: a master regulator of the pancreatic program. The review of diabetic studies: RDS *11*, 51.

Shah, R.G., Ghodgaonkar, M.M., Affar, E.B., and Shah, G.M. (2005). DNA vector-based RNAi approach for stable depletion of poly (ADP-ribose) polymerase-1. Biochemical and biophysical research communications *331*, 167-174.

Shall, S., and de Murcia, G. (2000). Poly(ADP-ribose) polymerase-1: what have we learned from the deficient mouse model? Mutation research *460*, 1-15.

Shapiro, A.J., Lakey, J.R., Ryan, E.A., Korbutt, G.S., Toth, E., Warnock, G.L., Kneteman, N.M., and Rajotte, R.V. (2000). Islet transplantation in seven patients with type 1 diabetes mellitus using a glucocorticoid-free immunosuppressive regimen. New England Journal of Medicine *343*, 230-238.

Shapiro, A.M., Pokrywczynska, M., and Ricordi, C. (2017). Clinical pancreatic islet transplantation. Nature reviews Endocrinology *13*, 268-277.

Sharma, S.S., Kumar, A., Arora, M., and Kaundal, R.K. (2009). Neuroprotective potential of combination of resveratrol and 4-amino 1,8 naphthalimide in experimental diabetic neuropathy: focus on functional, sensorimotor and biochemical changes. Free radical research *43*, 400-408.

Sharma, S.S., Kumar, A., and Kaundal, R.K. (2008). Protective effects of 4-amino1,8-napthalimide, a poly (ADP-ribose) polymerase inhibitor in experimental diabetic neuropathy. Life sciences *82*, 570-576.

Shevalye, H., Maksimchyk, Y., Watcho, P., and Obrosova, I.G. (2010). Poly(ADP-ribose) polymerase-1 (PARP-1) gene deficiency alleviates diabetic kidney disease. Biochimica et biophysica acta *1802*, 1020-1027.

Shi, Y., and Massagué, J. (2003). Mechanisms of TGF-β signaling from cell membrane to the nucleus. Cell *113*, 685-700.

Shih, H.P., Wang, A., and Sander, M. (2013). Pancreas organogenesis: from lineage determination to morphogenesis. Annual review of cell and developmental biology *29*, 81-105.

Sidney, L.E., Branch, M.J., Dunphy, S.E., Dua, H.S., and Hopkinson, A. (2014). Concise review: evidence for CD34 as a common marker for diverse progenitors. Stem Cells *32*, 1380-1389.

Simbulan-Rosenthal, C.M., Rosenthal, D.S., Luo, R., Samara, R., Espinoza, L.A., Hassa, P.O., Hottiger, M.O., and Smulson, M.E. (2003). PARP-1 binds E2F-1 independently of its DNA binding and catalytic domains, and acts as a novel coactivator of E2F-1-mediated transcription during re-entry of quiescent cells into S phase. Oncogene *22*, 8460-8471.

Slack, J. (1995). Developmental biology of the pancreas. Development *121*, 1569-1580.

Slattery, E., Dignam, J.D., Matsui, T., and Roeder, R.G. (1983). Purification and analysis of a factor which suppresses nick-induced transcription by RNA polymerase II and its identity with poly(ADP-ribose) polymerase. The Journal of biological chemistry *258*, 5955-5959.

Smukler, S.R., Arntfield, M.E., Razavi, R., Bikopoulos, G., Karpowicz, P., Seaberg, R., Dai, F., Lee, S., Ahrens, R., and Fraser, P.E. (2011). The adult mouse and human pancreas contain rare multipotent stem cells that express insulin. Cell stem cell *8*, 281-293.

Solar, M., Cardalda, C., Houbracken, I., Martín, M., Maestro, M.A., De Medts, N., Xu, X., Grau, V., Heimberg, H., and Bouwens, L. (2009). Pancreatic exocrine duct cells give rise to insulin-producing β cells during embryogenesis but not after birth. Developmental cell *17*, 849-860.

Soria, B., Skoudy, A., and Martin, F. (2001). From stem cells to beta cells: new strategies in cell therapy of diabetes mellitus. Diabetologia *44*, 407-415.

Soriano, F.G., Virag, L., and Szabo, C. (2001). Diabetic endothelial dysfunction: role of reactive oxygen and nitrogen species production and poly(ADP-ribose) polymerase activation. Journal of molecular medicine *79*, 437-448.

Sosa-Pineda, B. (2004). The gene Pax4 is an essential regulator of pancreatic beta-cell development. Molecules and cells *18*, 289-294.

Sosa-Pineda, B., Chowdhury, K., Torres, M., Oliver, G., and Gruss, P. (1997). The Pax4 gene is essential for differentiation of insulin-producing beta cells in the mammalian pancreas. Nature *386*, 399.

Srivastava, A., Bhatt, N.M., Patel, T.P., Dadheech, N., Singh, A., and Gupta, S. (2016). Anti-apoptotic and cytoprotective effect of Enicostemma littorale against oxidative stress in Islets of Langerhans. Pharmaceutical biology *54*, 2061-2072.

St-Laurent, J.-F., Gagnon, S.N., Dequen, F., Hardy, I., and Desnoyers, S. (2007). Altered DNA damage response in Caenorhabditis elegans with impaired poly (ADP-ribose) glycohydrolases genes expression. DNA repair *6*, 329-343.

Staels, W., De Groef, S., Heremans, Y., Coppens, V., Van Gassen, N., Leuckx, G., Van de Casteele, M., Van Riet, I., Luttun, A., Heimberg, H., *et al.* (2016). Accessory cells for beta-cell transplantation. Diabetes, obesity & metabolism *18*, 115-124.

Steil, G.M., Rebrin, K., Darwin, C., Hariri, F., and Saad, M.F. (2006). Feasibility of automating insulin delivery for the treatment of type 1 diabetes. Diabetes *55*, 3344-3350.

Sugiyama, T., and Kim, S.K. (2008). Fluorescence-activated cell sorting purification of pancreatic progenitor cells. Diabetes, obesity & metabolism *10 Suppl 4*, 179-185.

Sun, B., Roh, K.-H., Lee, S.-R., Lee, Y.-S., and Kang, K.-S. (2007a). Induction of human umbilical cord blood-derived stem cells with embryonic stem cell phenotypes into insulin producing islet-like structure. Biochemical and biophysical research communications *354*, 919-923.

Sun, Y., Chen, L., Hou, X.G., Hou, W.K., Dong, J.J., Sun, L., Tang, K.X., Wang, B., Song, J., Li, H., *et al.* (2007b). Differentiation of bone marrow-derived mesenchymal stem cells from diabetic patients into insulin-producing cells in vitro. Chin Med J (Engl) *120*, 771-776.

Sussel, L., Kalamaras, J., Hartigan-O'Connor, D., Meneses, J., Pedersen, R., Rubenstein, J., and German, M. (1998). Mice lacking the homeodomain transcription factor Nkx2. 2 have diabetes due to arrested differentiation of pancreatic beta cells. Development *125*, 2213-2221.

Sutherland, D.E., Gruessner, R.W., Dunn, D.L., Matas, A.J., Humar, A., Kandaswamy, R., Mauer, S.M., Kennedy, W.R., Goetz, F.C., and Robertson, R.P. (2001a). Lessons learned from more than 1,000 pancreas transplants at a single institution. Annals of surgery *233*, 463-501.

Sutherland, D.E., Gruessner, R.W., and Gruessner, A.C. (2001b). Pancreas transplantation for treatment of diabetes mellitus. World journal of surgery *25*, 487-496.

Suzuki, A., Nakauchi, H., and Taniguchi, H. (2004). Prospective isolation of multipotent pancreatic progenitors using flow-cytometric cell sorting. Diabetes *53*, 2143-2152.

Szabo, G., Bahrle, S., Stumpf, N., Sonnenberg, K., Szabo, E.E., Pacher, P., Csont, T., Schulz, R., Dengler, T.J., Liaudet, L., *et al.* (2002). Poly(ADP-Ribose) polymerase inhibition reduces reperfusion injury after heart transplantation. Circulation research *90*, 100-106.

Szkudelski, T. (2001). The mechanism of alloxan and streptozotocin action in B cells of the rat pancreas. Physiological research *50*, 537-546.

Takahashi, K., and Yamanaka, S. (2006). Induction of pluripotent stem cells from mouse embryonic and adult fibroblast cultures by defined factors. cell *126*, 663-676.

Takasawa, S., Akiyama, T., Nata, K., Kuroki, M., Tohgo, A., Noguchi, N., Kobayashi, S., Kato, I., Katada, T., and Okamoto, H. (1998). Cyclic ADP-ribose and inositol 1,4,5-trisphosphate as alternate second messengers for intracellular Ca2+ mobilization in normal and diabetic beta-cells. The Journal of biological chemistry *273*, 2497-2500.

Takasawa, S., Nata, K., Yonekura, H., and Okamoto, H. (1993). Cyclic ADP-ribose in insulin secretion from pancreatic beta cells. Science *259*, 370-373.

Takasawa, S., and Okamoto, H. (2002). Pancreatic beta-cell death, regeneration and insulin secretion: roles of poly(ADP-ribose) polymerase and cyclic ADP-ribose. International journal of experimental diabetes research *3*, 79-96.

Tamaki, M., Fujitani, Y., Uchida, T., Hirose, T., Kawamori, R., and Watada, H. Combination treatment of db/db mice with exendin-4 and gastrin preserves beta-cell mass by stimulating beta-cell growth and differentiation. J Diabetes Investig *1*, 172-183.

Tanaka, E.M., and Reddien, P.W. (2011). The cellular basis for animal regeneration. Developmental cell *21*, 172-185.

Tang, D.-Q., Cao, L.-Z., Burkhardt, B.R., Xia, C.-Q., Litherland, S.A., Atkinson, M.A., and Yang, L.-J. (2004a). In vivo and in vitro characterization of insulin-producing cells obtained from murine bone marrow. Diabetes *53*, 1721-1732.

Tang, D.Q., Cao, L.Z., Burkhardt, B.R., Xia, C.Q., Litherland, S.A., Atkinson, M.A., and Yang, L.J. (2004b). In vivo and in vitro characterization of insulin-producing cells obtained from murine bone marrow. Diabetes *53*, 1721-1732.

Thorel, F., Népote, V., Avril, I., Kohno, K., Desgraz, R., Chera, S., and Herrera, P.L. (2010). Conversion of adult pancreatic α-cells to β-cells after extreme β-cell loss. Nature *464*, 1149-1154.

Timper, K., Seboek, D., Eberhardt, M., Linscheid, P., Christ-Crain, M., Keller, U., Müller, B., and Zulewski, H. (2006). Human adipose tissue-derived mesenchymal stem cells differentiate into insulin, somatostatin, and glucagon expressing cells. Biochemical and biophysical research communications *341*, 1135-1140.

Tong, W.-M., Hande, M.P., Lansdorp, P.M., and Wang, Z.-Q. (2001). DNA strand break-sensing molecule poly (ADP-Ribose) polymerase cooperates with p53 in telomere function, chromosome stability, and tumor suppression. Molecular and cellular biology *21*, 4046-4054.

Totsuka, Y., Tabuchi, M., Kojima, I., Shibai, H., and Ogata, E. (1988). A novel action of activin A: stimulation of insulin secretion in rat pancreatic islets. Biochemical and biophysical research communications *156*, 335-339.

Trounson, A. (2002). Human embryonic stem cells: mother of all cell and tissue types. Reproductive BioMedicine Online *4*, 58-63.

Tulin, A., and Spradling, A. (2003). Chromatin loosening by poly (ADP)-ribose polymerase (PARP) at Drosophila puff loci. Science *299*, 560-562.

Tuomilehto, J., Lindström, J., Eriksson, J.G., Valle, T.T., Hämäläinen, H., Ilanne-Parikka, P., Keinänen-Kiukaanniemi, S., Laakso, M., Louheranta, A., and Rastas, M. (2001). Prevention of type 2 diabetes mellitus by changes in lifestyle among subjects with impaired glucose tolerance. New England Journal of Medicine *344*, 1343-1350.

Uchigata, Y., Yamamoto, H., Kawamura, A., and Okamoto, H. (1982). Protection by superoxide dismutase, catalase, and poly(ADP-ribose) synthetase inhibitors against alloxan- and streptozotocin-induced islet DNA strand breaks and against the inhibition of proinsulin synthesis. The Journal of biological chemistry *257*, 6084-6088.

Uzan, B., Figeac, F., Portha, B., and Movassat, J. (2009). Mechanisms of KGF mediated signaling in pancreatic duct cell proliferation and differentiation. PLoS One *4*, e4734.

van der Meulen, T., and Huising, M.O. (2015). Role of transcription factors in the transdifferentiation of pancreatic islet cells. Journal of molecular endocrinology *54*, R103-R117.

Van, P.P. (2011). Stem Cell Therapy for Islet Regeneration. In Stem Cells in Clinic and Research (InTech).

Venkatesan, V., Gopurappilly, R., Goteti, S.K., Dorisetty, R.K., and Bhonde, R.R. (2011). Pancreatic progenitors: The shortest route to restore islet cell mass. Islets *3*, 295-301.

Wada, M., Shintani, Y., Kosaka, M., Sano, T., Hizawa, K., and Saito, S. (1996). Immunohistochemical localization of activin A and follistatin in human tissues. Endocrine journal *43*, 375-385.

Wahlberg, G., Carlson, L.A., Wasserman, J., and Ljungqvist, A. (1985). Protective effect of nicotinamide against nephropathy in diabetic rats. Diabetes research *2*, 307-312.

Wang, R., Klöppel, G., and Bouwens, L. (1995). Duct-to islet-cell differentiation and islet growth in the pancreas of duct-ligated adult rats. Diabetologia *38*, 1405-1411.

Wang, T.C., Bonner-Weir, S., Oates, P.S., Chulak, M., Simon, B., Merlino, G.T., Schmidt, E.V., and Brand, S.J. (1993). Pancreatic gastrin stimulates islet differentiation of transforming growth factor alpha-induced ductular precursor cells. J Clin Invest *92*, 1349-1356.

Watabe, T., and Miyazono, K. (2009). Roles of TGF-β family signaling in stem cell renewal and differentiation. Cell research *19*, 103-115.

Watanabe, T., Yonemura, Y., Yonekura, H., Suzuki, Y., Miyashita, H., Sugiyama, K., Moriizumi, S., Unno, M., Tanaka, O., Kondo, H., *et al.* (1994). Pancreatic beta-cell replication and amelioration of surgical diabetes by Reg protein. Proceedings of the National Academy of Sciences of the United States of America *91*, 3589-3592.

Weinstein, D.C., i Altaba, A.R., Chen, W.S., Hoodless, P., Prezioso, V.R., Jessell, T.M., and Darnell, J.E. (1994). The winged-helix transcription factor HNF-3β is required for notochord development in the mouse embryo. Cell *78*, 575-588.

Wen, Y., Chen, B., and Ildstad, S.T. (2011). Stem cell-based strategies for the treatment of type 1 diabetes mellitus. Expert opinion on biological therapy *11*, 41-53.

Wilcox, N.S., Rui, J., Hebrok, M., and Herold, K.C. (2016). Life and death of β cells in Type 1 diabetes: A comprehensive review. Journal of Autoimmunity.

Wong, R.S. (2011). Extrinsic factors involved in the differentiation of stem cells into insulin-producing cells: an overview. Experimental diabetes research *2011*, 406182.

Wrighton, K.H., Lin, X., and Feng, X.-H. (2009). Phospho-control of TGF-β superfamily signaling. Cell research *19*, 8-20.

Wu, X.H., Liu, C.P., Xu, K.F., Mao, X.D., Zhu, J., Jiang, J.J., Cui, D., Zhang, M., Xu, Y., and Liu, C. (2007). Reversal of hyperglycemia in diabetic rats by portal vein transplantation of islet-like cells generated from bone marrow mesenchymal stem cells. World journal of gastroenterology *13*, 3342-3349.

Xia, M., and Laychock, S.G. (1993). Insulin secretion, myo-inositol transport, and Na(+)-K(+)-ATPase in glucose-desensitized rat islets. Diabetes *42*, 1392-1400.

Xiao, C.Y., Chen, M., Zsengeller, Z., and Szabo, C. (2004). Poly(ADP-ribose) polymerase contributes to the development of myocardial infarction in diabetic rats and regulates the nuclear translocation of apoptosis-inducing factor. The Journal of pharmacology and experimental therapeutics *310*, 498-504.

Xin, Y., Jiang, X., Wang, Y., Su, X., Sun, M., Zhang, L., Tan, Y., Wintergerst, K.A., Li, Y., and Li, Y. (2016). Insulin-Producing Cells Differentiated from Human Bone Marrow

Mesenchymal Stem Cells In Vitro Ameliorate Streptozotocin-Induced Diabetic Hyperglycemia. PloS one *11*.

Xu, G., Stoffers, D.A., Habener, J.F., and Bonner-Weir, S. (1999). Exendin-4 stimulates both beta-cell replication and neogenesis, resulting in increased beta-cell mass and improved glucose tolerance in diabetic rats. Diabetes *48*, 2270-2276.

Xu, X., D'Hoker, J., Stange, G., Bonne, S., De Leu, N., Xiao, X., Van De Casteele, M., Mellitzer, G., Ling, Z., and Pipeleers, D. (2008). β cells can be generated from endogenous progenitors in injured adult mouse pancreas. Cell *132*, 197-207.

Yagi, H., Soto-Gutierrez, A., Parekkadan, B., Kitagawa, Y., Tompkins, R.G., Kobayashi, N., and Yarmush, M.L. (2010). Mesenchymal stem cells: mechanisms of immunomodulation and homing. Cell transplantation *19*, 667.

Yamagami, T., Miwa, A., Takasawa, S., Yamamoto, H., and Okamoto, H. (1985). Induction of rat pancreatic B-cell tumors by the combined administration of streptozotocin or alloxan and poly(adenosine diphosphate ribose) synthetase inhibitors. Cancer research *45*, 1845-1849.

Yamamoto, H., and Okamoto, H. (1980). Protection by picolinamide, a novel inhibitor of poly (ADP-ribose) synthetase, against both streptozotocin-induced depression of proinsulin synthesis and reduction of NAD content in pancreatic islets. Biochemical and biophysical research communications *95*, 474-481.

Yang, L., Huang, K., Li, X., Du, M., Kang, X., Luo, X., Gao, L., Wang, C., Zhang, Y., and Zhang, C. (2013). Identification of poly (ADP-ribose) polymerase-1 as a cell cycle regulator through modulating Sp1 mediated transcription in human hepatoma cells. PloS one *8*, e82872.

Yasuda, H., Inoue, K., Shibata, H., Takeuchi, T., Eto, Y., Hasegawa, Y., Sekine, N., Totsuka, Y., Mine, T., and Ogata, E. (1993). Existence of activin-A in A-and D-cells of rat pancreatic islet. Endocrinology *133*, 624-630.

Ye, D.Z., Tai, M.H., Linning, K.D., Szabo, C., and Olson, L.K. (2006). MafA expression and insulin promoter activity are induced by nicotinamide and related compounds in INS-1 pancreatic beta-cells. Diabetes *55*, 742-750.

Yi, F., and Liu, G.-H. (2013). Stem Cell Therapy for Diabetes: A Call for Efficient Differentiation of Pancreatic Progenitors. Journal of Regenerative Medicine.

Yki-Järvinen, H. (2001). Combination therapies with insulin in type 2 diabetes. Diabetes care *24*, 758-767.

Yonemura, Y., Takashima, T., Miwa, K., Miyazaki, I., Yamamoto, H., and Okamoto, H. (1984). Amelioration of diabetes mellitus in partially depancreatized rats by poly(ADP-ribose) synthetase inhibitors. Evidence of islet B-cell regeneration. Diabetes *33*, 401-404.

Yoo, Y.D., Huang, C.T., Zhang, X., Lavaute, T.M., and Zhang, S.C. (2011). Fibroblast Growth Factor Regulates Human Neuroectoderm Specification Through ERK1/2-PARP-1 Pathway. Stem Cells *29*, 1975-1982.

Zalzman, M., Anker-Kitai, L., and Efrat, S. (2005). Differentiation of human liver-derived, insulin-producing cells toward the beta-cell phenotype. Diabetes *54*, 2568-2575.

Zampieri, M., Guastafierro, T., Calabrese, R., Ciccarone, F., Bacalini, M.G., Reale, A., Perilli, M., Passananti, C., and Caiafa, P. (2012). ADP-ribose polymers localized on Ctcf–Parp1–Dnmt1 complex prevent methylation of Ctcf target sites. Biochemical Journal *441*, 645-652.

Zanini, C., Bruno, S., Mandili, G., Baci, D., Cerutti, F., Cenacchi, G., Izzi, L., Camussi, G., and Forni, M. (2011). Differentiation of mesenchymal stem cells derived from pancreatic islets and bone marrow into islet-like cell phenotype. PloS one *6*, e28175.

Zarnegar, R., and Michalopoulos, G.K. (1995). The many faces of hepatocyte growth factor: from hepatopoiesis to hematopoiesis. J Cell Biol *129*, 1177-1180.

Zhang, C., Moriguchi, T., Kajihara, M., Esaki, R., Harada, A., Shimohata, H., Oishi, H., Hamada, M., Morito, N., and Hasegawa, K. (2005a). MafA is a key regulator of glucose-stimulated insulin secretion. Molecular and cellular biology *25*, 4969-4976.

Zhang, D., Jiang, W., Liu, M., Sui, X., Yin, X., Chen, S., Shi, Y., and Deng, H. (2009). Highly efficient differentiation of human ES cells and iPS cells into mature pancreatic insulin-producing cells. Cell research *19*, 429-438.

Zhang, L., Hong, T.P., Hu, J., Liu, Y.N., Wu, Y.H., and Li, L.S. (2005b). Nestin-positive progenitor cells isolated from human fetal pancreas have phenotypic markers identical to mesenchymal stem cells. World journal of gastroenterology *11*, 2906-2911.

Zhang, M., Lin, Q., Qi, T., Wang, T., Chen, C.-C., Riggs, A.D., and Zeng, D. (2016). Growth factors and medium hyperglycemia induce Sox9+ ductal cell differentiation into β cells in mice with reversal of diabetes. Proceedings of the National Academy of Sciences *113*, 650-655.

Zhang, P., Nakatsukasa, H., Tu, E., Kasagi, S., Cui, K., Ishikawa, M., Konkel, J.E., Maruyama, T., Wei, G., and Abbatiello, B. (2013). PARP-1 regulates expression of TGF-β receptors in T cells. Blood *122*, 2224-2232.

Zhang, Y.Q., Kritzik, M., and Sarvetnick, N. (2005c). Identification and expansion of pancreatic stem/progenitor cells. J Cell Mol Med *9*, 331-344.

Zhang, Y.Q., and Sarvetnick, N. (2003). Development of cell markers for the identification and expansion of islet progenitor cells. Diabetes/metabolism research and reviews *19*, 363-374.

Zhao, Y., Jiang, Z., Zhao, T., Ye, M., Hu, C., Yin, Z., Li, H., Zhang, Y., Diao, Y., and Li, Y. (2012). Reversal of type 1 diabetes via islet β cell regeneration following immune modulation by cord blood-derived multipotent stem cells. BMC medicine *10*, 3.

Zheng, L., Szabo, C., and Kern, T.S. (2004). Poly(ADP-ribose) polymerase is involved in the development of diabetic retinopathy via regulation of nuclear factor-kappaB. Diabetes *53*, 2960-2967.

Zhu, H., Xie, N., Adesanya, T., Kilic, A., Tan, T., and Xie, X. (2016). Transplantation of Mesenchymal Stem Cells Enhances Angiogenesis after Ischemic Limb Injury in Mice. Biophysical Journal *3*, 141a.

Zimmet, P., Tuomi, T., Mackay, I., Rowley, M., Knowles, W., Cohen, M., and Lang, D. (1994). Latent autoimmune diabetes mellitus in adults (LADA): the role of antibodies to glutamic acid decarboxylase in diagnosis and prediction of insulin dependency. Diabetic Medicine *11*, 299-303.

Zimmet, P.Z. (1995). The pathogenesis and prevention of diabetes in adults. Genes, autoimmunity, and demography. Diabetes Care *18*, 1050-1064.

Zingarelli, B., Salzman, A.L., and Szabo, C. (1998). Genetic disruption of poly (ADP-ribose) synthetase inhibits the expression of P-selectin and intercellular adhesion molecule-1 in myocardial ischemia/reperfusion injury. Circulation research *83*, 85-94.

Zulewski, H., Abraham, E.J., Gerlach, M.J., Daniel, P.B., Moritz, W., Muller, B., Vallejo, M., Thomas, M.K., and Habener, J.F. (2001). Multipotential nestin-positive stem cells isolated from adult pancreatic islets differentiate ex vivo into pancreatic endocrine, exocrine, and hepatic phenotypes. Diabetes *50*, 521-533.