Bibliography

1. Aronowski J, Strong R, Grotta JC. Reperfusion injury: demonstration of brain damage produced by reperfusion after transient focal ischemia in rats. J. Cereb. Blood Flow Metab. 1997; 17:1048 - 1056.
2. Verma S, Fedak PWM, Weisel RD, Butany J, Rao V, Maitland A, Ren-Ke Li, Dhillon B, Yau TM. Fundamentals of reperfusion injury for the clinical cardiologist. Circulation. 2002;105: 2332-2336.
3. Piper HM., Abdallah Y., Schafer C. The first minutes of reperfusion: a window of opportunity for cardioprotection. Cardiovas Res 2004;61:365– 371.
4. Kloner RA, Jennings RB. Consequences of brief ischemia: stunning, preconditioning, and their clinical implications: part I. Circulation. 2001;104:2981–2989.
5. Kloner RA, Jennings RB. Consequences of brief ischemia: stunning, preconditioning, and their clinical implications: part II. Circulation. 2001;104:3158 –3167.
6. Yellon DM, Baxter GF. Protecting the ischaemic and reperfused myocardium in acute myocardial infarction: distant dream or near reality? Heart. 2000;83:381–387.
7. Ambrosio G, Tritto I. Reperfusion injury: experimental evidence and clinical implications. Am Heart J. 1999;138:S69 –S75.
8. Hademenos GJ., Massoud TF. Biophysical Mechanisms of Stroke. Stroke 1997;28:2067-2077.
9. Mendis M. et al. Stroke and cardiovascular diseases: secondary management, a WHO study. Current sciences 2004;9;126-134.
10. Thomas Carmichael S. Rodent Models of Focal Stroke: Size, Mechanism, and Purpose. NeuroRx. 2005;2: 396–409.
11. Haffner S. Diabetes and the metabolic syndrome—When is it best to intervene to prevent? Atherosclerosis Supplements 2006;7:3–10.
12. Hammoud T, Tanguay JF, Bourassa MG. Management of coronary artery disease: therapeutic options in patients with diabetes. J Am Coll Cardiol 2000;36:355– 65.
13. Andersen B, Goldsmith GH, Spagnolo PJ. Neutrophil adhesive dysfunction in diabetes mellitus: the role of cellular and plasma factors. J Lab Clin Med 1988;111:275 – 85.
14. McDonagh P, Hokoma JY. Microvascular perfusion and transport in the diabetic heart. Microcirculation 2000;7:163– 81.
15. Mildenberger RR, Bar-Shlomo B, Druck MN, et al. Clinically unrecognized ventricular dysfunction in young diabetic patients. J Am Coll Cardiol 1984;4:234 –8.
16. Paulson DJ. The diabetic heart is more sensitive to ischemic injury. Cardiovasc Res 1997;34:104–12.
17. Feuvray D, Lopaschuk GD. Controversies on the sensitivity of the diabetic heart to ischemic injury: the sensitivity of the diabetic heart to ischemic injury is decreased. Cardiovasc Res 1997;34:113–20.

18. Kannel WP, Castelli WP, Gordon T, McNamara P. Serum cholesterol, lipoproteins and risk of coronary heart disease: the Framingham study. Ann Intern Med 1971;74:1 –12.
19. Gilligan DM, Guetta B, Panza J, Garcia CE, Quyyumi AA, Cannon RO. Selective loss of microvascular endothelial function in human hypercholesterolemia. Circulation 1994;90:35 –41.
20. White CR, Darley-Usmar V, Berrington WR, et al. Circulating plasma xanthine oxidase contributes to vascular dysfunction in hyper-cholesterolemic rabbits. Proc Natl Acad Sci U S A 1996;93:8745– 9.
21. Cambien F, Poirier O, Lecerf L, et al. Deletion polymorphism in the gene for angiotensin-converting enzyme is a potent risk factor for myocardial infarction. Nature 1992;359:641– 4.
22. Klein L, O'Connor CM, Gattis WA, et al. Pharmacologic therapy for patients with chronic heart failure and reduced systolic function: review of trials and practical considerations. Am J Cardiol 2003; 91(9A):18F – 40F.
23. Teo KK, Yusuf S, Pfeffer M, et al. ACE Inhibitors Collaborative Group. Effects of long-term treatment with angiotensin-converting-enzyme inhibitors in the presence or absence of aspirin: a systematic review. Lancet 2002;360:1037– 43.
24. Schmid-Schonbein GW, Seiffge D, DeLano FA, Shen K, Zweifach BW. Leukocyte counts and activation in spontaneously hypertensive and normotensive rats. Hypertension 1995;17:323– 30.
25. Boughner DR, Roach MR. Effect of low frequency vibration on the arterial wall. Circ Res. 1971;29:136-144.
26. Siegmund B, Klietz T, Schwartz P, Piper HM. Temporary contractile blockade prevents hypercontracture in cardiomyocytes. Am J Physiol 1991;260:H426 – 35.
27. The Cardiovascular disease management. Merck Med. Library, Updates 2006.
28. Hassan A and Markus HS.Genetics and ischemic stroke. Brain 2000;123:1784-1812.
29. Wilkins RH. Cerebral vasospasm. Contemp Neurosurg. 1988;10:1-66.
30. Molina CA and Saver JL. Extending Reperfusion Therapy for Acute Ischemic Stroke: Emerging Pharmacological, Mechanical, and Imaging Strategies. Stroke. 2005;36:2311-2320.
31. Marczin N., El-Habashi N., Hoare GS., Bundy RE., Yacouba M.Antioxidants in myocardial ischemia–reperfusion injury: therapeutic potential and basic mechanisms. Arc of Biochem Biophy 2003; 420:222–236.
32. Cuzzocrea S., Riely DP, Caputi AP., Salvemini D. Antioxidant therapy : A new pharmacological approach in shock, inflammation, and ischemia/reperfusion injury. Pharmacol. Rev. 2001; 53:135-159.
33. Thiyagarajan M, Sharma SS. Neuroprotective effect of curcumin in middle cerebral artery occlusion induced focal cerebral ischemia in rats. Life Sciences 2004;74: 969-85.
34. Du Y. and Ko KM. Effect of pharmacological preconditoning by emodin/oleanolic acid treatment and /or ischemic precondtioning on mitochondrial antioxidant components as well as the susceptibility to ischemia-reperfusion injury in rat hearts. Mol Cell Biochem, 2006; 288:135-142.

35. Zhai P.,Eureli TE., Cotthaus R., Jeffery EH., Bahr JM., Gross DR. Effect of estrogen on global myocardial ischemia-reperfusion injury in female rats.Am J Physiol Heart Circ Physiol 2000; 279:H2766-H2775.
36. Roof, R.L., Duvdevani, R., Stein, DG. Gender influences outcome of brain injury; progesterone plays a protective role. Brain Res 1993;607,333-336.
37. Panigrahi M, Sadguna Y, Shivakumar BR, et al. alpha-lipoic acid protects against reperfusion injury following cerebral ischemia in rats. Brain Res 1996;717:184-188.
38. Ferrara A., Quesenberry CP., Karter AJ., Njoroge CW., Jacobson AS., Selby JV. Current use of unopposed estrogen and estrogen plus progestin and the risk of acute myocarfdial infarction among women with diabetes. Circulation 2003; 107:43-48.
39. Manson JE, Judith Hsia PH, Johnson KC., Rossouw JE, Assaf AR,.Lasser NL, Trevisan M,.Black HR,.Heckbert SR, Detrano R, Strickland OL,.Wong ND, Crouse JR , Stein E, Cushman M. Estrogen plus Progestin and the Risk of Coronary Heart Disease. N Engl J Med2003; 349(6):523-534.
40. Toung TJ, Traystman RJ, Hurn PD. Estrogen-mediated neuroprotection after experimental stroke in male rats. Stroke 1998;29:1666-1670.
41. Carswell HV, Bingham D, Wallece K, Nilsen M, Graham DI, Dominiczak AF, macrae IM. Differencial effects of 17 beta-estradiol upon estrogen upon stroke damage in stroke prone and normotensive rats. J Cereb Blood Flow Metab 2004,24:298-304.
42. Harukuni I, Hurn PD, Crain BJ. Deleterious effect of beta-estradiol in rat model of transient forebrain ischemia. Brain Res 2001:900:137-142.
43. Stefanick ML. Estrogen, progestogens and cardiovascular risk. J Reprod Med. 1999; 44(2 Suppl):221-226.
44. Grady D, Herrington D, Bittner V, Blumenthal R, Davidson M, Hlatky M, Hsia J, Hulley S, Herd A, Khan S, Newby LK, Waters D, Vittinghoff E, Wenger N; HERS Research Group. Cardiovascular disease outcomes during 6.8 years of hormone therapy: Heart and Estrogen/progestin Replacement Study follow-up (HERS II). JAMA 2002; 288:49-57.
45. Rosano GM and Panina G.Cardiovascular pharmacology of hormone replacement therapy. Drugs Aging. 1999;15(3):219-234.
46. Cerquetani E, Leonardo F, Pagnotta P, Galetta P, Onorati D, Fini M, Rosano GM. Anti-ischemic effect of chronic oestrogen replacement therapy alone or in combination with medroxyprogesterone acetate in different replacement schemes. Maturitas. 2001; 39(3):245-51.
47. Murphy SJ, TRaystman RJ, Hurn PD. Progesterone exacerbates striatal stroke injury in progesterone-deficient female animals. Stroke 2000;31:1173-1178.
48. Love S. Oxidative stress in brain ischemia. Brain Patholo 1999; 9:119-131.
49. Halliwell B. Free radicals, antioxidants and human disease: curiosity, cause, or consequence? Lancet. 1994; 344:721- 4.

50. Poltronieri R, Cevese A, Sbarbati A. Cardioscience 1992;3:155-160.
51. Cao X and Philis JW. The free radical scavenger alpha- lipoic acid protects against cerebral ischemia-reperfusion injury in gerbils. Free Radic res 1995; 23:365-370.
52. Freisleben HJ. Lipoic acid reduces ischemia-reperfusion injury in animal models. Toxicology. 2000;148(2-3):159-71.
53. Maulik N, Yoshida T,. Engelman RM, Bagchi D, Otani H, Das DK. Dietary coenzyme Q10 supplement renders swine hearts resistant to ischemia-reperfusion injury. Am J Physiol Heart Circ Physiol 2000; 278: H1084–H1090.
54. Ostrowski RP. Effect of coenzyme Q10 on biochemical and morphological changes in experimental ischemia in the rat brain.Brain Res Bull 2000;53:399–407, 2000.
55. Schaper W, Schaper J. Reperfusion injury: an opinionated view. J Thrombos Thromboly 1997;4:113-116.
56. Hearse DJ. Myocardial protection during ischemia and reperfusion. Mol Cell Biochem 1998;186: 177–184.
57. Kloner RA, Ganote CE, Whalen D. Effect of a transient period of ischemia on myocardial cells, II: fine structure during the first few minutes of reflow. Am J Pathol 1974;74:399 –422.
58. Jennings RB, Murry CE, Steenbergen C Jr, et al. Development of cell injury in sustained acute ischemia. Circulation. 1990;82:II-2–II-12.
59. Park JW., ChunYS., Kim MS., Park YC., Kwak, SJ., and Park SC. Metabolic modulation of cellular redox potential can improve cardiac recovery from ischemia-reperfusion injury. Int. J. Cardiol 1998;65:139–147.
60. Rosenkranz ER, Buckberg GD. Myocardial protection during surgical coronary reperfusion. J Am Coll Cardiol 1983; 1:1235-1246.
61. ennings RB, Sommers HM, Smyth GA, Flack HA, Linn H. Myocardial necrosis induced by temporary occlusion of a coronary artery in the dog. Pathology 1960; 70:68-78.
62. Maxwell SRJ, Lip GYH. Reperfusion injury: a review of the pathophysiology, clinical manifestations and therapeutic options.International Journal of Cardiology 1997;58:95–117.
63. Jennings RB, Schaper J, Hill ML, et al Jr. Effect of reperfusion late in the phase of reversible ischemic injury: changes in cell volume, electrolytes, metabolites, and ultrastructure. Circ Res 1985;56:262–278.
64. Jennings RB, Reimer KA, Hill ML. Total ischemia, in dog hearts, in vitro, I: comparison of high energy phosphate production, utilization and depletion and of adenine nucleotide catabolism in total ischemia in vitro vs severe ischemia in vivo. Circ Res 1981;49:892–900.
65. Fleet WF, Johnson TA, Graebner CA, et al. Effect of serial brief ischemic episodes on extracellular K^+, pH, and activation in the pig. Circulation. 1985;72:922–932.
66. Jennings RB, Reimer KA, Steenbergen C Jr. Myocardial ischemia revisited: the osmolar load, membrane damage, and reperfusion. J Mol Cell Cardiol 1986;18:769 –780.

67. Rovetto MJ, Lamberton WF, Neely JR. Mechanisms of glycolytic inhibition in ischemic rat hearts. Circ Res 1975;37:742–751.

68. Jennings RB, Steenbergen C Jr. Nucleotide metabolism and cellular damage in myocardial ischemia. Annu Rev Physiol. 1985;47:727–749.

69. Kusumdevi V, Rehman F. Neutraceutical Antioxidants- An overview. Ind J Pharm Edu 2002; 36;67-74.

70. Mimic-Oka J, Simic DV, Simic V. Free Radicals In Cardiovascular Diseases, Med Biol 1999;6(1):11–22.

71. Hess ML, Manson NH. Molecular oxygen: friend and foe. The role of the oxygen free radical system in the calcium paradox, the oxygen paradox and ischemia/reperfusion injury. J Mol Cell Cardiol 1984;16(11):969 –985.

72. Hiraishi H, Terano A, Razandi M, et al. Reactive oxygen metabolite-induced toxicity to cultured bovine endothelial cells: status of cellular iron in mediating injury. J Cell Physiol 1994;160(1):132–134.

73. Grune T, Reinheckel T, Davies KJ. Degradation of oxidized proteins in mammalian cells. FASEB J 1997 ;11(7):526 – 34.

74. Bolli R, Jeroudi MO, Patel BS, et al. Direct evidence that oxygen-derived free radicals contribute to postischemic myocardial dysfunction in the intact dog. Proc Natl Acad Sci U S A. 1989;86:4695–4699.

75. Jordan JE, Zhao ZQ, Vinten-Johansen J. The role of neutrophils in myocardial ischemia-reperfusion injury. Cardiovasc Res. 1999; 43: 860–878.

76. Hwang, Y. C., Kaneko, M., Bakr, S., Liao, H., Lu, Y., Lewis, E. R., Yan, S., Ii, S., Itakura, M., Rui, L., Skopicki, H., Homma, S., Schmidt, A. M., Oates, P. J., Szabolcs, M., Ramasamy, R. Central role for aldose reductase pathway in myocardial ischemic injury. FASEB J. 2004;18, 1192–1199.

77. Hwang, Y. C., Bakr, S., Ellery, C. R., Oates, P. J., and Ramasamy R. () Sorbitol dehydrogenase: a novel target for adjunctive protection of ischemic myocardium. FASEB J. 2003;17: 2331–2333.

78. Stanley, W. C., Lopaschuk, G. D., Hall, J. L., and McCormack, J. G. (1997) Regulation of myocardial carbohydrate metabolism under normal and ischemic conditions. Potential for pharmacological interventions. Cardiovasc. Res.1997;33:243–257.

79. Taegtmeyer, H., King, L. M., and Jones, B. E. (1998) Energy substrate metabolism, myocardial ischemia, and targets for pharmacotherapy. Am J Cardiol 1998; 82: 54K–60K.

80. Stanley WC and Marzilli M. Metabolic therapy in the treatment of ischemic heart disease: the pharmacology of trimetazidine. Fundam Clin Pharmacol 2003; 17:133-145.

81. Trueblood, N. A., Ramasamy, R., Wang, L. F., and Schaefer, S. Niacin protects the isolated perfused heart from isch-emia/reperfusion injury: the role of lactate/proton co-trans-port. Am. J. Physiol.2000;279:H764–H77.
82. Trueblood, N. A., and Ramasamy, R. (1998) Aldose reductase inhibition improves the altered glucose metabolism of isolated diabetic rat hearts. Am J Physiol 1998; 275: H75–H83.
83. Ramasamy, R., Trueblood, N. A., and Schaefer, S. Metabolic effects of aldose reductase inhibition during low-flow ischemia and reperfusion. Am J Physiol 1998;275: H195–H203.
84. Hwang, Y. C., Sato, S., Tsai, J. Y., Bakr, S., Yan, S. D., Oates, P. J., and Ramasamy, R. Aldose reductase activation is a key component of myocardial response to ischemia. FASEB J. 2002;16:243–245.
85. Braunwald E, Kloner RA. The stunned myocardium: prolonged, postischemic ventricular dysfunction. Circulation. 1982; 66: 1146–9.
86. Di Wang H, Hope S, Du Y, et al. Paracrine role of adventitial superoxide anion in mediating spontaneous tone of the isolated rat aorta in angiotensin IIinduced hypertension. Hypertension 1999; 33: 1225-32.
87. Griendling KK, Minieri CA, Ollerenshaw JD,Alexander RW. Angiotensin II stimulates NADH and NADPH oxidase activity in cultured vascular smooth muscle cells. Circ Res 1994;74: 1141-8.
88. Rajagopalan S, Kurz S, Munzel T, et al. Angiotensin II-mediated hypertension in the rat increases vascular superoxide production via membrane NADH/NADPH oxidase activation. Contribution to alterations of vasomotor tone. J Clin Invest1996; 97: 1916-23.
89. Wildhirt SM, Dudek RR, Suzuki H et al. Expression of nitric oxide synthase isoforms after myocardial infarction in humans. Endothelium 1995;3:209–224.
90. Sears CE, Bryant SM, Ashley EA, et al. Cardiac neuronal nitric oxide synthase isoform regulates myocardial contraction and calcium handling. Circ Res 2003;92:e52 – e59.
91. Bolli R, Jeroudi MO, Patel BS, et al. Direct evidence that oxygen-derived free radicals contribute to postischemic myocardial dysfunction in the intact dog. Proc Natl Acad Sci U S A. 1989;86:4695–4699.
92. Bing RJ, Yamamoto T, Yamamoto M, Kakar R, Cohen A. New look at myocardial infarction: toward a better aspirin. Cardiovasc Res 1999;43:25–31.
93. Salvemini D, Misko TP, Masferrer JL et al. Nitric oxide activates cyclooxygenase enzymes. Proc Natl Acad Sci USA 1993;90:7240– 7244.
94. Davidge ST, Baker PN, McLaughlin MK, Roberts JM. Nitric oxide produced by endothelial cells increases production of eicosanoids through activation of prostaglandin H synthase. Circ Res 1995;77:274–283.
95. Vane J. Towards a better aspirin. Nature 1994;367:215–216.
96. Meldrum DR, Dinarello CA, Cleveland JCJ, Cain BS, Shames BD, Meng X, Harken AH. Hydrogen peroxide induces tumor necrosis factor alpha-mediated

cardiac injury by a p38 mitogen activated protein kinase-dependent mechanisms. Surgery. 1998;124:291–296.

97. Von Knethen A, Callsen D, Brune B. Superoxide attenuates macrophage poptosis by NF-kappa B and AP-1 activation that promotes cyclooxy-genase-2 expression. J Immunol. 1999;163:2858 –2866.

98. Oliver MR, Kurien VA, Greenwook TW. Relation between serum-free-fatty acids and arrhythmia and death after myocardial infarction. Lancet 1968;i:710-715.

99. Mueller HS, Ayres ST. Metabolic responses of the heart in acute myocardial infarction in man. Am J Cardiol 1978;42:363–371.

100. Lopaschuk GD, Belke DD, Gamble J, Itoi T, Scho¨ nekess BO. Regulation of fatty acid oxidation in the mammalian heart in health and disease. Biochim Biophys Acta 1994;1213:263–276.

101.Liu B.,Clanachan AS, Schulz R, Lopaschuk GD. Cardiac efficiency is improved following ischemia by altering both the source and fate of protons. Circ Res 1996;79:940 –948.

102.Karmazyn M. Mechanisms of protection of the ischemic and reperfused myocardium by sodium-hydrogen exchange inhibition. J Thromb Thrombolysis. 1999; 8: 33-8.

103.Mann DL, Kent RL, Parsons B, Cooper G 4th. Adrenergic effects on the biology of the adult mammalian cardiocyte. Circulation 1992;85:790–804.

104.McCormack JG, Denton RM. The role of mitochondrial Ca 2+ transport and matrix Ca 2+ in signal transduction in mammalian tissues. Biochim Biophys Acta 1990;1018:287–91.

105.Silverman HS. Mitochondrial free calcium regulation in hypoxia and reoxygenation: relation to cellular injury. Basic Res Cardiol 1993;88: 483–94.

106.Shen AC, Jennings RB. Myocardial calcium and magnesium in acute ischemic injury. Am J Pathol 1972;67:417–40.

107.Crompton M. The mitochondrial permeability transition pore and its role in cell death. Biochem J 1999;341:233–49.

108. Di Lisa F, Menabo R, Canton M, Barile M, Bernadi P. Opening of the mitochondrial permeability transition pore causes depletion of mito-chondrial and cytosolic NAD+ and is a causative event in the death of myocytes in postischemic reperfusion of the heart. J Biol Chem 2001;276:2571–5.

109.Peitsch MC, Polzar B, Stephan H, Crompton T, MacDonald HR, Mannherz HG, Tschopp J. Characterization of the endogenous deox-yribonuclease involved in nuclear DNA degradation during apoptosis (programmed cell death). EMBO J 1993;12:371–7.

110.Granger DN. Ischemia-reperfusion: mechanisms of microvascular dysfunction and the influence of risk factors for cardiovascular disease. Microcirculation. 1999; 6: 167–178.

111. Fujita, A., Kurachi, Y., 2000. Molecular aspects of ATP-sensitive K^+ channels in the cardiovascular system and K^+ Channel openers. Pharmacology and Therapeutics 85, 39–53.

112. DiDiego, J.M., Antzelevitch, C., 1993. Pinacidil-induced electrical heterogeneity and extrasystolic activity in canine ventricular tissues. Does activation of ATP-regulated potassium current promote phase II reentry? Circulation 88, 1177–1189.

113. Fagbemi, S.O., Chi, L., Lucchesi, B.R., 1993. Antifibrillatory and profibrillatory actions of selected class I antiarrhythmic agents. Journal of Cardiovascular Pharmacology 21, 709–719.

114.Auchampach, J.A., Gross, G.J., 1994. Reduction in myocardial infarct size by the new potassium channel opener bimakalim. Journal of Cardiovascular Pharmacology 23, 554–561.

115.Yao, Z., Gross, G.J., 1994. Effects of the KATP channel opener bimakalim on coronary blood flow, monophasic action potential duration, and infarct size in dogs. Circulation 89, 1769–1775.

116.Ganote CE. Contraction band necrosis and irreversible myocardial injury. J Mol Cell Cardiol 1983;15:67 –73.

117.Barrabe's J, Meana M, Piper HM, et al. Myocardial segment shrinkage during coronary reperfusion "in situ". Relation to hypercontracture and myocardial necrosis. Pflugers Arch-Eur J Physiol 1995;431:519– 26.

118.Altschuld RA, Wenger WC, Lamka KG, et al. Structural and functional properties of adult rat heart myocytes lysed with digitonin. J Biol Chem 1985;260:14325–34.

119.Nichols CG, Lederer WJ. The role of ATP in energy-deprivation contractures in unloaded rat ventricular myocytes. Can J Physiol Pharmacol 1990;68:183– 94.

120.Schluter KD, Jakob G, Ruiz-Meana M, Garcýa-Dorado D, Piper HM. Protection of reoxygenated cardiomyocytes against osmotic fragility by NO donors. Am J Physiol 1996;271:H428– 34.

121.Ladilov Y, Efe O., Schafer C, et al. Reoxygenation-induced rigor-type contracture. J Mol Cell Cardiol 2003;35:1481– 90.

122.Garcýa-Dorado D, Inserte J, Ruiz-Meana M, et al. Gap junction un-coupler heptanol prevents cell-to-cell progression of hypercontracture and limits necrosis during myocardial reperfusion. Circulation 1997;96:3579–86.

123.Ruiz-Meana M, Garcýa-Dorado D, Hofstaetter B, Piper HM, Soler-SolerJ. Propagation of cardiomyocyte hypercontracture by passage of Na^+ through gap junction. Circ Res 1999;85:280–87.

124.Bauldry SA, Bass DA, Cousart SL, et al. Tumor necrosis factor-alpha priming of phospholipase D in human neutrophils. Correlation between phosphatidic acid production and superoxide generation. J Biol Chem 1991;266:4173 –79.

125. Richter J, Ng-Sikorski J, Olsson I, et al. Tumor necrosis factor-induced degranulation in adherent human neutrophils is dependent on CD11b/CD18-integrin-triggered oscillations of cytosolic free Ca^{2+}. Proc Natl Acad Sci U S A 1990;87:9472– 6.

126.Sullivan GW, Luong LS, Carper HT, et al. Methylxanthines with adenosine alter TNF alpha-primed PMN activation. Immunopharmacology 1995;31:19 – 29.

127.Frangogiannis NG, Lindsey ML, Michael LH, et al. Resident cardiac mast cells degranulate and release preformed TNF-a, initiating the cytokine cascade in experimental canine myocardial ischemia/reperfusion. Circulation 1998;98:699– 710.

128.Frangogiannis NG, Smith CW, Entman ML. The inflammatory response in myocardial infarction. Cardiovasc Res 2002;53:31 – 47.

129.Shimizu Y, Minatoguchi S, Hashimoto K, et al. The role of serotonin in ischemic cellular damage and the infarct size-reducing effect of sarpogrelate, a 5-hydroxy-tryptamine-2 receptor blocker, in rabbit hearts. J Am Coll Cardiol 2002;40:1347–55.

130.Gaboury JP, Johnston B, Niu XF, et al. Mechanisms underlying acute mast cell-induced leukocyte rolling and adhesion in vivo. J Immunol 1995;154:804–13.

131.Chandrasekar B, Freeman GL. Induction of nuclear factor κB and activation protein 1 in postischemic myocardium. FEBS Lett 1997;401:30 – 4.

132.Fan H, Sun B, Gu Q, et al. Oxygen radicals trigger activation of NF- kB and AP-1 and upregulation of ICAM-1 in reperfused canine heart. Am J Physiol (Heart Circ Physiol) 2002;282:H1778– 86.

133.Zhao ZQ, Nakamura M, Wang N-P, Wilcox JN, Shearer ST, Katz-markS, et al. Reperfusion-induced apoptosis progressively develops in later stages of reflow. FASEB J 1999;13:A816.

134.Zhao ZQ, Velez DA, Wang N-P, et al. Progressively developed myocardial apoptotic cell death during late phase of reperfusion. Apoptosis 2001;6:279 – 90.

135.Chatelain P, Latour J-G, Tran D, et al. Neutrophil accumulation in experimental myocardial infarcts: relation with extent of injury and effect of reperfusion. Circulation 1987;75:1083–90.

136.Kaiser L, Sparks Jr HV. Endothelial cells: not just a cellophane wrapper. Arch Intern Med 1987;147:569 – 73.

137.Clancy RM, Leszczynska-Piziak J, Abramson SB. Nitric oxide, an endothelial cell relaxation factor, inhibits neutrophil superoxide anion production via a direct action on the NADPH oxidase. J Clin Invest 1992;90:1116– 21.

138.Fujii H, Ichimori K, Hoshia K, et al. Nitric oxide inactivates NADPH oxidase in pig neutrophils by inhibiting its assembling process. J Biol Chem 1997;272:32773–8.

139.Barankiewicz J, Jimenez R, Uyesaka J, et al. Endogenous adenosine formation can regulate human neutrophil function. Adv Exp Med Biol 1994;370:417–21.

140. Tsao PS, Ma X-L, Lefer AM. Activated neutrophils aggravate endothelial dysfunction after reperfusion of the ischemic feline myocardium. Am Heart J 1992;123:1464 –71.

141.Lefer AM, Tsao PS, Lefer DJ, et al. Role of endothelial dysfunction in the pathogenesis of reperfusion injury after myocardial ischemia. FASEB J 1991;5:2029– 34.

142.Buja LM, Eigenbrodt ML, Eigenbrodt EH. Apoptosis and necrosis: basic types and mechanisms of cell death. Arch Pathol Lab Med. 1993;117:1208-1214.

143.James TN. Normal and abnormal consequences of apoptosis in the human heart: from postnatal morphogenesis to paroxysmal arrhythmias. Circulation. 1994;90:556-573.

144.Martin SJ, Green DR, Cotter TG. Dicing with death: dissecting the components of the apoptosis machinery. Trends Biochem Sci. 1994;19:26-30.

145. Majno G, Joris I. Apoptosis, oncosis, and necrosis: an overview of cell death. Am J Pathol. 1995;146:3-15.

146.Hockenbery D. Defining apoptosis. Am J Pathol. 1995;146:16-19.

147.von Harsdorf R, Li PF, Dietz R. Signaling pathways in reactive oxygen species-induced cardiomyocyte apoptosis. Circulation 1999; 99: 2934-41.

148.Alnemri ES, Livingston DJ, Nicholson DW, Salvesen G, Thornberry NA, Wong WW, Yuan J. Human ICE/CED-3 protease nomenclature. Cell. 1996;87:171.

149. Nicholson DW, Thornberry NA. Caspases: killer proteases. Trends Biochem Sci. 1997;22:299 – 306.

150.Porter AG, Ng P, Janicke RU. Death substrates come alive. Bioessays. 1997;19:501–507.

151. Liu X, Zou H, Slaughter C, Wang X. DFF, a heterodimeric protein that functions downstream of caspase-3 to trigger DNA fragmentation during apoptosis. Cell. 1997;89.175–184.

152. Suda T, Takahashi T, Golstein P, Nagata S. Molecular cloning and expression of the Fas ligand, a novel member of the tumor necrosis factor family. Cell. 1993;75:1169 –1178.

153.Pitti RM, Marsters SA, Ruppert S, Donahue CJ, Moore A, Ashkenazi A. Induction of apoptosis by Apo-2 ligand, a new member of the tumor necrosis factor cytokine family. J Biol Chem. 1996;271:12687–12690.

154.Longthorne VL, Williams GT. Caspase activity is required for commitment to Fas-mediated apoptosis. EMBO J. 1997;16:3805–3812.

155.Hsu H, Shu HB, Pan MG, Goeddel DV. TRADD-TRAF2 and TRADD-FADD interactions define two distinct TNF receptor 1 signal transduction pathways. Cell. 1996;84:299 –308.

156. Ludwig RL, Bates S, Vousden KH. Differential activation of target cellular promoters by p53 mutants with impaired apoptotic function. MolCell Biol. 1996;16:4952– 4960.

157.Polyak K, Xia Y, Zweier JL, Kinzler KW, Vogelstein B. A model for p53-induced apoptosis. Nature. 1997;389:300 –305.

158.Enari M, Sakahira H, Yokoyama H, Okawa K, Iwamatsu A, Nagata S. A caspase-activated DNase that degrades DNA during apoptosis, and its inhibitor ICAD. Nature. 1998;391:43–50.

159.Adams, H.P., Jr., Bendixen, B.H., Kappelle, L.J., Biller, J., Love, B.B., Gordon, D.L., and Marsh, E.E., III. Classification of subtype of acute ischemic stroke. Definitions for use in a multicenter clinical trial. TOAST. Trial of Org 10172 in acute stroke treatment. Stroke 1993; 24:35–41.

160.Caplan, L.R. Caplan's Stroke: A Clinical Approach, 2000 Butterworth-Heinemann, Boston.

161. Lipton P. Ischemic Cell Death in Brain Neurons. Physiol Rev . 1999; 79(4):1431-1568.

162. Ginsberg, M. D., And R. Busto. Rodent models of cerebral ischemia. Stroke , 198920: 1627–1642.

163. Belayev, L., W. Zhao, R. Busto, And M. D. Ginsberg. Transient middle cerebral artery occlusion by intraluminal suture. I. Three-dimensional autoradiographic image analysis of local cerebral glucose metabolism-blood flow interrelationships during ischemia and early recirculation. J. Cereb. Blood Flow Metab. , 1997;17: 1266–1280.

164.Koizumi, J., Y. Yoshida, T. Nakazawa, And G. Ooneda. Experimental studies of ischemic brain edema. I. A new experi-mental model of cerebral embolism in rats in which recirculation can be introduced in the ischemic area. Jpn. J. Stroke 1986;8: 1–8,

165.Li, Y., M. Chopp, Z. G. Zhang, R. L. Zhang, And J. H. Garcia. Neuronal survival is associated with 72-kDa heat shock protein expression after transient middle cerebral artery occlusion in the rat. J. Neurol. Sci. 1993;120: 187–194.

166.Aspey, B. S., S. Cohen, Y. Patel, M. Terruli, And M. J. G. Harrison. Middle cerebral artery occlusion in the rat: consistent protocol for a model of stroke. Neuropathol. Appl. Neurobiol. 24: 487–497, 1998.

167. Nishigaya, K., Y. Yoshida, M. Sasuga, H. Nukui, And G. Ooneda. Effect of recirculation on exacerbation of ischemic vascular lesions in rat brain. Stroke 1991;22: 635–642,

168.Gido, G., T. Kristian, And B. K. Siesjo. Extracellular potassium in a neocortical core area after transient focal ischemia. Stroke 1997; 28: 206–210,

169.Mergenthaler P, Dirnagl U, Meisel A. Pathophysiology of Stoke: Lessons from animal models. Metab. Brain. Dis. 2004;19(314):151-67.

170.Ginsberg MD and Pulsinelli WA. The ischemic penumbra, injury thresholds and the therapeutic window for acute stroke. Ann. Neurol. 1994; 36: 553–554,

171.Nagasawa H and Kogure K. Correlation between cerebral blood flow and histological change in a new rat model of cerebral artery occlusion. Stroke 1989;20: 1037–1043,

172.Folbergova, J., H. Memezawa, M.-L. Smith, And B. K. Siesjo. Focal and perifocal changes in tissue energy state during middle cerebral artery occlusion in normo- and hyperglycemic rats. J. Cereb. Blood Flow Metab. 1992;12: 24–33.

173.Ikonomidou, C., J. L. Mosinger, K. S. Salles, J. La-Bruyere, And J. Olney. Sensitivity of the developing rat brain to hypobaric/ischemic damage parallels sensitivity to N-methyl-D-aspartate neurotoxicity. J. Neurosci. 1989;9: 2809–2816,

174.Klienman R, G. BANKER, AND O. STEWARD. Differential sub-cellular localization of particular mRNAs in hippocampal neurons in culture. Neuron 1990.5: 821–830.

175.Luo A, Hattori J, Munoz ZH, Qin, Roth GS. Intrastriatal dopamine injection induces apoptosis through oxidation-involved activation of transcription factors AP-1 and NF-kappaB in rats. Mol Pharm 1999; 56(2): 254-64.

176.Betz A. L., J. Randall, And D. Martz. Xanthine oxidase is not a major source of free radicals in focal cerebral ischemia. Am. J.Physiol.1991; 260:H563–H568.

177. Iadecola C. Bright and dark sides of nitric oxide in ischemic brain injury. Trends Neurosci. 1997;20: 132–139.

178.Dawson T. M., V. L. Dawson, And S. H. Snyder. Molecular mechanisms of nitric oxide actions in the brain. Ann. NY Acad. Sci. 1994;738: 76–85.

179.Katsuki H.. and Okuda S. Arachidonic acid as a neurotoxic and neurotrophic substance. Prog. Neurobiol. 1995;46: 607–636.

180.Kontos HA. Oxygen radicals in cerebral vascular responses. In: Physiology of Oxygen Radicals, edited by A. E. Taylor, S. Matalan, and P. A. Ward. Bethesda, MD: Am. Physiol. Soc., 1986, p. 207–216.

181.Ikeda Y and Long DM. The molecular basis of brain injury and brain edema: the role of oxygen free radicals. Neurosurgery 1990; 27: 1–11.

182.Au, A. M., P. H. Chan, And R. A. Fishman. Stimulation of phospholipase A2 activity by oxygen-derived free radical in iso-lated brain capillaries. J. Cell. Biochem.1985; 27: 449–453.

183.Nakano, S., K. Kogure, K. Abe, And T. Yae. Ischemia-induced alterations in lipid metabolism of the gerbil cerebral cortex. I. Changes in free fatty acid liberation. J. Neurochem 1990; 54: 1911–1916.

184.Beckman, J. S. Peroxynitrite versus hydroxyl radical: the role of nitric oxide in superoxide-dependent cerebral injury. Ann. NY Acad. Sci. 1994; 738: 69–75.

185. Freeman, B. A., And J. D. Crapo. Biology of disease: free radicals and tissue injury. Lab. Invest. 1982;47: 412–426.

186. Dirnagl, U., Iadecola, C., and Moskowitz, M.A. Pathobiology of ischaemic stroke: An integrated view.Trends Neurosci. 1999; 22:391–397.

187. Beckman, J. S., T. W. Beckman, J. Chen, P. A. Marshall, And B. A. Freeman. Apparent hydroxyl radical production by peroxynitrite: implications for endothelial injury from nitric oxide and superoxide. Proc. Natl. Acad. Sci. USA 1990.87: 1620–1624.

188.Meyer, F. B., R. E. Anderson, T. M. Sundt, Jr., And T. L. Yaksh. Intracellular brain pH, indicator tissue perfusion and electroencephalography and histology in severe and moderate focal cortical ischemia. J. Cereb. Blood Flow Metab. 1986; 8: 71–78, 1986.

189.Gevers W. Generation of protons by metabolic processes in heart cells. J. Mol. Cell. Cardiol. 1977.11: 867–874.

190.Katsura, K., B. Asplund, A. Ekholm, And B. K. Siesjo. Extra- and intracellular pH in the brain during ischemia, related to tissue lactate content in normo- and hypercapnic rats. Eur.J. Neurosci. 1992. 4: 166–176.

191.Chen, Q., M. Chopp, G. Bodzin, And H. Chen. Temperature modulation of cerebral depolarization during focal cerebral ischemia in rats: correlation with ischemic injury. J. Cereb. BloodFlow Metab. 1993:13: 389–394.

192.Kristian, T., K.-I. Katsura, G. Gido, And B. K. Siesjo. The influence of pH on cellular calcium influx during ischemia. Brain Res.1994; 641: 295–302.

193.Lipton, P., And T. S. Whittingham. Energy metabolism and brain slice function. In: Brain Slices, edited by R. Dingledine. New York: Academic, 1984, p. 113–154.

194.Young, J. N., P. G. Aitken, And G. G. Somjen. Calcium, magnesium, and long-term recovery from hypoxia in hippocampal tissue slices. Brain Res.1991; 548: 343–345.

195.Goldman, D. E. Potential, impedance and rectification in mem-branes. J. Gen. Physiol.1944; 27: 37–48.

196.Del Zoppo, G., Ginis, I., Hallenbeck, J.M., Iadecola, C., Wang, X., and Feuerstein, G.Z. Inflammationand stroke: Putative role for cytokines, adhesion molecules and iNOS in brain response to ischemia. BrainPathol. 2000; 10:95–112.

197.Bogdan, C., Paik, J., Vodovotz, Y., and Nathan, C. Contrasting mechanisms for suppression of macrophage cytokine release by transforming growth factor-beta and interleukin-10. J. Biol. Chem.1992; 267:23301–23308.

198.Abe, K., M. Yoshidomi, And K. Kogure. Arachidonic acid metabolism in ischemic neuronal damage. Ann. NY Acad. Sci1991..559: 259–268.

199.Zhang, Y.-P., And G. Y. Sun. Free fatty acids, neutral glycerides and phosphoglycerides in transient focal ischemia. J. Neurochem.1995;64: 1688–1695.

200.Rao, A. M., Hatcher, J. F., Kindy, M. S. and Dempsey, R. J. Arachidonic acid and leukotriene C4: role in transient cerebral ischemia of gerbils. Neurochem. Res.1999; 24, 1225-1232.

201.Savci, V., And R. J. Wurtman. Effect of cytidine on membrane phospholipid synthesis in rat brain slices. J. Neurochem. 1995;64: 378–384.

202.Romanic, A. M., R. F. White, A. J. Arleth, E. H. Ohlstein, And F. C. Barone. Matrix metalloproteinase expression in-creases after cerebral focal ischemia in rats: inhibition of matrix metalloproteinase-9 reduces infarct size. Stroke1998; 29: 1020–1030.

203.Dietrich WD, Halley M, Valdes I, Busto R. Interrelationships between increased vascular permeability and acute neuronal damage following temperature-controlled brain ischemia in rats. Acta.Neuropathol.Berl. 1991; 81: 615-25.

204.Ferrer, I., and Planas, A.M. Signaling of cell death and cell survival following focal cerebral ischemia: Life and death struggle in the penumbra. J. Neuropathol. Exp. Neurol. 2003;62:329–339.

205.Friedlander, R.M.. Apoptosis and caspases in neurodegenerative diseases. N.Engl.J.Med 2003;348:1365–1375.

206.Love, S. Apoptosis and brain ischaemia. Prog. Neuropsychopharmacol. Biol. Psychiatry 2003;27:267–282.

207.Yu SW., Wang H., Poitras MF., Coombs C., Bowers WJ., Federoff HJ., Poirier GG., Dawson TM., and Dawson VL. Mediation of poly(ADP-ribose) polymerase-1-dependent cell death by apoptosis-inducing factor. Science 2002;297:259–263.

208.Ferrari R, Ceconi C, Curello S, et al. Role of oxygen free radicals in ischemic and reperfused myocardium. Am J Clin Nutr 1991;53: 215S– 22S.

209.Okado-Matsumoto, A. and Fridovich, I. Subcellular distribution of superoxide dismutases (SOD) in rat liver. J. Biol. Chem.2001;276:38388-38393.

210.Yang G., Chan PH., Chen J., Carlson E., Chen S., Weinstein P., Epstein CJ. and Kamii, H. Human copper-zinc superoxide dismutase transgenic mice are highly resistant to reperfusion injury after focal cerebral ischemia. Stroke 1994; 25:165-170.

211.Marklund, SL. Extracellular superoxide dismutase in human tissues and human cell lines. J. Clin. Invest. 1984;74:1398-1403.

212.Tibell, L., Hjalmarsson, K., Edlund, T., Skogman, G., Engstom, A. and Marklund, SL. Expression of human extracellular superoxide dismutase in Chinese hamster ovary cells and characterization of the product. Proc. Natl. Acad. Sci. USA1987;84:6634-6638.

213.Warner DS, Sheng H Batinic′-Haberle I. Oxidants, antioxidants and the ischemic brain. The J Exptal Biol 2004; 207:3221-3231.

214. Halliwell, B., And Gutteridge MC. Oxygen toxicity, oxygen radicals, transition metals and disease. Biochem. J.1984;219: 1–14.

215. Namba, K., Takeda, Y., Sunami, K. and Hirakawa, M.Temporal profiles of the levels of endogenous antioxidants after four-vessel occlusion in rats. J. Neurosurg. Anesthesiol.2001;13:131-137.

216. Park, EM., Choi, JH., Park, JS., Han, MY. and Park, YM. Measurement of glutathione oxidation and 8-hydroxy-□¢-deoxyguanosine accumulation in the gerbil hippocampus following global ischemia. Brain. Res. Brain Res. Protoc.2000; 6: 25-32.

217.Vanella, A., Di Giacomo, C., Sorrenti, V., Russo, A , Castorina, C.,Campisi, A., Renis, M. and Perez-Polo, J R. Free radical scavenger depletion in post-ischemic reperfusion brain damage. Neurochem.Res.1993;18:1337-1340.

218.Shivakumar BR., Kolluri VRS. and Ravindranath V. Glutathione and protein thiol homeostasis in brain during reperfusion following cerebral ischemia, J. Pharmacol. Exp. Ther.1995;274:167-1173.

219.Ravindranath, V. and Reed, DJ., Glutathione depletion and formation of glutathione-protein mixed disulfide following exposure of brain mitochondria to oxidative stress. Biochem. Biophys. Res. Commun.1990;169:1075-1079.

220.Nakamura M., Thourani VH., Ronson RS., Velez DA., Ma XL., Katzmark S., Robinson J., Schmarkey LS., Zhao ZQ., Wang NP., et al. Glutathione reverses endothelial damage from peroxynitrite, the byproduct of nitric oxide degradation, in cristaloid cardioplegia. Circulation 2000; 102:III332-III338.

221.Fullerton HJ., Ditelberg JS., Chen SF., Sarco DP., Chan PH., Epstein CJ. and Ferriero DM. Copper/zinc superoxide dismutase transgenic brain accumulates hydrogen peroxide after perinatal hypoxia ischemia. Ann. Neurol 1998; 44:357-364.

222.Oliver MF, Opie LH. Effects of glucose and fatty acids on myocardial ischemia and arrhythmias. Lancet 1994;343:155–158.

223.Opie LH. Proof that glucose–insulin–potassium provides metabolic protection of ischaemic myocardium? Lancet 1999;353:768–769.

224.Van der Vusse GJ, Van Bilsen M, Glatz JFC. Cardiac fatty acid uptake and transport in health and disease. Cardiovasc Res 2000;45:279–293.

225.Van Bilsen M, De Vries JE, Van der Vusse GJ. Long-term effects of fatty acids on cell viability and gene expression of neonatal cardiac myocytes. Prostagl Leukotr Essent Fatty Acids 1997;57:39–45.

226.Floyd RA, Carney JM. Free radical damage to protein and DNA: Mechanism involved and relevant observations on brain undergoing oxidative stress. Ann. Neurol. 1992; 32:s22-s27.

227.Dhalla AK, Hill MF,Singal PK. Role of oxidative stress in transition of hypertrophy to heart failure. J Am Coll Cardiol 1996;28: 506-14.

228.Wojtczak, L. Effect of long-chain fatty acids and acyl-CoA on mitochondrial permeability, transport, and energy-coupling processes. J.Bioenerg. Biomembr1976; 8: 293-311.

229.Gaudet, RJ., Alam, I. and Levine, L. Accumulation of cyclooxygenase products of arachidonic acid metabolism in gerbil brain during reperfusion after bilateral common carotid artery occlusion. J.Neurochem 1980;35:653-658.

230.Imaizumi, S., Tominaga, T., Uenohara, H., Yoshimoto, T., Suzuki, J. And Fujita, Y. Initiation and propagation of lipid peroxidation in cerebral infarction models. Experimental studies. Neurol. Res.1986;8: 214-220.\

231.Kruman, I., Bruce-Keller, AJ., Bredesen, D., Waeg, G. and Mattson, MP. Evidence that 4-hydroxynonenal mediates oxidative stress-induced neuronal apoptosis. J. Neurosci.1997; 1: 5089-5100.

232.Mattson, MP., Culmsee, C. and Yu, ZF. Apoptotic and antiapoptotic mechanisms in stroke. Cell Tissue Res.2000;301:173-187.

233.Slater TF, Sawyer BC. The stimulatory effects of carbon tetrachloride and other halogenoalkanes or peroxidative reactions in rat liver fractions in vitro. Biochemical Journal 1971;123: 805-14.

234. Kjekshus JK., Sobel, BE. Depressed myocardial creatine phosphokinase activity following experimental myocardial infarction in rabbit. Circulation Res 1970;27:403-414.

235. Saks VA., Khuchua ZA., Vasilyeva EV., Belikova OY., Kuznetsov AV. Metabolic compartmentation and substrate channeling in muscle cells: role of coupled creatine kinases in in vivo regulation of cellular respiration: a synthesis. Mol Cell Biochem 1994;133: 155-192.

236. Watts DC. Creatine kinase (adenosine 5'-triphosphate-creatine phosphotransferase). In: Boyer PD, ed. The Enzyme. New York, NY: Academic Press 1973;8, 383-455.

237.Dreyfus JC., Schapira G., Rasnais J., Scebat L. La creatine-kinase sérique dans le diagnostic de l'infarctus myocardique, Revue Française d'études Cliniques et Biologiques 1960;5, 386-389.

238.Kleber, AG. ST-segment elevation in the electrocardiogram: a sign of myocardial ischemia. Cardiovas Res 2000; 45, 111-118.

239. Sobel BE., Bresnahan GF., Shell WE., Yoder RD., 1972. Estimation of infarct size in man and its relation to prognosis. Circulation 1972; 46, 640-648.

240.Mlynárik V, Kaš parová S., Liptaj T., Dobrota D., Horecký J. and Belan V. Creatine kinase reaction rates in rat brain during chronic ischemia.Magn Reso Mat in Bio Phy and Med 1998; 7:162-165.

241.Wang, Q.D., Pernow, J., Sjoquist, P., Ryden, L., 2002. Pharmacological possibilities for protection against myocardial reperfusion injury. Cardiovascular Research 55,25-37.

242.Tienfenbacher, CP., Kaptiza, J., Dietz, V., Lee, CH., Niroomand, F. Reduction of myocardial infarct size by fluvastatin. The Am J Physiolo Heart and Circulatory Physiolo 2003; 285, H59-H64.

243.Baldus S., Heitzer T., Eiserich, JP., Lau YD., Mollnau H., Ortak M., Petri S., Goldmann B., Duchstein HJ., Berger,J., Helmchen U., Freeman BA., Meinertz T., Nzel TM. Myeloperoxidase enhances nitric oxide catabolism during myocardial ischemia and reperfusion. Free Radl Biol & Med 2004;37,902-911.

244.Gumina RJ., Schultz Jo El., Zhenhai Yao, KennyD., Warltier DC., Newman PJ., Gross GJ. Antibody to platelet/endothelial cell adhesion molecule-1 reduces myocardial infarct size in a rat model of ischemia-reperfusion injury._Circulation. 1996;94:3327-3333.

245. Stendahi O., Coble BI, Dahigren C., Hed J., Molin L. Myeloperoxidase modulates the phagocytic activity of polymorpho neutrophil leukocytes. Studies with cells from a myeloperoxidase deficient patient. J Clin Invent 1984;73:366-373.

246.Matsuo Y, Onodera H, Shiga Y, Nakamura M, Ninomiya M, Kihara T, Kogure K. Correlation between myeloperoxidase-quantified neutrophil accumulation and ischemic brain injury in the rat. Effects of neutrophil depletion. Stroke. 1994; 25:1469-75.

247.Kleber AG., Janse MJ., Van Capelle FJC., Durrer, D. Mechanism and time course of S-T and T-Q segment changes during acute regional myocardial ischemia in the pig heart determined by extracellular and intracellular recordings. Circulation Res1978;42,603-613.

248. Wlhelmsen K, Svardsudd K, Koraan-Bengaten K, Lars-son B, Wolln L, Tlbblln Q. Rbrinogen as a risk factor for stroke and myocardial Infarction. N Engl J Med 1984; 311:501-505.

249. Stone MC, Thorpe JM. Plasma fibrinogen a major coronary risk factor. J R Coll Gen Pract 1985;35:565-569.

250. Welln L, Svardsudd K, Wllhelmsen L, Larsson B, Tlbblln G. Analysis for risk factors of stroke In a cohort of men bom In 1913. N Engl J Med 1987:317:521-526

251.Guarnieri C. and Muscari C. Effect of trimetazidine on mitochondrial function and oxidative damage during reperfusion of ischemic hypertrophied rat myocardium. Pharmacology. 1993; 46:324-31.

252. Opie LH and Boucher F. Trimetazidine and myocardial ischemic contracture in isolated rat heart. Am J Cardiol 1995;286:97-110.

253. Levine B, Kalman J, Mayer L, Fillit HM, and Packer M. Elevated circulating levels of tumor necrosis factor-_ in severe heart chronic failure. N Engl J Med 1990;323: 236–241.

254. Bozkurt B, Kribbs SB, Clubb FJ, Michael LH, Didenko V, Hornsby PJ, Seta Y, Spinale FG, and Mann DL. Pathophysiologically relevant concentrations of tumor necrosis factor-α promote progressive left ven-tricular dysfunction and remodeling in rats. Circulation 1998; 97: 1382–1391.

255.Bryant D, Becker L, Richardson J, Shelton J, Franco F, Peshock R, Thompson M, and Giroir B. Cardiac failure in transgenic mice with myocardial expression of tumor necrosis factor-α. Circulation 1998;97: 1375– 1381.

256.Berthonneche C., Sulpice T., Boucher F., Gouraud L., de Leiris J., O'Connor SE., Herbert JM. and Janiak P. New insights into the pathological role of TNF- α in early cardiac dysfunction and subsequent heart failure after infarction in rats. Am J Physiol Heart Circ Physiol 2004; 287: H340–H350.

257. Rajagopalan S, Kurz S, Munzel T, et al. Angiotensin II-mediated hypertension in the rat increases vascular superoxide production via membrane NADH/NADPH oxidase activation. Contribution to alterations of vasomotor tone. J Clin Invest1996; 97: 1916-23.

258.Feuerstein G, Yue TL, Ma X, Ruffolo RR. Novel mechanisms in the treatment of heart failure: inhibition of oxygen radicals and apoptosis by carvedilol. Prog Cardiovasc Dis 1998; 41: 17-24.

259.Feuerstein G, Ruffolo RRJ. Beta-blockers in congestive heart failure: the pharmacology of carvedilol, a vasodilating beta-blocker and antioxidant, and its therapeutic utility in congestive heart failure. Adv Pharmacol 1998; 42: 611-5.

260.Lerman A, Burnett JCJ, Higano ST, McKinley L. J,Holmes DR J. Long-term L-arginine supplementation improves small-vessel coronary endothelial function in humans. Circulation 1998;97: 2123- 8.

261.Rector TS, Bank AJ, Mullen KA, et al. Randomized, double-blind, placebocontrolled study of supplemental oral L-arginine in patients with heart failure. Circulation 1996;93: 2135-41.

262.Heller R, Munscher-Paulig F, Grabner R, Till U. L-Ascorbic acid potentiates nitric oxide synthesis in endothelial cells. J Biol Chem 1999;274: 8254-60.

263.Hirashima O, Kawano H, Motoyama T, et al. Improvement of endothelial function and insulin sensitivity with vitamin C in patients with coronary spastic angina: possible role of reactive oxygen species. J Am Coll Cardiol 2000;35: 1860-6.

264.Wilkinson IB, Megson IL, MacCallum H, Sogo N, Cockcroft JR,Webb DJ. Oral vitamin C reduces arterial stiffness and platelet aggregation in humans. J Cardiovasc Pharmacol 1999; 34: 690-93.

265.Wall SR., Fliss H and Korecky B. Role of catalase in myocardial protection against ischemia in heat shocked rats. Mol Cell Biol 1993; 129:187-194.

266.Margaill I , Plotkine M, Lerouet D. Antioxidant strategies in the treatment of stroke. Free Radical Biology & Medicine 2005;39: 429 – 443.

267.Marro, PJ., McGowan JE., Razdan B., Mishra OP. and Delivoria-Papadopoulos, M. Effect of allopurinol on uric acid levels and brain cell membrane Na + ,K (+) -ATPase activity during hypoxia in newborn piglets.Brain Res.1994; 650, 9-15.

268.Nihei, H., Kanemitsu, H., Tamura, A., Oka, H. and Sano, K. Cerebral uric acid, xanthine, and hypoxanthine after ischemia: the effect of allopurinol. Neurosurgery 1989;25, 613-617.

269.Williams GD., Palmer C., Heitjan DF. and Smith MB. Allopurinol preserves cerebral energy metabolism during perinatal hypoxia-ischemia:a 31P NMR study in unanesthetized immature rats. Neurosci. Lett. 1992;144, 103-106.

270.van Bel, F., Shadid, M., Moison, RM., Dorrepaal, CA., Fontijn, J., Monteiro, L., Van De Bor, M. and Berger, HM. Effect of allopurinol on postasphyxial free radical formation, cerebral hemodynamics, and electrical brain activity. Pediatrics 1998;101, 185-193.

271.Mohanty I., Arya DS., Dinda A., Joshi S., Talwar KK., Gupta SK. Protective effects of Curcuma longa on ischemia-reperfusion induced myocardial injuries and their mechanisms. Life Sciences 2004;75: 1701–1711.

272. Clark WM., Rinker LG., Lessov NS., Lowery SL., Cipolla MJ. Efficacy of antioxidant therapies in transient focal ischemia in mice. Stroke. 2001;32:1000-1004.

273.Yang, G., Chan, P. H., Chen, J., Carlson, E., Chen, S., Weinstein, P., Epstein, C. J. and Kamii, H.Human copper-zinc superoxide dismutase transgenic mice are highly resistant to reperfusion injury after focal cerebral ischemia. Stroke 1994;25, 165-170.

274.Chan, P. H., Kamii, H., Yang, G. Y., Gafni, J., Epstein, C. J., Carlson, E.

and Reola, L. Brain infarction is not reduced in SOD-1 transgenic mice after a permanent focal cerebral ischemia. NeuroReport 1993;5, 293-296.

275. Fujimura, M., Morita-Fujimura, Y., Copin, J., Yoshimoto, T. and Chan, P. H. Reduction of copper,zinc-superoxide dismutase in knockout mice does not affect edema or infarction volumes and the early release of mitochondrial cytochrome c after permanent focal cerebral ischemia. Brain Res. 2001;889, 208-213.

276. Kim GW., Kondo T., Noshita N. and Chan PH. Manganese superoxide dismutase deficiency exacerbates cerebral infarction after focal cerebral ischemia/reperfusion in mice: implications for the production and role of superoxide radicals. Stroke2002;33, 809-815.

277. Murakami K., Kondo T., Kawasr M., Li Y., Sato S., Chen SF. and Chan PH. Mitochondrial susceptibility to oxidative stress exacerbates cerebral infarction that follows permanent focal cerebral ischemia in mutant mice with manganese superoxide dismutase deficiency. J. Neurosci.1998;18, 205-213.

278. Sheng, H., Batinic-Haberle, I. and Warner, DS. Catalytic antioxidants as novel pharmacological approaches to treatment of ischemic brain injury. Drug News Perspect. 2002;15, 654-665.

279. Sheng, H., Brody, T., Pearlstein, RD., Crapo, J. and Warner, DS. Extracellular superoxide dismutase deficient mice have decreased resistance to focal cerebral ischemia. Neurosci. Lett 1999: 267, 13-17.

280. Mollace V., Iannone M., Muscoli C., Palma E., Granato T., Modesti A., Nisticò R., Rotiroti D. and Salvemini D. The protective effect of M40401,a superoxide dismutase mimetic, on post-ischemic brain damage in Mongolian gerbils. BMC Pharmacology 2003, 3:1-9.

281. Muller, A., Cadenas, E., Graf, P. and Sies, H. A novel biologically active seleno-organic compound-I. Glutathione peroxidase-like activity in vitro and antioxidant capacity of PZ 51 (Ebselen). Biochem. Pharmacol.1984;15, 3235-3239.

282. Schewe, T. Molecular actions of ebselen – an antinflammatory antioxidant. Gen. Pharmacol. 1995;26, 1153-1169.

283. Saito, I., Asano, T., Sano, K., Takakura, K., Abe, H., Yoshimoto, T., Kikuchi, H., Ohta, T. and Ishibashi, S. Neuroprotective effect of an antioxidant, ebselen, in patients with delayed neurological deficits after aneurysmal subarachnoid hemorrhage. Neurosurgery 1998; 42, 269-277; Discussion 277-278.

284.Yamaguchi, T., Sano, K., Takakura, K., Saito, I., Shinohara, Y., Asano,T. and Yasuhara, H. Ebselen in acute ischemic stroke: a placebo-controlled, double-blind clinical trial. Ebselen Study Group. Stroke 1998;29, 12-17.

285. Lo, E. H., Hara, H., Rogowska, J., Trocha, M., Pierce, A. R., Huang, P. L., Fishman, M. C., Wolf, G. L. and Moskowitz, MA. Temporal correlation mapping analysis of the hemodynamic penumbra in mutant mice deficient in endothelial nitric oxide synthase gene expression. Stroke 1996;27, 1381-1385.

286.Amin-Hanjani, S., Stagliano, NE., Yamada, M., Huang, PL., Liao, JK. and Moskowitz, MA. Mevastatin, an HMG-CoA reductase inhibitor, reduces stroke damage and upregulates endothelial nitric oxide synthase in mice. Stroke 2001;32, 980-986.

287.Suzuki, M., Tabuchi, M., Ikeda, M. and Tomita, T. Concurrent formation of peroxynitrite with the expression of inducible nitric oxide synthase in the brain during middle cerebral artery occlusion and reperfusion in rats. Brain Res. 2002;951, 113-120.

288.Eliasson, MJL., Sampei, K., Mandir, AS., Hurn, PD., Traystman, RJ., Bao, J., Pieper, A., Wang, ZQ., Dawson, TM., Snyder, SH.. Poly(ADP-ribose) polymerase gene disruption renders mice resistant to cerebral ischemia. Nature Med.1997; 3, 1089-1095.

289. Giovannelli, L., Cozzi, A., Guarnieri, I., Dolara, P. and Moroni, F. Comet assay as a novel approach for studying DNA damage in focal cerebral ischemia: differential effects of NMDA receptor antagonists and poly(ADP-ribose) polymerase inhibitors. J. Cereb. Blood Flow Metab.2002; 22, 697-704.

290. Mandir, AS., Poitras, MF., Berliner, AR., Herring, WJ., Guastella, DB., Feldman, A., Poirier, GG., Wang, ZQ., Dawson, TM. and Dawson, VL. NMDA but not non-NMDA excitotoxicity is mediated by Poly(ADP-ribose) polymerase. J. Neurosci. 2000;20, 8005-8011.

291.Yang, J., Klaidman, LK., Chang, ML., Kem, S., Sugawara, T., Chan, P. and Adams, JD. Nicotinamide therapy protects against both necrosis and apoptosis in a stroke model. Pharmacol. Biochem. Behav.2002; 73, 901-910.

292.Britigan, BE., Roeder, TL. and Buettner, G. R. Spin traps inhibit formation of hydrogen peroxide via the dismutation of superoxide: implications for spin trapping the hydroxyl free radical. Biochim. Biophys. Acta 1991;1075, 213-222.

293.Marshall, JWB., Duffin, KJ., Green, R. and Ridley, RM. NXY-059,a free radical-trapping agent, substantially lessens the functional disability resulting from cerebral ischemia in a primate species. Stroke2003;32, 190-198.

294.Yoshimoto, T., Kristian, T., Hu, B., Ouyang, H.-B. and Siesjo, BK. NYX-059 maintains Akt activation and inhibits release of cytochrome c after focal cerebral ischemia. Brain Res. 2002;947, 191-198.

295. Shimizu-Sasamata M., Kano T., Rogowska J, Wolf GL., Moskowitz MA., Lo EH. YM872, a highly water-soluble ampa receptor antagonist, preserves the hemodynamic penumbra and reduces brain injury after permanent focal ischemia in rats. Stroke. 1998;29:2141-2148.

296.Liu F, Kim JK, Li Y, Liu XQ, Li J, Chen X. An extract of Lagerstroemia speciosa L. has insulin-like glucose uptake–stimulatory and adipocyte differentiation–inhibitory Activities in 3T3-L1 cells. J. Nutr. 2001; 131: 2242–2247.

297.Lopchack PA, et al., Novel MMP inhibitor, batimastat, on BBB of mice. Stoke 2000;31:3034-3035.

298. Rossouw JE., Anderson GL., Prentice RL., LaCroxie AZ., Kooperberg C., Stefanick ML., Jackson RD., Beresford SA., Howard BV., Johnson KC., Kotchen JM., Ockene J. Risk and benefits of estrogen plus progestin in healthy postmenopausal women: principla results from Women's Health Initiative randomized controlled trial. JAMA 2002; 288:321-333.

299.Viscoli CM., Brass LM., Kernan WN., Sarrel PM., Suissa S., Horwitz RI. A clinical trial of estrogen-replacement therapy after ischemic stroke. N Engl J Med 2001; 345:1243-1249.

300.Sitruk-Ware, R.Progestins and cardiovascular risk markers. Steroids 2000.65, 651–658.

301.Delyani, J.A., Murohara, T., Nossuli, T.O., Lefer, A.M. Protection from Myocardial Reperfusion Injury by Acute Administration of 17b-Estradiol. J Mol and Cell Cardiol 1996.28, 1001–1008.

302.Stampfer, MJ., Coldifz, GA., Willett, WC. Post-menopausal estrogen therapy and cardiovascular disease. Ten-year follow-up from the nurses' health study. Nengl J Med 1991;325, 756-762.

303.Behl, C. Oestrogen as a neuroprotevctive hormone.Nature Rev 2002;3, 433-442.

304.Roof, R.L., Duvdevani, R., Heyburn, J.W., Stein, S.G. Progesterone rapidly decreases brain edema: treatment delayed up to 24 hours is still effective. Exptal Neurol 1996;138, 246-251.

305. Zhai, P., Eurell, E.T., Cottaus, R., Jeffery, E.H., Bahr, J.M., Gross, D.R., 2002. Effect of estrogen on global myocardial ischemia-reperfusion injury in female rats. The American Journal of Physiology-Heart and Circulatory Physiology 279, H2766-2775.

306.Roof RL., Duvdevani R., Stein DG., Gender influences outcome of brain injury; progesterone plays a protective role. Brain Res 1993.607,333-336.

307. Schumacher M., Guennoun R., Robert F. Local synthesis and dual actions of progesterone in the nervous system: neuroprotection and myelination. Growth Hormone IGF Research 2004;14, S18-S33.

308.Bramlett HM. Sex differences and the effect of hormonal therapy on ischemic brain injury. Pathophysiology 2005;12:17-27.

309.Butterworth J., Prielipp J.R., Cerese J., Livingston J., Burnett D, 2000. Female gender associates with increased duration of intubationand length of stay after coronary artery surgery. CABG clinical benchmarking database participants. Anesthesiology 2000; 92, 414 –424.

310.Abramov D., Tamariz MG., Sever JY. The influence of gender on the outcome of coronary artery bypass surgery. Ann Thorac Surg 2000;70,800 –805.

311.Carswell HV., Bingham D., Wallace K., Nilsen M., Graham DI., Dominiczak AF., Macrae IM. Diffenerntial effects of 17 beta-estradiol upon stroke damage in stroke prone and normotensive rats. J Cereb Blood Flow Metab 2004; 24:298-304.

312. Adashi EY, Hennebold JD. Mechanisms of disease: single-gene mu-tations resulting in reproductive dysfunction in women. N Engl J Med. 1999;340:709 –718.

313. Hall P.F. Role of cytochromes P-450 in the biosynthesis of steroid hormones,Vitam.Horm. 1985;42:315-368

314.Stocco, DM. Clinical disorders associated with abnormal cholesterol transport:mutations in the steroidogenic acute regulatory protein,Mol.Cell.Endocrinol. 2002;191:19 –25.

315.Waterman MR. Steroidogenesis,StAR and PBR:is there light at the end of the tunnel?,Proc.Soc.Exp.Biol.Med. 1998;217:121-122.

316.Miller WL., Strauss JF. Molecular pathology and mechanism of action of the steroidogenic acute regulatory protein,StAR. ,J.Steroid Biochem.Mol.Biol. 1999;69:131-141.

317. Luu-The V., Takahashi M., deLaunoit Y., .Dumont,Y.achance,F.Labrie,Evidence for distinct dehydrogenase and isomerase sites within a single 3 beta-hydroxysteroid dehydrogenase/5-ene-4-ene isomerase protein,Biochemistry 1991; 30: 8861-8865.

318.Gutai JP., Meyer WJ., Kowarski AA., Migeon CJ. Twenty-four hour integrated concentrations of progesterone,17-hydroxyprogesterone and cortisol in normal male subjects. J.Clin.Endocrinol.Metab. 1977;44:116-120.

319. Boomsma D, Paoletti J. A review of current research on the effects of Progesterone. Int. lour Pharm compound 2002;6(4):245-250.

320. Williams, G. D., Palmer, C., Heitjan, D. F. and Smith, M. B. (1992). Allopurinol preserves cerebral energy metabolism during perinatal hypoxia-ischemia: a 31P NMR study in unanesthetized immature rats. Neurosci. Lett.1992;144, 103-106.

321.Santos RL., Abreu GR., Bissoli NS., Moysés MR.Endothelial mediators of 17ß-estradiol-induced coronary vasodilation in the isolated rat heart. Braz J Med Biol Res 2004;37(4),569-575.

322.Hurn PD, Brass LM. Estrogen and Stroke: A Balanced Analysis. Stroke. 2003;34:338-341.

323.Pines A., Bornstein NM, Shapira I. Menopause and ischemic stroke: basic, clinical and epidermiological consideraqtions: The role of hormonal replacement. Hum Rep Update 2002;8:161-168.

324.Chidlow G., Schmidt KG., Wood JPM., Melena J.,. Osborne NN. Alpha-lipoic acid protects the retina against ischemia-reperfusion. Neuropharmacol 2002;43:1015–1025.

325.Packer L, Witt EH, Tritschler HJ. Alpha-lipoic acid as a biological antioxidant. Free Rad Biol Med 1995;19:227-250.

326.Sigel H, Prijs B, McCormick DB, Shih JCH. Stability of binary and ternary complexes of alpha-lipoate and lipoate derivatives with Mn2+, Cu2+, and Zn2+ in solution. Arch Biochem Biophys 1978;187:208-214.

327.Panigrahi M, Sadguna Y, Shivakumar BR, et al. alpha-lipoic acid protects against reperfusion injury following cerebral ischemia in rats. Brain Res 1996;717:184-188.

328.Haramaki N, Assadnazari H, Zimmer G, et al. The influence of vitamin E and dihydrolipoic acid on cardiac energy and glutathione status under hypoxia-reoxygenation. Biochem Mol Biol Int 1995;37:591-597.

329.Hendrickson SC, St. Louis JD, Lowe JE, Abdel-aleem S. Free fatty acid metabolism during myocardial ischemia and reperfusion. Mol Cell Biochem 1997;166: 85-94.

330.Passwater RA. Lipoic Acid: The Metabolic Antioxidant. New Canaan, CT: Keats Publishing, Inc; 1995:1-47.

331.Freisleben H-J. Lipoic acid reduces ischemia-reperfusion injury in animal models Toxicol 2000;148:159–171.

332.Sunamori, M., H. Tanaka, T. Maruyama, I. Sultan, T. Sakamoto, and A. Suzuki. Clinical experience of coenzyme Q10 to enhance intraoperative myocardial protection in coronary artery revascularization. Cardiovasc. Drugs Ther. 1991;5: 297–300.

333.Kagan, V. E., A. Arroyo, V. A. Tyurin, Y. Y. Tyurina, J. M. Villalba, and P. Navas. Plasma membrane NADH-coenzymeQ10 reductase generates semiquinone radicals and recycles vita-min E homologue in a superoxide-dependent reaction. FEBS Lett. 1998;428: 43–46.

334.Ernster, L. Ubiquinol as a biological antioxidant. In: Active Oxygens, Lipid Peroxides and Antioxidants, edited by K. Yagi. Tokyo: Japan Sci. Soc., 1993, p. 1–38.

335.Ernster L, Forsmark P. Ubiquinol: an endogenous antioxidant in aerobic organisms. Clin Invest 1993;71(Suppl):60-5.23. Konishi, T., Y. Nakamura, T. Konishi, and C. Kawai. Accentuated negative inotropism of verapamil after ischaemic interven-tion in isolated working rat heart. Cardiovasc. Res. 1984;19: 38–43.

336.Das, D. K., and N. Maulik. Protection against free radical injury in the heart and cardiac performance. In: Exercise and Oxygen Toxicity, edited by C. K. Sen, L. Packer, and O. Ha¨ nninen. Amsterdam, The Netherlands: Elsevier Science, 1995, p. 359–388.

337.Crestanello JA, KamelgardJ, Lingle D, Mortensen SA, Rhode M, Whitman GJ. Elucidation of a tripartite mechanism underlying the improvement in cardiac tolerance to ischemia by Coenzyme Q10 pretreatment. J Thorac Cardiovasc Surg 1996;111:443-50.

338.Beyer R. An analysis of Coenzyme Q in free radical generation and as an antioxidant. Biochem Cell Biol 1992;70:390-403.

339.Sugawara H, Yamamoto T, Scimizu SI, et al. Inhibition of ubiquinone synthesis in isolated rat heart under ischemic condition. Int J Biochem 1990;22:477-80.

340.Matthews RT., Yang L., Browne S., Baik M., and Flint Beal M. Coenzyme Q_{10} administration increases brain mitochondrial concentrations and exerts neuroprotective effects. PNAS 1998;95:8892-8897.

341.Frei B, Kim M, Ames B. Ubiquinol 10 is an effective lipid soluble antioxidant at physiological concentrations. Proc Natl Acad Sci USA 1987;87:4879-83.31.

342.Kutala VK, Khan M, Mandal R, Ganesan LP, Tridandapani S, Kalai T, Hideg K, Kuppusamy P. Attenuation of Myocardial Ischemia-reperfusion Injury by Trimetazidine Derivatives Functionalized with Antioxidant Properties. J. Pharmacol. Exp. Ther. 2006; 317(3): 921-8.

343.Lopaschuk GD, Barr R, Thomas PD, Dyck JR. Beneficial effects of trimetazidine in ex-vivo working ischemic hearts are due to a stimulation of glucose oxidation secondary to inhibition of long-chain 3-ketoacyl coenzyme a thiolase. Circ. Res. 2003; 93: e33-e37.

344.Kantor PF, Lucien A, Kozak R, Lopaschuk GD. The antianginal drug trimetazidine shifts cardiac energy metabolism from fatty acid oxidation to glucose oxidation by inhibiting mitochondrial long-chain 3-ketoacyl coenzyme A thiolase. Circ. Res. 2000; 86:580-8.

345.Guarnieri C. and Muscari C. Effect of trimetazidine on mitochondrial function and oxidative damage during reperfusion of ischemic hypertrophied rat myocardium. Pharmacology. 1993; 46:324-31.

346.Mironova OP, Zarubina IV, Krivoruchko BI, Smirnov AV. Antioxidant effects of amtizol and trimetazidine in brain ischemia. Patol. Fiziol. Eksp. Ter. 2001;7(3):13-6.

347.Levy DE, Brierly JB. Communication between vertebro-basilar and carotid arterial circulations in the gerbils. Exp. Neurol. 1974; 45: 503-8.

348.Maupoil V, Rochette L, Tabard A, Clauser P, Harpey C. Direct measurement of free radical generation in isolated rat heart by electron paramagnetic resonance spectroscopy. Effects of Trimetazidine. Adv Exp Med Biol 1990;264:373-6.

349.Harpey C, Clausen P, Labrid C, Freyria JL, Poirier JP. Trimetazidine, a cellular anti-ischemic agent, Cardiovasc Drug Rev 1989; 6:292-312.

350. Kay L, Finelli C, Aussedat J, Guarnieri C, Rossi A. Improvement of long term preservation of the isolated arrested rat heart by Trimetazidine: Effects on the energy state and mitochondrial function. Am. J. Cardiol. 1995; 76: 45B-49B.

351.Yamaguchi Y, Yamada K, Yoshikawa N, Nakamura K, Haginaka J, Kunitomo M. Corosolic acid prevents oxidative stress, inflammation and hypertension in SHR/NDmer-cp rats, a model of metabolic syndrome. Life Sciences. 2006; 79:2474-2479.

352.Liu F, Kim JK, Li Y, Liu XQ, Li J, Chen X. An extract of Lagerstroemia speciosa L. has insulin-like glucose uptake–stimulatory and adipocyte differentiation–inhibitory Activities in 3T3-L1 cells. J. Nutr. 2001; 131: 2242–2247.

353. Hattori K., Sukenobu N, Sasaki T, Takasuga S, Hayashi T, Kasai R, Yamasaki K, Hazeki O. Activation of Insulin Receptors by Lagerstroemin. J Pharmacol Sci. 2003; 93: 69 – 73.

354.Tienfenbacher CP, Kaptiza J, Dietz V, Lee CH, Niroomand F. Reduction of myocardial infarct size by fluvastatin. The American Journal of Physiology-Heart and Circulatory Physiology. 2003; 285: H59-H64.

355.Shinghala JR and Balaraman R. Antihypertensive effect of 5-HT1A agonist buspirone and 5HT2B antagonist in experimentally induced hypertension in rats. Pharmacology. 2005; 73(3):129-139.

356.Altavilla D, Squadrito F, Ioculano M et al. E- selectin in the pathogenesis of experimental myocardial reperfusion injury. Eur J Pharmacol 1994; 270(1): 45-51.

357.Alkayed, N.J., Harukuni, I., Kimes, A.S., London, E.D.,Traystman, R.J., Hurn, P.D., 1998. Gender-Linked Brain Injury in Experimental Stroke. Stroke 29,159-166.

358.Longa EZ, Weinstein PR, Carlson S, Cummins R. Reversible middle cerebral artery occlusion without craniectomy in rats. Stroke. 1989; 20(1): 84-91.

359.Maier M, Ahern K, Cheng M, Lee J, Yenari M, Steinberg G. Optimal depth and duration of mild hypothermia in a focal model of transient cerebral ischemia. Stroke. 1998;29:2171-80.

360.Slater TF, Sawyer BC. The stimulatory effect of carbon tetrachloride and other halogenoalkanes or peroxidative reactions in rat liver fractions in vitro. Biochemical Journal 1971; 123: 805-814.

361.Misra HP, Fridovich L. The role of superoxide anions in the auto-oxidation of epinephrine and simple assay for superoxide dismutase. Journal of Biological Chemistry. 1972; 247: 3170-3175.

362.Aebi H. Catalase in vitro. Methods in Enzymology. 1984; 105: 121-26.

363.Moron MS, Depierre JW, Mannervik B. Levels of glutathione, glutathione reductase and glutathione S-transeferase activities in rat lung and liver. Biochimica et Biophysica Acta. 1979; 582: 67-78.

364.Lowery OH, Rosenbrough NJ, Farr Al, Randall RJ. Protein measurement with the Folin phenol reagent. Journal of Biological Chemistry. 1951; 193: 265-75.

365.Bradley PP, Priebat DA, Christensen RD, Rothstein G. Measurement of cutaneous inflammation: estimation of neutrophil content with an enzyme marker. The Journal of Investigative Dermatology. 1982; 78: 206-209.

366.Simonian NA, Getz RL, Leveque JC, Konardi C, Coyle JT. Kainic acid induced apoptosis in neurons. Neuroscience. 1996; 75: 1047-1055.

367.Gursoy-Ozdemir Y, Bolay H, Saribas O, Dalkara T. Role of endothelial nitric oxide generation and peroxynitrite formation in reperfusion injury after focal cerebral ischemia. Stroke. 2000; 31: 1974- 81.

368.Das DK, Cordis GA, Rao PS, Liu XK, Maity S. High-performance liquid chromatographic detection of hydroxylated benzoic acids as an indirect measure of hydroxyl radical in heart: its possible link with the myocardial reperfusion injury. Journal of Chromatography 1991;536: 273-282.

369.Shankar RA, Hideg K, Zweier JL, Kuppusamy P. Targeted antioxidant properties of N-[(tetramethyl-3-pyrroline-3-carboxamido) propyl]phthalimide and its nitroxide metabolite in preventing postischemic myocardial injury. Journal of Pharmacology and Experimental Therapeutics 2000; 292: 838-845.

370. Xu, K.Y., Zweier, J.L., Becker, L.C., 1997. Hydroxyl radical inhibits sarcoplasmic reticulum Ca2+ -ATPase function by directattack on the ATP binding site. Circulation Research 80, 76–81.

371.Berne, R. M. Cardiac nucleotides in hypoxia: possible role in regulation of coronary blood flow. Am. J. Physiol. 204: 317–322,1963.

372.Berne, R. M., J. R. Blackmon, and T. H. Gardner. Hypoxemia and coronary blood flow. J. Clin. Invest. 36: 1101–1106, 1957.

373.Nuutinen, E. M., K. Nishiki, M. Erecinska, and D. F. Wilson. Role of mitochondrial oxidative phosphorylation in regulation of coronary blood flow. Am. J. Physiol. 243 (Heart Circ. Physiol. 12):H159–H169, 1982.

374.Rubio, R., and R. M. Berne. Regulation of coronary blood flow. Prog. Cardiovasc. Dis. 18: 105–122, 1975.

375.Wang, Q.D., Pernow, J., Sjoquist, P., Ryden, L., 2002. Pharmacological possibilities for protection against myocardial reperfusion injury. Cardiovascular Research 55,25-37.

376.Dhalla NS, Elmoselhi AB, Hata T, Makino N. Status of myocardial antioxidants in ischemia–reperfusion injury. Cardiovasc Res 2000; 47: 446-56.

377.Watts DC. Creatine kinase (adenosine 5'-triphosphate-creatine phosphotransferase). In: Boyer PD, ed. The Enzyme. New York, NY: Academic Press 1973. 8, 383-455.

378.Mannign AS, Coltant DJ. Ischemia and reperfusion-induced arrhythmias in the rat: effects of xanthine oxidase inhibition with allopurinol.Circ Res 1984;55:545 –8.

379.Merts TE, Kaplan HR. Pirmenol hydrochloride (CI-845) and reference antiarrhythmic agents: effects on early ventricular arrhythmias afteracute coronary artery ligation in anesthetized rats. J Pharmacol Exp Ther 1982;223:580– 6.

380.Zweier JL. Measurement of superoxide-derived free radicals in the reperfused heart: evidence for a free radical mechanism of reperfusion injury. J Biol Chem 1988;263:1353 – 7.

381.Brooks WW, Conarad CH, Morgan JP. Reperfusion induced arrhythmias following ischemia in intact rat heart: role of intracellular calcium. Cardiovasc Res 1995;29:536– 42.

382. Cheng XS, Shimokawa H, Momii H, et al. Role of superoxide anion in the pathogenesis of cytokine-induced myocardial dysfunction in dogs in vivo. Cardiovasc Res 1999;42: 651-9.

383. Fujiwara H, Onodera T, Tanaka M, et al. Acceleration of cell necrosis following reperfusion after ischemia in the pig heart without collateral circulation. Am J Cardiol 1989;63:14E– 8E.

384.Garcia-Dorado D, Theroux P, Duran JM, et al. Selective inhibition of the contractile apparatus. A new approach to modification of infarct size, infarct composition, and infarct geometry during coronary artery occlusion and reperfusion. Circulation 1992;85:1160–74.

385.Karas RH, Patterson BL, Mendelsohn ME. Human vascular smooth muscle cells contain functional estrogen receptor. Circulation 1994; 89:1943–50.

386.Mendelsohn ME, Karas RH. The protective effects of estrogen on the cardiovascular system. New Engl J Med 1999;340(23):1801–11.

387.The writing group for the PEPI trial. Effects of androgen or estrogen/progestin regimens on heart disease risk factors in postmenopausal women: The Postmenopausal Estrogen/Progestin (PEPI) Trial.JAMA 1995;273:199 –208.

388.Bradley DD, Wingerd J, Petitti DB, Krauss RM, Ramcharan S. Serum high-density-lipoprotein cholesterol in women using oral contraceptives, estrogens and progestins. N Engl J Med 1978;299:17–20.

389.Vakeva AP, Agah A, Rollins SA. Myocardial infarction and apoptosis after myocardial ischemia and reperfusion: role of the acute myocardial terminal complement components and inhibition by anti-C5 therapy. Circulation 1998; 97: 2259-2267.

390.Fliss H, Gattinger D. Apoptosis in ischemic and reperfused rat myocardium. Circulation Research 1996; 79: 949-956.

391.Biewenga, G.P., Haenen, G.R., Bast, A., 1997. The pharmacology of the antioxidant lipoic acid. General Pharmacology 29, 315–331.

392. Freisleben, H.-J., Kriege, H., Clarke, C., Beyersdorf, F., Zimmer,G., 1991. Hemodynamic and mitochondrial parameters during hypoxia and reoxygenation in working rat hearts. Arzneim. Forsch.:Drug Res. 41, 81–88.

393.Scartezzini P and Speroni E. Review on some plants of Indian traditional medicine with antioxidant activity. J of Ethanopharmacol 2000; 71:23-43.

394.Freisleben, H.J. Lipoic acid reduces ischemia-reperfusion injury in animal models. Toxicology 2000;148, 159–171.

395.Gottlieb, R.A., Burleson, K.O., Kloner, R.A., Babior, B.M., Engler, R.L. Reperfusion injury induces apoptosis in rabbit cardiomyocytes. The Journal of Clinical Investigations 1994; 94,1621-1628.

396.Gottlieb, R.A., Giesing, H.A., Engler, R.L., Babior, B.M., The acid deoxyribonuclease of neutrophils: a possible participant in apoptosis-associated genome destruction. Blood1995; 86, 2414-2418.

397.Galang, N., Sasaki, H., Maulik, N., 2000. Apoptotic cell death during ischemia / reperfusion and its attenuation by antioxidant therapy. Toxicology 148, 111-118.

398.Becker LB. New concepts in reactive oxygen species and cardiovascular reperfusion physiology. Cardiovas Resh 2004;61:461– 470.

399.Valverde MA, Rojas P, Amigo J, Cosmelli D, Patricio O, Bahamonde MI, Mann GE, Vergara C & Latorre R.. Acute activation of maxi-K channels (hSlo) by estradiol binding to the ß subunit. Science 1999;285: 1929-1931.

400.Nakajima T, Kitazawa T, Hamada E, Hazama H, Omata M & Kurachi Y. 17ß-Estradiol inhibits the voltage-dependent L-type Ca 2+currents in aortic smooth muscle cells. Euro Jourof Pharmacol 1995,294: 625-635.

401.Nagao T, Illiano S & Vanhoutte PM. Heterogenous distribu- tion of endothelium-dependent relaxation resistant to N G -nitro-L-arginine in rats. Am J Physiol, 1992; 263: H1090-H1094.

402.Cohen RA, Plane F, Najibi S, Huk I, Malinski T & Garland CJ. Nitric oxide is the mediator of both endothelium-dependent relaxation and hyperpolarization of the rabbit carotid artery. Proc of the Nat Acad Sci, USA,1997; 94: 4193-4198.

403.Garland CJ, Plane F, Kemp BK & Cocks TM. Endothelium-dependent hyperpolarization: a role in the control of vascular tone. Trends in Pharmacol Sci 1995;16: 23-30.

404. Weglicki WB, Mak IT. Antioxidant drug mechanism: transition metal-binding and vasodilation. Mol Cell Biochem 1991;1118: 105–111.

405.Wang M., Baker L, Tsai BM., Meldrum, KK., Meldrum, DR. Sex differences in the myocardial inflammatory response to ischemia-reperfusion injury. The Am Jourof Physiol- Endocrinol and Metab 2005; 288, E321–E326.

406.Jain M., Liao R., Podesser BK., Ngoy S., Apstein CS., Eberli, FR. Influence of gender on the response to hemodynamic overload after myocardial infarction. The American Journal of Physiology- Heart Circulatory Physiology 2002; 283, H2544–H2550.

407.Cannon, J.G. and Dinarello, C.A., 1985. Increased plasma interleukin-1 activity in women after ovulation. Science 227, 1247-1249.

408.Gandolfo, M.T., Verzola, D., Salvatore, L., Gianiorio, G., Procopio, V., Romagnoli, A., Giannoni, M., Garibotto, G.. Gender and progression of chronic renal diseases: does apoptosis makes a difference. Italian Jour of Urolo and Nephrol 2006;56, 1-14.

409.Makai K, Daifuku K, Yokoyama S, Nakano M, Stopped-flow investigation of antioxidant activity of estrogens in solution. Biochim Biophys Acta 1990;1035: 348– 352.

410.Taggart DP, Jenkins M, Hooper J, Hadjinikolas L, Kemp M, Hue D, Bennett G. Effects of short-term supplementation with coenzyme Q10 on myocardial protection during cardiac operations. Ann Thorac Surg 1996;62(4):1243-1244.

411.Chaplin ER., Free RG. and Goldstein GW. Inhibition by steroids of the uptake of potassium by capillaries isolated from rat brain, Biochem. Pharm. 1981 ; 30 :241-245.

412.Wolff JEA., Laterra J. and Goldstein GW. Steroid inhibition of neural microvessel morphogenesis in vitro: Receptor mediation and astroglial dependence, J. Neurochem., 58 (1992) 1023-32.

413.Muller, U., Krieglstein, J. Prolonged pretreatment with alpha-lipoic acid protects cultured neurons against hypoxic, glutamate-,or iron-induced injury. Jour Cereb Blood Flow and Metab , 1995.15, 624–630.

414.Zhao, W.; Richardson, J. S.; Mombourquette, M. J.; Weil, J. A.; Ijaz, Z.; Shuaib, A. Neuroprotective effects of hypothermia and U-78517F in cerebral ischemia are due to reducing oxygen-based free radicals: An electron paramagnetic resonance study with gerbils. J. Neurosci. Res. 1996;45:282–288.

415.Grieb, P.; Ryba, M. S.; Sawicki, J.; Chrapusta, S. J. Oral coenzyme Q10 administration prevents the development of ischemic brain lesions in a rabbit model of symptomatic vasospasm. Acta Neuropathol.1997;94:363–368.

416.Taylor, J.; Gatchalian, C. L.; Keen, G.; Rubin, L. L. Apoptosis in cerebellar granule neurones: Involvement of interleukin-1 b converting enzyme-like proteases. J. Neurochem. 1997;68:1598 – 1605.

417. Reed, J. C.; Jurgensmeyer, J. M.; Matsuyama, S. Bcl-2 family proteins and mitochondria. Biochim. Biophys. Acta 1998;1366:127–137.

418.Cai, J.; Yang, J.; Jones, D. P. Mitochondrial control of apoptosis: The role of cytochrome c. Biochim. Biophys. Acta 1998;1366:139 –149.

419.Atlante, A.; Gagliardi, S.; Marra, E.; Calissano, P. Neuronal apoptosis in rats is accompanied by rapid impairment of cellular respiration and is prevented by scavengers of reactive oxygen species. Neurosci. Lett. 1998;245:127–130.

420.Toran-Allerand, C.D. Novel mechanisms of estrogen action in the developing brain: role of steroid–neurotrophin intereactions. In Neurosteroids: A New Regulatory Function in theNervous System (Beaulieu, E-E. et al., eds), 1999; pp. 293–316, Humana Press.

421.Gumuslu S, Serteser M, Ozben T, Balkan S, Balkan E. Inhibitory role of N omega-nitro-L-arginine methyl ester (L-NAME), a potent nitric oxide synthase inhibitor, on brain malondialdehyde and conjugated diene levels during focal cerebral ischemia in rats. Clinical. Chim. Acta. 1997; 267(2): 213-23.

422.Serteser M, Ozben T, Gumuslu S, Balkan S, Balkan E. Lipid peroxidation in rat brain during focal cerebral ischemia: prevention of malondialdehyde and lipid conjugated dienes production by a novel antiepileptic, lamotrigine. Neurotoxicology. 2002;23(1):111-9.

423.Iqbal S, Baziany A, Hussain M, James S, Wright S, Zini R, Simon N, Morin C, d'Athis, P, Tillement JP. Inhibition of rat cerebral mitochondrial respiration by cyclosporins A, D and G and restoration with trimetazidine. C. R. Acad. Sci. 1996; III: 1087–92.

424.Goodman, Y. et al. Estrogens attenuate and corticosterone exacerbates excitotoxicity, oxidative injury, and amyloid beta-peptide toxicity in hippocampal neurons. J. Neurochem. 1996;66, 1836–1844 35.

425.Garcia-Segura, L.M. et al. Steroid effects on brain plasticity: role of glial cells and trophic factors. In Neurosteroids: A New Regulatory Function in the Nervous System (Beaulieu, E-E.et al., eds), pp. 255–268, Humana Press regulatory system that affects the brain. J. Neurobiol 1999;40, 484–496.

426.Rusa, R. et al. 17 beta-estradiol reduces stroke injury in estrogen-deficient female animals. Stroke1999; 30, 1665–1670.