Figure no.	TITLE	Page no.

Figure 57. Knockdown of $cFLIP_L$ induces p53 dependent ubiquitination of Beclin-1. 110
Figure 58. Knockdown of $cFLIP_L$ mitigates TNF-α response for cell survival. 111
Figure 59. Knockdown of $cFLIP_L$ induces p53 dependent ubiquitination of Beclin-1. 112
Figure 60. FADD regulates proIL-1β 114
Figure 61. FADD supresses Inflamasome activation. 115
Figure 62. Isolation of total RNA and cDNA synthesis. 116
Figure 63. Cloning of human FADD. 117
Figure 64. Expression of pET-FADD in *E. coli* BL21 (*DE3*)-pLys cells 118
Figure 65. Purification of His tagged human FADD (hFADD). 119
Figure 66. Characterization of purified FADD. 121
Figure 67. Mass determination of purified FADD by MALDI-TOF mass spectrometry. 121
Figure 68. Intrinsic fluorescence spectrum. 122
Figure 69. FT-IR analysis of CP-FADD 123
Figure 70. Cellular localization of CP-FADD inside the cells 124
Figure 71. Apoptotic potential of CP-FADD. 125
Figure 72. Anti-inflammatory potential of CP-FADD. 126
Figure 73. Comparative analysis of CP-FADD with conventional apoptotic inducers. 127

LIST OF TABLES

Table no.	TITLE	Page no.

Table 1. RT-qPCR primers. Following primers were used in this study. 61
Table 2. Quantitative estimation of RNA and cDNA. 116
Table 3. Quantification of RE digested insert and vector. 117
Table 4. IPTG dose standardization and induction profile. 118
Table 5. Purification yield of human FADD (hFADD) protein. 119
Table 6. Preliminary quantification of purified hFADD. 120
Table 7. Quantitative analysis of CP-FADD. 122