

LIST OF FIGURES

Chapter-1

| | |
|---|----|
| Figure 1. 1: Female reproductive organs (Hall 2010) | 1 |
| Figure 1. 2: Functional anatomy and developmental changes in adult ovary during an ovarian cycle. (Freeman 2001). | 4 |
| Figure 1. 3: Hypothalamus-pituitary ovarian axis (Erickson et al. 1985)..... | 4 |
| Figure 1. 4: Developmental stages of ovarian follicle. (Erickson et al. 1985)..... | 4 |
| Figure 1. 5: Steroidogenic pathway in the ovary, modified from. (Leão and Esteves 2014)..... | 7 |
| Figure 1. 6: The two cell two gonadotropin theory of ovarian steroidogenesis (Will et al. 2001; Boron and Boulpaep 2012)..... | 8 |
| Figure 1. 7 : The apoptotic pathway in a cell..... | 10 |
| Figure 1. 8: The negative feedback along the HPO axis ensures that steroidogenic signaling is tightly controlled. (Restuccia et al. 2012)..... | 11 |
| Figure 1. 9: Structure of insulin receptor, type I IGF receptor and type II IGF receptor (Livingstone 2014)..... | 14 |
| Figure 1. 10: Insulin and IGF signalling pathways. (White 2002)..... | 16 |
| Figure 1. 11: Cross talk between gonadotropins and insulin in granulosa cells. (Leung and Adashi 2003) | 17 |
| Figure 1. 12: Overview of energy metabolism during key developmental events in mammalian folliculogenesis modified from (Trounson et al. 2013). | 18 |
| Figure 1. 13: Pictorial explanation of normal cell and insulin resistant cell. Insulin resistant cell have less insulin receptors, excess free-floating insulin and glucose..... | 22 |
| Figure 1. 14: Mechanism of insulin resistance (Biddinger and Kahn 2006) | 24 |
| Figure 1. 15: Pictorial representation of cysts in PCOS as compared to normal ovary..... | 26 |
| Figure 1. 16: Overall signaling relation between Insulin resistance and PCOS..... | 26 |

Chapter-4

| | |
|--|----|
| Figure 4. 1: Effect of Cd, Dexamethasone and Dexamethasone + Cd treatment on (A) OGTT profile, (B) serum insulin levels, (C) FIRI and (D) protein expression of insulin receptor in granulosa cell. | 66 |
| Figure 4. 2: Effect of Dexamethasone and Cd alone or in combination on estrous cyclicity of rats..... | 67 |
| Figure 4. 3: Histological assessment of ovarian structure in Control, Cd, IR and IR+Cd treated rats. | 68 |
| Figure 4. 4: Effect of IR and Cd on % cell viability of luteinized granulosa cells of rats. | 68 |
| Figure 4. 5: Effect of IR and Cd either alone or in combination on cell death parameters. (A) mRNA expression of Bax and Bcl2 was done by semi-quantitative RT-PCR. (B). Densitometric analysis of Bax (C) Densitometric analysis of Bcl2.. | 69 |
| Figure 4. 6: Effect of IR and Cd either alone or in combination on protein expression of PARP and caspase-3. (A) western blot images of PARP and caspase-3. (B). Densitometric analysis of Activated caspase-3. (C) Densitometric analysis of Cleaved PARP-F2. | 70 |
| Figure 4. 7: Effect of Cd, IR alone and in combination on Annexin V/PI staining in granulosa cells. Red color indicates presence of PI, green color indicates Annexin V staining. | 70 |
| Figure 4. 8: Effect of IR and Cd either alone or in combination on serum (A) estradiol and (B). progesterone levels. (C) E2/P4 ratio. | 71 |

| | |
|--|----|
| Figure 4. 9: Effect of IR and Cd either alone or in combination on (A) mRNA expression of genes involved in steroidogenesis (B) densitometric analysis for mRNA (C) protein expression of genes involved in steroidogenesis (D) densitometric analysis for protein. | 71 |
| Figure 4. 10: Effect of IR and Cd either alone or in combination on A. 17 β -HSD activity and B. 3 β -HSD activity in rat luteinized granulosa cells. | 73 |
| Figure 4. 11: Effect of IR and Cd either alone or in combination on gonadotropin receptors. mRNA of FSH-R and LH-R was analysed by real time PCR. (A) FSH-R. (B) LH-R. | 73 |
| Figure 4. 12: Effect of IR and Cd either alone or in combination on genes involved in regulation of steroidogenesis. mRNA expression of GnRH, CYP19A1, FSH- β and LH- β was done by real time PCR. A. CYP19A1. B. GnRH. C. FSH- β D. LH- β . E. Ratio of LH/FSH. | 74 |
| Figure 4. 13: Schematic diagram summarizing potential effect of Cd on dexamethasone induced IR rat model | 80 |

Chapter-5

| | |
|---|----|
| Figure 5. 1: Schematic diagram of IVF. After the retrieval of oocyte, the remaining follicular fluid is discarded. It is this follicular fluid that is rich in granulosa cells and is used for research purposes. | 86 |
| Figure 5. 2: Characterization of hLGC's isolated from follicular fluid. Blue color indicates presence of nucleus and green color reflects the steroidogenic enzymes. | 88 |
| Figure 5. 3: IR in hLGC's from PCOS samples. A) Western blot of INR- β using β - actin as endogenous control. B) Densitometry analysis. | 88 |
| Figure 5. 4: Cd concentration as estimated in control and PCOS by AAS in few representative samples. | 89 |
| Figure 5. 5: A. Dose dependent study of Cd on hLGC's viability (in vitro). B. Effect of IR and Cd either alone or in combination on hLGC's viability. | 90 |
| Figure 5. 6: Effect of IR and Cd either alone or in combination on protein expression of A). activated caspase-3 B). Densitometry for activated caspase-3 C). PARP-F2 D) Densitometry for cleaved PARP-F2. | 90 |
| Figure 5. 7: Effect of IR and Cd alone or in combination on Annexin V and PI staining in hLGC's. | 91 |
| Figure 5. 8: Effect of IR and Cd either alone or in combination on steroid hormone concentrations in hLGC cell culture supernatant. A). estradiol and B). progesterone | 92 |
| Figure 5. 9: Effect of IR and Cd either alone or in combination on hLGC's. (A) mRNA expression of genes involved in steroidogenesis (B) protein expression of genes involved in steroidogenesis (C) densitometric analysis for protein. | 92 |
| Figure 5. 10: Effect of Cd and IR either alone or in combination on A. 17 β -HSD and B. 3 β -HSD activity in hLGC's. | 93 |
| Figure 5. 11: Effect of IR and Cd either alone or in combination hLGC's. Effect of IR and Cd either alone or in combination in hLGC's on (A) mRNA expression of FSH-R (B) mRNA expression of LH-R | 93 |

Chapter-6

| | |
|--|-----|
| Figure 6. 1: The diagnostic criteria for PCOS includes two or more of these features. (Diamanti-Kandarakis and Dunaif 2012). | 102 |
| Figure 6. 2: Human luteinized granulosa cell A). Protein expression of INSR- β . B). Densitometry. C). % cell viability. | 105 |
| Figure 6. 3: Expression of insulin signalling proteins in control, PCOS-IR and PCOS-NIR. A) Western blot B) Densitometry | 106 |

| | |
|--|-----|
| Figure 6. 4: Expression of fatty acid metabolism genes in control, PCOS-IR and PCOS-NIR. | 106 |
| Figure 6. 5: Expression of genes and proteins involved in steroidogenesis in control, PCOS-IR, PCOS-NIR. A. mRNA expression B. Western blot study using β - actin as endogenous control C. Densitometry analysis. | 107 |
| Figure 6. 6: Steroid hormone concentration in follicular fluid aspirates. A. estradiol, B. progesterone and C. testosterone. | 108 |
| Figure 6. 7: Expression of FSH-R and LH-R genes in PCOS-IR and PCOS-NIR. | 108 |
| Figure 6. 8: Expression of IGF's and their receptors in PCOS-IR and PCOS-NIR. A. IGF-1 and IGF-1R and B. IGF-2 and IGF-2R | 109 |
| Figure 6. 9: Schematic figure showing difference in signalling between control, PCOS-IR and PCOS-NIR luteinized granulosa cells. | 114 |

Chapter-7

| | |
|--|-----|
| Figure 7. 1: Structure of metformin. | 119 |
| Figure 7. 2: A. Picture of <i>Enicostemma Littorale</i> Blume. B. Molecular structure of swertiamarin. | 120 |
| Figure 7. 3: A. Picture of Curcumin Longa B. Powdered form used as spice. C. Molecular formula of curcumin | 120 |
| Figure 7. 4: Effect of swertiamarin, curcumin and metformin on cell viability in PCOS-IR and PCOS-NIR. The normalized expression values are represented as mean + SEM of three independent experiments. | 123 |
| Figure 7.5: Effect of swertiamarin, curcumin and metformin on expression of proteins involved in insulin signalling in hLGC's. A. western blot from PCOS-IR. B. Densitometric analysis for PCOS-IR. C. western blot for PCOS-NIR. D. Densitometric analysis for PCOS-NIR. | 124 |
| Figure 7. 6: Effect of swertiamarin, curcumin and metformin on mRNA expression of fatty acid metabolism genes in A. PCOS-IR. B. PCOS-NIR. The normalized expression values are represented as mean + SEM of three independent experiments. | 125 |
| Figure 7. 7: Effect of swertiamarin, curcumin and metformin on steroidogenic machinery in hLGC's from PCOS-IR. A. mRNA expression. B. western blot. C. Densitometric analysis. E. Enzyme activity. | 126 |
| Figure 7. 8: Effect of swertiamarin, curcumin and metformin on steroidogenic machinery in hLGC's from PCOS-NIR. A. mRNA expression. B. western blot. C. Densitometric analysis. E. Enzyme activity. | 126 |
| Figure 7. 9: Effect of swertiamarin, curcumin and metformin on enzyme activity in hLGC's from PCOS-IR and PCOS-NIR A. 3β -HSD and B. 17β -HSD. The values are represented as mean + SEM of three independent experiments. | 127 |
| Figure 7. 10: Effect of swertiamarin, curcumin and metformin on steroid hormone levels in hLGC's from. A. PCOS-IR. B. PCOS-NIR. The normalized expression values are represented as mean + SEM of three independent experiments. | 128 |
| Figure 7. 11: Effect of swertiamarin, curcumin and metformin on mRNA expression of A.FSH-R and B. LH-R in hLGC's from PCOS-IR B. and NIR. The values are represented as mean + SEM of three independent experiments. | 128 |
| Figure 7. 12: Effect of swertiamarin, curcumin and metformin on mRNA expression of genes involved in IGF system. A. PCOS-IR. B. PCOS-NIR. The normalized expression values are represented as mean + SEM of three independent experiments. | 129 |
| Figure 7. 13: Schematic diagram summarizing potential effect of swertiamarin, curcumin and metformin in PCOS-IR and PCOS-NIR hLGC's. | 133 |
