

# References

## REFERENCES

- ♣ Abreu PM, Luis MH (1996) Constituents of *Tephrosia uniflora*. *Natural Product Letters* 9:81-86.
- ♣ Acharya L, Mukherjee AK, Pandya PC (2004) Genome relationship among nine species of Millettieae (Leguminosae; Papillonoideae) based on Random Amplified polymorphic DNA (RAPD). *Z.Naturforsch* 59:868-873.
- ♣ Agarwal K, Gupta PK (1983) Cytological studies in the genus *Tephrosia* Pers. *Cytologia* 48:795- 801.
- ♣ Ahluwalia KS and Smith AR (1967) A new species of *Tephrosia* pres. (leguminosae) from India, *Kew Bulletin*, 21(2):311-313.
- ♣ Ahmad VU, Ali Z, Hussaini SR, Iqbal F, Zahid M, Abbas M, Saba N (1999) Flavonoids of *Tephrosia purpurea*. *Fitoterapia*, 70(4):443-445.
- ♣ Ahmedullah M and Nayar MP (1986) *Endemic plant of the Indian region Vol-1*, Pub. in Botanical Survey of India:102.
- ♣ Ali M (2011) *Pharmacognosy (Pharmacognosy and Phytochemistry) Vol 1*, Pubs: CBS Publishers and distributor Pvt. Ltd.:1-32.
- ♣ Andrei CC, Vieira PC, Fernandes JB, da Silva MDG, Fo ER (2002) New spirorotenoids from *Tephrosia candida*. *Zeitschrift. Fur Naturforschung C* 57:418-422.
- ♣ Andrei CC, Vieira PC, Fernandes JB, Da Silva MFGF (1997) Dimethylchromene rotenoids from *Tephrosia candida*. *Phytochemistry* 46:1081-1085.
- ♣ Angiosperm Phylogeny Group (2009) An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants: APG III, *Botanical Journal of the Linnean Society* 161 (2):105–121.
- ♣ Anonymous (2001) *Threatened biodiversity of Gujarat-Baseline information* Pub: Gujarat Ecology Commission Vadodara.
- ♣ Anonymous (2002) *Quality control methods for medicinal plants materials*, Pub: World Health Organization, Geneva. A.I.T.B.S. New Delhi:115.
- ♣ Anonymous (2003) *Rare and endangered plant and animal of Gujarat*, Pub: Gujarat Ecology Commission Vadodara:37–38.
- ♣ Anonymous (2008) *Indian Pharmacopoeia Commission* Pub: Central Indian Pharmacopoeia Laboratory, Govt. Of India Ministry of Health &family Welfare, Raj Nagar, Ghaziabad.

- ♣ Anonymous (2014) *Conservation assessment and Management Plan(CAMP) Workshop*, Pub: Gujarat ecological education and research Foundation:109
- ♣ AOAC (1995) *Official method of analysis of AOAC International*, 16<sup>th</sup> edition AOAC International, suite 400, 2200 Wilson Boulevard, Arlington, Virginia, USA.
- ♣ AOSA (1978) Seed vigor committee report. Vigour test “referee” Program, *Association of official seed analyst Newsletter* 52(4):31-42.
- ♣ Arriaga AM, Magalhães FE, Feitosa EM, Malcher GT, Andrade-Neto M, Nascimento RF (2005) Composition of the Essential Oil of *Tephrosia egregia* Sandw *J Essential Oil Res* 17(4):451-452.
- ♣ Arriaga AMC, Malcher GT, Lima JQ, Magalhães FE, Gomes TM, Conceição M D, Santiago GMP (2008) Composition of the essential oil of *Indigofera microcarpa* from the Northeast of Brazil *J Essential Oil Res* 20(5):450-451.
- ♣ Ashok DB, Thomas PV, Devasagayam A (2007) Current status of herbal drugs in India: An Overview. *J Clin Biochem Nutr. plants: a compendium of 500 species. Health and Fitness* 41(1):1-11.
- ♣ Asuntha GI, Prasannaraju Y, Sujatha D, Prasad K (2010) Assessment of effect of ethanolic extract of *Tephrosia purpurea* (L.) Pers., Fabaceae, activity on lithium-pilocarpine induced Status epilepticus and oxidative stress in Wistar ratsfarmacogn. *Revista Brasileira de Farmacognosia Brazilian Journal of Pharmacognosy* 20 (5):767–772.
- ♣ Atchison E (1951) Studies in the Leguminosae. VI. Chromosome Numbers among Tropical Woody Species. *American Journal of Botany*:538-546.
- ♣ Augustine KT (2000) The seedling morphology of *Tephrosia purpurea* (L) Pers., *T. maxima* (L) Pers., *T. pumila* (Lam) Pers., and *T. villosa* (L) Pers. From Thumba, Kerala, *J. Econ. Taxon. Bot.* 24 (1):243–250.
- ♣ Azimova SS and Yunusov MS (2013) *Natural compounds* Pub: Springer New York: 308.
- ♣ Babayemi OJ, Daniel IO, Bamikole MA, Ogungbesan A, Oduguwa BO (2003) Preliminary studies on *Tephrosia* species: Effect of seed treatments on germination, *Nigerian Journal of Animal Production*, 30 (2):209–216.
- ♣ Baruah PB and Swain T (1959) In anthocyanins the sugar imparts stability to the aglycones. *J. s. Food Agri.* 10:125.

- ♣ Belmain SR, Amoah BA, Nyirenda SP, Kamanula JF, Stevenson PC (2012) Highly variable insect control efficacy of *Tephrosia vogelii* chemotypes. *J Agric Food Chem.* 60(40):10055–63.
- ♣ Bentham G and Hooker JD (1862–1883) *Genera Plantarum, Vols. 1-3.* L. Reeve & Co., London:1862–1883.
- ♣ Berlyn, GP and Miksche, JP (1976) *Botanical Microtechnique and Cytochemistry.* Iowa State University Press, Iowa.
- ♣ Bhandari MM, Gehlot MS, Anand SK (1985) Identification of Indian desert species of *Tephrosia* (L.) Pers. by seed characters. *J Pl Morph* 2:41–50.
- ♣ Bisky F (1994) *Phytochemical dictionary of the leguminosae vol 1* Chapman and Hall, :888
- ♣ Bole PV and Pathak JM (1988) *Flora of Saurashtra. Part II and III.* Botanical Survey of India, Calcutta: 545
- ♣ Bramwell D (1972) *Endemism in the flora of the Canary Islands.* In: Valentine DH (ed) *Taxonomy, Phytogeography and Evolution*, Academic press, London: 141-159
- ♣ Bridson D and Forman L (1992) *The herbarium handbook.* Pub: Royal Botanic Gardens, Kew, London.
- ♣ Brummitt RK (1981) Reconsideration of the genera *Ptychobium*, *Caulocarpus*, *Lupinophyllum* and *Requena* in relation to *Tephrosia*. *Kew Bull* 35:459–473.
- ♣ Bubna GA, Lima RB, Zanardo DY, Santos WDD, Ferrarese MLL, Ferrarese Fillcho (2011) Exogenous caffeic acid inhibits the growth and enhances the lignification of the roots of soybean (*Glycine max*), *Journal of plant physiology* 168(14):1627-33.
- ♣ Cai YZ, Sun M, Xing J, Luo Q, Corke H (2006) Structure-radical scavenging activity relationships of phenolic compounds from traditional Chinese medicinal plants. *Life Sci.* 78:2872–2888.
- ♣ Catherine A, Rceo-Evans, Miller NJ, Paganga G (1997) Antioxidant properties of phenolic compounds. *Trends plant sci.* 2(4):152-159.
- ♣ Chadha YR (1972) *The wealth of India: raw materials series IX (Rh-So):*151-157.
- ♣ Chang LC, Chávez D, Song LL, Farnsworth NR, Pezzuto JM, Kinghorn AD (2000) Absolute configuration of novel bioactive flavonoids from *Tephrosia purpurea*. *Organic letters* 2:515-518.

- ♣ Chang LC, Gerhäuser C, Song L, Farnsworth NR, Pezzuto JM, Kinghorn AD (1997) Activity guided isolation of constituents of *Tephrosia purpurea* with the potential to induce the phase II enzyme, quinone reductase. *Journal of Natural Products* 60:869–87.
- ♣ Chaudhari TB, Tambe DA, Chaudhari SR (2012) Phytopharmacology of *Tephrosia purpurea* Pers (Fabaceae)-A review. *IJPI's Journal of pharmacology and herbal formulation* 2(8):1-13.
- ♣ Chen Y, Yan T, Gao C, Cao W, Huang R (2014) Natural products from the genus *Tephrosia* *Molecules* 19:1432–1458.
- ♣ Chenin L and Yein CH (1950) Turbidimetri determination of available sulphates *Soil Sci. Soc.Am. Proc.*15:149–151.
- ♣ Chopra GL (1987) "*Angiosperms*", Pradeep Publications. Sitla Mandir, Jalandhar. India.
- ♣ Coe EH, Neuffer MG, Hoisington DA (1988) *The genetics of corn*. In: Corn and corn improvement. American Society of Agronomy, Madison, WI. Eds. Sprague GF, Dudley JW:81.
- ♣ Dagne E, Yenesew A, Gray AI, Waterman PG (2012). Praecansone A: Evidence for the existence of 8, 9-(E) and 8, 9-(Z) isomers in extracts from *Tephrosia pumila*. *Bulletin of the Chemical Society of Ethiopia* 4:141-145.
- ♣ Dai J, Mumper RJ (2010) Plant Phenolics: Extraction, Analysis and Their Antioxidant and Anticancer Properties. *Molecules*. 15: 7313-7352.
- ♣ Damery S (2011) The herbal medicines by people with cancer: a cross sectional survey. *British Journal of cancer* 104:927-933.
- ♣ Damre AS (2003) Studies on the Immunomodulatory activity of flavonoidal fraction of *Tephrosia purpurea*. *Fitoterapia*. 74(3):257-261.
- ♣ Daniel M (1991) *Methods in Plant Chemistry and Economic Botany*, Kalyani Publishers, New Delhi, India.
- ♣ Darbar S, Bhattacharya A, Chakraborty MR, Chattopadhyay S (2010) Livina, a polyherbal preparation protects liver against aceclofenac-induced hepatic insult in sprague-dawley rats: a comparison with silymarin. *Pharmacologyonline* 2:889-907.
- ♣ Decandolle AP (1825) *Prodromus systematis naturalis regni vegetabilis*. 2nd Pub: International Plant Names Index. Paris:248–256.
- ♣ Decandolle, AP (1855) *Geographie botanique Raisonnee Vols I &II*. Geneva.
- ♣ Despande SE, Shah GB, Parmar NS (2003) Antiulcer activity of *Tephrosia purpurea* in rats. *Indian journal of Pharmacology* 35:168–172.

- ♣ Dharmalingam C, Rao SM and Sundaraj D (1971) Pre-germination treatment of kolinji seed (*Tephrosia purpurea*) to improve germination. *Seed Res.*1:58–62.
- ♣ Dharmalingam C, Rao SM and Sundaraj D (1973) Pre-germination treatment of kolinji seed (*Tephrosia purpurea*) to improve germination. *Seed Res.*1:58–62.
- ♣ Doyle JJ, Doyle JL (1987) A rapid DNA isolation procedure for small quantities of fresh leaf material. *Phytochemical Bulletin* 19:11–15.
- ♣ Dunnet CW (1964) New tables for multiple comparisons with a control. *Biometrics* 20:482–491.
- ♣ Dutt SK, Chibber SS. (1983). Candidol, a flavonol from *Tephrosia candida*. *Phytochemistry* 22:325–326.
- ♣ Dzenda T, Ayo JO, Adelaiye AB, Auda AO (2007) Ethnomedical and veterinary uses of *Tephrosia vogelii* Hook F (Fabaceae): a review. *Niger Vet J* 28:24–39.
- ♣ [ensembl-sigenae.jouy.inra.fr/META/NEWAGE?type=ORGANISM&object=TAX-1157238](http://ensembl-sigenae.jouy.inra.fr/META/NEWAGE?type=ORGANISM&object=TAX-1157238)
- ♣ Erakar SR and Murumkar CV (1995) Proline accumulation in *Tephrosia purpurea* Pers, *Biologia Plantarum* 37(2):301–304.
- ♣ European Food Safety Authority (2009) EFSA Compendium of botanicals that have been reported to contain toxic, addictive, psychotropic or other substances of concern, *EFSA Journal* 7(9):281.
- ♣ European Food Safety Authority(2013) Scientific Opinion on safety evaluation of Ephedra species for use in food1 EFSA Panel on Food Additives and Nutrient Sources added to Food (ANS), *EFSA Journal* 11(11):3467.
- ♣ Evans WC (2002) *Trease and Evans Pharmacognosy*, 15 W.B.Saunders; Edinburgh.
- ♣ Fahn A (1967) *Plant anatomy* Pregamon Press in Oxford:534.
- ♣ Falak AH, Shoeb A. (1987) A new epoxyflavanone from *Tephrosia hamiltonii*. *Planta Medica* 53:220–221.
- ♣ Fang N and Casida JE (1998) Anticancer action of cubé insecticide: correlation for rotenoid constituents between inhibition of NADH: ubiquinone oxidoreductase and induced ornithine decarboxylase activities. *Proc Natl Acad Sci U S A.* 31; 95(7):3380–4.
- ♣ Feng Xu, Hongmei W, Xiangpei W, Yang Y, Yuanmin W, Haibing Q, Yanyan Z (2014) RP-HPLC Characterization of Lupenone and  $\beta$ -Sitosterol in Rhizoma Musae and Evaluation of the Anti-Diabetic Activity of Lupenone in Diabetic Sprague-Dawley Rats *Molecules* 19(9):14114 – 14127.

- ♣ Firm R (2010) *Nature's Chemicals*. Oxford University Press, Oxford: 74–75.
- ♣ Gajera PL, Dalal MV (2011). Mast cell stabilizing potential of the ethanolic extract of *Tephrosia purpurea* (Linn.) in the management of asthma. *IJRAP* 2:1308–1312.
- ♣ Ganapaty S, Lakshmi P, Bobbarala V (2010). Chemical and biological examination of the leaves of *Tephrosia tinctoria* Pers. *International Journal of Chemical and Analytical Science* 1:14–1.
- ♣ Ganapaty S, Lakshminarayana K, Lakshmi P, Thomas PS (2009) Antiprotozoal and Cytotoxicity Assays of the Isolates of *Tephrosia tinctoria*. *Asian Journal of Chemistry* 21:1007–1010.
- ♣ Ganapaty S, Nyamathulla S, Srilakshmi GVK, Prasad KVNMR (2008). Chemical and Antimicrobial Studies of the Roots of *Tephrosia villosa* (L) Pers. *Asian Journal of Chemistry* 20:4498–4502.
- ♣ Ganapaty S, Pannakal ST, Srilakshmi GV, Lakshmi P, Waterman PG, Brun R (2008) Pumilanol, an antiprotozoal isoflavanol from *Tephrosia pumila*. *Phytochemistry Letters* 1:175–178.
- ♣ Ganapaty S, Srilakshmi GVK, Pannakal ST, Laatsch H (2008) A pyranochalcone and prenylflavanones from *Tephrosia pulcherrima* (Baker) drumm. *Nat. Prod. Commun.* 3, 49:52.
- ♣ Gennaro AR (1995) Remington: *The science and practice of pharmacy*, 19<sup>th</sup> edition: 111–117.
- ♣ Ghosh AB, Bajaj JC, Hasan R, Singh D (1983) *Soil and water testing Methods-A laboratory Manual*. Pub: Division of soil science and Agricultural chemistry, IARI, New Delhi.
- ♣ Girish C, Koner BC, Jayanthi S, Rao KR, Rajesh B, Padhan SC (2009) Hepatoprotective activity of six polyherbal formulations in paracetamol induce liver toxicity in mice, *Indian J Med Res* 129:569–578.
- ♣ Gokhale AB, Dikshit VJ, Damre VS (2000) Influence of ethanolic extract of *Tephrosia purpurea* Linn. on mast cells and erythrocytes membrane integrity. *Indian J Exp Biol.* 38(80):837–40.
- ♣ Gopalakrishnan S, Vadivel E and Dhanalakshmi K (2009) Phytochemical and Pharmacognostical Studies of “*Tephrosia purpurea*” Linn. Aerial and Root Parts. *Journal of Herbal Medicine and Toxicology* 3 (2):73–78.

- ♣ Gopalakrishnan S, Vadivel E, Dhanalakshmi K (2010) Anti-inflammatory and analgesic activities of *Tephrosia purpurea* Linn. aerial and root extracts *Journal of Pharmacy Research*.3(5):6.
- ♣ Gupta YK, Sharma M, Chaudhary G, Katiyar CK (2004) Hepatoprotective effect of New Livfit, a polyherbal formulation, is mediated through its free radical scavenging activity. *Phytother Res*.18(5):362–4.
- ♣ Han XZ, Shen T, Lou HX (2007) Dietary polyphenols and their biological significance. *Int J Mol Sci*. 8:950–988.
- ♣ Harborne JB (1984) *Phytochemical Methods*, Chapman a hall, London.
- ♣ Harborne JB (1998) *Phytochemical Methods: A Guide to Modern Techniques of Plant Analysis*, 3rd ed.; Chapman and Hall: London,
- ♣ Harborne JB (2007) *Phytochemical Methods- A Guide to Modern techniques of Plant analysis*. Third ed. Springer Private Ltd., Delhi.
- ♣ Hegazy MEF, Mohamed AEHH, El-Halawany AM, Djemgou PC, Shahat AA, Paré PW (2011) Estrogenic activity of Chemical Constituents from *Tephrosia candida*. *Journal of natural products* 74:937–942.
- ♣ Hegazy MF, Abd MH, Nagashima F, Asakawa Y, Pare PW (2009) Rare prenylated flavonoids from *Tephrosia purpurea*, *Phytochemistry* 70(11-12):1474–1477.
- ♣ Hill AF (1979) "*Economic Botany*". Mc Graw Hill, New York.
- ♣ Hisham A, John S, Al-Shuaily W, Asai T, Fujimoto Y (2006) (+)-Apollineanin: a new flavanone from *Tephrosia apollinea*. *Natural Product Research* 20:1046–1052.
- ♣ Hosler. D.M. & Mikita, M.A. (1987) Ethnobotany: the chemist's source for the identification of useful natural products. *Journal of Chemistry Education*. 64:328–332.
- ♣ Hu J, Lavin M, Wojciechowski MF, Sanderson M J (2000) Phylogenetic Systematics Of The Tribe Millettieae (Leguminosae) Based On Chloroplast Trnk/MatK Sequences And Its Implications For Evolutionary Patterns In Papilionoideae, *American Journal of Botany* 87(3):418–430.
- ♣ Hu J, Peng P, Chivas AR (2009) Molecular biomarker evidence of origins and transport of organic matter in sediments of the Pearl River estuary and adjacent *South China Sea*. *Appl. Geochem*. 24: 1666 – 1676.
- ♣ Ibrahim RK, Tower GHN (1960) Studies of Hydrangenol In *Hydrangea macrophylla* Ser.: I. Isolation, Identification, And Biosynthesis From C14-Labelled Compounds *Canadian journal biochemistry and Physiology*. 38(7):627–634.

- ♣ Irvine E, Freyre HR (1959) Occurrence of rotenoids in some species of the genus *Tephrosia*, *Agriculture and Food chemistry* (7):2.
- ♣ Jain A, Lodhi S, Singhai AK (2009). Simultaneous estimation of quercetin and rutin in *Tephrosia purpurea* Pers by high performance thin-layer chromatography. *Asian Journal of Traditional Medicine* 4:104–109.
- ♣ Jain A, Singhai AK, Dixit VK (2006) A comparative study of ethanol extract of *Tephrosia purpurea* Pers and Flavonoids extract for Hepatoprotective, *Indian Journal of Pharmaceutical Science* 28 (6):740–743.
- ♣ Jan TBJ (2007) Structural studies of some natural products form *Kaempferia praviiflora* *Iris Germanica* Linn and *Andrographis paniculata* Nees, Thesis submitted at University Sain Malaysia:63-66.
- ♣ Jayaraman I, Ghanim A, Khan HA (1980) A new prenylated flavanone [7-methylglabranin] from *Tephrosia villosa*. *Phytochemistry*:19.
- ♣ Jin Y, Kim HP, Chi M, Ifedigbo E, Ryter SW, Choi AMK (2008) Deletion of Caveolin-1 Protects against Oxidative Lung Injury via Up Regulation of Heme Oxygenase-1. *Am J Respir Cell Mol Biol* 39:171–179.
- ♣ Johansen DA (1940) *Plant Microtechnique*. Pub: Tata Mc-Graw Hill limited, New Delhi.
- ♣ Joshi SS, Shrivastava RK, Shrivastava DK (1979) Chemical examination of *Tephrosia purpurea* seeds. *Journal of the Indian Chemical Society*, 56(3):303–5.
- ♣ Joshi, JV (1980) A study of Surat and Its Environs. Thesis submitted to The M.S. University of Baroda, Vadodara.
- ♣ Juma WP, Akala HM, Eyase FL, Muiva LM, Heydenreich M, Okalebo FA, Yenesew A (2011) Terpurinflavone: An antiplasmodial flavone from the stem of *Tephrosia purpurea*. *Phytochemistry Letters*, 4:176–178.
- ♣ Kar A (2007) *Pharmaocgnosy and Pharmacobiotechnology* (Revised-Expanded Second Edition). New Age International Limited Publishers New Delhi:332–600.
- ♣ Kashaw V, Nema AK, Agarwal A (2011) Hepatoprotective Prospective Of Herbal Drugs and Their Vesicular Carriers– A Review, *International Journal of Research in Pharmaceutical and Biomedical Sciences* 2(2):360–374.
- ♣ Khalid J, Qadir MI, Jan A, Gilani AH (2013) Anti-diarrheal activity of methanolic extract of *Tephrosia purpurea*. *Acta poloniae pharmaceutica* 70(2):345–7.

- ♣ Khalivulla SI, Reddy BAK, Gunasekar D, Blond A, Bodo B, Murthy MM, Rao TP (2008) A new diO-prenylated isoflavone from *Tephrosia tinctoria*. *Journal of Asian natural products research* 10: 953–955.
- ♣ Khan N, Sharma S, Alam A, (2001) *Tephrosia purpurea* Ameliorates N-Diethylnitrosamine and Potassium Bromate- Mediated Renal oxidative stress and Toxicity in Wistar Rats. *Pharmacology & Toxicology* 88:294–299.
- ♣ Khatria A, Arun G, Shyam S, Agrawal C (2009) Evaluation of hepatoprotective activity of aerial parts of *Tephrosia purpurea* and stem bark of *Tecomella undulate*. *Journal of Ethnopharmacology* 122(1):1–5.
- ♣ Kirkbride JH, Gunn CR, Weitzman AL (2003) *Fruits and seeds of genera in the subfamily Faboideae (Fabaceae), vol 1*. United Seed morphology of *Tephrosia* (Leguminosae): 469,123.
- ♣ Kishore K, Roy D (2011) *Tephrosia purpurea* Pers.(Fabaceae)A common winter weed of Chhattisgarh, India-As a source of anticancer drug. *Indian J.Applied & Pure Bio.1* 26(1):53–55.
- ♣ Kokate CK (1994) *Practical pharmacognosy*, published by M.K. Jain for Vallabh Prakashan, Pitampura, Delhi, India: 152.
- ♣ Kokila AP, Patel AN (2010) Preliminary Phytochemical Screening and study of antiviral activity and antibacterial activity of *Tephrosia purpurea* flower *Life sciences Leaflets* 1:7–13.
- ♣ Kole RK, Satpathi C, Chowdhury A, Ghosh MR, Adityachaudhury N (1992) Isolation of amorpholone, a potent rotenoid insecticide from *Tephrosia candida*. *J. Agric. Food Chem.* 40: 1208–1210.
- ♣ Krupadanam GLD, Srimannarayana G, Rao NVS (1977) New C-6 oxygenated rotenoids from *Tephrosia villosa*—villosin, villosone, villol, and villinol. *Tetrahedron Letters* (24): 2125–8.
- ♣ Kumar GS (2007) Antimicrobial effects of Indian medicinal plants against acne-inducing bacteria. *Tropical Journal of Pharmaceutical Research.* 6(2): 717–723.
- ♣ Kumar KJ, Prasad AGD (2011) Identification and comparison of biomolecules in medicinal plants of *Tephrosia tinctoria* and *Atylosia albicans* By Using Ftir. *Romanian J. Biophys.* 21(1): 63–71.
- ♣ Kumar S, Sane PV (2003) *Legumes of South Asia: a Check list*. Royal Botanical Gardens- Science :543.

- ♣ Lakshmi P, Khan AA, Reddy PN, Lakshminarayana K, Ganapaty S (2008) Genetic Relationship among *Tephrosia* species as revealed by RAPD analysis. *Asian Journal of Biological Sciences* 1 (1):1–10.
- ♣ Lavate SM, Shendkar CD, Deshpande NR (2014) GC-MS Study of a Steam Volatile Matter from *Aglaia lawii* leaves. *International Journal of ChemTech Research* 6(7): 3903–3908.
- ♣ Leon LJG, Bertiller MB (1982) Aspectos fenológicos, de dos comunidals dal pastizal de la depression dal salado (Pgai des Buenos Aires), *Bulletin de la sociadat Argentina Botanic*, 20:320–347.
- ♣ Lewis G, Schrire B, MacKinder B, Lock M (2005) *Legumes of the world*. Royal Botanical Gardens, Kew, UK.
- ♣ Mabberley DL (2008) *Mabberley's Plant-book: A Portable Dictionary of Plants, Their classification and uses*, third edition pub: Cambridge:476.
- ♣ Mabry TJ, Mabry H, Markham H (1970) *The Systematic Identification of Flavonoids*. Springer, Berlin.
- ♣ Madhusudhana J, Reddy RVN, Reddy BAK, Reddy MVB, Gunasekar D, Deville A, Bodo B (2010) Two new geranyl flavanones from *Tephrosia villosa*. *Natural Product Research* 24(8): 743–749.
- ♣ Madlener S, Svacinová J, Kitner M, Kopecky J, Eytner R, Lackner A, Vo TP, Frisch R, Grusch M, De Martin R, Dolezal K, Strnad M, Krupitza G (2009) In vitro antiinflammatory and anticancer activities of extracts of *Acalypha alopecuroidea* (Euphorbiaceae). *Int. J. Oncol.* 35(4): 881–91.
- ♣ Magoma G (2014) Introduction to Biochemical Pharmacology and Drug Discovery In *Drug Discovery*:475–515.
- ♣ Mahajan RT, Chaudhari GS, Chopada MZ (1999) Screening of some indigenous plants for their possible antibacterial activity *Environmental Bulletin*. 15(2):61–62.
- ♣ Manjula RR, Spandana U, Joshi TA, Sudheer M (2013) *In vitro* anthelmintic activity of aqueous and methanolic leaf extract of *Tephrosia purpurea* linn. *International Journal Of Research In Pharmacy And Chemistry* 3(1):12–14.
- ♣ Manoharan S, Kavitha K, Senthil N, Renju GL (2006) Evaluation of anticarcinogenic effects of *Clerodendron inerme* on 7,12-dimethylbenz(a) anthracene-induced hamster buccal pouch carcinogenesis. *Singapore Med J* 47(12):1038–43.

- ♣ Martinez MJA, Lazaro RM, DelOlmo LMB, Benito PB (2008) Anti-infectious activity in the anthemideae tribe. In: Atta-ur- (Ed.) *Studies in Natural Products Chemistry*. 35: 445–516.
- ♣ Mathews RR, Sujith K and Christina AJM (2012) Basic Research on the herb *Tephrosia purpurea* (l) Pers- the Translation Challenges-A Review. *International Journal of Pharmaceutical and Chemical sciences* 1:466-46.
- ♣ Misra KC (1980) *Manual of Plant Ecology*. Oxford and IBH Publishing Co., New Delhi:107–117.
- ♣ Mitra SK (1999) Anticholestatic activity of HD-03(an herbal formulation) in thioacetamide induced experimental cholestasis. *Indian journal of Expt. Biology*. 37(4):409–410.
- ♣ Mohammad S, Alam A (1999) *Tephrosia purpurea* Ameliorates Benzoyl Peroxide–induced Cutaneous Toxicity in Mice: Diminution of Oxidative stress. *Pharmacy and Pharmacology Communications* 5(7):455–461.
- ♣ Mosmann T (1983) Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. *J Immunol Methods* 65:55–63.
- ♣ Muiva LM, Yenesew A, Derese S, Heydenreich M, Peter MG, Akala HM, Eyase F, Waters NC, Mutai C, Keriko JM, *et al.*, (2009) Antiplasmodial beta-hydroxydihydrochalcone from seedpods of *Tephrosia elata*. *Phytochem. Lett.* 2:99–102.
- ♣ Mukarram M, Ahmad S, Ahmad I, Ahmad M (1987) Chemical screening of seed oils. *Journal of the Oil Technologists' Association of India (Mumbai, India)*, 19(2): 48–9.
- ♣ Mukherjee P (2007) *Quality control of Herbal Drugs-An approach to evaluation of Botanicals*. Second ed. Business Horizons, New Delhi.
- ♣ Murthy MSR and Srinivasan M (1993) Hepatoprotective effect of *Tephrosia purpurea* in experimental animals *Journal of Pharmacology*. 25(1):34–36.
- ♣ Nadaf M, Nasrabadi M, Halimi M (2012) GC-MS Analysis of n-Hexane Extract from Aerial Parts of *Salvia nemorosa*. *Middle-East Journal of Scientific Research* 11 (8): 1127–1130.
- ♣ Nagar PS (2007) A little known plant species of Gujarat *Tephrosia collina* var *lanuginocarpa* Sharma in *J. Bombay Nat. Hist. Soc.* (3):104.
- ♣ Nagar PS, Sata SJ, Pathak SJ (2003) Rediscovery of *Tephrosia jamnagarensis* (Fabaceae), An endangered and narrow endemic plant species of Saurashtra, *Sida* 20(4):1701–1705.

- ♣ Nayar MP (1980) Endemic Flora of peninsular India and Its Significance. *Bull. Bot. Surv. India* 22:12–23.
- ♣ Nayar MP, Sastry ARK (1988) *Red Data Book of Indian Plants, Vol. 2* BSI, Calcutta: 214–218.
- ♣ Nile HS, Khobragade CN (2011) Phytochemical analysis, antioxidant and xanthine oxidase inhibitory activity of *Tephrosia purpurea* Linn. Root extract. *Indian Journal Of Natural Products And Resources*. 2(1):52–58.
- ♣ OECD. Acute Oral Toxicity-Acute Toxic Class Method- Series on testing and assessment No. 423. Paris: Organisation for Economic Co-operation and Development:2001.
- ♣ Parmar NS, Parmar S (1998) Antiulcer potential of flavonoids. *Ind J Physiol Pharmacol*. 48:343–51.
- ♣ Parmar VS, Jain R, Gupta SR, Boll PM, Mikkelsen JM (1988) Phytochemical Investigation of *Tephrosia candida*: Hplc Separation of Tephrosin and 12a-Hydroxyrotenone. *Journal of Natural Products* 51:185–185.
- ♣ Parmar VS, Rathore JS, Jain R, Henderson DA, Malone JF (1989) Occurrence of pongamol as the enol structure in *Tephrosia purpurea*. *Phytochemistry* 28:591–593.
- ♣ Patel A, Patel A, Patel A, Patel NM (2010) Determination of polyphenols and free radical scavenging activity of *Tephrosia purpurea* linn leaves (Leguminosae). *Pharmacognosy Res*. 2(3):152–8.
- ♣ Patil VP, Hugar S, Nanjappaiah HM, Kalyane N, Chowdhary M, Pandarinath (2011) Phytopharmacology of *Tephrosia purpurea* Linn: An Overview. *Pharmacologyonline* 3: 1112–1140.
- ♣ Pavana P, Sethupathy S, Manoharan S (2007) Antihyperglycemic and Antilipidperoxidative effects of *Tephrosia purpurea* seed extract in streptozotocin induced diabetic rats. *Indian Journal of Clinical Biochemistry*. 22(1): 77–83.
- ♣ Perur NG, Subramanian CK, Muhr GR, Roy HE (1973) *Soil fertility evaluation to serve Indian farmers* Mysore Department of Agriculture, Mysore University of Agriculture science and U. S Agency for international development Bangalore: 40–44.
- ♣ Prashant A, Krupadanam GLD (1993) A new prenylated dehydrorotenoid from *Tephrosia villosa* seeds. *Journal of Natural Products*. 56(5):765–6.

- ♣ Queiroz RTD, Tozzi AMGD, Lewis GP (2013) Seed morphology: an addition to the taxonomy of *Tephrosia* (Leguminosae, Papilionoideae, Millettieae) from South America *Plant Syst Evol* 299:459–470.
- ♣ Qureshi R, Bhatti GR, Memon RA (2010) Ethnomedicinal uses of herbs from northern part of NARA desert, Pakistan. *Pak. J. Bot* 42, 839–851.
- ♣ Raghavan RS, Wadhwa BM, Ansari MY and Rao RS (1981), A checklist of Gujarat plants in *Rec. Bot. Surv. India*. 21(2):28.
- ♣ Raghavan SR, Wadhwa BM (1968) On the identity of *Tephrosia jamnagarensis* Santapau and *Tephrosia axillaris* A.R.Smith from Gujarat. *Curr. Sci.* 18:536.
- ♣ Raina SN, Srivastav PR, Ramarao S (1985) Nuclear DNA Variation in *Tephrosia*. *Genetica*. 69:27–33.
- ♣ Rajabudeen E, Ganthi AS, Subramanian MPS (2014) Affiliations Pharmacognostical Studies on *Tephrosia villosa* (L.) Pers. (Fabaceae) *Research Journal Of Pharmacognosy And Phytochemistry* 6(4):190–194.
- ♣ Rajani P, Sarma N (1988) A coumestone from the roots of *Tephrosia hamiltonii*. *Phytochemistry* 27: 648–649.
- ♣ Rao EV, Raju NR (1979) Occurrence of (-)-isolonchocarpin in the roots of *Tephrosia purpurea*. *Phytochemistry* 18(9): 1581–2.
- ♣ Rao EV, Raju NR (1984) Two flavonoid from *Tephrosia purpurea* Lin. *Phytochemistry* 23(10): 2339–2342.
- ♣ Rao EV, Sridhar P (1999) Chemical examination of *Tephrosia strigosa* *Indian Journal of Chemistry* 38B(7): 872–873.
- ♣ Rao KC, Geetha BL, Suresh G (2003) *Red list of threatened Vascular plant in India*, pub-Botanical survey of India: 59.
- ♣ Rao PP, Srimannarayana G (1980) Tephrosol, a new coumestone from the roots of *Tephrosia villosa*. *Phytochemistry*, 19(6): 1272–3.
- ♣ Rao PP, Srimannarayana G (1981) Tephtrinone - a new flavanone from *Tephrosia villosa*, *Cur. Sci.* 50(7): 319–20.
- ♣ Rao S, Timsina B, Nadumane VK (2013) Antimicrobial effects of medicinal plants and their comparative cytotoxic effect of HepG2 cell line. *International J. Of Pharmacy and Pharm. Sci.* 6(1): 101–105.
- ♣ Riahi L, Elferchichi M, Ghazghazi H, Jebali J, Ziadi S, Aaiodhi C, Chograni H, Zaovali Y, Zoghiami N, Mliki A (2013) Phytochemistry, antioxidant and antimicrobial activity of

the essential oil of *Mentha rotundifolia* L. In Tunisia. *Industrial Crops and Product* 49: 883–889.

- ♣ Richardson IBK (1978) *Endemic Taxa and The Taxonomist*, In: Street, HE(ed) *Essays in Plant Taxonomy*, Academic Press:245–262.
- ♣ Rivera AG, Mariscal HA, Romero-Ceronio N, Fuente LFRL, Garcia CEL (2013) Synthesis and antiinflammatory activity of three nitrochalcones *Bioorganic and Medicinal chemistry letters* 23:5519–5522.
- ♣ Ross MSF, Brain KR (1977) "*An Introduction to Phytopharmacy*", Pitman Medical. Kent: 84.
- ♣ Roy M, Bhattacharya PK, Pal S, Chowdhuri A, Adityachaudhury N (1987) Dehydrodihydroteneone and flemichapparin-B in *Tephrosia candida*. *Phytochemistry* 26:2423–2424.
- ♣ Roy M, Mitra SR, Bhattacharyya A, Adityachaudhury N (1986) Candidone, a flavanone from *Tephrosia candida*. *Phytochemistry* 25:961–962.
- ♣ Ruppel EG, Freyre RH, Barnes DK (1967) Hot Water Immersion Effective for Scarifying Seeds of *Tephrosia vogelii*. *Crop Science* 7(3):273–275.
- ♣ Rutunga V, Karanja NK, Gachene CKK, Palm C (1999) Biomass production and nutrient accumulation by *Tephrosia vogelii* (Hemsley) A. Gray and *Tithonia diversifolia* Hook F. fallows during the six-month growth period at Maseno, Western Kenya *Biotechnol. Agron. Soc. Environ.* 13 (4): 237–246.
- ♣ Sadashiva CT, Naidoo Y, Naidoo J R, Naidoo G (2013) Chemical composition of the essential oil from the leaves of *Endostemon obtusifolius* (E.Mey. ex Benth.) N.E.Br. *Biochem. & Pharmacol.* 2:4.
- ♣ Sahayaraj K, Kombiah P, Dikshit AK, Rathi JM, (2014) Chemical constituents of essential oils of *Tephrosia purpurea* and *Ipomoea carnea* and their repellent activity against *Odoiporus longicollis* J. *Serb. Chem. Soc.* 79 (0): 1–13.
- ♣ Sairam RK, Tyagi A (2004) Physiology and molecular biology of salinity stress tolerance in plants *Curr. Sci.* 86(3): 407–421.
- ♣ Saleem M, Ahmad Su, Alam A (2001) *Tephrosia purpurea* alleviates phorbol ester – induced tumor protection response in murine skin *Pharmacol Res.* 43(2):135–144.
- ♣ Sanchez I, Gomez-Garibay F, Taboada J, Ruiz BH (2000) Antiviral effect of flavonoids on the dengue virus. *Phytother. Res.* 14:89–92.

- ♣ Sandhya S, Venkatramana K, Vinod KR, Sunitha CH, Murali K (2011) Pharmacognostical Standardization of *Tephrosia maxima* Pers Root. *Pharmacognosy Journal* 3(26):25–27.
- ♣ Sangeetha B, Krishnakumari S (2010) Folk Medicinal plant *Tephrosia purpurea* ameliorates carbon tetrachloride induced hepatic damage in rats. *International Journal of Pharma and Biosciences* 1(2).
- ♣ Sanjappa M (1992) *Legumes of India*, Dehra Dun: Bishen Singh Mahendra.
- ♣ Santapau H (1958) Addition and corrections to the Indo-Nepalese Flora. *Proc. Natl. Sci. Inst. India* 24B:133.t.1
- ♣ Santapau H (1962) *The flora of Saurashtra Part-I*. (Ranunculaceae to Rubiaceae) Saurashtra Research Society, Rajkot:270.
- ♣ Santos AFD, Fonseca SA, César FA, Albuquerque MCPA, Santana JV, Santana AEG (2014) A penta-substituted pyridine alkaloid from the rhizome of *Jatropha elliptica* (Pohl) Muell. Arg. is active against *Schistosoma mansoni* and *Biomphalaria glabrata*. *Parasitol Res.* 113(3):1077–84.
- ♣ Santram L, Pawar RS, Jain AP, Singhai AK (2006) Woundhealing potential of *Tephrosia purpurea* (Linn.) Pers in rats. *Journal of Ethnopharmacology* 108(2):204–210.
- ♣ Sarker SD, Nahar L (2007) *Chemistry for Pharmacy Students General, Organic and Natural Product Chemistry*. England: John Wiley and Sons:283–359.
- ♣ Sasidharan N (2006) Illustrated manual on tree flora of Kerala supplemented with computer aided identification. Pub: Kerala Forest Research Institute India:462.
- ♣ Sasikumar G (1985) Studies on leguminous weeds and a holoparasite- *Orobancha* Linn. Thesis submitted in The Maharaja Sayajirao University of Baroda:69.
- ♣ Saxena VK, Choubey A (1997) A novel neoflavonoid glycoside from *Tephrosia purpurea* stem. *Fitoterapia* 68:359–360.
- ♣ Sayad M (2012) Comparative study of *Tephrosia purpurea* (Linn) leaves and Lovastatin on cholesterol level of hyperlipidemic wistar rats. *IOSR Journal of Pharmacy & Biological Science* 2(1):25–30.
- ♣ Schalk M, Cabello-Hurtado F, Atanossova MPR, Saindrenan P, Werck-Reichhart D (1998) Piperonylic Acid, a Selective, Mechanism-Based Inactivator of the trans-Cinnamate 4-Hydroxylase: A New Tool to Control the Flux of Metabolites in the Phenylpropanoid Pathway *Plant Physiol.* 118(1):209–218.

- ♣ Schrire BD (2005) Milletieae. In G Lewis, D. Schrire B. Mackinda and M. Locks (eds) *Legumes of the world*, Kew UK Royal botanical Garden:367-387.
- ♣ Sethupathy S, Pavana P, Manoharan S (2007) Antihyperglycemic and Antilipidperoxidative effects of *Tephrosia purpurea* seed extract in streptozotocin induced diabetic rats. *Indian Journal of Clinical Biochemistry* 22(1):77–83.
- ♣ Shah GL (1978) *Flora of Gujarat Part I*, Pub:- Sardar Patel University Vallabh Vidyanagar: 248
- ♣ Shah GL (1983) Rare species with restricted distribution in south Gujarat: 50–54, In Jain SK and RR Rao (Eds:). *An Assessment of Threatened Plants Of India*. Botanical Survey of India. Calcutta.
- ♣ Shan B, Cai YZ, Sun M, Corke H (2005) Antioxidant capacity of 26 spice extracts and characterization of their phenolic constituents. *J Agric Food Chem*. 53: 7749–7759.
- ♣ Sharma A, Shing RT, Sehgal V and Handa SS (1991) Antihepatotoxic activity of some plants used in herbal formulations. *Fitoterapia*. 62:131–8.
- ♣ Sharma AK (2008) Phytochemical examination of some plants widely used in Ayurveda, in thesis submitted at University of Rajasthan:, chapter 1 Introduction:1–28.
- ♣ Sharma P, Rastogi S, Bhatnagar S (2003) Antileishmanial action of *Tephrosia purpurea* linn, extract and its fractions against experimental visceral leishmaniasis *Drug Development Research*.60(4):285–293.
- ♣ Sharma R, Mehan S, Kalra S, Khanna D (2013) *Tephrosia purpurea*- A magical herb with blessing in human biological System. *Int. J. Of Recent advances in Pharm, Res*. 3(3): 12–22.
- ♣ Sharma SK, Vasudeva N, Rathi P (2008) Isolation and identification of a new phytosterol ester from *Tephrosia purpurea* (Linn.) Pers. Root. *Int. J. Chem. Sci* 6:1734–1741.
- ♣ Sharma VS (1963) Description of *Tephrosia collina* var *lanuginocarpa* Sharma and two new varieties. *J. Bombay Nat. Hist. Soc*. 60(3):754–758.
- ♣ Sing JS, Yagava PS (1974) Seasonal variation in composition, plant biomass and net primary productivity of a torpical grassland at Kurushedra, India. *Ecological Monograph* 44:351–376.
- ♣ Singh NP, Karthikeyan S (2000) *Flora of Maharashtra State Dicotyledons. Vol.1*. Botanical Survey of India, Kolkata.

- ♣ Sinha B, Natu AA, Nanavati DD (1982) Prenylated flavonoids from *Tephrosia purpurea* seeds *Phytochemistry* 21:1468–1470.
- ♣ Soni K, Suresh KP, Saraf MN (2003) Free radical scavenging activity of *Tephrosia purpurea* linn *Indian Journal of Pharmaceutical Science*. 65(1): 27–30.
- ♣ Sreenivasulu B, Sarma PN (1998) Chemical examination of the whole plant of *Tephrosia strigosa*. *Indian Journal of Chemistry section-B*. 37(11): 1217–1218.
- ♣ Sudhi KS (2012) Redlist has 132 species of plants, animal from India, *The Hindu*.
- ♣ Takhtajan A. (1997) *Diversity and Classification of Flowering Plants*.
- ♣ Thomas K, Coleman P (2004) Use of complementary or alternative medicine in a general population in Great Britain. Results from the National Omnibus survey, *Journal of Public Health* 26:152–157.
- ♣ Touqeer S, Saeed AM, Ajaib M (2013) A review on phytochemistry and pharmacology of genus *Tephrosia*. *Phytopharmacology* 4(3): 598–637.
- ♣ Tropicos.org. Missouri Botanical Garden. 13 Apr 2015 <http://www.tropicos.org>
- ♣ Valli G, Vasanthi A, Vijayalakshmi R, Thirupathi AT (2011) Antipyretic and anti-inflammatory activities of *Tephrosia purpurea* root extracts. *IJPRD*. 3(6):211–217.
- ♣ Verpoorte R, Schripsema (2009) Isolation, identification, and structure elucidation of Alkaloids A general overview. In: Linskens H.F. and Jackson J.F.(ed) *Alkaloids* Springer India,; 1–22.
- ♣ Vyas KJ (1973) *A Contribution To The Florist And Phytosociology Of The River Narmada In Gujarat State*. Thesis submitted to S.P. University, Vallabh Vidyanagar.
- ♣ Wagner H, Bladt S (1996) *Plant Drug Analysis, A Thin Layer Chromatography Atlas*, second ed. Verlag, Berlin, Heidelberg, Germany, 1996.
- ♣ Wagner H, Wolff P. (1977) *New Natural Products and Plant Drugs with Pharmacological, Biological or Therapeutical Activity*, Springer-Verlag. Berlin: 6.
- ♣ Wallis TE (1957) *Textbook of pharmacognosy*. 5<sup>th</sup> edition J. and A. Churchill, London.
- ♣ Walter KS, Gillet HJ (1997) *IUCN Red list of threatened plants* Pub International.
- ♣ Waterman PG, Khalid SA (1980) The major flavonoids of the seed of *Tephrosia apollinea*. *Phytochemistry* 19: 909–915.
- ♣ WCMC (1994) Status reports as of 24 November 1994. Gujarat. Print out from plant database BG-BASE. World conservation monitoring centers.
- ♣ Weingartner O, Bohm M, Laufs U (2008) Plant sterols as dietary supplement for the prevention of cardiovascular disease *Dtsch. Med. Wochenschr*. 133:1201–1204.

- ♣ Willaman JJ, Schubert B (1961) *Alkaloid-bearing Plants and Their Contained Alkaloids*. Pub in Washington 25 DC:126
- ♣ Wu YH, YZ Cai, Zhang Y (2009) Natural Phenolic Compounds From Medicinal Herbs and Dietary Plants: Potential Use for Cancer Prevention. *Nutr. Cancer*. 62(1): 1–20.
- ♣ [www.icar.org.in/files/ar0506/cs.pdf](http://www.icar.org.in/files/ar0506/cs.pdf)
- ♣ [www.imdahm.gov.in](http://www.imdahm.gov.in)
- ♣ Xin Y, Yan Z, Xiaohua L, Huang X, Zeng X (2009) Analysis of physical characteristic of *Tephrosia vogelii* Hook F. Seeds and its oil component *Acta Scientiarum Naturalium Univer. Sunyatseni* 1: 37–41.
- ♣ Yadav A (2000) Rediscovery of new location of *Tephrosia collina* var. *lanuginocarpa* in Rajasthan. *Journal of Phytological Research* 13(1):103–104.
- ♣ Yang DY, Fushimi H, Cai SQ (2004) Molecular analysis of Rheum species used as Rhei Rhizoma based on the chloroplast *matK* gene sequence and its application for identification[J]. *Biol Pharm Bull* 27(3):375–383.
- ♣ Yenesew A, Dagne E, Waterman PG (1989) Flavonoids from the seed pods of *Tephrosia pumila* *Phytochemistry* 28:1291–1292.
- ♣ Yoshiki Y, Kim JH, Okubo K (1994) Saponins conjugated with 2,3-dihydro-2,5-dihydroxy-6-methyl-4H-pyran-4-one from *Phaseolus coccineus*. *Phytochemistry* 36(4):1009–12.
- ♣ Young TP (1984) Comparative demography of semelparous *Lobelia telekii* and iteroparous *Lobelia keniensis* on Mount Kenya. *Journal of Ecology* 72: 637–650.