

LIST OF FIGURES

Figure 1	Treatment methods for the removal of textile dyes.	7
Figure 2	Wood and its cell wall components.	12
Figure 3	Corresponding phenylpropanoid H, G and S units.	17
Figure 4	Structure of lignin formed by the polymerization of phenyl propanoid units.	18
Figure 5	Pie chart representation of percent (G:H:S) units of lignin composition in different wood types: a. Hardwood, b. Softwood and c. Monocot.	19
Figure 6	Schematic representation of overall processes involved in lignin degradation by of white rot fungi.	22
Figure 7	Schematic diagram representing mechanism of MnP action.	24
Figure 8	Schematic diagram representing mechanism of LiP action.	27
Figure 9	Schematic diagram representing mechanism of Lac action.	29
Figure 10	Application of ligninolytic enzymes in different industrial sectors.	31
Figure 11	Solid State fermentation A) Growth of <i>Trametes hirsuta</i> on 1 mm wheat straw after 12 days of incubation B) Growth of <i>Trametes versicolor</i> on Pigeon pea (<i>Cajanus cajan</i>) pod shells after 9 days of incubation.	64
Figure 12-13	Optimization of agro-industrial waste used as substrates for ligninolytic enzyme production by <i>Trametes hirsuta</i> and <i>Trametes versicolor</i> .	65
Figure 14-15	Optimization of incubation time for ligninolytic enzyme production with respect to their specific substrates by <i>Trametes hirsuta</i> and <i>Trametes versicolor</i> .	66
Figure 16-17	Effect of pH on MnP and MnIP enzyme activity produced by <i>Trametes hirsuta</i> and <i>Trametes versicolor</i> .	67

Figure 18-19	Effect of pH on laccase enzyme activity produced by <i>Trametes hirsuta</i> and <i>Trametes versicolor</i> .	68
Figure 20-21	Effect of pH on LiP enzyme activity produced by <i>Trametes hirsuta</i> and <i>Trametes versicolor</i> .	69
Figure 22-23	Effect of temperature on ligninolytic enzyme production by <i>Trametes hirsuta</i> and <i>Trametes versicolor</i> .	70
Figure 24-25	Production profile of ligninolytic enzymes after partial purification with ammonium sulphate precipitation produced by <i>Trametes hirsuta</i> and <i>Trametes versicolor</i> .	71
Figure 26	SDS-PAGE of the purified Laccase from fungal strains a) marker b) <i>Trametes hirsuta</i> and c) <i>Trametes versicolor</i> .	72
Figure 27	Decolourization of dye Reactive Red HE8B in liquid medium by (left to right) <i>Trametes hirsuta</i> , Control and <i>Trametes versicolor</i> .	73
Figure 28	Decolourization of dye Reactive Orange R in liquid medium by (left to right) <i>Trametes hirsuta</i> , Control and <i>Trametes versicolor</i> .	73
Figure 29	Decolourization of dye Reactive Black B in liquid medium by (left to right) <i>Trametes hirsuta</i> , Control and <i>Trametes versicolor</i> .	74
Figure 30	Decolourization of dye Reactive Red ME4BL in liquid medium by (left to right) <i>Trametes hirsuta</i> , Control and <i>Trametes versicolor</i> .	74
Figure 31	Decolourization of dye Reactive Yellow FG in liquid medium by (left to right) <i>Trametes hirsuta</i> , Control and <i>Trametes versicolor</i> .	75
Figure 32	Percent decolourization of dye Reactive Red HE8B in liquid medium by <i>Trametes hirsuta</i> (TH) and <i>Trametes versicolor</i> (TV).	76
Figure 33	Percent decolourization of dye Reactive Orange 2R in liquid medium by <i>Trametes hirsuta</i> (TH) and <i>Trametes versicolor</i> (TV).	76
Figure 34	Percent decolourization of dye Reactive Black B in liquid medium by <i>Trametes hirsuta</i> (TH) and <i>Trametes</i>	77

	<i>versicolor</i> (TV).	
Figure 35	Percent decolourization of dye Reactive Red ME4BL in liquid medium by <i>Trametes hirsuta</i> (TH) and <i>Trametes versicolor</i> (TV).	77
Figure 36	Percent decolourization of dye Reactive Yellow FG in liquid medium by <i>Trametes hirsuta</i> (TH) and <i>Trametes versicolor</i> (TV).	78
Figure 37	FTIR spectra of a) control dye Reactive Red HE8B and b) its degradation product treated by <i>Trametes hirsuta</i> .	80
Figure 38	FTIR spectra of a) control dye Reactive Red HE8B and b) its degradation product treated by <i>Trametes versicolor</i> .	80
Figure 39	FTIR spectra of a) control dye Reactive Orange 2R and b) its degradation product treated by <i>Trametes hirsuta</i> .	81
Figure 40	FTIR spectra of a) control dye Reactive Orange 2R and b) its degradation product treated by <i>Trametes versicolor</i> .	82
Figure 41	FTIR spectra of a) control dye Reactive Black B and b) its degradation product treated by <i>Trametes hirsuta</i> .	83
Figure 42	FTIR spectra of a) control dye Reactive Black B and b) its degradation product treated by <i>Trametes versicolor</i> .	83
Figure 43	FTIR spectra of a) control dye Reactive Red ME4BL and b) its degradation product treated by <i>Trametes hirsuta</i> .	84
Figure 44	FTIR spectra of a) control dye Reactive Red ME4BL and b) its degradation product treated by <i>Trametes versicolor</i> .	85
Figure 45	FTIR spectra of a) control dye Reactive Yellow FG and b) its degradation product treated by <i>Trametes hirsuta</i> .	86
Figure 46	FTIR spectra of a) control dye Reactive Yellow FG and b) its degradation product treated by <i>Trametes versicolor</i> .	86
Figure 47	Average percentage of weight loss in <i>Eucalyptus globulus</i> after different incubation periods of the woods with a) <i>T. hirsuta</i> and b) <i>T. versicolor</i> .	91

Figure 48	Transverse (a, i, m and n), tangential (b, d-h, j-l, o and p) and radial longitudinal (c) sections of <i>Eucalyptus globulus</i> wood blocks infected with <i>Trametes hirsuta</i> for 30 (a-d), 60 (e-k), 90 (l and m) and 120 (n-p) days.	113
Figure 49	Transverse (a, l, p and t) tangential (b-k, m, n, o, q and r) and radial longitudinal (s and u) sections of <i>Eucalyptus globulus</i> wood blocks infected with <i>Trametes versicolor</i> for 30 (a-k), 60 (l-o), 90 (p-s) and 120 (t and u) days.	114
Figure 50	Confocal images of transverse sections from the wood of <i>Eucalyptus globulus</i> infected with <i>Trametes hirsuta</i> (a-d) and <i>Trametes versicolor</i> (e-h).	115
Figure 51	Average percentage of weight loss in <i>Azadirachta indica</i> after different incubation periods of the woods with a) <i>T. hirsuta</i> and b) <i>T. versicolor</i> .	94
Figure 52	Transverse (a, f and g) and tangential longitudinal (b-e and h) sections of <i>Azadirachta indica</i> wood blocks infected with <i>Trametes hirsuta</i> for 30 (a-h) days.	116
Figure 53	Transverse (a, b, h and i) and tangential longitudinal (c, d, e, f, g and j) sections from the wood of <i>Azadirachta indica</i> infected with <i>Trametes hirsuta</i> for 60 (a-e), 90 (f and g) and 120 (h-j) days.	117
Figure 54	Transverse (a,b, d-g and k) and tangential longitudinal (c and h-j) and radial longitudinal (l) sections from the wood of <i>Azadirachta indica</i> infected with <i>Trametes versicolor</i> for 30 (a-d), 60 (e-i), 90 (j) and 120 (k and l) days.	118
Figure 55	Confocal laser scanning images from the transverse sections of <i>Azadirachta indica</i> wood blocks infected with <i>Trametes hirsuta</i> (a-f) and <i>Trametes versicolor</i> (g-j).	119
Figure 56	Average percentage of weight loss in <i>Tectona grandis</i> after different incubation periods of the woods with a) <i>T. hirsuta</i> and b) <i>T. versicolor</i> .	97
Figure 57	Transverse (a, b, k, m and o) tangential (c-j, l and n) and radial longitudinal (p and q) sections of <i>Tectona grandis</i> wood blocks infected with <i>Trametes hirsuta</i> for 30 (a), 60 (b-e), 90 (f-m) and 120 (n-q) days.	120

Figure 58	Transverse (a, e and f) tangential (c-j, l and n) sections of <i>Tectona grandis</i> wood blocks infected with <i>Trametes versicolor</i> for 30 (a), 60 (b-d), 90 (e) and 120 (f and g) days.	121
Figure 59	Confocal laser scanning images from the transverse sections of <i>Tectona grandis</i> wood blocks infected with <i>T. hirsuta</i> (a-e) and <i>T. versicolor</i> (f-m).	122
Figure 60	Average percentage of weight loss in <i>Leucaena leucocephala</i> after different incubation periods of the woods with a) <i>T. hirsuta</i> and b) <i>T. versicolor</i> .	99
Figure 61	Transverse (f and i) and tangential longitudinal (a-e, g, h and j) sections from the wood of <i>Leucaena leucocephala</i> infected with <i>Trametes hirsuta</i> for 30 days (a and b), 60 (c-f), 90 (g and h) and 120 (i and j) days.	123
Figure 62	Transverse (a and i), tangential (b-h and j) and radial longitudinal (k and l) section from the <i>Leucaena leucocephala</i> wood infected with <i>Trametes versicolor</i> for 30 days (a and b), 60 (c-i), 90 (j) and 120 (k and l) days.	124
Figure 63	Confocal images from transverse sections of <i>Leucaena leucocephala</i> wood infected with <i>Trametes versicolor</i> (a-f) and <i>Trametes hirsuta</i> (g-k).	125