



4. References

- Agrawal A. (2001). The regulatory community: Decentralization and the environment in the Van Panchayats (Forest Councils) of Kumaon, India. *Mountain research and development*, 21(3): 208-211.
- Agrawal A. (2001). Common Property Institutions and Sustainable Governance of Resources. *World Development* 29(10): 1649-72.
- Ahmad F., Khan M. A., Ahmad M., Zafar M., Nazir A. and Marwat S. K. (2009). Taxonomic studies of grasses and their indigenous uses in the salt range area of Pakistan. *African Journal of Biotechnology*, 8(2): 231-249.
- Ajtay G.L., Ketner P. and Duvigneaud P. (1979) Terrestrial primary production and phytomass. In Bolin, B., Degens, E., Kempe, S. and Ketner, P. (Eds): *The Global Carbon Cycle. SCOPE* edn., Vol. 13. Wiley, Chichester, pp. 129-182.
- Akçin ÖE, Şenel G and Akçin Y (2013). Leaf epidermis morphology of some *Onosma* (Boraginaceae) species from Turkey. *Turkish Journal of Botany* 37: 55–64
- Banerjee S.K., and Chauhan K.P.S. (1981) Studies on the evolution of seed coat pattern in wheat by scanning electron microscopy identification. *Seed Science and Technology*. 9(3) : 819-822.
- Barthlott W. (1981). Epidermal and seed surface characters of plants: systematic applicability and some evolutionary aspects. *Nordic Journal of Botany*, 1(3):345-355.
- Barthlott W. (1990). Scanning electron microscopy of the epidermal surface in plants. *Scanning electron microscopy in taxonomy and functional morphology Systematics association special volume*,(41): 69-94.
- Bengoechea G. and Gomez Campo C. (1975). Algunos caracteres de la semilla en la tribu Brassiceae. *An. Inst. Bot. AJ Cavanilles*, 32(2): 793-841.
- Blatter E. and McCann C. (1935). *The Bombay grasses*. Imperial Council of Agriculture research, Scientific monograph.
- Bockenstedt, P. J. (2001) Lower Wells Creek Natural Resource Inventory.
- Bogdan A. V. (1965). Vegetative propagation of fodder and pasture grasses. *Kenya Farmer*, 105:33-34.
- Bogdan A. V. (1966). Seed morphology of some cultivated African grasses. In *Proceedings of the International Seed Testing Association*, 31: 789-799.
- Bole P. V., and Pathak J. M. (1988). *Flora of saurashtra*. Botanical Survey of India.
- Bor, N. L. (1960). *The grasses of Burma. Ceylon, India and Pakistan*. Pergamon Press
- Bradley, K. W., and Fishel, F. (2010). *Integrated pest management: Identifying grass seedlings*; University of Missouri Extension
- Briske D. D. (1991). Developmental morphology and physiology of grasses. *Grazing management: an ecological perspective*. Timber Press, Portland 85-108.

- Brisson J. D. and Peterson R. L. (1976). Critical review of the use of scanning electron microscopy in the study of the seed coat. In *Proceedings of the Scanning Electron Microscope Symposium*. 2: 477-495.
- Brisson J. D. and Peterson, R. L. (1977). The scanning electron microscope and X-ray microanalysis in the study of seeds: a bibliography covering the period of 1967-1976. *Scanning Electron Microscopy*, 2: 697-712.
- Brochmann C. and Elven R. (1992). Ecological and genetic consequences of polyploidy in arctic *Draba* (Brassicaceae). *Evolutionary Trends in Plants*.6(2): 111-124.
- Brown R. (1814). General remarks, geographical and systematical, on the botany of *Terra Australis*, 2: 533-613.
- Burr S. and Turner D. M. (1933). *British economic grasses, their identification by the leaf anatomy*. Edward Arnold and Co., London.
- Buth G. M. and Ara R. (1983). Seed coat anatomy of some cultivated Brassicas. *Phytomorphology*.31: 69-78.
- Campbell A. G., Phillips D. S. M., and O'reilly E. D. (1962) "An electronic instrument for pasture yield estimation." *Grass and Forage Science* 17(2): 89-100.
- Carrier L. (1917). *The identification of grasses by their vegetative characters* (No. 461). US Dept. of Agriculture. Caryopsis. *SIDA* 2002, 20: 525-548.
- Chavan A. R., and Mehta A. R. (1959). *Grasses of Pavagarh*. Journ. M.S. Uni. Baroda 7(2): 171-185.
- Chuang T.I. and Heckard L. R. (1972). Seed coat morphology in *Cordylanthus* (Scrophulariaceae) and its taxonomic significance. *American Journal of Botany*, 258-265.
- Chuang T. I. and Heckard L. R. (1983). Systematic significance of seed-surface features in *Orthocarpus* (Scrophulariaceae-Subtribe Castillejinae). *American Journal of Botany*, 877-890.
- Clark L. G. and Fisher J. B. (1987). Vegetative morphology of grasses: shoots and roots. Soderstrom TR, and others ed (s). *Grass systematics and evolution: an International Symposium held at the Smithsonian Institution, Washington, DC, 27-31 July 1986*. Washington, DC London, Smithsonian Institution Press.
- Clayton W. D. and Renvoize S. A. (1986). *Genera graminum. Grasses of the World*. Royal Botanical Garden, Kew. pp.389.
- Colledge S. (1988). Scanning electron studies of the pericarp layers of some wild wheats and ryes: methods and problems. In *Scanning electron microscopy in archaeology* (pp. 225-236). British Archaeological Reports.
- Cook C. D. K. (1996) *Aquatic and wetland plants of India*, Oxford University Press, New Delhi, 95-104 and 280-316
- Cooke T. (1901-1908). *The flora of the Presidency of Bombay*. Taylor and Francis, London. Vol. I and II.

- Cope T.A. (1982). *Poa* L. In: Nasir E, Ali SI (eds). Flora of Pakistan, Univ. of Karachi, Karachi, Pakistan. 143: 559-583.
- Costea, M. and Tardif, F. J. (2002). Taxonomy of the most common weedy European *Echinochloa* species (Poaceae: Panicoideae) with special emphasis on characters of the lemma and caryopsis. *SIDA*. 20(2): 525 – 548.
- Crow G. E. and Storks I. M. (1980). Rare and endangered plants of New Hampshire: a phytogeographic viewpoint. *Rhodora*, 82(829): 173-189.
- Dabadghao P. M. and Shankarnarayan K. A. (1973). *Grass cover of India*. Indian Council of Agricultural Research; New Delhi.pp. 713.
- Desai R. J. and Raole V. M. (2012). *Spodiopogon aristatus*, a new species of Poaceae from Gujarat, India. *Kew Bulletin*, 67(1), 103-107.
- Desai R. J. and Raole V. M. (2013). Noteworthy grasses of Gujarat. *J. Indian bot. Soc.* Vol, 92 (3&4). 179-183.
- Durgan B. R. (1999). *Broadleaf and grass weed seedling identification keys*. University of Minnesota, Extension Service.
- Ellison A.M. (1987). Effect of seed dimorphism on the density-dependent dynamics of experimental populations of *Atriplex triangularis*(Chenopodiaceae). *Am. J. Bot.*, 74: 1280-1288.
- Etkin N.L. (2000). *Eating on the Wild Side: The Pharmacologic, Ecologic, and Social of Using Noncultigens*. Tucson: University of Arizona Press.
- FAOSTAT.1999. Database online, [http://:apps.fao.org/](http://apps.fao.org/)
- Fenner M. (1987). Seedlings. *New phytologist*. 106 (1): 35-47.
- Fishel F. 2004. Identifying Grass seedling, In: *Integrated Pest Management Manual*. University of Missouri Extension Columbia. 5–18.
- G. Yadaiah , Bindiya Oza , Parul Joshi (2011)Ethno-medico-botany of Godhra Forest Division of Panchmahal District, Gujarat State 137(8) : 948-961
- Gaertner J. (1788). *Josephus Gärtne: De fructibus et seminibus plantarum*.Vol. 1. Typis Academiae Carolinae.
- Gaertner J. (1791). *De fructibus et seminibus plantarum*. U.S. Department of Agriculture, National Agricultural Library
- Gandhi D. J., Albert S., Pandya N. R. and Panchal, K. R. (2011). Socioeconomic Study of Grasses and Legumes in Baria and Godhra Forest Division, Gujarat. *Notulae Scientia Biologicae*, 3(3): 53.
- Gandhi D., Albert S.,and Pandya N. (2016). *Handbook on the Morphology of Common Grasses: Identification and Characterization of Caryopses and Seedlings*. CRC Press. pp. 314.
- Gohil T. G. and Patel D. D. (2006). New records for Gujarat State from Dharampur forests of Valsad district in South Gujarat. *J. Econ. Taxon. Bot*, 30(3): 715-716.
- Gómez-Campo C. (1980). Morphology and morpho-taxonomy of the tribe Brassiceae. *Morphology and morpho-taxonomy of the tribe Brassiceae*. 32: 3-31.

- Barker, N. P., Clark, L. G., Davis, J. I., Duvall, M. R., Guala, G. F. and Mathews, S. Y. (2001). Phylogeny and subfamilial classification of the grasses (Poaceae). *Annals of the Missouri Botanical Garden*. 88: 373-457.
- Guest E. (1966). *Flora of Iraq*, Vol 1: Introduction. Baghdad: Ministry of Agriculture, Republic of Iraq.
- Haber A., and Runyon R. P. (1969). *General statistics*. Addison-Wesley.
- Harper J. L. (1977). *Population biology of plants*. Academic press, Newyork. pp. 892.
- Harries, W. F. (2010) *The identification of some of the more common native Oklahoma grasses by vegetative characters*, Oklahoma native plant record. 10.
- Hazra C. R. (2014) Feed and forage resources for sustainable livestock development. *Range management and Agroforestry* 35(1):1-14.
- Henning, E. (1930) *Table for the identification of grasses and legumes in the nonflowering condition*, Translated from the Swedish by F. V. Meissner, Springer. Berlin.
- Heywood V.H. (1971). *Scanning electron microscopy: Systematic and evolutionary applications*. Academic press, Newyork. London. pp. 331
- Hill R. J. (1976). Taxonomic and phylogenetic significance of seed coat microsculpturing in *Mentzelia* (Loasaceae) in Wyoming and adjacent western states. *Brittonia*, 28(1), 86-112.
- Hillman F. H. (1916). *Distinguishing characters of the seeds of Sudan grass and Johnson grass*. USDA, Bulletin 406 (Washington, D. C.).
- Hilu K. W. (2007). A century of progress in grass systematics. *Kew Bulletin*. 62: 355-373.
- Hilu K. W. and Alice L. A. (2001). A phylogeny of Chloridoideae (Poaceae) based on matK sequences. *Systematic Botany*, 26(2): 386-405.
- Hilu K. W. and Wright K. (1982). Systematics of Gramineae: a cluster analysis study. *Taxon*. 31: 9-36.
- Hitchcock A. S. (1936). *Manual of the grasses of the West Indies*. United State Department of Agriculture. 240-250.
- Hoagland R. E. and Paul R. N. (1978). A Comparative SEM Study of Red Rice and Several Commerical Rice (*Oryza sativa*) Varieties. *Weed Science*, 26(6): 619-625.
- Hufford L. (1995) Seed morphology of Hydrangeaceae and its phylogenetic implications. *International Journal of Plant Science* 156: 555–580.
- Jacobs S. W. L., Everett J., Barkworth M. E., and Hsiao C. (2000). Relationships within the stipoid grasses (Gramineae). *Grasses: Systematics and Evolution*. CSIRO, Melbourne, 75-82.
- Jadhav R. N., Sastry K. L. N., Thakker P. S. and Chavan S. A. (1998). *Bio-diversity threat through exotic sp., using remotely sensed data*. A case study of Banni (Kachchh) Gujarat. ENVIS bulletin. 17: 1-22.

- Jain S. K. and Rao R. R. (1977). *A handbook of field and herbarium methods*. New Delhi: Today and Tomorrow's Printers and Publishers. pp. 157.
- Jensen L. A., Johnston W. R. and Pierpoint M. (1957). Seed characteristics of certain wild barley, *Hordeum* spp. In *Proceedings of the International Seed Testing Association* 7: 87-91.
- Jian-Guo H., Clifford H. T. and Jia Shen-Xiu W. P. (1993) A cluster analysis of seedling characters of Gramineae. *Acta Pytotax. Sin* 31(6):517–532.
- Jones S. B. and Luchsinger A.E (1986) *Plant systematics. McGraw Hill Book Co., New York*, 15: 3-24.
- Kanwal D., Abid R. and Qaiser M. (2009). The Seed Atlas of Pakistan-I Aizoaceae. *Pakistan Journal of Botany*, 41(4): 1557-1564.
- Karez R., Engelbert S. and Sommer, U. (2000). 'Co-consumption' and 'protective coating': two new proposed effects of epiphytes on their macroalgal hosts in mesograzed-epiphyte-host interactions. *Marine Ecology Progress Series* 205: 85-93.
- Karihaloo J. L. and Malik S. K. (1994). Systematic relationships among some species of *Solanum* L. sect. *Melongena* L.—evidence from seed characters. *Indian Journal of Plant Genetic Resources*, 7(1): 13-21.
- Karthikeyan.S., S.K., Jain, M. P. Nayar and M. Sanjappa. (1989). *Florae Indicae Enumerate Monocotyledonae*. FI. Ind. Ser. 4. Botanical Surey of India. Calcutta.
- Kaul V., Koul A. K. and Sharma M. C. (2000). The underground flower. *Current Science*, 78(1): 39-44.
- Koul K. K., Nagpal R. and Raina S. N. (2000). Seed coat microsculpturing in *Brassica* and allied genera (subtribes Brassicinae, Raphaninae, Moricandiinae). *Annals of Botany*, 86(2): 385-397.
- Kumar V, Kodandaramaiah J and Rajan M.V (2012). Leaf and anatomical traits in relation to physiological characteristics in mulberry (*Morus* sp.) cultivars. *Turkish Journal of Botany* 36: 683–689.
- Laird S. A. and Kate, K. (2002). Linking biodiversity prospecting and forest conservation. 2002). *Selling forest environmental services—Market-based mechanisms for conservation and development*, 151-172.
- Langer, R.H.M. (1979). *How grasses grow*. 2nd ed. Edward Arnold (Publishers) Ltd., London.
- Liu Q., Zhao N. X., and Hao, G. (2005). Inflorescence structures and evolution in subfamily Chloridoideae (Gramineae). *Plant Systematics and Evolution*, 251(2-4): 183-198.
- Liu Q., Zhao N. X., Hao G., Hu X. Y., and Liu Y. X. (2005). Caryopsis morphology of the Chloridoideae (Gramineae) and its systematic implications. *Botanical Journal of the Linnean Society*, 148(1): 57-72.
- Looman J. (1982). *Prairie grasses identified and described by vegetative characters*. Research Station, Swift Current, Saskatchewan, Agriculture Canada.

- Maden K. (2004). Plant collection and herbarium techniques. *Our Nature*, 2(1): 53-57.
- Matsutani A. (1986). Identification of Italian millet from Esashika site by means of scanning electron microscope. *Journal of the Anthropological society of Nippon*, 94(1): 111-118.
- Matthews J. F. and Levins P. A. (1986). The systematic significance of seed morphology in *Portulaca* (Portulacaceae) under scanning electron microscopy. *Systematic Botany*. 11: 302-308
- McAlpine A. N. (1890). How to know grasses by their leaves. *Standard Cyc. Mod. Agri*, 6: 1-153.
- Meena S. L. (2004). Some new plants to the flora of Gujarat, India-I. *Journal of Economic and Taxonomic Botany*, 28(2): 387-388.
- Meena S. L. (2005). Some new plants to the flora of Gujarat, India-II. *Journal of Economic and Taxonomic Botany*, 29(2):375-377.
- Meena S. L., and Pandey R. P. (2004). A reassessment of the phytodiversity of Gujarat state: Floristic composition and floristic analysis, vegetation, threatened and rare taxa and their conservation strategies. *Journal of Economic and Taxonomic Botany*, 28(4): 867.
- Metcalfe C. R. (1960). *Anatomy of the monocotyledons. 1. Gramineae*. Oxford: Clarendon Press.
- Meyer M. H. and Gaynor V. (2000). A field guide and key to fifteen grass seedlings. *Journal of Natural Resources and Life Sciences Education*, 29:141-147.
- Moazzeni H., Zarre S., Al-Shehbaz I. A. and Mummenhoff K. (2007). Seed-coat microsculpturing and its systematic application in *Isatis* (Brassicaceae) and allied genera in Iran. *Flora-Morphology, Distribution, Functional Ecology of Plants*, 202(6): 447-454.
- Moulik S. (1997). *Grasses and bamboos of India*. Scientific Publishers.
- Mulligan G. A. and Bailey L. G. (1976). Seed coats of some Brassica and Sinapis weedy and cultivated in Canada. *Economic Botany*, 30(2): 143-148.
- Nelson D. M., Hu F. S., Scholes D. R., Joshi N. and Pearson, A. (2008). Using SPIRAL (Single Pollen Isotope Ratio AnaLysis) to estimate C3-and C4-grass abundance in the paleorecord. *Earth and Planetary Science Letters*, 269(1-2): 11-16.
- Nowosad F. S., Woodward J. C., and Wright L. E. (1942). The Effect of Fertilizer Treatments on the Yield and Chemical Composition of Pasture Species. *Scientific Agriculture*, 22(12):733-745.
- Olsson P., Folke C. and Berkes F. (2004). Adaptive comanagement for building resilience in social–ecological systems. *Environmental management*, 34(1):75-90.
- Osborne C.P., Visser V. Chapman S. Baker A. Freckleton R.P., Salamin N., and Simpson D. (2011). Uren V. GrassPortal: an online ecological and

evolutionary data facility. [WWW document]. URL <http://www.grassportal.org..>

- Osborne C.P and Freckleton R.P. (2009). Ecological selection pressures for C4 photosynthesis in the grasses. *Proc. R. Soc. Lond. B*, 276:1753–1760.
- Osman A.K, Mohammed A.Z, Hamed S.T and Hussein NR (2012). Fruit morphology of annual grasses *Egypt. Asian Journal of Plant Sciences* 11(6):268-284.
- Osman A. K., Zaki M. A., HamedS. T., and Hussein N. R. (2012). Fruits Morphology of Annual Grasses from Egypt. *Asian Journal of Plant Sciences*, 11(6): 268.
- Pandey R. P. (2001). Additions to the flora of Gujarat: 2. *J. Econ. Taxon. Bot*, 25(3): 623-625.
- Pandey R. P. (2002). Additions to the flora of Gujarat-III. *Journal of Economic and Taxonomic Botany*, 26(2): 505-508.
- Pandey R. P., & Padhye, P. M. (1997). Additions to the flora of Gujarat: 1. *J. Econ. Taxon. Bot*, 21(1), 103-106.
- Pandey, R. P. and Padhye P. M. (2005). Additions to the flora of Gujarat-IV. *Journal of Economic and Taxonomic Botany*, 29(2):346-351.
- Pandey R. P. and Padhye P. M. (2006). Additions to the Flora of Gujarat-IV. *J. Econ. Taxon. Bot*, 30(3): 603-608.
- Patel R. M. (1971). The flora of Bulsar and its environ. Ph. D. thesis, Sardar Patel University, Vallabh Vidyanagar, Gujarat.
- Peterson P. M. and Vega I. S. (2007). Eragrostis (Poaceae: Chloridoideae: Eragrostideae: Eragrostidinae) of Peru. *Annals of the Missouri Botanical Garden*, 94(4):745-790.
- Raghavan, R. S., Wadhwa B. M, Ansari M. Y. and. Rao R. S.(1981). A checklist of the plants of Gujarat. *Records of the Botanical Survey of India*. 21 (2): 1-127.
- Raole V. M. and Desai R. J. (2008). *Desmostachyapingalaiae* sp. nov. from Gujarat, India. *Nordic Journal of Botany*, 26(3-4): 196-198.
- Raole V. M., Desai R. J. and Veldkamp J. F. (2011). *Ischaemum sayajiraoi*, a new species of Poaceae from Gujarat, India. *Kew Bulletin*, 66(2):303.
- RedfordK. H. and Padoch C. (1992). *Conservation of neotropical forests: working from traditional resource use*. Columbia University Press.
- Reeder J. R. (1957). The embryo in grass systematics. *American Journal of Botany*. 44 (9): 756-768.
- Reimer E. (2006). *A phylogenetic study of Danthonia DC.(Poaceae) in North America* (Doctoral dissertation).
- Robson M. J., Ryle G. J. A. and Woledge J. (1988). The grass plant—its form and function. In *The grass crop* Springer, Dordrecht. (pp. 25-83).
- Santapau H. and Janardhanan K. P. (1966). The flora of Saurashtra. *Bulletin of the Botanical Survey of India*, 8: 1-58.

- Schmink M., Redford K. H. and Padoch C. (1992). Traditional peoples and the biosphere: framing the issues and defining the terms. *Conservation of neotropical forests: Working from traditional resource use*, 3-10.
- Schulz OE (1936). *Cruciferae*. In: A Engler & H Harms (ed.). Die Natürlichen Pflanzenfamilien ed. 2, 17B: 227-658. Leipzig: Verlag von Wilhelm Engelmann.
- Segarra J. G. and Mateu I. (2001). Seed morphology of *Linaria* species from eastern Spain: identification of species and taxonomic implications. *Botanical Journal of the Linnean Society*, 135(4): 375-389.
- Shah G. L. (1978). Flora of Gujarat state. Sardar Patel University. Pp. 1074.
- Shavvon, R. S. Mehrvarz, S. S. and Golmohammadi, N. (2012). Evidence from micromorphology and gross morphology of the genus *Loranthus* (Loranthaceae) in Iran. *Turkish Journal of Botany*, 36(6): 655-666.
- Singh J. S. (1988) The environmental problems: Issues and challenges. In: K.S. Valdiya (Eds.), *Kumaun: Land and People*. Gyanodaya Prakashan, Nainital, India. pp. 347-351
- Singh J.S., Hanxi Yang and Sajise P.F. (1985). *Structure and functional aspects of Indian and south east Asia savanna ecosystems. Ecology and management of world's savanna* (ed. by J. C. Tothill and J. J. Mott), pp. 34-55. Australian Academy of Sciences, Canberra.
- Spearman, P. J., Alexander, R. W. and Brodie, I. D. S. (2000). The use of microscale field mapping in a study of the ox-eye daisy (*Leucanthemum vulgare* L.) as a component of wild flower meadows. Pages 41–52 in R. W. Alexander A. C. and Millington, eds. *Vegetation mapping: From patch to planet*. John Wiley & Sons, Chichester, UK. 339 pp.
- Stanton M.L. (1984) Seed Variation in Wild Radish: Effect of Seed Size on Components of Seedling and Adult Fitness. *Ecology*, 65:1105-1112.
- Surrency D., Owsley M. and Kirkland M. (2006). *Seeling ID guide for native grasses in the Southeast*, United state of Agriculture. pp. 35.
- Terrell E. E. and Peterson P. M. (1993). Caryopsis morphology and classification in the Triticeae (Pooideae: Poaceae). *Smithsonian Contribution to Botany*. 1993, 83, 1–25.
- Tillich H. J. (2007). Seedling diversity and the homologies of seedling organs in the order Poales (Monocotyledons). *Annals of Botany*, 100(7): 1413-1429.
- Tzvelev N. (1989). The system of grasses (Poaceae) and their evolution. *The Botanical Review*, 55(3) :141-203.
- Undersander D., Casler M. D., Cosgrove D., and Martin N. E. (1996). *Identifying pasture grasses*. University of Wisconsin--Extension.
- Venable D L (1985) The evolutionary ecology of seed heteromorphism. *Am. Nat.* 126: 577-595.

- Wang S. and Henwood M. J. (1999). The taxonomic utility of micromorphological characters in Australian and New Zealand *Elymus* species (Poaceae). *Telopea*, 8(3): 351-362.
- Ward H. M. *Grasses*. (1901). Cambridge Univ. Press, London,
- Warwick, S. I., & Black, L. D. (1993). Molecular relationships in subtribe Brassicinae (Cruciferae, tribe Brassiceae). *Canadian Journal of Botany*, 71(7), 906-918.
- Warwick S. I. and Black L. D. (1994). Evaluation of the subtribes Moricandiinae, Savignyinae, Vellinae, and Zillinae (Brassicaceae, tribe Brassiceae) using chloroplast DNA restriction site variation. *Canadian Journal of Botany*, 72(11):1692-1701.
- Wenger K. F. (1984). *Forestry handbook* 84(1) eds John Wiley & Sons. John Wiley & Sons Inc.
- Whiffin T. and Tomb A. S. (1972). The systematic significance of seed morphology in the neotropical capsular-fruited Melastomataceae. *American Journal of Botany*. 59: 411-422.
- White E. G. (1984). A multispecies simulation model of grassland producers and consumers II. Producers. *Ecological modelling*, 24(3-4):241-262.
- Whittaker R.H. and Likens G.E. (1973) Primary productivity: The biosphere and man. *Human Ecol.* 1: 357-369.
- Whittaker R.H. and Likens G.E. (1975) The biosphere and man. In Lieth, G.E. and Whittaker, R.H. (Eds): Primary Productivity of the Biosphere. Ecological Studies edn., Vol. 14. Springer, New York, pp. 305-328.
- Whyte J. H. (1930). The recognition of some agricultural grasses by their vegetative characters. In *Transactions of the Botanical Society of Edinburgh* (Vol. 30, No. 3, pp. 206-208). Taylor & Francis Group.
- Wintle B., Moore G., Nichols P. (2009) *Identifying sub-tropical grass seedlings*. Bulletin No 4775, Department of Agriculture and Food Western Australia, South Perth, Australia.
- Yousaf Z., Shinwari Z.K., Qureshi R.A. and Perveen A. (2008) Leaf epidermal anatomy of selected Allium species, family Alliaceae from Pakistan. *Pak. J. Bot.*, 40(1): 77-90.
- Yusufzai A. E. and Gandhi A. P. (1999). *Grasses of Gujarat*. Gujarat Agricultural University. Directorate of Research, Gujarat Agricultural University India. pp. 277.
- Zhang Y., Xiaoying H. U., Yunxiao L. and Qing L. (2014) Caryopsis morphological survey of the genus *Themeda* (Poaceae) and allied spathaceous genera in the Andropogoneae. *Turkish Journal of Botany*, 38: 665–676.
- Zhang Z. Y., Yang D. Z., Lu A. M. and Knapp S. (2005). Seed morphology of the tribe Hyoscyameae (Solanaceae). *Taxon*, 54(1):71-83