

4. References



- Abbasi M. and Assadi M. (2014). Evaluation of generic concept in *Colpodium* Trin. and *Catabrosella* Tzvel. (Tzvel.): a new subspecies in Poaceae family in Iran. *International Journal of Biosciences*. 5: 50–57.
- Abbasi M., Assadi M., Attar, F. and Nejadsttari T. (2010) Micromorphological and anatomical features of *Puccinellia dolicholepis* (Poaceae), a new record for the flora of Iran. *Iranian Journal of Botany*. 16: 267–272.
- Aliscioni S. S., Ospina, J. C. and Gomiz, N. E. (2016). Morphology and leaf anatomy of *Setaria* (Poaceae: Panicoideae: Paniceae) and its taxonomic significance. *Plant systematics and evolution*. 302(2): 173-185.
- Allen P. S., Benech-Arnold R. L., Batlla D., Bradford K. J., and Nonogaki, H. (2007). *Seed development, dormancy and germination*. Oxford: Wiley-Blackwell. pp. 392.
- Alvarez J. M., Rocha J. F. and Machado, S. R. (2008). Bulliform cells in *Loudetiopsis chrysothrix* (Nees) Conert and *Tristachya leiostachya* Nees (Poaceae): structure in relation to function. *Brazilian Archives of Biology and Technology*. 51(1): 113-119.
- Alvarez, J. M., Rocha, J. F. and Machado S. R. (2005). Estrutura foliar de *Loudetiopsis chrysothrix* (Nees) Conert and *Tristachya leiostachya* Nees (Poaceae). *Revta. brasil. Bot.* 28:23-37.
- America O. T. N., Stebbins, G. L. and Crampton B. (1961). The natural classification of the gramineae. *Recent advances in botany*. 1: 133-145.
- Arber A. (1918). The phyllode theory of the monocotyledonous leaf, with special reference to anatomical evidence. *Annals of botany*. 32(128): 465-501.
- Arber A. (1923). Leaves of the Gramineae. *Bot. Gaz.* 76: 374-388.
- Arber A. (1937). The interpretation of the flower: a study of some aspects of morphological thought. *Biological Reviews*. 12(2): 157-184.
- Arber A. (2010). *The Gramineae: a study of cereal, bamboo and grass*. Cambridge University Press. pp. 506.
- Armstrong J. M. (1937). A cytological study of the genus *Poa* L. *Canadian Journal of Research*. 15(6): 281-297.
- Armstrong J., Jones R. E. and Armstrong W. (2006). Rhizome phyllosphere oxygenation in *Phragmites* and other species in relation to redox potential, convective gas flow, submergence and aeration pathways. *New Phytologist*. 172(4): 719-731.
- Artschwager E., Brandes E. W. and Starrett R. C. (1929). Development of flower and seed of some varieties of sugar cane. *J. agr. Res.* 39(1): 1-30.
- Avdulov N. P. (1931). Karyo-systematische Untersuchungen der Familie Gramineen. (In Russian; German summary.) *Bull. Appl. Bot. Suppl.* 44. pp. 428.

- Barker N. P., Clark L. G., Davis J. I., Duvall M. R., Guala G. F and Mathews, S. Y. (2001). Phylogeny and subfamilial classification of the grasses (Poaceae). *Annals of the Missouri Botanical Garden*. 88: 373-457.
- Barker N. P., Linder H. P. and Harley E. H. (1999). Sequences of the grass-specific insert in the chloroplast rpoC2 gene elucidate generic relationships of the Arundinoideae (Poaceae). *Systematic Botany*. 23: 327-350.
- Bechtel D. B. and Pomeranz Y. (1977). Ultrastructure of the mature ungerminated rice (*Oryza sativa*) caryopsis: The caryopsis coat and the aleurone cells. *American Journal of Botany*. 64 (8): 966-973.
- Bews, J. W. (1929). *World'S Grasses: Their differentiation, distribution, economic and ecology*. Longmans: Green and Co. Ltd., London. pp. 408.
- Blackman E. (1969). Observations on the development of the silica cells of the leaf sheath of wheat (*Triticum aestivum*). *Canadian Journal of Botany*. 47(6): 827-838.
- Bor. N. L. (1960). *The Grasses o f Burma, Ceylon, India and Pakistan*. London: Pergamon Press. pp. 767.
- Bowen C. C. (1956). Freezing by liquid carbon dioxide in making slides permanent. *Stain Technology*. 31(2): 87-90.
- Breakwell E. (1914). A study of the leaf anatomy of some native species of the genus *Andropogon*, N.O. Gramineae. *Proc. Linn. Soc. N.S.W.* 39: 385-394.
- Bremer K. (2002). Gondwanan evolution of the grass alliance of families (Poales). *Evolution*. 56(7): 1374-1387.
- Brown W. V. (1960). A cytological difference between the Eupanicoideae and the Chloridoi- deae (Gramineae). *Southwest. Nat.* 5:7-11.
- Brown W. V. (1974a). Another cytological difference among the Kranz Sub-families of the Gramineae. *Bulletin of Torrey Botanical Club*. 101(3): 120-123.
- Brown W. V. (1974b). Morphological variation and chromosome numbers of North American populations of *Koeleria cristata*. *Bulletin of Torrey Botanical Club*. 101(3): 124-128.
- Brown W. V. (1975). Variations in anatomy, associations, and origins of kranz tissue. *Am J Bot.* 62: 395-402.
- Brown W. V. (1977). The Kranz syndrome and its subtypes in grass systematics. *Memoirs of the Torrey Botanical Club*. 23(3): 1-97.
- Brown W. V. and Emery W. H. (1957). Persistent nucleoli and grass systematics. *American Journal of Botany*. 44: 585-590
- Brown W. V., Heimsch C. and Emery W. H. P. (1957). The organization of the grass shoot apex and systematics. *American Journal of Botany*. 44: 590-595.
- Brown, W. V. (1958). Leaf anatomy in grass systematics. *Botanical Gazette*. 119(3): 170-178.
- Bugnon P. (1921). *La feuille chez les Graminees*. Paris.

- Burton R. A. and Fincher G. B. (2014). Evolution and development of cell walls in cereal grains. *Frontiers in plant science*. 5: 456.
- Calderón C. E. and Soderstrom T. R. (1973). Morphological and anatomical considerations of the grass subfamily Bambusoideae based on the genus *Maclurolyra*. *Smith. Contr. Bot.* 11: 1-54.
- Canfield R. H. (1934). Stem structure of grasses on the *Jornada* experimental Range. *Botanical Gazette*. 95(4): 636-648.
- Carlquist S. and Schneider E. L. (1997). Origins and nature of vessels in monocotyledons. I. *Acorus*. *International Journal of Plant Sciences*. 158(1): 51-56.
- Cenci C. A., Grando S. and Ceccarelli, S. (1984). Culm anatomy in barley (*Hordeum vulgare*). *Canadian journal of botany*. 62(10): 2023-2027.
- Chaffey N. J. (1984). Epidermal structure in the ligules of four species of the genus *Poa* L. (Poaceae). *Botanical Journal of Linnean Society*. 89: 341–354.
- Chaffey N. J. (1994). Structure and function of the membranous grass ligule – a comparative study. *Botanical Journal of Linnean Society*. 116: 53–69.
- Chaffey N. J. (2000). Research review: Physiological anatomy and function of the membranous grass ligule. *New Phytologist*. 146: 5–21.
- Chakravarty A. K. and Verma C. M. (1966). Anatomical study in arid zone grasses of Western Rajasthan. *J. Ind. bot. Soc.* 44: 506-511.
- Choi H., Kim W. and Shin M. (2004). Properties of Korean amaranth starch compared to waxy millet and waxy sorghum starches. *Starch-Stärke*. 56(10): 469-477.
- Chonan N. (1978). Comparative anatomy of mesophyll among the leaves of gramineous crops. *Jpn. Agric. Res. Q.*, 12: 128-131.
- Clayton W. D. (1989). *Melinis* Beauv.–Pp. 115–128 in: Launert E. and Pope G.(ed.), *Flora zambesiaca* 10 (3). London: *Flora Zambesiaca Managing Committee*.—Published at: <http://apps.kew.org/efloras/search.do>.
- Clayton W. D. and Renvoize S. A. (1986). *Genera Graminum*. Kew Bulletin Adicional Series XIII, London. pp. 389.
- Clayton, W. D., & Renvoize, S. A. (1982). *Flora of Tropical East Africa: Gramineae* (Part 3). Royal Botanic Gardens, Kew. pp. 451.
- Colmer T. D. (2003). Long-distance transport of gases in plants: a perspective on internal aeration and radial oxygen loss from roots. *Plant, Cell and Environment*. 26(1): 17-36.
- Columbus J. T. (1996). *Lemma micromorphology, leaf anatomy and phylogenetics of Bouteloua, Hilaria and relatives (Gramineae: Chlorideae: Boutelouinae)*. Ph.D. dissertation. Univeristy of California. Berkeley. pp. 258.
- Crookston R. K. and Moss D. N. (1973). A variation of C4 leaf anatomy in *Arundinella hirta* (Gramineae). *Plant physiology*. 52(5): 397-402.
- De Agrasar Z. E. R. and Rodríguez M. F. (2002). Cauline anatomy of native woody bamboos in Argentina and neighbouring areas: epidermis. *Botanical Journal of the Linnean Society*. 138(1): 45-55.

- DE Wet J. M. J. (1960). Leaf anatomy in six South African grass genera. *Bothalia*. 7: 299-301.
- Decker H. F. (1964). Affinities of the grass genus *Ampelodesmos*. *Brittonia*. 16(1): 76-79.
- Dengler N. G., Dengler R. E., Donnelly P. M. and Hattersley P. W. (1994). Quantitative leaf anatomy of C3 and C4 grasses (Poaceae): bundle sheath and mesophyll surface area relationships. *Annals of Botany*. 73(3): 241-255.
- Dengler N. G., Donnelly P. M. and Dengler R. E. (1996/4) Differentiation of bundle sheath, mesophyll, and distinctive cells in the C₄ grass *Arundinella hirta* (Poaceae). *American Journal of Botany*. 83(11): 1391-1405.
- Deshpande B. D. and Sarkar S. (1962). Nodal anatomy and the vascular system of some members of the Gramineae. *In Proc. Nat. Inst. Sci. India Bulletin*. 28: 1-12.
- Duval-Jouve J. (1875). Histotaxie des feuilles de Graminees. *Ann. Sci. Nat. Bot. Ser.* 6(1): 294-371.
- Eichenberg M. T. and Scatena V. L. (2013). Morphology and anatomy of the diaspores and seedling of *Paspalum* (Poaceae, Poales). *Anais da Academia Brasileira de Ciências*. 85(4): 1389-1396.
- Eleftheriou E. P. and Noitsakis B. (1978). A comparative study of the leaf anatomy of the grasses *Andropogon ischaemum* and *Chrysopogon gryllus*. *Phyton*. 19: 27-36.
- Ellis R. P. (1974). Anomalous vascular bundle sheath structure in *Alloteropsis semialata* leaf blades. *Bothalia*. 11(3): 273-275.
- Ellis R. P. (1976). A procedure for standardizing comparative leaf anatomy in the Poaceae. I. The leaf-blade as viewed in transverse section. *Bothalia*. 12(1): 65-109.
- Ellis R. P. (1987). A review of comparative leaf blade anatomy in the systematics of the Poaceae: the past twenty-five years. Soderstrom T R, and others ed (s). *Grass systematics and evolution: an International Symposium held at the Smithsonian Institution, Washington, DC, 27-31 July 1986*. Smithsonian Institution Press, Washington, DC, London.
- Esau K. (1965). *Plant anatomy* (2nd ed.). John wiley and sons, New York. Pp. 767.
- Esau, K. (1953). *Plant anatomy*. John wiley and sons, New York. Pp. 407.
- Fahn A. (1990). *Plant anatomy* (4th edn). Oxford, Pergamon.
- Fisher B. S. (1939). A contribution to the leaf anatomy of Natal grasses, Series I: *Chloris* Sw. and *Eustachys* Desv. *Ann. Natal Mus.* 9: 245-267.
- Garnier E. and Laurent G. (1994). Leaf anatomy, specific mass and water content in congeneric annual and perennial grass species. *New Phytologist*. 128(4), 725-736.
- Gasser M., Vegetti A. C. and Tivano J. A. (1994). Anatomía de las estructuras foliares y caulinares senescentes productivos de *Eleusine indica*

- (L.) Gaertner y E. tristachya (Lam.) Lam. In *Resúmenes VI Congreso Latinoamericano de Botánica. Mar del Plata*. pp. 39.
- Ghahraman A., Alemi M., Atar F., Hamzeh B. and Columbus T. (2006). Anatomical studies in some species of *Bromus* L.(Poaceae) in Iran. *Iran. Journ. Bot.* 12 (1): 1-14.
 - Gibson D. J. (2009). *Grasses and grassland ecology*. Oxford University Press.
 - Gibson D. J. (2009). *Grasses and grassland ecology*. Oxford University Press.
 - Goossens A. P. (1938). A study of the South African species of *Sporobolus* with special reference to leaf anatomy. *Trans. R. Soc. S. Afr.* 26: 173-223.
 - Gopal B. H. and Ram H. M. (1985). Systematic significance of mature embryo of bamboos. *Plant systematics and evolution*. 148(3-4): 239-246.
 - Grosser D. and Liese W. (1971). On the anatomy of Asian bamboos, with special reference to their vascular bundles. *Wood Science and technology*. 5(4): 290-312.
 - Guerin M. P. (1898). On the development of seminal integuments and pericarp of Gramineae. *Bulletin of the Botanical Society of France*. 45 (5): 405-411.
 - Haberlandt G. (1914) *Physiological plant anatomy*. Macmillan and Co, London.
 - Hameed M., Ashraf M. and Na, N. (2009). Anatomical adaptations to salinity in cogon grass [*Imperata cylindrica* (L.) Raeuschel] from the Salt Range, Pakistan. *Plant and soil*. 322(1-2): 229-238.
 - Hands P., Kourmpetli S., Sharples D., Harris R. G. and Drea S. (2012). Analysis of grain characters in temperate grasses reveals distinctive patterns of endosperm organization associated with grain shape. *Journal of experimental botany*. 63(17): 6253-6266.
 - Harrington G. T. and Crocker W. and Plant P. (1923). Structure, physical characteristics, and composition of the pericarp and integument of Johnson grass seed in relation to its physiology. *Journal of Agricultural Research*. 23(3): 193-222.
 - Hattersley P. W. (1984). Characterization of C₄ type leaf anatomy in grasses (Poaceae). Mesophyll: bundle sheath area ratios. *Annals of Botany*. 53(2): 163-180.
 - Hattersley P. W. Wong S. C., Perry S., and Roksandic Z. (1986). Comparative ultrastructure and gas exchange characteristics of the C₃-C₄ intermediate Neurachne minor ST Blake (Poaceae). *Plant, Cell & Environment*. 9(3): 217-233.
 - Hinton J. J. and Shaw B. (1954). The distribution of nicotinic acid in the rice grain. *Brit. J. Nutrition*. 8: 65-71.
 - Hinton J. J. C. (1948). The Distribution of Vitamin B 1 in the Rice Grain. *British Journal of Nutrition*. 2(3): 237-241.
 - Hisham H. N., Othman S., Rokiah H., Latif M. A., Ani S. and Tamizi M. M. (2006). Characterization of bamboo *Gigantochloa scortechinii* at different ages. *Journal of Tropical Forest Science*. 18: 236-242.

- Hitch P. A. and Sharman B. C. (1971). The vascular pattern of festucoid grass axes, with particular reference to nodal plexi. *Botanical Gazette*. 132(1): 38-56.
- Hitch P. A. and Shiarman B. C. (1968). Initiation of procambial strands in leaf primordia of *Dactylis glomerata* L. as an example of a temperate herbage grass. *Ann. Bot.* 32: 153-164.
- Hitchcock A. S. (1914). *A text-book of grasses*. The Macmillan Company, New York.
- Hitchcock A. S. (1922). *A textbook of grasses*. Macmillan Co., New York.
- Hodkinson T. R., Chase M. W., Lledó D. M., Salamin N. and Renvoize S. A. (2002). Phylogenetics of *Miscanthus*, *Saccharum* and related genera (Saccharinae, Andropogoneae, Poaceae) based on DNA sequences from ITS nuclear ribosomal DNA and plastid trnL intron and trnL-F intergenic spacers. *Journal of plant research*. 115(5): 381-392.
- Hossain M. A. (2010). *Culm Internode and Leaf Anatomy of Two Economic Grass of Bangladesh* [Napier and Kas]. M.S. Thesis, Rajshahi University, Bangladesh.
- Hsiao C., Jacobs S. W. L., Barker N. P. and Chatterton N. J. (1998). A molecular phylogeny of the subfamily Arundinoideae (Poaceae) based on sequences of rDNA. *Australian Systematic Botany*. 11(1): 41-52.
- Huang X. Y., Qi J. Q., Xie J. L., Hao J. F., Qin B. D. and Chen S. M. (2015). Variation in anatomical characteristics of bamboo, *Bambusa rigidula*. *Sains Malays*. 44(1): 17-23.
- Hubbard C. E. (1934). Gramineae. J. Hutchinson: *The families of Flowering plants*. 2: 199-299.
- Irving D. W. (1983). Anatomy and histochemistry of *Echinochloa turnerana* (channel millet) spikelet. *Cereal Chemistry*. 60(2): 155-160.
- Izaguirre de Artucio, P. and Laguardia A. (1987). Un nuevo enfoque hacia la definicion del fruto de las gramineas. *Uruguay Fac. Agron. Bot. Invest.* 3: 1-16
- Jane W. N. (2004). Ultrastructure of endosperm development in *Arundo formosana* Hack. (Poaceae) from differentiation to maturity. *Botanical Bulletin of Academia Sinica*. 45: 69-85.
- Jane W. N. and Chiang S. H. T. (1991). Morphology and development of bulliform cells in *Arundo formosana* Hack. *Taiwania*. 36: 85-97.
- Jauhar P. P. and Joshi A. B. (1967). Cytotaxonomic Investigations in the *Panicum maximum* Jacq. Complex II. Studies on Epidermal Pattern. *Nelumbo*. 9(1-4): 59-62.
- Joarder N., Roy A. K., Sima S. N. and Parvin K. (2010). Leaf blade and midrib anatomy of two sugarcane cultivars of Bangladesh. *Journal of Bio-Science*. 18: 66-73.
- Jung M. J., Veldkamp J. F. and Kuoh C. S. (2008). Notes on *Eragrostis wolf* (Poaceae) for the flora of Taiwan. *Taiwania*. 53(1): 96-102.

- Justin S. H. F. W. and Armstrong W. (1987). The anatomical characteristics of roots and plant response to soil flooding. *New Phytologist*. 106(3): 465-495.
- Kanai R. and Kashiwagi M. (1975). *Panicum milioides*, a Gramineae plant having Kranz leaf anatomy without C4-photosynthesis. *Plant and Cell Physiology*. 16(4): 669-679.
- Kar R.K., Misra N.M. and Kabi T. (2002). *Textbook on fundamentals of botany*. 4th Ed. Kalyani publishers. India.
- Kar R.K., Misra N.M. and Kabi T. (2002). *Textbook on fundamentals of botany*. 4th Ed. Kalyani publishers. India.
- Kaufman B. P. (1956). Cytological studies of changes induced in cellular materials by ionizing radiations. *Annals of New York Academic Sciences*. 59: 1-53.
- Kawai M., Samarajeewa P. K., Barrero R. A., Nishiguchi M. and Uchimiya H. (1998). Cellular dissection of the degradation pattern of cortical cell death during aerenchyma formation of rice roots. *Planta*. 204(3): 277-287.
- Kawai M., Umeda M. and chimiya H. (1998). Stimulation of adenylate kinase in rice seedlings under submergence stress. *Journal of plant physiology*. 152(4-5): 533-539.
- Keshavarzi M., Khaksar M. and Seifali M. (2007). Systematic study of annual weed *Phalaris minor* Retz (Poaceae) in Iran. *Pakistan Journal of Biological Sciences*. 10(8): 1336-1342.
- Keshavarzi M., Rahiminejad M.R. and Kheradmandnia M. (2002). Anatomical and morphological variation of *Aegilops triuncialis* L. of Iran. *Pagoheesh Sazandegi*. 5: 14-20.
- Keshavarzi M., Seifali M. and Babaii K. (2007). A Morphological and Anatomical Study of an annual grass *Eremopyrum* (Poaceae) in Iran. *Pakistan journal of Biological sciences*. 10(1): 32-40.
- Kiesselbach T. A. (1949). The structure and reproduction of corn. University of Nebraska, College of Agriculture, Agricultural Experiment Station. *Research Bulletin*, 161. University of Nebraska College of Agriculture, Agricultural Experiment Station, Lincoln. pp. 96.
- Kok P. D. F and Van Der Schijff, H. P. (1973). A key based on epidermal characteristics for the identification of certain highveld grasses. *Koedoe*. 16(1): 27-43.
- Korzun V., Malyshev S., Voylovkov A. and Borner A. (1997). RFLP-based mapping of three mutant loci in rye (*Secale cereale*) and their relation to homoeologous loci within the Gramineae. *Theoretical and Applied Genetics*. 95: 468-473.
- Kumar N. and Nautiyal S. (2017). Leaf anatomy of two genera of tribe Eragrostideae (Poaceae) from Mandal forest of Kedarnath wildlife sanctuary, Uttarakhand, India. *International Journal of Botany Studies*. 2(5): 50-55.

- Kuo J., O'Brien T.P., Canny M.J., (1974) Pit-field distribution, plasmodesmatal frequency, assimilate flux in the mestome sheath cells of wheat leaves. *Planta*. **121**: 97–118.
- Lambers J. H. R., Harpole W. S., Tilman D., Knops J. and Reich P. B. (2004). Mechanisms responsible for the positive diversity–productivity relationship in Minnesota grasslands. *Ecology Letters*. **7**(8): 661–668.
- Liese W. (1980). Anatomy of bamboo. In *Bamboo research in Asia*. In Proceedings of a workshop held in Singapore. International Development Research Centre Singapore: Ottawa. pp. 161–164.
- Liese W. (1985). Anatomy and Properties of Bamboo. In International Bamboo Workshop (INBAR). October 6-14. China. pp. 196-208.
- Liese W. (1998). *The anatomy of bamboo culms*. Technical report: International network for Bamboo and Rattan, Beijing. pp. 207.
- Londoño X., Camayo G. C., Riaño N. M. and López, Y. (2002). Characterization of the anatomy of *Guadua angustifolia* (Poaceae: Bambusoideae) culms. *Bamboo Science and Culture: The Journal of the American Bamboo Society*. **16**(1): 18-31.
- Longhi-Wagner H. M. (2001). Tribo Arundinelleae. In: Wanderley, M. G. L.; Shepherd, G. J. and Giulietti, A. M. *Flora fanerogâmica do Estado de São Paulo, Poaceae*. Fapesp - Hucitec, São Paulo. Vol.1, pp. 119-123.
- Lundgren M. R., Osborne C. P. and Christin P. A. (2014). Deconstructing Kranz anatomy to understand C4 evolution. *Journal of experimental botany*. **65**(13): 3357-3369.
- Makino Y. and Ueno O. (2018). Structural and physiological responses of the C4 grass *Sorghum bicolor* to nitrogen limitation. *Plant Production Science*. **21**(1): 39-50.
- Mann A. and Harlan H. V. (1916). Morphology of the Barley Grain with Reference to its Enzyme-Secreting Areas. *Journal of the Institute of Brewing*. **22**(2): 73-108.
- Martínez-Sagarra G., Abad P. and Devesa J. A. (2017). Study of the leaf anatomy in cross-section in the Iberian species of *Festuca* L. (Poaceae) and its systematic significance. *PhytoKeys*. (83): 43-74.
- Matumura M. and Nakajima N. (1988). Comparative ecology of intraspecific variation of the Chigaya, *Imperata cylindrica* Var. *koenigii* (Alono-along). III. Annual growth of the 3rd year communities originated from the seedlings. *J. Jap. Soc. Grassl.* **34**: 77-84.
- Mauseth J. D. (1988). *Plant anatomy*. Benjamin Cummings, California.
- Mendes M. M., Gazarini L. C., Rodrigues M. L. (2001) Acclimation of *Myrtus communis* to contrasting Mediterranean light environments effects on structure and chemical composition of foliage and plant water relations. *Environ Exp Bot.* **45**:165–178.
- Metcalfe C. R. (1960). *Anatomy of the monocotyledons*. 1. *Gramineae*. Oxford: Clarendon Press. pp. 731.

- Mohmod A. L. and Mustafa M. T. (1992). Variation in anatomical properties of three Malaysian bamboos from natural stands. *Journal of Tropical Forest Science*. 5(1): 90-96.
- Mojzes A. and Kalapos T. (2005). Leaf anatomical plasticity of *Brachypodium pinnatum* (L.) Beauv. growing in contrasting microenvironments in a semiarid loess forest-steppe vegetation mosaic. *Community ecology*. 6(1): 49-56.
- Moreno M. A., Harper L. C., Krueger R. W., Dellaporta S. L. and Freeling M. (1997). Liguleless 1 encodes a nuclear-localized protein required for induction of ligules and auricles during maize leaf organogenesis. *Genes and Development* 11: 616–628.
- Moulia B. (2000). Leaves as shell structures: double curvature, auto-stresses, and minimal mechanical energy constraints on leaf rolling in grasses. *J. Pl. Growth Reg.* 19: 19-30.
- Narayanaswami S. (1953). The structure and development of the caryopsis in some Indian millets. 1. *Pennisetum typhoideum* Rich. *Phytomorphology*. 3: 98-112.
- Narayanaswami S. (1954). The Structure and Development of the Caryopsis in Some Indian Millets. III. *Paspalum scrobiculatum* L. *Bulletin of the Torrey Botanical Club*. 81: 288-299.
- Negbi, M. (1984). The structure and function of the scutellum of the Gramineae. *Botanical journal of the Linnean Society*. 88(3): 205-222.
- Neumann H. (1938). Zur Kenntnis der Anatomie und ersten Anlage der Graminenligula. *Beitrage zur Biologie der Pflanzen*. 25: 1–22.
- Oguchi R., Hikosaka K. and Hirose T. (2003). Does the photosynthetic light-acclimation need change in leaf anatomy?. *Plant, Cell & Environment*. 26(4): 505-512.
- Ogundipe O. T. and Olatunji O. A. (1992). Systematic anatomy of *Brachiaria* (Trin.) Griseb. (Poaceae). *Feddes Repertorium*. 103(1-2): 19-30.
- Olsen J. T., Caudle K. L., Johnson L. C., Baer S. G. and Maricle B. R. (2013). Environmental and genetic variation in leaf anatomy among populations of *Andropogon gerardii* (Poaceae) along a precipitation gradient. *American Journal of Botany*. 100(10): 1957-1968.
- Olsen O. A. (2001). Endosperm development: cellularization and cell fate specification. *Annual review of plant biology*. 52(1): 233-267.
- Olsen O. A., Linnestad, C. and Nichols S. E. (1999). Developmental biology of the cereal endosperm. *Trends in plant science*. 4(7): 253-257.
- Osman A. K., Zaki M. A., Hamed S. T. and Hussein N. R. (2012). Comparative anatomical study on fruits of some tribes of family Gramineae from Egypt. *Pak. J. Bot.* 44(2): 599-618.
- Paliwal G. S. (1969). Stomatal ontogeny and phylogeny. I. Monocotyledons. *Acta botanica neerlandica*. 18(5): 654-668.
- Parameswaran N. and Liese W. (1981). Torus-like structures in interfibre pits of *Prunus* and *Pyrus*. *IAWA Journal*. 2(2-3): 89-93.

- Patel S. R. (1976). *Thesis: Morpho taxonomic studies in some Gramineae*. Sardar Patel University.
- Pathak S., Kar S., and Singh P. (2013). Ligules as aid to Identification of Grasses. *Pleione* 7(1): 241 - 246.
- Picket-Heaps J. D. and Northcote D. H. (1966). Cell division in the formation of the stomatal complex of the young leaves of wheat. *J. Cell. Sci.* 1 : 121-128.
- Prat, H. (1932). L'epiderme des Graminées. Étude anatomique et systematique. *Ann. Sci. Nat. Bot., Ser.* 10(14): 117-324.
- Prat, H. (1936). La Sistématique des Graminées. *Ann. Sci. Nat. Bot., Ser.* 10(18): 165-257.
- Ramos J. C., Tivano J. C. and Vegetti A. C. (2002). Estudioanatomico de vastagos reproductivosen Bromus auleticus Trin.ex Ness (Poaceae). *Gayana. Botánica.* 59(2): 51-60.
- Reeder J. R. (1953). Affinities of the grass genus Beckmannia Host. *Bulletin of the Torrey Botanical Club.* 80: 187-196.
- Reeder J. R. (1957). The embryo in grass systematics. *American Journal of Botany.* 44 (9): 756-768.
- Reichert E. T. (1913). Differentiation and specificity of starches in relation to genera, species, etc. *Stereo-Chemistry Applied to Protoplasmic Processes and Products, and as a Strictly Scientific Basis for the Classification of Plants and Animals* by Edward Tyson Reichert. 41. pp. 686-689
- Renvoize S. (2002). Grass anatomy. *Flora of Australia.* 43: 71-132.
- Renvoize S. A. (1982). A Survey of Leaf-Blade Anatomy in Grasses III. Garnotieae *Kew Bulletin.* 37(3): 497-500.
- Renvoize S. A. (1986). A Survey of Leaf-Blade Anatomy in Grasses IX. Centothecoideae. *Kew Bulletin.* 41(2): 339-342
- Renvoize S. A. (1982). A survey of Leaf-Blade anatomy in Grasses II. Arundinelleae. *Kew bulletin.* 37(3): 489-495.
- Renvoize S. A. (1982). A survey of leaf-blade anatomy in grasses. I. Andropogoneae. *Kew Bulletin.* 37(2): 315-321.
- Renvoize S. A. (1983). A survey of leaf-blade anatomy in grasses IV. Eragrostideae. *Kew bulletin.* 38(3): 469-478.
- Renvoize S. A. (1986). A Survey of Leaf-Blade Anatomy in Grasses VIII. Arundinoideae *Kew Bulletin.* 41(2): 323-338.
- Renvoize S. A. (1987). A survey of leaf-blade anatomy in grasses XI. Paniceae. *Kew Bulletin.* 42: 739-768.
- Renvoize S. A. (1987). A survey of leaf-blade anatomy in grasses X: Bambuseae. *Kew Bulletin.* 42(1): 201-207.
- Rost T. L. (1971). Fine structure of endosperm protein bodies in *Setaria lutescens* (Gramineae). *Protoplasma.* 73(3-4): 475-479.
- Rost T. L. (1973). The anatomy of the caryopsis coat in mature caryopses of the yellow foxtail grass (*Setaria lutescens*). *Botanical Gazette.* 134(1): 32-39.

- Rost T. L., De Artucio P. I., and Risley E. B. (1984). Transfer cells in the placental pad and caryopsis coat of *Pappophorum subbulbosum* Arech.(Poaceae). *American journal of botany*. 71(7): 948-957.
- Row H. C. and Reeder J. R. (1957). Root-hair development as evidence of relationships among genera of Gramineae. *American Journal of Botany* 44: 596-601.
- Rudall P. J., Stuppy W., Cunniff J., Kellogg E. A. and Briggs B. G. (2005). Evolution of reproductive structures in grasses (Poaceae) inferred by sister-group comparison with their putative closest living relatives, Ecdociaceae. *American Journal of Botany*. 92(9): 1432-1443.
- Sage T. L. and Sage R. F. (2009). The Functional Anatomy of Rice Leaves: Implications for Refixation of Photorespiratory CO₂ and Efforts to Engineer C4 Photosynthesis into Rice. *Plant and Cell Physiology*. 50(4): 756-772.
- Sanchez E., Arriaga M. O. and Ellis R. P. (1989). Kranz distinctive cells in the culm of *Arundineua* (Arundinelleae; Panicoideae; Poaceae). *Bothalia*. 19(1): 45-52.
- Sanchez E., Arriaga M. O. and Ellis R. P. (1990). Kranz distinctive cells in the culm of *Arundineua* (Arundinelleae; Panicoideae; Poaceae). *Bothalia*. 20(1): 45-52.
- Sanders E. H. (1955). Developmental morphology of the kernal of grain sorghum. *Cereal Chem*. 32: 12-25.
- Sangster A. G. (1985). Silicon Distribution and Anatomy of the Grass Rhizome, with Special Reference to *Miscanthus sacchariflorus* (Maxim.) Hackel. *Annals of botany*. 55(5): 621-634.
- Sangster A. G. and Parry D. W. (1969). Some factors in relation to bulliform cell silicification in the grass leaf. *Annals of Botany*. 33(2): 315-323.
- Sargent E., Robertson A. and Hill Q. (1905). The anatomy of the scutellum in *Zea mais*. *Annals of Botany*. 19(73): 115-123.
- Schlickum A. (1896). Morphologischer und anatomischer Vergleich der Kotyledonen und ersten Laubblätter der Keimpflanzen der Monokotylen, *Bibliotheca Botanica*. 35:1-88.
- Schwendener S. (1890). Die Mestomscheiden der Gramineenblätter. *S.B. Akad. Wiss. Berlin*. 22: 405-426.
- Seago J. L., Marsh L. C., Stevens K. J., Soukup A. and Votrubova O. (2005). A re-examination of the root cortex in wetland flowering plants with respect to aerenchyma. *Ann. Bot.* 96: 565–579.
- Sekar T. and Balasubramanian A. (1994). Culm anatomy of *Guadua* and its systematic position. *BIC, Indian Bulletin*. 4: 6-9.
- Shah C. K. and Sreekumari S. B. (1980). Developmental morphology and homologies in grass embryo. *Current Science*. 49(7): 284-285.

- Shapter F. M., Henry R. J. and Lee L. S. (2008). Endosperm and starch granule morphology in wild cereal relatives. *Plant Genetic Resources*. 6(2): 85-97.
- Shields L. M. (1951). The involution mechanism in leaves of certain grasses. *Phytomorphology*. 1: 225-251.
- Sima S. N., Roy A. K. and Joarder N. (2015). Erect culm internodal anatomy and properties of sun ecotype of *Imperata cylindrica* (L.) P. Beauv. *Bangladesh Journal of Botany*. 44(1): 67-72.
- Siqueiros Delgado M. E. and Herrera Arrieta Y. (1996). Taxonomic value of culm anatomical characters in the species of *Bouteloua lagasca* (Poaceae: Eragrostideae). *Phytologia*. 81(2): 124-141.
- Siqueiros-Delgado M. (2007). Anatomía del tallo de *Bouteloua* y géneros relacionados (Gramineae: Chloridoideae: Boutelouinae). *Acta botánica Mexicana*. (78): 39-54.
- Siqueiros-Delgado M. E., Ainouche M., Columbus J. T. and Ainouche A. (2013). Phylogeny of the *Bouteloua curtispindula* complex (Poaceae: Chloridoideae) based on nuclear ribosomal and plastid DNA sequences from diploid taxa. *Systematic botany*. 38(2): 379-389.
- Stebbins G. L. (1956). Taxonomy and the evolution of genera, with special reference to the family Gramineae. *Evolution*. 10(3): 235-245.
- Stebbins G. L. and Crampton B. (1961). A suggested revision of the grass genera of temperate North America. *Recent Advances in Botany*. 1: 133-144.
- Stebbins G. L. and Jain S. K. (1960). A study of stomatal development in *Allium*, *Rhoeo* and *Commelina*. *Developmental Biol.* 1: 409-426.
- Stebbins G. L. and Khush G. S. (1961). Variations in the organization of the stomatal complex in the leaf epidermis of monocotyledons and its bearing on their phylogeny. *Amer. J. Bot.* 48: 51-59.
- Stebbins G. L. and Shah S. S. (1960). Developmental studies of cell differentiation in the epidermis of Monocotyledons. II. Cytological figures of stomatal development in the Gramineae. *Developm. Biol.* 2: 477-500.
- Sulthoni A. (1989). *Bamboo: physical properties, testing methods and means of preservation*. pp. 1-15 in Bassiii A.V. and Davies W.G. (Eds.) Proceedings of a Workshop on Design and Manufacture of Bamboo and Rattan Furniture. March 3-14.
- Swift J. G. and O'brien T. P. (1972). The fine structure of the wheat scutellum before germination. *Australian Journal of Biological Sciences*. 25(1): 9-22.
- Szabo Z. K., Papp M. and Daroczi L. (2006). Ligule anatomy and morphology of five *Poa* species. *Acta Biologica Cracoviensia Series Botanica*. 48(2): 83-88.
- Taiz L. and Zeiger E. (2002). Photosynthesis: physiological and ecological considerations. *Plant physiology*. Massachusetts: Sinauer Associates, 171-192.

- Tateoka T. (1956). The place of the genus *Phyllorachis* in the system of gramineae. *Bot. Mag.* 69: 83-86.
- Tateoka T. (1957). Miscellaneous papers on the phylogeny of Poaceae, proposition of a new phylogenetic system of Poaceae. *Jour. Jap. Bot.* 32: 275-287.
- Thornton M. L. (1966). Seed dormancy in tall wheat grass (*Agropyron longatum*). In *Proceedings of the Association of Official Seed Analysts.* 56: 116-119.
- Tillich H. J. (2007). Seedling diversity and the homologies of seedling organs in the order Poales (Monocotyledons). *Annals of Botany.* 100(7): 1413-1429.
- Vartapetian B. B. and Jackson M. B. (1997). Plant adaptations to anaerobic stress. *Annals of Botany,* 79: 3-20.
- Vecchia F. D., Asmar T. E., Calamassi R., Rascio N. and Vazzana C. (1998) Morphological and ultrastructural aspects of dehydration and rehydration in leaves of *Sporobolus stapfianus*. *Pl. Growth Reg.* 24: 219-228.
- Vickery J. W. (1935). Leaf anatomy and vegetative characters of the indigenous grasses of New South Wales. *Proc. Linn. Soc. N.S.W.* 60: 340-373.
- Wang S. G., Pu X. L., Ding Y. L., Wan X. C. and Lin S. Y. (2011). Anatomical and chemical properties of *Fargesia yunnanensis*. *Journal of Tropical Forest Science.* 23(1): 73-81.
- Weatherwax P. (1930). The endosperm of *Zea* and *Coix*. *American Journal of Botany.* 17: 371-380.
- Wolf M. J., Buzan C. L., Mac Masters M. M. and Rist C. E. (1952). Structure of the mature corn kernel. Gross anatomy and structural relationships. *Cereal Chemistry.* 29(5): 321-333.
- Yakovlev M. S. (1950). Structure of endosperm and embryo in cereals as a systematic feature. *Inst. Akad. Sci. S.S.S.R.* 1: 121-218.
- Yang C., Zhang X., Zhou C. and Seago J. L. (2011). Root and stem anatomy and histochemistry of four grasses from the *Jiangnan* Floodplain along the Yangtze River, China. *Flora-Morphology, Distribution, Functional Ecology of Plants.* 206(7): 653-661.
- Yao X., Yi T., Ma N., Wang Y. and Li Y. (2002). *Bamboo culm anatomy of China* (Vol. 187). Science Press: Beijing.
- Zeleznak K. and Varriano-Marston E. (1982). Pearl millet (*Pennisetum americanum* (L.) Leeke) and grain sorghum (*Sorghum bicolor* (L.) Moench) ultrastructure. *American Journal of Botany.* 69(8): 1306-1313.
- Zuloaga F., Morrone O. and Dubcovsky J. (1989). Exomorphological, anatomical, and cytological studies in *Panicum validum* (Poaceae: Panicoideae: Paniceae): its systematic position within the genus. *Systematic Botany.* 14: 220-230.