REFERENCES

1. Aakif, A. and Khan, M. F. (2015). Automatic classification of plants based on their leaves. *Biosystems Engineering*, 139, 66-75
2. Adie, P. A., Torsabo, S. T., Uno, U. A. and Ajegi, J. (2014). *Funaria hygrometrica* moss as bioindicator of atmospheric pollution of heavy metals in Makurdi and environs, North Central Nigeria. *Research Journal of Chemical Sciences*, 4(10), 10-17
3. Agrawal, C. G. (1986). Structure and tectonics of exposed rocks between Narmada and Kim rivers in South Gujarat. *Journal of Geological Society of India*, 27, 531- 542
4. Ahmad, S. (1942). Three new species of *Riccia* from India. *Current Science*, 11: 433–434
5. Akiyama, H. (1993). A submerged rheophytic moss, *Fissidens dalamair* sp. nov. (Fissidentaceae) from West Kalimantan, Indonesia. *Acta phytotaxonomica et geobotanica*, 44(1), 21-26
6. Alam, A. (2011). Diversity and altitudinal distribution of terrestrial liverworts (Hepaticae) in Nilgiri hills, Tamil Nadu, India. *Proceedings National Academy of Sciences, India, Sect., B*, 81, 206-217
7. Alam, A. (2013). Moss flora of Western Himalayas, India - An updated Checklist. *Archive for bryology*,168,1-28
8. Alam, A. (2015). *Moss flora of India: An updated checklist*. Grain Verlag Publication, Germany. 196 p
9. Alam, A., Rawat, K. K., Verma, P. K., Sharma, V. and Gupta, D. S. (2015). Moss flora of central India. *Plant science today*, 2(4), 159-171
10. Alam, A., Pandey, S., Singh, V., Sharma, S. C. and V. Sharma. (2014). Moss flora of Mount Abu (Rajasthan), India: An updated checklist. *Tropical Plant Research*, 1(1), 8-13
11. Alam, A., Sharma, V. and Sharma, S. C. (2011). *Bryoflora of Ranthambhore Tiger Reserve*, *Rajasthan (India). Archive for Bryology*, 106, 1-8
12. Alexander, D. C., Levett, P. N. and Turenne, C. Y. (2015). Molecular taxonomy. In: *Molecular Medical Microbiology*, Academic Press. 369-379 pp

13. Amandita, F. Y., Rembold, K., Vornam, B., Rahayu, S., Siregar, I. Z., Kreft, H. and Finkeldey, R. (2019). DNA barcoding of flowering plants in Sumatra, Indonesia. *Ecology and Evolution*, 9(4), 1858-1868
14. Anon. (2020) *Lucid.* http://www.lucidcentral.org. Accessed on 17th March 2020.
15. Anon. (2020). *1225 Weeds of the 48 states& Adjacent Canada*. http://www. weed-id.com/.Accessed on 12th Feb 2020
16. Anon. (2020). *ActKey*. http://ucjeps.berkeley.edu/mea cham/meka/index03.html. Accessed on 18th Feb 2020
17. Anon. (2020). *Navikey*. http://www.huh.harvard.edu/databases/legacy /navikey Accessed on 17th March 2020
18. Asakawa, Y. (1990). Biologically Active Substances from Bryophytes. In: *Bryophyte Development: Physiology and Biochemistry*, (Eds) Chopra, R.N and Bhatla, S.C., Boca Raton, CRC Press. 259 p
19. Asakawa, Y. (1995). Chemical constituents of the bryophytes. In *Progress in the chemistry of organic natural products*, Springer, Vienna. 618 p
20. Asakawa, Y. (1999). Phytochemistry of bryophytes. In Romeo, J. T. (Eds) *Phytochemicals in human health protection, nutrition, and plant defense*. Springer, Boston, MA. 319-342 pp
21. Asrani, K. and Jain, R. (2013). Contour based retrieval for plant species. *International Journal of Image, Graphics and Signal Processing*, 5(9), 29-35
22. Asthana, A. K. and Gupta, R. (2015). Hornwort diversity at Pachmarhi Biosphere Reserve (Madhya Pradesh), India. *Plant Science Today*, 2(4),145-150
23. Asthana, A. K., Nath V. and Sahu, V. (2005). *Phaeoceros kashyapii* Asthana et Sriv-new to Eastern Himalayas. *Current Science*, 88(6), 864-865
24. Asthana, A. K., Omar, I. and Srivastava, P. (2015). *Lejeunea minutiloba* A. Evans, new to bryoflora of South India with a note on its interesting morphoform. *Indian Journal of Forestry*, 38(3), 245-248
25. Asthana, A. K. and Omar, I. (2016). *Archidium acanthophyllum* Snider, a new recordto moss flora of India. *Phytotaxonomy*, 16(1), 38-41

26. Asthana, A. K and Sahu, V. (2010). Growth response of moss *Brachymenium capitulatum* (Mitt.) Par. in different culture media. *National Academy Science letters*, 34(1), 23-25
27. Asthana, A. K. and Sahu, V. (2013). Bryophyte diversity in Mukteshwar (Uttarakhand): an overview. *Archive for bryology*, 154, 1-11
28. Asthana, A. K., Sahu, V. and Srivastava, A. (2015). Invitro propagation of three species of *Bryum* Hedw.:A comparative study. *Geophytology*, 45(2), 215-220
29. Asthana, A. K., Sahu, V. and Gupta, D. (2019). A new species of genus Cololejeunea (Spruce) Schiffn. (Lejeuneaceae) from India. *Journal of Bryology*, 41(3), 274-277
30. Asthana, A. K., Singh T., V. J., Pathak I. and Sahu, V. (2018). In vitro conservation strategy for endemic and endangered Himalayan liverwort *Stephensoniella brevipedunculata* Kashyap (Marchantiophyta). *International Journal of plant and Environment*, 4(2), 81-87
31. Asthana A. K. and Srivastava S. C. (1991). *Indian Hornworts – A taxonomic study*. Bryophytorum Bibliotheca , Berlin, Stuttgart. 158 p.
32. Attigala, L., De Silva, N. I. and Clark, L. G. (2016). Simple Web-based interactive key development software (WEBiKEY) and an example key for Kuruna (Poaceae: Bambusoideae). *Applications in plant sciences*, 4(4), 1-7
33. Awasthi, V., Asthana, A. K. and Nath, V. (2012a). In vitro propagation of an epiphytic pleurocarpous moss *Erythrodontium julaceum* (Schwaegr.) Par. *Journal of Bryology*, 34(2), 140-144
34. Awasthi, V., Nath, V. and Asthana, A. K. (2010). Effect of some physical factors on reproductive behaviour of selected bryophytes. *The International Journal of Plant Reproductive Biology*, 2(2), 141-145
35. Awasthi, V., Nath, V., Pande, N. and Asthana, A. K. (2012b). Morphogenetic Studies and" In vitro" Propagation of two Mosses: *Philonotis thwaitesii* Mitt. and *Brachythecium plumosum* (Hedw.) BSG. *Taiwania*, 57(1), 27-36
36. Ayob, M. Z. andKadir, M. K. A. (2016). Case report Computer Aided Taxonomy: A case study on the automated identification of invasive ladybirds in the UK. *Asian Journal of Scientific Research*, 9, 266-272

37. Aziz, M. N. and Vohra, J. N. (2008). *Pottiaceae (Musci) of India*. Bishen Singh & Mahendra Pal Singh, Dehra Dun, India. 366p
38. Backes, A. R. and Bruno, O.M. (2009). *Plant leaf identification using multi-scale fractal dimension*. In Foggia P., Sansone C., and Vento, M.(Eds)Image analysis and processing,Springer, Berlin, Heidelberg. 143-150 p
39. Backes, A. R., Casanova, D. and Bruno, O. M. (2009). Plant leaf identification based on volumetric fractal dimension. *International Journal of Pattern Recognition and Artificial Intelligence*, 23(6), 1145-1160
40. Baena-Bejarano, N. and Heads, S. W. (2015). Three new species of the genus *Ripipteryx* from Colombia (Orthoptera, Ripipterygidae). *ZooKeys,* 502, 129-143
41. Bahuguna, Y. M., Gairola, S., Semwal, D. P., Uniyal, P. L. and Bhatt, A. B. (2013). *Bryophytes and ecosystem*. InGupta R. K. and Kumar M. (Eds), *Biodiversity of Lower Plants*, IK International publishing house Pvt. Ltd. 279-296 pp
42. Bansal, P. and Nath, V. (2012). A new record of *Bryum coronatum* Schwaegr.(Bryophyte) in Meghalaya, India. *Taiwania,* 57(3), 294-299
43. Bansal, P. and Nath, V. (2014). Present status of genus *Bryum* Hedw., (Bryophyta) in Rajasthan, India. G-*Journal of Environmental Science and Technology*, 1(6), 118-122
44. Bansal, P., Nath, V. and Pande, N. (2012). *Bryum dunense* Smith et Whitehouse (Bryophyta: Bryaceae): New to Central India. *Proceeding of National Academy of SciencesIndia Sect B Biological Sciences*, 82(3), 453- 454
45. *Bapna, K. R. (1975). On *Funaria (Enthostodon) nutans* (Mitt.) Broth. from Rajasthan and its distribution. *Journal ofHattori Botanical Laboratory*, 54, 234–237
46. Bapna, K. R. (1980). Occurrence of *Fissidens* Hedw. in Rajasthan*. Journal of Indian Botanical Society*, 59, 20-26
47. Bapna, K. R. and Choudhary, B. L. (1989). *Mosses of Rajasthan-II.* Dicranales and Pottiales. *Journal of Indian Botanical Society*, 68, 379-388
48. Bapna, K. R. and Kachroo, P. (2000). *Hepaticology in India I and II*. Himanshu publications, Udepur. 446 p
49. Basile, A., Sorbo, S., Bassi, P., Napolitano, E., Cogoni, A. E. and Castaldo Cobianchi, R. (2008). Effects of heavy metals on protonemal development and ultrastructure in

populations of the moss *Funaria hygrometrica* Hedw.(Bryophyta) from a mine and an unpolluted site. *Fresenius Environmental Bulletin*, 17, 1956-1963

50. *Bauer, L. and Mohr, H. (1959). Der Nachweis des reversiblen Hellrot-Dunkelrot-Reaktionssystems bei Laubmoosen. *Planta*, 54(1), 68-73
51. Bell, D., Long, D. G., Forrest, A. D., Hollingsworth, M. L., Blom, H. H. and Hollingsworth, P. M. (2012). DNA barcoding of European *Herbertus* (Marchantiopsida, Herbertaceae) and the discovery and description of a new species. *Molecular Ecology Resources*, 12(1), 36-47
52. *Beneke, W. (1903). *Uber die Keimung der Brutknospen von Lunularia cruciata*. Botanische Zeitung, 61, 19-46
53. Benny, M. E. and Mathew, A. (2020). *Taxonomic studies on leafy liverworts and mosses from Ernakulam district, Kerala, India*. In AIP Conference Proceedings, 2263 (1):030013
54. Berg, A., Gardenfors, U., Hallingback, T., and Noren, M. (2002). Habitat preferences of red-listed fungi and bryophytes in woodland key habitats in southern Sweden-analyses of data from a national survey. *Biodiversity & Conservation*, 11(8), 1479-1503
55. Bergamini, A. and Peintinger, M. (2002).Effects of light and nitrogen on morphological plasticity of the moss *Calliergonella cuspidata*. *Oikos*, 96: 255–363
56. Beutelmann, P. and Bauer, L. (1977). Purification and identification of a cytokinin from moss callus cells. *Planta*, 133(3), 215-217
57. Bhadauriya, G., Rathore, K. S. and Singh, S. (2018). Phytochemical screening and total phenolic content in the extract of bryophyte *Plagiochasma appendiculatum* and *Dicranum scoparium. Environment Conservation Journal*, 19(1&2), 175-181
58. Bharadwaj, D. C. (1950). Studies in Indian Anthocerataceae I. The morphology of *Anthoceros crispulus* (Montin.) Douin. *Journal of Indian Botanical Society*, 29: 145-163.
59. Bharadwaj, D. C. (1960). Studies in Indian Anthocerotaceae II.The morphology of *Anthoceros* cf *gemmulosus* (Hattori) Pande. *Journal of Indian Botanical Society*, 37, 75-92
60. Bhatla, S. C. and Chopra, R. N. (1981). Hormonal regulation of gametangial formation in the moss *Bryum argenteum* Hedw. *Journal of Experimental Botany*, 32(6), 1243-1256

61. Bhattacharyya, T., Pal, D. K., Mandal, C., Chandran, P., Ray, S. K., Sarkar, D., Velmourougane, K., Srivastava, A., Sidhu, G. S., Singh, R. S, Sahoo, A. K., Dutta, D., Nair, K., Srivastava, R., Tiwari, P., Nagar, A. P. and Nimkhedkar, S. S. (2013). *Soils of India: historical perspective, classification and recent advances*.104 (10), 1308-1323
62. Bhattarai, K. R. and Vetaas, O. R. (2003). Variation in plant species richness of different lifeforms along a subtropical elevation gradient in the Himalayas, east Nepal. *Global Ecology and Biogeography*, 12, 327–340
63. Bhattarai, K. R. and Vetaas, O. R. (2006). Can Rapoport's rule explain tree species richness along the Himalayan elevation gradient, Nepal? *Diversity and Distributions*, 12, 373–378
64. Bhattarai, K. R., Vetaas, O. R. and Grytnes, J. A. (2004). Fern species richness along a Central Himalayan elevational gradient, Nepal. *Journal of Biogeography*, 31, 389–400
65. Bijelovic, A., Sabovljevic M., Grubisic D. and Konjevic, R. (2004). Phytohormone influence on the morphogenesis of two mosses (*Bryum argenteum* Hedw. and *Atrichum undulatum* (Hedw.) P. Beuav.). *Israel Journal of Plant Sciences*, 52(1), 31-36
66. Bittrich, V.; Souza, C. S. D.; Coelho, R. L.; Martins, M.V.; Hopkins, M. J. andAmaral, M.C. (2012). An interactive key (Lucid) for the identifying of the genera of seed plants from the Ducke Reserve, Manaus, AM, Brazil. *Rodriguesia*, 63(1), 55-64
67. Blocher, R. and Frahm, J. P. (2002). A comparison of the moss floras of Chile and New Zealand. Studies in austral temperate rain forest bryophytes. *Tropical Bryology*, 21, 81–92
68. Bopp, M. (1983). Developmental physiology of bryophytes. In: Schuster R. M., (Eds). *New manual of bryology*, Volume I. The Hattori Botanical Laboratory, Nichinan, Japan.276-324 pp
69. Bopp, M. and Knop, B (1984) Culture methods for bryophytes. In Vasil, I. (Eds),*Cell Culture and Somatic Cell Genetics of Plants - Laboratory Procedures and their Applications* Vol – 1, Academic Press, 96-105 pp
70. Botanical Survey of India (BSI). (2019) *Plant Discoveries 2018-New Genera, species and New Records.* Mao A.A. and Dash S.S. (Eds), Botanical Survey of India, Kolkatta, 183 p

71. Boughey, A. S., Bridges, K. and Wand Ikeda, A. G. (1968). *An automated biological identification key. Research series no.2*. Museum of Systematic Biology, University of California, Irvine. 36 p

72. Bramley-Alves, J., King, D. H., Robinson, S. A., and Miller, R. E. (2014). Dominating the Antarctic environment: bryophytes in a time of change. *Advances in Photosynthesis and Respiration*, 37, 309–324

73. *Braun, A. (1864). Uebersicht des naturlichen systems nach der Anordnung desselben. In Ascherson, P. *Fl. Brandenburg*, 1(2), 22-67 pp

74. BSI ENVIS. (2021). *Status of plant diversity in india: An overview*. http://www.bsienvis.nic.in/Database/Status_of_Plant_Diversity_in_India_17566.aspx. Accessed on 22ndJanuary 2021

75. Buck, W. R. and Goffinet, B. (2000). *Morphology and classification of mosses*. Bryophyte Biology (Eds.) A. J. Shaw and B. Goffinet. Cambridge University Press, University of Cambridge, UK .71-123 pp

76. Buckley, P. M. and Reed, B. M. (1994). Antibiotic susceptibility of plant associated bacteria. *HorticulturalScience*, 29, 434-434

77. Buryova, B, and Shaw, A. J. (2005) Phenotypic plasticity in Philonotis fontana (Bryopsida: Bartramiaceae). *Journal of Bryology*, 27, 13–22

78. Cano, M, J., Ros, R. M. and Guerra, J. (1996). In vitro culture of *Pterygoneurum compactum* (Musci: Pottiaceae): Control of Taxonomical Characters. *Cryptogamie Bryologie Lichenologie*, 17(1), 67-70

79. Cano, M. J., Werner, O. and Guerra, J. (2005). A morphometric and molecular study in *Tortula subulata* complex (Pottiaceae, Bryophyta). *Botanical Journal of the Linnean Society*, 149(3), 333-350

80. CBOL Plant Working Group (2009). A DNA barcode for land plants. *Proceedings of National Academy of Sciences*, 106, 2794–2797

81. Cerretti, P., Tschorsnig, H. P., Lopresti, M. andDi Giovanni, F. (2012). MOSCHweb—a matrix-based interactive key to the genera of the Palaearctic Tachinidae (Insecta, Diptera). *ZooKeys*, 205, 5-18

82. Chase, M. W., Salamin, N., Wilkinson, M., Dunwell, J. M., Kesanakurthi, R.P., Haidar, N., Savolainen, V. (2005). Land plants and DNA barcodes: short-term and long-term

goals. *Philosophical Transactions of the Royal Society B-Biological Sciences* ,360, 1889–1895

83. Chaudhary, B. L. and Deora, G. S. (1993). *Moss Flora of Rajasthan (India)*, Himanshu Publications, Udepur.127 p
84. Chaudhary, B.L. and Sharma, T.P. (2002) Occurrence of epiphytic *Stereophyllum* Mitt. on Girnar hill, Gujarat (India). P*hytomorphology*,46: 273–277
85. Chaudhary, B. L., Sharma, T. P. and Bhagora, F. S. (2008). B*ryophyte Flora of North Konkan Maharashtra, India*. Himanshu Publication, Udepur and New Delhi, 342p
86. Chaudhary, B. L., Sharma, T. P. and Charu, S. (2006). *Bryophyte Flora of Gujarat (INDIA),* Himanshu Publicatons, Udepur, 198 p
87. Chavan, A. R. (1937a).A morphological study of *Cyathodium barodae.American Journal of Botany*, 24(8), 484 – 492
88. Chavan, A. R. (1937b). A new species of *Cyathodium* from India. *The Bryologist*, 15(3), 57-60
89. Chavan, A. R.and Mahabale, T. S. (1945). Distribution of Liverworts in Gujarat. *Proc. 32nd Indian Science Congress*, Nagpur, 70
90. Chavan, A. R., Mehta, S. and Mehta, A. R. (1961). A preliminary report of mosses from Pavagadh in Gujarat. *Journal of M. S. University of Baroda*, 10 (3), 35-42
91. Chen, Y. Y., Lou, Y. X., Guo, S. L. and Cao, T. (2009). Successful tissue culture of medicinal moss *Rhodobryum giaganteum* and factors influencing proliferation of its protonemata. *Annales Botanici Fennici*, 46, 516-524
92. Cho, S. Y. (2012). Content-based structural recognition for flower image classification. *In: 2012 7th IEEE conference on industrial electronics and applications* (ICIEA), 541–546 pp
93. Chopra, R. S. (1943). A census of Indian Hepatics. *Journal of Botanical society*, 22, 237-259
94. Chopra, R. S. (1975). *Taxonomy of Indian Mosses. Botanical Monograph, No.10*, Council of Scientific and Industrial Research (CSIR), New Delhi. 631 pp

95. Chopra, R.N. and Kumra, S. (1984). Regulation of growth and gametangial formation in *Riccia gangetica* Ahmad by some amino acids. *Journal of experimental botany*, 35(10), 1537-1546

96. *Chopra, R. N and Rawat, M. S. (1977). Studies on the intitation of sexual phase in the moss *Leptobryum pyriforme*. *Beitrage zur Biologie der Pflanzen*., 53, 353-357

97. Chopra, R. N and Sood, S. (1973). In vitro studies in Marchantiales. I. Effects of some carbohydrates, agar, pH, light and growth regulators on the growth and sexuality in *Riccia crystallina*. *Phytomorphology*, 23, 230-44

98. Chopra. R. S. (1938). Notes on Indian Hepatics I. South India. *Proceedings of Indian Academic of ScienceSection B.* 8(6), 239-251

99. Christianson, M. L. (2000). *Control of morphogenesis in bryophytes*. In Shaw A. J.and Goffinet, B. (Eds.), *Bryophyte Biology* Cambridge University Press, University of Cambridge, UK. 199-224 pp

100. Costa, D. P. (1999). Epiphytic bryophyte diversity in primary and secondary lowland rainforests in southeastern Brazil. *The Bryologist*, 102(2), 320-326

101. Costa, D. P., Nadal, F. and da Rocha, T. C. (2020). The first botanical explorations of bryophyte diversity in the Brazilian Amazon mountains: high species diversity, low endemism, and low similarity. *Biodiversity and Conservation*, 29, 2663-2688

102. Costa, J. L., Paulsrud, P., Rikkinen, J. and Lindblad, P. (2001). Genetic diversity of *Nostoc* symbionts endophytically associated with two bryophyte species. *AppliedEnvironmental Microbiology*, 67, 4393–4396

103. Cove, D. J., Knight, C. D. and Lamparter, T. (1997). Mosses as model systems. *Trends in Plant Science*, 2(3), 99-105

104. Cove, D. J., Perroud, P. F., Charron, A. J., McDaniel, S. F., Khandelwal, A. and Quatrano, R. S. (2009). *Culturing the moss Physcomitrella patens*. Cold Spring Harbor Protocols, 4(2), 1-6

105. Cox, C. J., Goffinet , B., Newton,A. E., Shaw, A .J. and Hedderson, T. A. (2000). Phylogenetic relationships among the diplolepideous-alternate mosses (Bryidae) inferred from nuclear and chloroplast DNA sequences. *Bryologist*, 103(2), 224-241

106. Crandall-Stotler B., Stotler, R. E. and Long, D. A. (2008). Morphology and Classification of the Marchantiophtyta, In: *Bryophyte bryology*. 2nd Ed., Goffinet B. and Shaw A. J (Eds), University Press, Cambridge, 1-54 pp
107. Cruz, T. E. E., Pangilinan, M. V. B. and Litao, R. A. (2012). Printed identification key or web-based identification guide: an effective tool for species identification?. *Journal of Microbiology & Biology Education*,13(2), 180-182
108. Cuenoud, P., Savolainen, V., Chatrou, L.W., Powell, M., Grayer, R.J. and Chase, M.W. (2002). Molecular phylogenetics of Caryophyllales based on nuclear 18S rDNA and plastid rbcL, atpB, and matK DNA sequences. *American Journal of Botany*, 89(1), 132-144
109. Dabhade, G. T. (1969). Investigation of the genus *Bryum* (Hedwig) Schimper from Western India. *Maharashtra Vig Mandir Patrika*, 4, 13-21
110. Dabhade, G .T. (1970). New records of Funaria (Entosthodon) nutans (Mitt.) Broth. from Western India.*Bulletin of the Botanical Society of Bengal*, 24, 83-85
111. Dallwitz, M. J. (1993). DELTA and Intkey. In Fortner,R. (Ed.),*Advances in computer methods for systematic biology: artificial intelligence, databases, computer vision*. The Johns Hopkins University Press, Baltimore, Maryland. 287-296 pp
112. Dallwitz, M. J. (1997) onwards. Desirable attributes for interactive identification programs. delta-intkey.com. Accessed on 10th April 2020
113. Dallwitz M.J. (2018). *Overview of Delta system*. Southern IPM Centre. http://delta-intkey.com/www/overview.htm. Accessed on 13th March 2020
114. Dallwitz, M. J., Paine, T. A. and Zurcher, E. J. (1993).Conditions of use for the Delta System. https://www.delta-intkey.com/www/use.htm. Accessed on March 2020
115. Dallwitz, M.J; Paine, T.A, and Zurcher E.J. (2011). Principles of interactive keys. http://delta-intkey.com. Accessed on 21stMarch 2020
116. Dallwitz M. J, Paine T. A, and Zurcher E. J. (2016). Conditions of use for the Delta System. https://www.delta-intkey.com/www/use.pdf. Accessed on 15th March 2020
117. Dandotiya, D., Govindapyari, H., Suman, S. andUniyal, P. L. (2011). Checklist of the bryophytes of India. *Archive forBryology*, 88(1), 1-126
118. Daniels, A.E.D. (2010). Checklist of the bryophytes of Tamil Nadu, India. *Archieve for bryology*, 60, 1-117

119. Daniels, A. E. D. and Kariyappa, K. C. (2007). Bryophyte diversity along a gradient of human disturbance in the southern Western Ghats. *Current science*, 93(7), 976-982
120. Davey, M. L. and Currah R. S. (2006). Interactions between mosses (Bryophyta) and fungi. *Canadian Journal of Botany*, 84, 1509–1519
121. De Mattia, F., Bruni, I., Galimberti, A., Cattaneo, F., Casiraghi, M. and Labra, M. (2011). A comparative study of different DNA barcoding markers for the identification of some members of Lamiacaea. *Food Research International*, 44(3), 693-702
122. Decker, E. L., Frank, W., Sarnighausen, E. and Reski, R. (2006). Moss systems biology en route: Phytohormones in *Physcomitrella* development. *Plant Biology*, 8, 397-406
123. *Dellinger-Johnston, R. A. (2015). *A New Method for Creating a Visual Plant Identification Key*. Ph.D thesis submitted to University of North Carolina at Greensboro. 88 p
124. Deora, G. S. and Chaudhary, B. L. (1996). Occurrence of *Bryum* Hedw. in Rajasthan. *Phytomorphology*, 46, 299- 304
125. Deora, G. S. and Suhalka, D. (2016). Phytochemical composition and fungicidal potential of moss *Philonotis revoluta* against spore germination process of fungus *Helminthosporium turcicum*. *Journal of Pharmacy and Biological Sciences*, 11(6), 38-43
126. Dey M. and Singh, D. K. (2012). *Epiphyllous liverworts of eastern Himalaya*. Botanical Survey of India. 422 p
127. *Dillenius, J. J. (1741). *Historia Muscorum*. Oxford. 218 pp
128. Directorate of Census Operations (DCO). (2011a).*District CensusHandbook-Dahod. Directorate of Census Operations*, Gandhinagar, Gujarat, 460 p
129. Directorate of Census Operations (DCO). (2011b).*District CensusHandbook-Narmada. Directorate of Census Operation*s, Gandhinagar, Gujarat, 410p
130. Directorate of Census Operations (DCO). (2011c).*District CensusHandbook-Panchmahal. Directorate of Census Operations*, Gandhinagar, Gujarat,712p
131. Directorate of Economics and Statistics (DES). (2019).*Statistical Abstract of Gujarat State-2019.Directorate of Economics and Statistics*, Gandhinagar, 231p
132. Dixon, H. N. (1914). Report on the mosses collected by C.F.C Fischer and others from South India and Ceylon. *Records of Botanical Survey of India*, 6, 75-89

133. Dixon, H. N. (1921). On collection of mosses from the Kanara district. *Journal of Indian Botanical society,* 2, 174-188
134. Dixon, H. N. (1937). Mosses collected in Assam. *Journal of Bombay Natural History Society*, 39, 769-795
135. Du, J. X., Huang, D. S., Wang, X. F. and Gu, X. (2006). Computer-aided plant species identification (CAPSI) based on leaf shape matching technique. *Transactions of the Institute of Measurement and Control*, 28(3), 275-285
136. Duckett, J. G., Burch, J., Fletcher, P. W., Matcham, H. W., Read, D. J., Russell, A. J. and Pressel, S. (2004). In vitro cultivation of bryophytes: a review of practicalities, problems, progress and promise. *Journal of bryology*, 26(1), 3-20
137. Duckett, J. G., Goode, J. A. and Matcham, H. W. (2001). Studies of protonemal morphogenesis in mosses. VIII. The gemmiferous protonemata of *Orthodontium* and *Dicranoweisia. Journal of Bryology*, 23, 181–194
138. Duff, R. J., Villarreal, J. C., Cargill, D. C. and Renzaglia, K. S. (2007). Progress and challenges toward developing a phylogeny and classification of the hornworts. *Bryologist,* 110, 214–43
139. Edwards, M. and Morse, D.R. (1995). The potential for computer-aided identification in biodiversity research. *Trends in ecology & evolution*, 10(4), 153-158
140. *Eichler A. W. (1883). Sylalbus der Vorlesungenuber Phanerogamenkunde, (Eds) Borntrager 3rd , Berlin, 36 p
141. Erdag,A. andKurschner, H. (2017). A reference list of Turkish bryophytes. The state of knowledge from 1829 until 2017. *Anatolian Bryology*, 3(2), 81-102
142. Fataniya, B. M. (2019). *Microscopic Image Based Classification Algorithms for few Herbal Plants Identification using Machine Learning Approaches*. Ph.D thesis submitted to Nirma University. 144 p
143. Fazekas, A. J., Kesanakurti, P. R., Burgess, K. S., Percy, D. M., Graham, S. W., Barrett, S. C. H, Newmaster, S. G., Hajibabaei, M. and Husband, B. C. (2009). Are plant species inherently harder to discriminate than animal species using DNA barcoding markers? *Molecular Ecology Resources*, 9(1), 130–139
144. Feldberg, K., Hentschel, J., Wilson, R., Rycroft, D. S., Glenny, D. and Heinrichs, J. (2007). Phylogenetic biogeography of the leafy liverwort *Herbertus* (Jungermanniales,

Herbertaceae) based on nuclear and chloroplast DNA sequence data: correlation between genetic variation and geographical distribution. *Journal of biogeography*, 34(4), 688-698

145. Fensham, R. J. and Streimann, H. (1997). Broad landscape relations of the moss flora from inland dry rainforest in north Queensland, Australia. *The Bryologist*, 100, 56–64
146. Fischer, J. (2013). *Fish identification tools for biodiversity and fisheries assessments: review and guidance for decision-makers*. FAO Fisheries and Aquaculture Department. Rome, Italy, 118 p
147. Foreau, G. (1961). The moss Flora of the Palni hills. *Journal Bombay Natural History Society*, 58, 13-47
148. Forest Survey of India (FSI). (2019a) *Indian state of forest Report 2019. Volume I*.Forest Survey of India, Ministry of Environment, Forest and Climate change, Dehradun, Uttarakhand, India, 185 p
149. Forest Survey of India (FSI). (2019b). *Indian state of forest Report 2019. VolumeII*: Gujarat.Forest Survey of India, Ministry of Environment, Forest and Climate change, Dehradun, Uttarakhand, India, 71-81p
150. Forrest, L. L., Schuette, S. W., Crandall-Stotler, B.J. and Stotler, R. E. (2005). A molecular study of the simple thalloid liverwort *Jensenia* (Marchantiophyta, Pallaviciniaceae). *The Bryologist,* 108(2), 204-212
151. Fox, G. E., Stackebrandt, E., Hespell, R. B., Gibson, J., Maniloff, J., Dyer, T. A., Wolfe, R. S., Balch, W. E., Tanner, R. S., Magrum, L. J. and Zablen, L. B. (1980). The phylogeny of prokaryotes. *Science*, 209(4455), 457-463
152. French, J. C. and Paollillo Jr, D. J. (1976). Effect of light and other factors on capsule expansion in *Funaria hygrometrica*. *The Bryologist*, 79(4), 457-465
153. Gabriel, R. and Bates, J. W. (2005) Bryophyte community composition and habitat specificity in the natural forests of Terceira, Azores. *Plant Ecology*, 177 (1), 125-144.
154. *Gabriel, R. M. and De Almeida. (2000). *Ecophysiology of Azorean Forest Bryophytes*. Ph.D. thesis submitted to Department of Biology, Imperial College of Science, Technology and Medicine, Silwood Park, England
155. Gangulee, H. C. (1959). *Mosses of Eastern India II. Eubryiidae. Series II*. Dicranales. Family: Ditrichaceae. *Bulletin ofBotanicalSociety ofBengal*, 13, 1-9

156. Gangulee, H. C. (1960). *Mosses of Eastern India II. Eubryiidae. SeriesII*. Dicranales. Family: Dicranaceae. *Bulletin of BotanicalSociety ofBengal*, 14, 10-57

157. Gangulee, H. C. (1961). Mosses of Eastern India II. Eubryiidae. SeriesII. Dicranales. Family: Leucobryaceae and Series III Pottiales: Family Calymperaeceae. *Journal Bombay Natural History Society*, 60, 606-637

158. Gangulee, H. C. (1969-1980). *Mosses of Eastern India & Adjacent regions*, Fascicles 1-8. Calcutta. 1- 2145 p

159. Gangulee, H. C. (1969-72). *Mosses of Eastern India and Adjacent Regions. Vol. I*. Books and Allied (P) Ltd., Kolkata, India. 830 pp.

160. Gangulee, H. C. (1974-78). *Mosses of Eastern India and Adjacent Regions. Vol. II.* Books and Allied (P) Ltd., Kolkata, India. 1546 p

161. Gangulee, H. C. (1978-80). *Mosses of Eastern India and Adjacent Regions. Vol. III.* Books and Allied (P) Ltd., Kolkata, India. 2145 p

162. Gibson, S. I. (2004). Sugar and phytohormone response pathways: navigating a signalling network. *Journal of Experimental Botany*, 55, 253-264

163. Glime, J.M. (2017), Chapter 1 - Field Taxonomy and Collection Methods, *Bryophyte Ecology Volume 3: Methods*. 3.https://digitalcommons.mtu.edu/bryophyte-ecology3/3. Accessed on 14th August 2013

164. Glime, J. M. and Knop, B. C. (1986). Spore germination and protonemal development in *Fontinalis squamosa*. *Journal of the Hattori Botanical Laboratory*, 61, 487–497

165. Glime, J. M. and Wagner, D. H. (2013). Laboratory Techniques: Equipment. Chapt. 2-1. In: Glime, J. M. Bryophyte Ecology. Volume 2-1-1 3. Methods. Ebook sponsored by Michigan Technological University and the International Association of Bryologists. Ebook last updated 20 July 2020 and available at https://digitalcommons.mtu.edu/bryophyte-ecology3/3. Accessed on 19th September 2013

166. Glime, J.M., 2019. Chapter 4 Aquaria in *Bryophyte ecology, vol 5. Uses.* https://digitalcommons.mtu.edu/bryophyte-ecology1/..Accessed on 5 May 2020

167. Goffinet, B. and Shaw, A. J. (2008). *Bryophyte biology*. 2nd Ed. Cambridge University Press. Cambridge 565 p

168. Goffinet, B., Buck, W. R., and Shaw, A. J. (2008). Morphology, anatomy and classification of the Bryophyta. In *Bryophyte Biology*, 2nd Ed. (Eds.) Goffinet, B and Shaw, A. J. Cambridge University Press: England. 55–138 pp
169. Gonzalez, M. L., Mallon, R., Reinoso, J. and Rodriguez-Oubina, J. (2006). In vitro micropropagation and long-term conservation of the endangered moss *Splachnum ampullaceum*. *Biologia plantarum*, 50(3), 339-345
170. Goode, J. A., Duckett, J. G. and Stead, A. J. (1993). Protonemal morphogenesis in *Sphagnum*: the cytoskeleton and effects of growth regulators. *Advances in Bryology*, 5, 129–151
171. Google Earth pro. (2019). Elevation Information.https://www.google.com/intl/en_in/earth/versions/. Accessed on 22 December 2019.
172. Gradstein S. R. and Culmsee, H. (2010). Bryophyte diversity on tree trunks in montane forests of Central Sulawesi, Indonesia. *Tropical Bryology* 31, 95–105
173. Gradstein, S. R. (1992). Threatened bryophytes of the Neotropical rain forest: a status report. *Tropical Bryology*, 6, 83–93
174. Gradstein, S. R. and Allen, N. S. (1992). Bryophyte diversity along an altitudinal gradient in Dárien National Park, Panama. *Tropical Bryology*, 5, 61-71
175. Gradstein, S. R. and Costa, D. P. (2016). A new species of *Syzygiella* (subg. *Cryptochila*) from Brazil. *Nova Hedwigia*, 103, 13–16
176. Gradstein, S. R. and Costa, D. P. (2018) *Plagiochila lamyana*, a new liverwort species from the Guayana Highland of Brazil. *Cryptogamie Bryologie*, 39, 147–153
177. Grau, O., Grytnes, J. A. and Birks, H. J. B. (2007). A comparison of altitudinal species richness patterns of bryophytes with other plant groups in Nepal, Central Himalaya. *Journal of biogeography*, 34(11), 1907-1915
178. *Griffith, W. (1849). *Notulae ad Plantas Asiaticas II*. Government of Bengal, Calcutta
179. Gujar, R. V. and Shah D. G. (2019). Bryodiversity of Wilson Hill and Surrounding Area of Valsad, Gujarat. In Afroz Alam, (Ed.), *Recent Advances in Botanical Science*., Bentham Science Publishers, 1, 81-86 pp
180. Gujar, R. V., Shah, D. G. and Singh, A. P. (2016). The diversity and distribution of mosses in Gujarat: current status. *Cryptogam Biodiversity and Assessment*, 1(2), 83-86

181. Gujarat Ecological Commission (GEC). (2012). *State of Environment Report On Coastal and Marine Environment, Gujarat State*. Gujarat Ecological Commission, Gandhinagar, 13 p
182. Gujarat Forest Department (GFD). (2011). *Management plan: Jambughoda Wildlife Sanctuary*. Wildlife Division, Vadodara Circle, Gujarat Forest Department, Vadodara
183. Gujarat Forest Department (GFD). (2015a). *Management plan: Ratanmahal Wildlife Sanctuary*. Wildlife Division, Vadodara Circle. Gujarat Forest Department, Vadodara
184. Gujarat Forest Department (GFD). (2015b). *Management plan: Shoolpaneshwar Wildlife Sanctuary*. Gujarat, Surat Circle, Narmada Forest Division, Gujarat Forest Department, Surat
185. Gujarat Forest Department (GFD). (2018). *Gujarat Forest Statistics year 2018-19*. Gujarat Forest Department, Gandhinagar, 130 p
186. Gupta, R., Nath, V. and Asthana, A. K. (2013). Present scenario of moss diversity at Tamia hills and Patalkot valley (Madhya Pradesh) India. *National Academy of Science Letter*, 36(6), 629- 634
187. Gupta, R., Nath, V. and Asthana, A. K. (2016). Genus *Brachymenium* Schwagr.(Bryophyta) at Pachmarhi Wildlife Sanctuary (Madhya Pradesh), India. *Tropical Plant Research*, 3(2), 283-288
188. Gupta, R., Nath, V. and Asthana, A. K. (2017). Two species of *Fissidens* (Fissidentaceae): new to Indian bryoflora. *National Academy science letters*, 40(4), 295-299
189. Gupta, R., Nath, V., Pandey, N. and Asthana, A. K. (2018). Diversity and species composition of family Hypnaceae (Bryophyta) at Pachmarhi Sanctuary (India). *Indian Journal of Forestry*, 41(4), 337-351
190. Hagedorn, G, Rambold, G and Martellos, S. (2010). Types of identification keys.In Nimis, P L and Vignes-LebbeR. (Eds), *Tools for identifying biodiversity: progress and problems*, Trieste: EUT Edizioni Universita di Trieste. 59-64 pp
191. Hajibabaei, M., Smith, M. A., Janzen, D. H., Rodriguez, J. J., Whitfield, J. B. and Hebert, P. D. (2006). A minimalist barcode can identify a specimen whose DNA is degraded. *Molecular Ecology Notes,* 6(4), 959-964

192. Hallingback, T. and Hodgetts, N. (2000). *Mosses, Liverworts, and Hornworts. Status Survey and Conservation Action Plan for Bryophytes*. IUCN, Gland, Switzerland and Cambridge, UK. 106 p
193. Hassel, K., Pedersen, B. and Soderstrom, L. (2005). Changes in life-history traits in an expanding moss species: Phenotypic plasticity or genetic differentiation? A reciprocal transplantation experiment with *Pogonatum dentatum*. *Ecography*, 28, 71–80
194. Hassel, K., Segreto, R. and Ekrem, T. (2013). Restricted variation in plant barcoding markers limits identification in closely related bryophyte species. *Molecular ecology resources*, 13(6), 1047-1057
195. Hassid, W. Z. and Putman, E. W. (1950). Transformation of sugars in plants. *Annual Review of Plant Biology*, 1, 109-124
196. Hebert, P. D. N., Cywinska, A., Ball, S. L., de Waard, J. R. (2003) Biological identifications through DNA barcodes. *Proceedings of Royal Society B: Biological Sciences*, 270, 313–321
197. Hebert, P. D. N., Cywinska, A., Ball, S. L., DeWaard, J. R. (2003). Biological identifications through DNA barcodes. *Proceedings of Royal Society B: Biological Sciences*. 270, 313–322
198. Hebert, P. D., Stoeckle, M. Y., Zemlak, T. S. and Francis, C. M. (2004). Identification of birds through DNA barcodes. *Plos biology*, 2(10), 1657-1663
199. Hentschel, J., Zhu, R. L., Long, D. G., Davison, P. G., Schneider, H., Gradstein, S. R. and Heinrichs J. (2007). A phylogeny of *Porella* (Porellaceae, Jungermanniopsida) based on nuclear and chloroplast DNA sequences. *Molecular Phylogenetics and Evolution*, 45(2), 693-705
200. Herdiyeni, Y., Nurfadhilah, E., Zuhud, E. A.Damayanti, E. K., Arai, K. and Okumura, H. (2013). A computer aided system for tropical leaf medicinal plant identification. *International Journal on Advanced Science, Engineering and Information Technology*, 3(1), 23-27.
201. Higuchi, M. (2011). Endemic species of bryophytes in Japan. *Bulletin of the National Museum of Nature and Science, SeriesB*, 37(3), 117-126

202. *Hodgetts, N. G. (2015). Checklist and country status of European bryophytes – towards a new red list for Europe. Ireland: National Parks and Wildlife Service natova EA. 2006. Checklist of mosses of East Europe and North Asia. *Arctoa*, 15, 1–130
203. Hofbauer, W. K., Forrest, L. L., Hollingsworth, P.M. and Hart, M. L. (2016). Preliminary insights from DNA barcoding into the diversity of mosses colonising modern building surfaces. *Bryophyte Diversity and Evolution*, 38(1), 1-22
204. Hollingsworth, M. L., Andra Clark, A. L. E. X., Forrest, L. L., Richardson, J., Pennington, R. T., Long, D. G., Cowan, R., Chase, M. W., Gaudeul, M. and Hollingsworth, P. M. (2009). Selecting barcoding loci for plants: evaluation of seven candidate loci with species-level sampling in three divergent groups of land plants. *Molecular ecology resources*, 9(2), 439-457
205. Holmes, S. (1983). *Outline classification of plants*. Longman group limited, Harlow, England. 182p
206. Holzenthal, R. W. and Rázuri-Gonzales, L. E. (2011). A new species of *Amphoropsyche* (Trichoptera, Leptoceridae) from Ecuador, with a key to the species in the genus. *ZooKeys*, 111, 59-65
207. *Hooker, W. J. (1808). XXIV. Musci Nepalenses; or Descriptions of several new Mosses from Nepal. *Transactions of the Linnean Society of London*, 9(1), 307-322
208. Howe, M. A. (1899). The hepaticae and anthocerotes of California. *Memoirs of the Torrey Botanical Club*, 7, 1-208
209. Hueber, F. M. (1961). Hepaticites devonicus, a new fossil liverwort from the Devonian of New York. *Annals of the Missouri Botanical Garden*, 48, 125–132
210. Ignatov, M. S., Afonina, O. M. and Ignatova, E. A. (2006). Check-list of mosses of East Europe and North Asia. *Arctoa,* 15, 1–130
211. *Jaeger, A. and Sauerbeck A. W. (1879). *Genera et Species Muscorum,* St. Gallen : Zollikofer'sche Buchdruckerei, 825 p
212. Johri, M. M. and Desai, S. (1973). Auxin regulation of caulonema formation in moss protonema. *Nature New Biology*, 245(146), 223-224
213. Kajita, M., Takio, S., Takami, S. and Hino, S. (1987). Establishment and growth characterization of suspension culture of cells from the moss, *Sphagnum imbricatum*. *Physiologia Plantarum* 70, 21–26

214. Kallio, P. and Heinonen, S. (1973). *Ecology of Rhacomitrium lanuginosum (Hedw.) Brid.* Report of Kevo Subartic Research. Station. 10, 43-54
215. Kapoor, R., Nath, V. and Asthana, A. K. (2011). Bryophytes of Kanha national Park (Madhya Pradesh), India. *Nelumbo*, 53, 140-144
216. Kashyap, S. R. (1914). Morphological and biological notes on new and little known West Himalayan liverworts II. *New Phytologist*, 13, 308-323
217. Kashyap, S. R. (1915). Morphological and biological notes on new and little known western Himalayan Liverworts. III. *New Phytologist*, 14(1), 1-48
218. Kashyap, S. R. (1917). Liverworts of the Western Himalayas and the Punjab, with notes on known species and descriptions of new species. *Journal of the Bombay Natural History Society*, 25, 279-281
219. Kashyap, S. R. (1929). *Liverworts of western Himalayas and Panjab plains. Vol. 1*. University of Panjab, Lahore. 129 p
220. Kashyap, S. R., and Chopra, R. N. (1932).*Liverworts of western Himalayas and Panjab plains. Vol. 2* ¸University of Panjab, Lahore. 137 p
221. Kaur, S. and Kaur, P. (2019). Plant species identification based on plant leaf using computer vision and machine learning techniques. *Journal of Multimedia Information System*, 6(2), 49-60
222. Klavina, L., Springe, G., Nikolajeva, V., Martsinkevich, I., Nakurte, I., Dzabijeva, D. and Steinberga, I. (2015). Chemical composition analysis, antimicrobial activity and cytotoxicity screening of moss extracts (moss phytochemistry). *Molecules,* 20(9), 17221-17243
223. Kranz, H. D., Miks, D., Siegler, M. L., Capesius, I., Sensen, C. W. and Huss, V. A. (1995). The origin of land plants: phylogenetic relationships among charophytes, bryophytes, and vascular plants inferred from complete small-subunit ribosomal RNA gene sequences. *Journal of Molecular Evolution*, 41(1), 74-84
224. Kress, W. J. and Erickson, D. L., 2007. A two-locus global DNA barcode for land plants: the coding rbcL gene complements the non-coding trnH-psbA spacer region. *PLoS one*, 2(6), 1-10
225. Kress, W. J. and Erickson, D. L. (2012). *DNA barcodes: methods and protocols*. In DNA Barcodes. Humana Press, Totowa, NJ. 3-8 pp

226. Kress, W. J., Erickson, D. L., Jones, F. A., Swenson, N. G., Perez, R., Sanjur, O. and Bermingham, E. (2009). Plant DNA barcodes and a community phylogeny of a tropical forest dynamics plot in Panama. *Proceedings of the National Academy of Sciences*, 106(44), 18621-18626
227. Kress, J. W., Wurdack, K. J., Zimmer, E. A., Weigt, L. A.and Janzen, D. H. (2005). Use of DNA barcodes to identify flowering plants. *Proceedings of the National Academy of Sciences of the United States of America*, 102, 8369–8374
228. Lal, J. and Parihar, N. S. (1979). Contributions to the bryoflora of central Indian Zone I, Liverworts. *Journal of Indian Botanical Society*, 58, 110-114
229. Lang, A. S., Bocksberger, G. and Stech, M. (2015). Phylogeny and species delimitations in European *Dicranum* (Dicranaceae, Bryophyta) inferred from nuclear and plastid DNA. *Molecular Phylogenetics and Evolution*, 92, 217-225
230. Lang, A. S., Kruijer, J. D. and Stech, M. (2014). DNA barcoding of Arctic bryophytes: an example from the moss genus *Dicranum* (Dicranaceae, Bryophyta). Polar biology, 37(8), 1157-1169
231. Lavate, R. A. (2020). Bryodiversity, Threats and Conservation of Liverworts and Hornworts of Kolhapur District (Maharashtra), India. *In Contemporary Research on Bryophytes*, 1,134-158
232. *Lebbe, J. (1984). *Manuel d'utilisation du logiciel XPER*. Micro Application, Paris.148 p.
233. Legg, C. J. (1992). Random-access identification guides for a microcomputer. *Field Studies*, 8(1), 1-30
234. Leutner, B., Horning, N., Rocchini, D, and Wegmann, M (2016) *Spatial data and software in Remote Sensing and GIS for Ecologists: Using Open Source Software*,(Eds). Wegmann, M., Leutner, B and Dech,S., Pelagic Publishing, UK, 324 p
235. Levin, R. A., Wagner, W. L., Hoch, P. C., Nepokroeff, M., Pires, J. C., Zimmer, E. A. and Sytsma, K. J. (2003). Family-level relationships of Onagraceae based on chloroplast rbcL and ndhF data. *American Journal of Botany*, 90(1), 107-115
236. Lewis, L. A., Mishler, B. D.and R. Vilgalys, R.(1997). Phylogenetic relationships of the liverworts (Hepaticae), a basal embryophyte lineage, inferred from nucleotide sequence data of the chloroplast gene rbcL. *Molecular phylogenetics and evolution*, 7, 377–393

237. Lewis, L. R., Ickert-Bond, S. M., Biersma, E. M., Convery, P., Goffinet, B., Hassel, K., Kruijer, H. J. D., La Farge, C., Metzgar, J., Stech, M., Villarreal, J. C., and McDaniel, S.F. (2017). Future directions and priorities of Arctic bryophyte research. *Arctic Science*, 3, 475–497
238. Liang, S. F., Sun, Y. and Zhu, R. L. (2010). In vitro micropropagation of *Bryum argenteum* Hedw. *Cryptogamie Bryologie*, 31(3), 233-239
239. *Linnaeus, C. (1753). *Species Plantarum*. (2nd Ed.) *Vol. II, Musci. Holmiae*. 1200 pp
240. Liu ,Y., Cao, T., and Ge, X. J. (2011) A case study of DNA barcoding in Chinese Grimmiaceae and a moss recorded in China for the first time. *Taxon*, 60, 185–193
241. Liu, Y., Yan, H. F., Cao, T. and Ge, X. J. (2010) Evaluation of 10 plant barcodes in Bryophyta (Mosses). *Journal of Systematics and Evolution*, 48, 36–46
242. Lodge, E. (1959). Effects of certain cultivation treatments on the morphology of some British species of *Drepanocladus*. *Botanical Journal of the Linnean Society*, 56(366), 218-224
243. Loreti, E., Bellis, L.D., Alpi, A. and Perata, P. (2001). Why and how do plant cells sense sugars? *Annals of Botany*, 88(5), 803-812
244. Machlis, L. and Doyle W, T. (1962). Submerged growth of pure cultures of the liverwort *Sphaerocarpos donnellii*. *Physiologia Plantarum*, 15, 351–353
245. Magdum, S. M., Lavate, R. A., Patil, S. M. andDongare, M. M.(2017a) Studies on corticolous Mosses from Panhalgad in Western Ghats of Maharashtra, India. *Bioscience Discovery*, 8(4), 796-801
246. Magdum, S. M., Patil, S. M.; Lavate, R. A. and Dongare, M. M. (2017b). Checklist of mosses from the Western Ghats of Maharashtra, India. *Bioscience Discovery*, 8(1), 73-81
247. Magill, R. E. (2010). Moss diversity: new look at old numbers. *Phytotaxa*, 9, 167–174
248. Mahabale, T. S. (1941). A long-lost Liverwort from south India. *Aspiromitus* A rare member of the Anthocerotes.*Currrent Science*, 10, 530-533
249. Maiden, M. C. J. (2006). Multi-locus sequence typing of bacteria. *Annual Review of Microbiology*, 60, 561-588

250. Mallon, R., Reinoso, J., Rodriguez-Oubina, J. and Gonzalez, M. L. (2006). In vitro development of vegetative propagules in *Splachnum ampullaceum*: brood cells and chloronematal bulbils. *The Bryologist*, 109(2), 215-223
251. Manju, C. N., Rajesh, K. P. and Madhusoodanan, P. V. (2009). Contribution to the bryophyte flora of India: Agasthyamalai Biosphere Reserve in Western Ghats. *Taiwania*, 54(1), 57-68
252. *Marchal, L. E.and Marchal, E. (1906). Recherches physiologiques sur l'amidon chez les Bryophytes. *Bulletin de la Société royale de botanique de Belgique*., 43, 115-214
253. Marshall, E. (2005). Taxonomy: will DNA bar codes breathe life into classification? *Science*, 307, 1037
254. Mathur, L. P., Rao, K. L. N. and Chaube, A. N. (1968) Tectonic Framework of Cambay Basin. *Bulletin of Oil and Natural Gas Commission*, 5(1), 7-28
255. MBG (1995-2021). MBG Research: Bryology Home Page http://www.mobot.org/MOBOT/tropicos/most/welcome.shtml. Accessed on 21st January 2021
256. Mehta, D. N., Gujar, R. V. and Shah, D. G. (2016). Status of family Fissidentaceae in Gujarat. *Cryptogam Biodiversity and Assessment*, 1(1), 84-91
257. Merh S. S. (1995). *Geology of Gujarat*. Geological society of India, Bangalore. 222p
258. Micro Small Medium Enterprise (MSME). (2016a). *Brief Industrial Profile of Chhota Udepur District.* MSME Development Institute, Government of Gujarat, Ahmedabad, 65p .
259. Micro Small Medium Enterprise (MSME). (2016b).*Brief Industrial Profile of Dahod District*. MSME Development Institute, Government of Gujarat, Ahmedabad, 11p
260. Micro Small Medium Enterprise (MSME). (2016c). *Brief Industrial Profile of NarmadaDistrict*. MSME Development Institute, Government of Gujarat, Ahmedabad, 16p
261. Micro Small Medium Enterprise (MSME). (2016d). *Brief Industrial Profile of Panchmahal District*. MSME Development Institute, Government of Gujarat, Ahmedabad, 9 p .
262. Micro Small Medium Enterprise (MSME). (2016e) *Brief Industrial Profile of Vadodara District.* MSME Development Institute, Government of Gujarat, Ahmedabad, 42 p

263. *Mishler, B. D. (1985) Biosystematic studies of the *Tortula ruralis* complex. I. Variation of taxonomic characters in culture. *Journal of the Hattori Botanical Laboratory*, 58, 225–253
264. Mishler, B. D. and Churchill, S. P. (1984). A cladistic approach to the phylogeny of the bryophytes. *Brittonia*, 36, 406–424
265. Mishra, R., Vijay, P. and Ramesh, C. (2014). Potential of Bryophytes as Therapeutics, *International Journal of Pharmaceutical Sciences and Research*, 5(9), 3584-3593
266. *Mitten, W. (1859). *Musci Indiae Orientalis: an enumeration of the mosses of the East Indies (Vol. 1)*. Longman, Brown, Green, Longmans & Roberts, and Williams and Norgate, London,1-96 pp
267. *Mitten, W. (1860). Hepaticae Indiae Orientalis: an enumeration of the Hepaticae of the East Indies. *Botanical Journal of the Linnean Society*, 5(18), 89-108
268. *Mitten, W. (1861). Hepaticæ Indiae Orientalis: an Enumeration of the Hepaticæ of the East Indies.(continued). *Botanical Journal of the Linnean Society*. 5(19), 109-128
269. Mogensen, G. S. (1978). Spore development and germination in Cinclidium Sw. (Mniaceae Mitt., Bryophyta), with special reference to spore mortality and false anisospory. *Canadian Journal of Botany*, 56, 1032-1060
270. *Montagne, C. J. F. (1842). Cryptogamme Nilgheriensis. *Annales des Sciences NaturellesBotanique*, 17, 213-256
271. Morris, J. L., Puttick, M. N., Clark, J. W., Edwards, D., Kenrick, P., Pressel, S. and Donoghue, P. C. (2018). The timescale of early land plant evolution. *Proceedings of the National Academy of Sciences*, 115(10), 2274-2283
272. Morse, L. E. (1974). *Computer programs for specimen identification, key construction and description printing*. Publ. Mus. Michigan State University, Biology
273. *Mota de Oliveira S. (2010). Diversity of epiphytic bryophytes across the Amazon. Ph.D. thesis submitted toUniversiteit Utrecht, Netherlands
274. Mower, J. P., Stefanovic, S., Young, G. J. and Palmer, J. D. (2004). Gene transfer from parasitic to host plants. *Nature*, 432(7014), 165-166
275. Muller, S. J., Gutle, D. D., Jacquot, J. P. and Reski, R. (2016). Can mosses serve as model organisms for forest research? *Annals of forest Science*, 73(1), 135-146

276. Nair M. C. Rajesh K. P. and Madhusoodanan P. V. (2005). *Bryophytes of Wayanad in Western Ghats.* Malabar Natural History Society, Calicut. 284 p
277. Nakanishi, K. (2001). Floristic diversity of bryophyte vegetation in relation to island area. *The Journal of the Hattori Botanical Laboratory*, 91, 301-316
278. Narendran, T. C. (2008). Taxonomy and its Relevance. *Govt. Arts and Science College Research Journal*. 1,9-14
279. Narmada, Water resources, water supply and Kalpsar Department (NWRWS). (2019a). Geology Map of Gujarat. https://gujnwrws.gujarat.gov.in/showpage.aspx?contentid=1459&lang=English. Accessed on 15th November 2019
280. Narmada, Water resources, water supply and KalpsarDepartment (NWRWS). (2019b). River Map of Gujarat. https://guj-nwrws.gujarat.gov.in/showpage.aspx?contentid=1524&lang=english . Accessed on 15th November 2019.
281. Narmada, Water resources, water supply and Kalpsar Department (NWRWS). (2019c). Soil Map of Gujarat https://gujnwrws.gujarat.gov.in/showpage.aspx?contentid=1518&lang=english. Accessed on 15th November 2019.
282. Nath, V. and Asthana, A. K. (1998). Diversity and distribution of genus *Frullani radii* in South India. *Journal of Hattori Botanical Laboratory*, 85, 63-82
283. Nath, V. and Bansal, P. (2009). Bryophyte diversity of Bhimbetka world heritage site, Madhya Pradesh (India). *Journal of Indian Botanical Society*, 88(3&4), 129-140
284. Nath, V. and Gupta, R. (2009). *Anoectangium strachyanum* Mitt. – A moss new to central *India. Geobios*, 36, 5-8
285. Nath, V. and Singh, A. P. (2006). *Frullania udarii* sp. nov. – A new species from Meghalaya, India. *Current science*, 91(6), 744-746
286. Nath, V., Asthana, A. K. and Gupta, R. (2011a). An overview of family Pottiaceae (Bryopsida) in central India with special reference to Pachmarhi Biosphere Reserve (PBR). *Lindbergia*, 34, 30-39

287. Nath, V., Asthana, A. K. and Gupta, R. (2011b). Genus Fissidens Hedw.(Fissidentaceae, Bryopsida) at Pachmarhi Biosphere Reserve (Madhya Pradesh), India. *Taiwania*, 56(1), 71-80
288. Nath, V., Asthana, A. K. and Kapoor, R. (2005). A study on the genus *Fissidens* Hedw. (Musci) of Achanakmar sanctuary (Chhatisgarh). *IndianJournal ofForestry*, 28(4), 433-438
289. Nath, V., Asthana, A. and Kapoor, R. (2007). Enumeration of the Mosses in Amarkantak (Madhya Pradesh), India - I. *Taiwania*, 52(2), 168-176
290. Nath, V., Pande, N., Asthana A. K. and Gupta, R. (2012). Epiphytic moss flora of Pachmarhi Biosphere Reserve (M.P.): An important aspect of Bryophyte diversity.*National Academy Science Letter*, 35(3), 195-200
291. Nath, V., Pande, N., Asthana, A. K.and Gupta, R. (2013). Some acrocarpus mosses new to central India. *Geophytology*, 42(1), 49-57
292. Negbi, M. (1989). Theophrastus on geophytes. *Botanical Journal of the Linnean Society*, 100, 15-43
293. Negi, H. R. (2001). Diversity and dominance of liverworts of Chopta Tunaganathinthe Garhwal Himalaya. *Journal of Ecology and Evironment Science*, 27, 13-21
294. Nemejc, F. and Pacltova, B. (1974). Hepaticae in the Senonian of South Bohemia. *Paleobotanist*, 21, 23–26
295. Newmaster, S. G., Fazekas, A. J. and Ragupathy, S. (2006). DNA barcoding in land plants: evaluation of rbcL in a multigene tiered approach. *Botany*, 84(3), 335-341
296. *Noguchi, A. (1987). *Illustrated Moss Flora of Japan. Part 1*. Hattori Botanical Laboratory, Nichinan.1-242 pp
297. *Noguchi, A. (1988). *Illustrated Moss Flora of Japan. Part 2*. Hattori Botanical Laboratory, Nichinan. 243–491 pp.
298. *Noguchi, A. (1989). *Illustrated Moss Flora of Japan. Part 3.* Hattori Botanical Laboratory, Nichinan. 493-742 pp.
299. *Noguchi, A. (1991). *Illustrated Moss Flora of Japan. Part 4*. Hattori Botanical Laboratory, Nichinan. 743-1012 pp.
300. *Noguchi, A. (1994). *Illustrated Moss Flora of Japan. Part 5*. Hattori Botanical Laboratory, Nichinan. 1013-1253 pp.

301. Ogilvie, B. W. (2003). The many books of nature: Renaissance naturalists and information overload. *Journal of History of Ideas*, 64(1), 29-40
302. Olesen, P. and Mogensen, G. S. (1978). Ultrastructure, histochemistry and notes on germination stages of spores in selected mosses. *The Bryologist*, 81, 493- 516
303. Oliveira, B. A., Pereira, A. F. D. N., Porto, K. C. and Maciel-Silva, A. S. (2017). Spore germination and young gametophyte development of the endemic Brazilian hornwort *Notothylas vitalii* Udar & Singh (Notothyladaceae-Anthocerotophyta), with insights into sporeling evolution. *Acta Botanica Brasilica*, 31(2), 313-318
304. Palani, R., Sathish, S. S., Thamizharasi, T. and Vijayakanth, P. (2017). Checklist of Mosses (Bryophyta) of Bodamalai Hills in Eastern Ghats, Tamil Nadu. *Plant Science Today*, 4(1), 49-54
305. Palhares, R. M., Drummond, M. G., Brasil, B. D. S. A. F., Cosenza, G. P., Brandão, M. D. G. L. and Oliveira, G. (2015). Medicinal plants recommended by the world health organization: DNA barcode identification associated with chemical analyses guarantees their quality. *PloS one*, 10(5), 1-29
306. Palmer, J. D., Adams, K. L., Cho, Y., Parkinson, C . L., Qiu, Y. L. and Song, K. (2000). Dynamic evolution of plant mitochondrial genomes: mobile genes and introns and highly variable mutation rates. *Proceedings of the National Academy of Sciences*, 97(13), 6960-6966
307. Pandey, B. P. (2005). *College Botany Vol. 1*. S. Chand Publishing house, New Delhi. 221 p
308. Pandey, V. J., Mishra, R. and Chandra, R. (2014). Invitro culture of moss B*ryumcoranatum* Schwaegr. (Bryaceae) and its phytochemical analysis. *International Journal of Pharmacy and Pharmacuetical Sciences*, 6(9), 307-311
309. Pankhrust, R. J. (1975). *Biological Identification with computers*. Academic Press Inc. 343 p
310. Pankhurst, R. J. (1991). *Practical taxonomic computing*. Cambridge University Press. 202 p
311. *Paris, E. G. (1898).*Index bryologicus sive, Enumeratio muscorum hucusque cognitorum adjunctis synonymia distributioneque geographica locupletissimus*, Parisiis Apud Paul Klincksieck, 1380 p

312. Pasarell, L (1999) *Computer assisted programs or references available for the mycologists or microbiologist*. http://www.doctorfungus.org. Accessed on 16th April 2020

313. Passinho-Soares, H., Felix, D., Maria Auxiliadora Kaplan, M. A., Margis-Pinheiro, M. and Margis, R. (2006). Authentication of medicinal plant botanical identity by amplified fragmented length polymorphism dominant DNA marker: inferences from the Plectranthus genus. *Planta Medica*, 72(10), 929-931

314. Patel, R. J. (1977a). On *Riella affinis*, Hepaticeae from India. *Aquatic Botany*,3(3), 287-290

315. Patel, R. J. (1977b). On *Riella cossoniana* New record from Gujarat state.*Journal of Indian Botanical Society*, 56(3), 237-239

316. Pedersen, N. and Hedenas, L. (2003). Phylogenetic investigations of a well supported clade within the acrocarpous moss family Bryaceae: evidence from seven chloroplast DNA sequences and morphology. *Plant Systematics and Evolution*, 240(1),115-132

317. Pedersen, N. and Hedenas, L. (2005). Taxonomic and nomenclatural implications of phylogenetic studies of the Bryaceae based on molecular data and morphology. *The Bryologist*, 108(1), 123-128

318. Pedersen, N., Holyoak, D.T. and Newton, A. E. (2007). Systematics and morphological evolution within the moss family Bryaceae: A comparison between parsimony and Bayesian methods for reconstruction of ancestral character states. *Molecular phylogenetics and evolution*, 43(3), 891-907

319. Pedersen, N., Russell, S. J., Newton, A. E. and Ansell, S. W. (2006). A novel molecular protocol for the rapid extraction of DNA from bryophytes and the utility of direct amplification of DNA from a single dwarf male. *The Bryologist*, 109(2), 257-264

320. Pennisi, E. (2007). Taxonomy – Wanted: a barcode for plants. *Science,* 318, 190–191

321. Piippo, S. (1992). On the phytogeographical affinities of temperate and tropical Asiatic and Australasiatic hepatics. *Journal of the Hattori Botanical Laboratory*, 71, 1–35

322. Post, A, R., McCall, D. S.and Askew, S. D. (2016). Pre emergence control of silvery threadmoss (*Bryum argenteum*) grown from spores and bulbils in Axenic culture. *Weed Technology*, 30,198-206

323. Pressel, S. Matcham, H.W., Duckett, J. G. (2007). Studies of protonemal morphogenesis in mosses. XI. *Bryum* and allied genera: a plethora of propagules. *Journal of Bryology*, 26, 241-258

324. Priya, C. A., Balasaravanan, T. and Thanamani, A. S. (2012).An efficient leaf recognition algorithm for plant classification using support vector machine. *In International conference on pattern recognition, informatics and medical engineering (PRIME-2012) IEEE*, 482-834 pp

325. Proctor, M. C. F. (2014). The diversification of bryophytes and vascular plants in evolving terrestrial environments. In Hanson, D. T. and Rice, S. K. (Eds.)., *Photosynthesis in Bryophytes and Early Land Plants. Advances in Photosynthesis and Respiration*, 37, 59-77

326. Proskauer, J. (1957). Studies on Anthocerotales V. *Phytomorphology*, 7, 113-135

327. Prosperi, J., Ramesh, B. R., Grard, P., Jayatissa L. P., Aravajy, S. and Depommier, D. (2005).*Mangroves v.1.0 - A multimedia identification system of mangrove species*. Available in DVD

328. Raghavan, R. and Wadhwana, B. M. (1970). Mosses of Agumbe (Shimoga district Mysore state) II key to species and new records for India. *Bulletin of Botanical survey of India*, 5, 28-35

329. Raghavan, R.S. and Wadhwa, B.M. (1968). Mosses of Agumbe-Hulical Ranges, Shimoga District. *Nelumbo*, 10(3-4), 344-347

330. Ramasubbu, R., Sathya, S. and Manikandan, G. (2015). Computer aided alternative tool for the identification of angiosperm families (dicots) of south India. In :Recent Advances in Computer Science and Applications. International Conference on Evolution in Engineering & Management (ICEEM-2015), 359-362 pp

331. Ramesh, B. R., Ayyappan, N., Grard, P., Prosperi, J., Aravajy, S. and Pascal, J. P. (2010). *Western Ghats v.1.0 - A multimedia identification system of evergreen species of the Western Ghats*, India. Available in DVD & online

332. Ramsay, H. P. and Cairns, A. (2004). Habitat, distribution and the phytogeographical affinities of mosses in the Wet Tropics bioregion, north-east Queensland, Australia.*Cunninghamia,* 8, 371–408

333. Rana, S. (2020). Genus *Physcomitrium* (brid.) Fuernr (bryophyta) in district Banswara of south Rajasthan. *Plant Archives*, 2(2), 4697-4700

334. Rao, G. N. and Dora, S. (2019). Some ecological studies on distribution of bryophytes and its application in Vishakhapatnam district Eastern Ghats of India. *Bionature*, 39(2), 99-105

335. Rashid, A. (2007). *An Introduction to Bryophyta (Diversity, Development and Differentiation)* 1st edition, Vikas Publishing House Pvt Ltd, New Delhi, 225 p

336. Ratnasingham, S. and Hebert, P. D. (2007). BOLD: The Barcode of Life Data System (http://www. barcodinglife. org). *Molecular ecology notes*, 7(3), 355-364

337. Rawat, K. K., Alam, A. and Verma, P. K. (2015). Moss flora of Rajasthan and Punjab plains. *Plant Science Today*, 2(4), 154-158

338. Redfearn, P. L. (1990). Tropical component of the moss flora of China. *Tropical Bryology*, 2, 201–222

339. Reid, G. M. (2009). Carolus Linnaeus (1707–1778): His life, philosophy and science and its relationship to modern biology and medicine. *Taxon,* 58(1), 18-31

340. Renzaglia K. S., Duff R. J., Nickrent D. L. and Garbary D. J. (2000). Vegetative and reproductive innovations of early land plants: implications for a unified phylogeny. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 355, 769–793 Report of Kevo Subartic Research. Station. 10, 43-54

341. Reski, R. andAbel, W. O. (1985). Induction of budding on chloronemata and caulonemata of the moss, *Physcomitrella patens*, using isopentenyladenine. *Planta*, 165, 354-358

342. Rice, S. K. (2009). Mosses (Bryophytes). In: *Encyclopedia of Inland Waters*. (Eds), Likens G.E., Academic press. 88-96 pp

343. Richardson, J. B. (1985). Lower Palaeozoic sporomorphs: their stratigraphical distribution and possible affinities. *Philosophical Transactions of the Royal Society B: Biological Sciences*. 309, 201–205

344. Rivera-Jimenez, H., Rossini, B. C., Tambarussi, E. V., Veasey, E. A., Ibanes, B. and Marino, C. L. (2017). DNA barcode regions for differentiating *Cattleya walkeriana* and C. *loddigesii*. *Acta Scientiarum*, 39, 45–52

345. Rodgers, W. A. and Panwar, H. S. (1988) *Planning a Protected Area Network in India, Vol. 1: Technical Report*. Wildlife Institute of India, Dehradun, 267 p

346. Ros, R. M., Mazımpaka, V.,Abou-Salama, U., Aleffi, M., Blockeel, T. L.,Brugues, M., Cros, R. M., Dia, M. G.,Dirkse, G. M., Draper, I.and El-Saadawi, W. (2013). Mosses of the mediterranean, an annotated checklist. *Cryptogamie Bryologie*,34, 99–283

347. *Rothmaler, W. (1951). *Die Abteilungen und Klassen der Pflanzen- Field Report*. Repertorium specierum novarum regni vegetabilis. 54, 256-266

348. Rowntree, J. K. (2006). Development of novel methods for the initiation of in vitro bryophyte cultures for conservation. *Plant cell, tissue and organ culture*, 87(2), 191-201

349. *Rudolph, H. J. (1978). *Jahre Kultur von Sphagnum inter definierten Bedingungen: Eine Übersicht über Resultate, Probleme und Prespectiven*. In: Suire C, ed. Bryophytorum Bibliotheca 13: 279–309

350. Sabovljevic, A. andSabovljevic, M. (2008). *Bryophytes, a source of bioactive and new compounds.* In: Phytopharmacology and Therapeutic Values IV, the Series "Recent Progress in Medicinal Plants". (Ed.): J.N. Govil. Studium Press, Houston, Texas, USA. 9-25 pp

351. Sabovljevic, M. S., Bijelovic, A. and Dragicevic, I. C. (2002). Effective and easy way of establishing in vitro culture of mosses, *Bryum argenteum* Hedw. and *Bryum capillare* Hedw, Bryaceae. *Archives of Biological Sciences*, 54(1-2): 7-8

352. Sabovljevic, M.,Bijelovic A. andDragicevic I. (2003). In vitro culture of mosses: *Aloina aloides* (K.F. Schultz) Kindb., *Brachythecium velutinum* (Hedw.) B.S. & G., *Ceratodon purpureus* (Hedw.) Brid., *Eurhynchium praelongum* (Hedw.) B.S.& G. and *Grimmia pulvinata* (Hedw.) Sm. *Turkish Journal of Botany*, 27, 441-446

353. Sabovljevic, A., Sabovljevic, M. and Grubisic, D. (2010). Gibberellin influence on the morphogenesis of the moss *Bryum argenteum* Hedw. in in vitro conditions. *Archives of Biological Sciences*, 62(2), 373-380

354. Sabovljevic, A., Sabovljevic, M., Grubisic, D. and Konjevic, R. (2005). The effect of sugars on development of two moss species (*Bryum argenteum* and *Atrichum undulatum*) during in vitro culture. *Belgian journal of botany*, 138(1), 79-84

355. Sabovljevic, M. S., Papp, B., Sabovljevic, A., Vujicic, M., Szurdoki, E. and Segarra-Moragues, J. G. (2012). In vitro micropropagation of rare and endangered moss *Entosthodon hungaricus* (Funariaceae). *Bioscience Journal*, 28(4), 632-640
356. Sahu, V. andAsthana, A.K. (2013). An observation on growth response of *Anomobryum filiforme* var. *concinnatum* (Spruce) Aman. (Bryaceae) in different Culture Media. *National Academy Science Letters*. 36(6), 587–589
357. Sahu, V. and Asthana A. K. (2013). An observation on growth response of *Anomobryum filiforme* var. *concinnatum* (Spruce) Aman. (Bryaceae) in different culture media. *National Academy Science Letters*, 36(6), 587–589
358. Sahu, V. and Asthana, A. K. (2015). Bryophyte diversity in Terai regions of Uttar Pradesh, India with some new additions to the state. *Tropical Plant Research*. 2(3), 180-191
359. Sahu, V. and Asthana, A.K. (2018). A study on photomorphogenesis of protonema and bud formation in *Pohlia ludwigii* (Spreng. ex Schwägr.) Broth. *Indian Forester*, 144(8), 781-783
360. Sasidharan, N. (2006). *Illustrated manual on tree flora of Kerala supplemented with computer aided identification*. Kerala Forest Research Institute, Peechi, Kerala. 698p.
361. Saucede, T.,Eleaume, M., Jossart, Q.,Moreau, C., Downey, R.,Bax, N., Sands, C., Mercado, B., Gallut, C. and Vignes-Lebbe, R. (2021). Taxonomy 2.0: computer-aided identification tools to assist Antarctic biologists in the field and in the laboratory. *Antarctic Science*, 33(1), 39-51
362. Saxena D. and Harinder K. (2004). Uses of bryophytes. *Resonance*, 9, 56-65
363. Schoch, C.L., Seifert, K. A., Huhndorf, S., Robert, V., Spouge, J. L., Levesque, C. A., Chen, W. and Fungal Barcoding Consortium (2012). Nuclear ribosomal internal transcribed spacer (ITS) region as a universal DNA barcode marker for Fungi. *Proceedings of the National Academy of Sciences*, 109(16), 6241-6246
364. Schussler, A. (2000). *Glomus claroideum* forms an arbuscular mycorrhiza-like symbiosis with the hornwort *Anthoceros punctatus*. *Mycorrhiza*, 10, 15–21
365. Schuster, R. M. (1958). Boreal Hepaticae, a manual of the liverworts of Minnesota and adjacent regions III. Phytogeography. The American Midland Naturalist, 59(2), 257-332

366. Schwarz, U. (2013). An updated checklist of Bryophytes of Karnataka. *Archive for Bryology*, 181,1-42
367. *Servettaz C. (1913). Researches experimentales sur le development et la nutrition des mousses en milieu sterilise. *Annales des sciences naturelles Botanique*, 17, 111-223
368. Shah, D. G. and Gujar, R. (2016). Diversity and Distribution of Liverworts in Gujarat–Current Status. *Plant Science Today*, 3(2),124-128
369. Shah, D. G. and Gujar, R. V.(2014). *A pilot study on Bryophyte diversity and distribution in Gujarat*. Gujarat Biodiversity Board, Gandhinagar, 117 p
370. Sharkey, M. J., Clutts, S., Tucker, E. M., Janzen, D., Hallwachs, W., Dapkey, T. and Smith, M. A. (2011). *Lytopylus forster* (Hymenoptera, Braconidae, Agathidinae) species from Costa Rica, with an emphasis on specimens reared from caterpillars in Area de Conservacion Guanacaste. *ZooKeys*, (130), 379-419
371. Sharma, K., Agrawal, V., Gupta, S., Kumar, R. and Prasad, M. (2008). ISSR marker-assisted selection of male and female plants in a promising dioecious crop: jojoba (Simmondsia chinensis). *Plant Biotechnology Reports*, 2(4), 239-243
372. Sharma, O. P. (2014). *Diversity of microbes and cryptogams: Bryophyta*. McGraw Hill Education, India. 416 p
373. Shaw A. J. and Renzaglia K. J. (2004). Phylogeny and diversification of bryophytes. *American Journal of Botany*, 91, 557-1581.
374. Shaw, J. (1986). A new approach to the experimental propagation of bryophytes. *Taxon*, 35, 671-675
375. Shevock, J. R., MA, W. Z. and Akiyama, H. (2017). Diversity of the rheophytic condition in bryophytes: field observations from multiple continents. *Bryophyte Diversity and evolution*, 39(1), 75-93
376. Silva, A. S. M. D., Porto, K. C. and Simabukuro, E. A. (2010). Effects of light and nutrients on different germination phases of the cosmopolitan moss *Bryum argenteum* Hedw. (Bryaceae). *Brazilian Archives of biology and technology*, 53(4), 763-769
377. Silva-e-Costa, J. D. C., Luizi-Ponzo, A. P., Resende C. F. D. and Peixoto, P. H. P. (2016). Spore germination, early development and some notes on the effects of in vitro

culture medium on *Frullania ericoides* (Nees) Mont. (Frullaniaceae Marchantiophyta).*ActaBotanica Brasilica*, 31(1), 19-28

378. Simola, L. K. (1975). The effects of several protein amino acids and some inorganic nitrogen sources on the growth of *Sphagnum nemoreum*. *Physiologia Plantarum*, 35, 195–199
379. Singh, D. K. (2002).*Notothylaceae of India and Nepal* (*A morpho-taxonomic revision).* Bishen Sing h Mahendra Pal Singh, Dehra Dun. 271 pp.
380. Singh, D. K. and Singh, S. K. (2003). *Conocephalum japonicum* (Thumb.) Groll (Bryophyta: Conocephalaceae)-A liverwort rediscovered in India. *Indian Journal of Forestry*, 26, 442-444
381. Singh, S. K. (1999). Indian hepaticae: status and strategies. *Annals of Forest Science*, 7, 199-211
382. Singh, S. K., Singh D. and Singh, D. K. (2004).On two noteworthy liverworts form Western Himalaya India. *Annals of Forest Science*, 12, 56-60
383. Singh, S.K. and Kumar, S. (2016). A preliminary study on liverworts and hornworts of Tripura, North-East India. *Nelumbo*, 58, 130-151
384. Slipiko, M., Myszczynski, K., Buczkowska, K., Bączkiewicz, A., Szczecinska, M. and Sawicki, J. (2020). Molecular delimitation of European leafy liverworts of the genus *Calypogeia* based on plastid super-barcodes. *BMC Plant Biology*, 20, 1-15
385. Smeekens, S. (2000). Sugar-induced signal transduction in plants. *Annual Review of Plant Physiology and Plant Molecular Biology*, 51, 49-81
386. Smith, G. M.(1955). *Cryptogamic Botany: Bryophytes and Pteridophytes*. 2nd Ed. Tata McGraw-Hill Publishing, London.380 p
387. Smith, G.M. (1938). *Cryptogamic botany. Volume II. Bryophytes and pteridophytes* Tata McGraw-Hill Publishing Co. Ltd, London.380 p
388. Soderstrom, L., Hagborg, A., von Konrat, M., Bartholomew- € Began, S., Bell, D., Briscoe, L., Brown, E., Cargill, D. C., Costa, D. P., Crandal-Stotler, B. J., Cooper, E. D., Dauphin, G., Engel, J. J., Feldberg, D. G., Gradstein, S. R., He, X., Heinrichs, J., Hentschel, J., Ilkiu-Gorges, A. L., Katagiri, T., Konstantinova, N. A., Larraın, J., Long, D. G., Nebel, M., Pocs, T., Puche, F., Reiner-Drehwald, E., Renner, M. A. M. m Sass-Gyarmati, A., Schafer-

Verwimp, A., Moragues, J. G. S., Stotler, R. E., Sukkharak, P., Thiers, B. M., Uribe, J., Vana, J., Villarreal, J. C., Wigginton, M., Zhang, L., and Zhu R.-L. (2016). World checklist of hornworts and liverworts. *PhytoKey*s, 59, 1–828

389. Soni, A. and Kumar, A. (2009). Protocol for improved extraction and PCR amplification of genomic DNA from liverwort, *Plagiochasma appendiculatum*. *IndianJournal of Experimental Biology*, 47, 921-924
390. Soni, A., Kumar, A., Nath, V. and Niveden, A., (2009). Genetic diversity of Indian liverwort *Plagiochasma appendiculatum* revealed by RAPD marker. *Research Journal of Botany*, 4(3), 89-100
391. Sreenath, A., Kumar, M. A., Anjaneyulu, P. and Boyina, R. P. R. (2020). Three moss families (Bryopsida: Calymperaceae, Hyopterygiaceae, & Pterobryaceae): new distribution records to bryoflora of Andhra Pradesh, India. *Journal of Threatened Taxa*, 12(4), 15481-15488
392. Srivastava, K. P. (1964). Bryophytes of India I. Ricciaceae. *Bulletin of National Botanical Garden, Lucknow*. 104, 1-103
393. Srivastava, S.C. and Alam A. (2005). Family Scapaniaceae, new to south India Bryoflora. Indian*Journal ofForestry*, 28(3), 291-294
394. Srivastava, A., Vinay Sahu and Asthana, A. K. (2018). An observation on morphogenetic response of *Marchantia polymorpha* subsp *ruderalis* Bischl & Boissel.-Dub in different culture media. *International Journal of Plant and Environment*, 4(1), 64-69
395. Stagg, B. C., Donkin, M. E. and Smith, A .M. (2015). Bryophytes for Beginners: The usability of a printed dichotomous key versus a multi-access computer-based key for bryophyte identification. *Journal of Biological Education*, 49(3), 274-287
396. Staples, G. W., Imada, C. T., Hoe, W. J. and Smith, C. W. (2004). A revised checklist of Hawaiian mosses. *Tropical Bryology*, 25, 35–69
397. Stech, M. and Frey, W. (2008). A morpho-molecular classification of the mosses (Bryophyta). *Nova hedwigia*, 86, 1-21
398. Stech, M. and Quandt, D. (2014). 20,000 species and five key markers: the status of molecular bryophyte phylogenetics. *Phytotaxa*, 9(1), 196-228

399. Stech, M., Osman, S., Sim-Sim, M. and Frey, W. (2006). Molecular systematics and biogeography of the liverwort genus *Tylimanthus* (Acrobolbaceae) Studies in austral temperate rain forest bryophytes 33. *Nova Hedwigia*, 83, 17-30
400. Stech, M., Quandt, D., Lindlar, A. and Frahm, J. P. (2003). The systematic position of *Pulchrinodus inflatus* (Pterobryaceae, Bryopsida) based on molecular data. Studies in austral temperate rainforest bryophytes 21. *Australian Systematic Botany*, 16(5), 561-568
401. *Stephani, F. (1906-1924). *Species Hepaticarum* 2-6, Geneva.
402. Stevenson, L. A., Gonzalez-Orozco, C. E., Knerr, N., Cargill, D. C. and Miller, J. T. (2012). Species richness and endemism of Australian bryophytes. *Journal of Bryology*, 34(2), 101-107
403. Suesatpanit, T., Osathanunkul, K., Madesis, P. and Osathanunkul, M. (2017). Should DNA sequence be incorporated with other taxonomical data for routine identifying of plant species?. *BMC complementary and alternative medicine*, 17(1),1-10
404. Sun, S.Q., Wu, Y.H., Wang, G.X., Zhou, J., Yu, D., Bing, H.J. and Luo, J. (2013). Bryophyte species richness and composition along an altitudinal gradient in Gongga Mountain, China. *PloS one*, 8(3), 1-10
405. Suzuki, T., Inoue, Y. and Tsubota, H. (2018). Molecular phylogeny of the genus *Fissidens* (Fissidentaceae, Bryophyta) and a refinement of the infrageneric classification. *Molecular phylogenetics and evolution*, 127, 190-202
406. Taberlet, P., Gielly, L., Pautou, G. and Bouvet, J. (1991). Universal primers for amplification of three non-coding regions of chloroplast DNA. *Plant molecular biology*, 17(5), 1105-1109
407. Takhtajan, A. L. (1953). Phylogenetic principles of the system of higher plants. *The Botanical Review*, 19(1), 1-45
408. Taylor, W. A. (1995). Spores in earliest land plants. *Nature*, 373, 393–392
409. Techen, N., Parveen, I., Pan, Z. and Khan, I.A. (2014). DNA barcoding of medicinal plant material for identification. *Current Opinion in Biotechnology*, 25,103-110
410. Tewari, S. D., Joshi, P., Bhandari, M., Singh, D. and Joshi, K. (2020). An assessment of bryophyte diversity of Nagar Nigam areas at Haldwani in Kumaun region of northwest Himalaya. *Journal of Pharmacognosy and Phytochemistry*, 9(5): 365-372

411. Thamizharasi, T.,Sahaya Sathish, S., Palani, R., Vimala, A. andVijayakanth, P. (2018). Enumeration and phytogeographical pattern of mosses (Bryopsida) in Kalrayan Hills of Eastern Ghats of Tamilnadu, India. *International Journal of Development Research*, 8(1), 18212-18216
412. Thomas, R. and De Franceschi, D. (2013). Palm stem anatomy and computer-aided identification: the Coryphoideae (Arecaceae). *American Journal of Botany*, 100(2), 289-313
413. Thornton, L.E., Keren, N., Ohad, I. and Pakrasi, H. B. (2005). *Physcomitrella patens* and *Ceratodon purpureus*, mosses as model organisms in photosynthesis studies. *Photosynthesis research*, 83(1), 87-96
414. *Tixier, P. (1979). *Bryogeographie du Mont Bokor (Cambodge*). Bryophyt. Biblioth, 18, 1-121
415. Tofilski, A. (2018). DKey software for editing and browsing dichotomous keys. *ZooKeys,* 735, 131-140
416. Turetsky, M. R. (2003). The role of bryophytes in carbon and nitrogen cycling. *The bryologist*, 106(3), 395-409
417. Tyagi, S. N., Joshi, P. M., Oza, B. D. and Dalal, H. N. (2007). *Forest Resource Survey Chhotaudepur Forest Division (Vadodara District) Series I & II. Tree Enumeration,* Forest Department, Gandhinagar
418. Udar, R. and Singh, D. K. (1979b). *Notothylas pfleiderei* sp.nov. from Mangalore, Indian. *Lindbergia*, 5, 28-30
419. Udar, R. and Singh, D. K. (1979a). *Notothylas disseta* a hornwort new to India. *TheBryologist*, 82, 625-628
420. Udar, R. and Srivastava, S. C. (1975). Notes on South Indian HepatocaeI. *Journal of Bomabay Natural History Society*, 12, 401-406
421. Ung, V., Dubus, G., Zaragueta-Bagils, R. andVignes-Lebbe, R. (2010). Xper2: introducing e-taxonomy. *Bioinformatics*, 26(5), 703-704
422. Valente Valente, D., Aguiar Saraiva Câmara, P.E., Peralta, D.F. and Stech, M. (2019). Molecular tools to identify tropical mosses: a case study of the Brazilian species of Schlotheimia Brid. (Bryophyta, Orthotrichaceae). *Systematics and Biodiversity*, 17(6),609-621

423. Valliammal, N. (2014). *Computer aided plant identification through leaf recognition using enhanced image processing and machine learning algorithms*. Ph. D thesis submitted to Avinashilingam Deemed University for Women.

424. Vanderpoorten, A. and Goffinet, B. (2009). *Introduction to bryophytes*. Cambridge University Press. 303 p

425. Vashistha, B. D. (1987). Effect of some auxins and cytokinins on growth and archegonial formation in the liverwort *Riccia frostii* Aust. *Biochemie und Physiologie der Pflanzen.* 182, 309-321

426. Venugopal, M. and Nair, M.C.(2017). Bryophyte diversity of Thamarassery pass (Wayanad pass) in the Western Ghats of Kerala. *Plant Science Today*, 4(1), 41-48

427. Vignes-Lebbe, R., Bouquin, S., Kerner, A. andBourdon, E. (2017). Desktop or remote knowledge base management systems for taxonomic data and identification keys: Xper2 and Xper3*. Biodiversity Information Science and Standards*, 1, 1-3

428. Vitt, D. H., Li, Y. and Belland, R. J. (1995). Patterns of bryophyte diversity in peatlands of continental western Canada. *The Bryologist*, 98(2), 218-227

429. Von Crautlein, M., Korpelainen, H., Pietilainen, M. and Rikkinen, J.(2011). DNA barcoding: a tool for improved taxon identification and detection of species diversity. *Biodiversity and conservation*, 20(2), 373-389

430. Von Konrat, M., Shaw, A. J., and Renzaglia, K. S. (2010). A special issue of Phytotaxa dedicated to Bryophytes: the closest living relatives of early land plants. *Phytotaxa,* 9, 5–10

431. Vorontsova, M. S., Clayton, D. and Simon, B. K. (2015). Grassroots e-floras in the Poaceae: growing GrassBase and GrassWorld. *PhytoKeys*, 48, 73-84

432. Vujicic M., Sabovljevic, A.and Sabovljevic, M. (2009). Axenically culturing the bryophytes: a case study of the moss *Dicranum scoparium* Hedw. (Dicranaceae, Bryophyta). *Botanica Serbica*, 33(2), 137-140

433. Vujicic, M., Sabovljevic, A.and Sabovljevic, M. (2011). Axenically culturing the bryophytes: establishment and propagation of the moss *Hypnum cupressiforme* Hedw. (Bryophyta, Hypnaceae) in In vitro conditions. *Botanica Serbica*, 35(1), 71-77

434. Vujicic, M., Cvetic, T., Sabovljevic, A. and Sabovljevic, M. (2010). Axenically culturing the bryophytes: A case study of the liverwort *Marchantia polymorpha* L. ssp. *ruderalis*

Bischl. & Boisselier (Marchantiophyta, Marchantiaceae). *Kragujevac Journal of Science*, (32), 73-81

435. *Wada, K., Hirabayashi, Y. and Saito, W. (1984). Light germination of *Anthoceros miyabeanus* spores. *The botanical magazine= Shokubutsu-gaku-zasshi*, 97(3), 369-379
436. Wahrmund, U., Rein, T., Müller, K. F., Groth-Malonek, M. and Knoop, V. (2009). Fifty mosses on five trees: comparing phylogenetic information in three types of non-coding mitochondrial DNA and two chloroplast loci. *Plant systematics and evolution*, 282(3-4), 241-255
437. Waldchen, J. andMader, P. (2018). Plant species identification using computer vision techniques: A systematic literature review. *Archives of Computational Methods in Engineerin*g, 25(2), 507-543
438. Waldchen J. Rzanny M., Seeland M.and Mader P. (2018). Automated plant species identification. *PLOS Computional Biology*, 14(4), 1-19
439. Wang, D. H., Wang Y. F., Zuo, Q., Li, M., WU, W. Y., Huang, J. H. and Zhao, M. S. (2012). Bryophyte species diversity in seven typical forests of the West Tianmu Mountain in Zhejiang, China. *Chinese Journal of Plant Ecology*, 36(6), 550-559
440. Wang, W., Wu, Y., Yan, Y., Ermakova, M., Kerstetter, R. and Messing, J. (2010). DNA barcoding of the Lemnaceae, a family of aquatic monocots. *BMC Plant Biology*, 10(1), 1-11.
441. Wardlaw, C. W. (1955). *Embryogenesis in plants.*Metheun & Co., London.381 p.
442. Watson, L. and Dallwitz, M. J. (1992a). *The grass genera of the world*. C.A.B. International Wallingford, Oxfordshire, UK. 1038 p
443. Watson, L. and Dallwitz, M. J. (1992b). The families of flowering plants: descriptions, illustrations, identification, and information retrieval. Version: 1st March 2021.https://www.delta-intkey.com.Accessed on 22nd February 2020
444. Watson, L. and Dallwitz, M. J. (2008). *The families of gymnosperms*. Version: 5th August 2019. www delta-intkey.com. Accessed on 22nd February 2020
445. Watson, L., Macfarlane, T. D. and Dallwitz, M. J. (1992). *The grass genera of the world: descriptions, illustrations, identification, and information retrieval; including synonyms, morphology, anatomy, physiology, phytochemistry, cytology, classification, pathogens,*

world and local distribution, and references. Version: 15th December 2020. www.delta-intkey.com Accessed on 24th February 2020

446. Weis Gabriel, E., Wamper, D. and Santarius, K. A. (1986). Heat sensitivity and thermal adaptation of photosynthesis in liverwort thalli. *Oecologia*, 69, 134-139
447. Wen-Zhang, M., Wen-Yao, L. and Xing -Yang, L. (2009). Species composition and life forms of epiphytic bryophytes in old–growth and secondary forests in Mt. Ailao, SW China. *CryptogamieBryologie*, 30, 477-500
448. *Wijk, R., Van-Der, W. D. M. and Florschutz, P. A. (1959-67). *Index muscorum*, Utrech, 922 p
449. Wildlife Institute of India. (2019). National Wildlife Database. Ministry of Environment, Forest and climate change, Government of India. <https://wii.gov.in/national_wildlife_database. Accessed on 10thNovember 2019
450. Williams, P. G., Appavoo, M. R., Aravind, A. and Prakash, G. (2016). Antibacterial activity and phytochemical analysis of selected bryophytes against penicillin resistant bacterial pathogens. *South Indian Journal of Biological Sciences*, 2, 347-353
451. Woese, C. R. and Fox, G. E. (1977). Phylogenetic structure of the prokaryotic domain: the primary kingdoms. *Proceedings of the National Academy of Sciences*, 74(11), 5088-5090
452. Wolf, J. H. (1993). Diversity patterns and biomass of epiphytic bryophytes and lichens along an altitudinal gradient in the northern Andes. *Annals of the Missouri botanical garden*, 80(4), 928-960
453. World Conservation Monitoring Centre (WCMC). (1992) *Global Biodiversity: Status of the Earths Living Resources*. Chapman & Hall, London, 594p
454. Wu, S. G., Bao, F. S., Xu, E. Y., Wang, Y. X., Chang, Y. F. and Xiang, Q. L. (2007).A leaf recognition algorithm for plant classification using probabilistic neural network. *In 2007 IEEE international symposium on signal processing and information technology*, 11-16
455. Yang, H. P., Ma, C. S., Wen, H., Zhan, Q. B. andWang, X. L. (2015). A tool for developing an automatic insect identification system based on wing outlines. Scientific reports, 5(1), 1-11

456. Zander, R. (1993).*Genera of the Pottiaceae: Mosses of Harsh Environments*, Bulletin of the Baffalo Society of Natural Sciences, 32, 378 p
457. Zavala, A. M., Pérez, N. M., Becerra, A. A. and Lopez, M. A. V. (2012). Glucose inhibits the shoot bud formation in the moss *Bryum billarderi*. *Central European Journal of Biology*, 7(4), 648-654
458. Zeltyn, S. A. and Pchelkin, A. V. (1993) *Application of a computerized lichen identifier in background ecological monitoring systems*. In: Problems of Ecological Monitoring and Ecosystem Modelling, Gidrometeoizdat Publisher, St. Petersburg. 15, 264-273 pp.
459. Zuquim, G, Tuomisto, H. and Prado J. (2017) A free-access online key to identify Amazonian ferns. *PhytoKeys*, 78, 1-15

Note: References Marked with Asterik (*) were not seen.