LIST OF FIGURES

Figure No.	Figure Name	Page No.
1.1	Salient features of bryophytes	3
1.2	Morphological diversity of different groups of bryophytes	4
2.1	Figure showing the locations of Gujarat state visited by the Researchers in past study	19
2.2	Figure showing the locations of study area state visited by the Researchers in past study	20
2.3	Important components of Moss tissue culture experiments	22
3.1	Outline Map of Study area	43
3.2	Geology Map of Gujarat	44
3.3	River Map of Gujarat	46
3.4	Soil Map of Gujarat	47
4.1	Major objectives of the current study	53
4.2	Outline methodology of diversity and distribution study	54
4.4	Outline methodology of Culturing of bryophytes	61
4.5	Outline methodology of barcoding of bryophytes	64
4.6	Outline methodology of Computer aided identification study	67
4.7	Dialog box for entering the Taxon characters	70
4.8	Method of running the confor files for final ouput	71
5.1	The distribution of the locations across the study area.	72
5.2	Map showing the distribution of families in different districts of the study area	135
5.3	Distribution of groups in different district	137
5.4	Elevation distribution map of *Anthoceros* spp and *Notothylas* spp	141
5.5	Elevation distribution map of *Asterella angustata* and *Plagiochasma* spp	141
5.6	Elevation distribution map of *Cyathodium* spp, *Targiona hypophylla* and *Rectolejunea aloba*	142
5.7	Elevation distribution map of *Riccia* spp	142
5.8	Elevation distribution map of *Brachymenium turgidum, Bryum* spp. and *Ptychostomum capillare*	143
5.9	Elevation distribution map of *Fissidens* spp	143

Figure No.	Figure Name	Page No.
5.10	Elevation distribution map of *Funaria hygrometrica*, *Philonotis* spp and *Physcomitrium eurystomum*.	144
5.11	Elevation distribution map of *Gymnostomiella vernicosa*	144
5.12	Elevation distribution map of *Hydrogonium* spp, *Hyophila* spp. and *Semibarbula orientalis*	145
5.13	Elevation distribution map of *Stereophyllum* spp	145
5.16	Evolutionary analysis by Maximum Likelihood method of *Fissidens zollingeri* with different members of Fissidentaceae	158
5.17	Evolutionary analysis by Maximum Likelihood method of of *Bryum coronatum* with different members of Bryaceae	160
5.18	Evolutionary analysis by Maximum Likelihood method of different member of Ayotoniaceae	162
5.19	The INTKY Window Panel in Normal Mode	167
5.20	Subset of characters considered for the identification of sporophyte possessing members of Bryaceae	168
5.21	Difference in characters between the moss species *Bryum coronatum* and *Stereophyllum tavoyense*	169
5.22	Liverwort *Riccia discolor* identified using DELTA INTKEY	170
5.23	Hornwort *Notothylas khasiana* identified using DELTA INTKEY	171
5.24	Moss *Bryum coronatum* identified using DELTA INTKEY	171