

REFERENCES

REFERENCES

1. Abdugamatov, A., Ubaev, Kh. and S.Yu Yunusov, 1968. The alkaloids of Pedicularis olgae. Khim. Prir. Soedin (Tashk.) 4 (2) : 136-137.
2. Affifi-Yazar; Fatma, U. and O. Sticher 1980. Veproside, a new iridoid glycoside from Veronica officinalis. Helv. Chim. Acta 63 (7) : 1905-1907.
3. Affifi-Yazer; Fatma, U., Sticher, O., Uesato, S., Nagajima, K and H. Inouye. 1981. Ladroside, a new iridoid glycoside from Varonica officinalis. Helv. Chim. Acta 64 (1):16-24.
4. Agababyan, E.Yu., Arutyunyan, L.S., Minatsakanyan, V.A., Gach-Baitts, E. and L. Radich. 1982. Iridoids from Verbascum georgicum. Khim. Prir. Soedin (Tashk.) 0 (4): 446-451.
5. Agardh, J.G. 1858. Theoiria systematis plantarum; accedit familiarum phanerogamarum. Lund.
6. Agarwal, S.C., Bhaskar, A. and T.R. Seshadri 1973. Chemical compounds of the roots of Didymocarpus pedicellata Isolation and structure of pashanone, a new chalcone. Indian J. Chem. 11(1) : 9-12.
7. Agarwal, S.G., Thappa, R.K., Vashist, V.N., Atal C.K. and R. Gupta 1975. Chemical examination of the volatile oil of Limnophila rugosa (Roth) Merrill. Indian J. Pharm 37 (4): 99-100.

8. Agarwal, S.K. and R.P. Rastogi 1974. Celsiogenin-C, a new genin from celsiosides isolated from Celsia coromandeliana Indian J. Chem 12 (9) : 907-910.
9. Ahmad, K.J. 1974a. Cuticular studies on some Nelsonioideae (Acanthaceae) Bot. J.Linn.Soc. 68: 73-80.
10. Ahmed, K.J. 1974b. Cuticular studies in some species of Mendoncia and Thunbergia (Acanthaceae) Bot.J.Kinn.Soc. 69 : 53-63.
11. Akhmedov, S.G. and V.I. Litvinenko, 1968. Garashangin : New phenol compound from Scrophularia grossheimii Dokl. Akad. Nauk. Azerb. Ser. 24(9) : 89-92.
12. Akhmedov, S.G. and V.I. Litvinenko, 1969. Scrophulein : A new flavonoid from Scophularia grossheimii. Khim.Prir. Soedin (Tashk.) 5 (1): 65-55
13. Andary, C., Wylde, R., Laffite, C. Privat.G. and F. Winternitz 1982. Structure of Verbascoside and orobanchoside, caffeic acid, sugar esters from Orobanche rapum genistae Phytochemistry 21 (5): 1123-1128.
14. Arati, D. and P.C.Datta 1975. Floral vasculature of common Indian Scrophulariaceae. Acta Soc. Bot. Polon. 44:57-58.
15. Arati, D. and P.C. Datta 1977. Interrelations of Scrophulariaceae based on secondary xylem of common Indian species J. Indian Bot. Soc. 56 : 149-164.
16. Arekal, D., Rajeshwari,S. and S.N.R. Swamy 1971. Contribution to the embryology of Scoparia dulcis L. Bot. Notiser. 124 (2): 237-248.

17. Arutyunyan, L.S., Agababyan, E.Yu. and V.A.Minatsakanyan 1983. Verbascoside, B : A minor iridoid from Verbascum georgicum Khim Prir.Soedin (Tashk.) Q (5): 478-518.
18. Bachmann, E.T. 1882. Darstellung der Entwicklungs geschichte und des Baues der Samenschalen der Scrophularineen. Verhandl. K. Leop-Carol. Dent. Akad. Naturforscher 43 : 1-180.
19. Baillon, H. 1888. Histoire des Plantes IX. Libraire Hachette and Cie. Paris.
- *20. Barloy, J. 1963. Annali Physio-Veget. 5, 41.
21. Bate-Smith, E.C. 1972. Chemistry and Phylogeny of the Angiosperms. Nature 236 : 353-354.
22. Bate-Smith, E.C. and T.Swain 1966. The asperulosides and the aucubins - in Comparative Phytochemistry (Swain, T. Ed.) Academic Press. London, 159-174.
23. Baudouin, G. and R. Paris. 1975. Polyphenols of Digitalis V : The polyphenols of the flowers of the Canary Island foxglove. Plant Med. Phytoter. 9(4): 278-288.
24. Bellini, R. 1907. Criteri per una nuova classificazione della Personatae (Scrophulariaceae et Rhinanthaceae) Annali di Botanica 6 : 131-145.
25. Bendall, M.R., Clive, W.F. and D.M.Thomas 1979. The structure of procumbide. Aust.J.Chem. 32(9) : 2085-2092.
26. Benson, L. 1957. Plant classification, Boston.
27. Bentham, G. 1835. Scrophularineae Indicae. A synopsis of the East Indian Scrophulariaceae. London.

28. Bentham, G. 1846. Scrophulariaceae. In de Candolle. A.Prod.Syst.Nat.Regn.Veg.X Victoris Masson Paris.
29. Bentham, G. and J.D.Hooker. 1876. Scrophulariaceae, In Genera Plantarum 2 : 913-980. London.
30. Bentham, G. and J.D. Hooker 1862-1883. Genera Plantarum 3 Vols. London.
31. Berg, T., Soren, D., Soren, R., Jenson, B., Juhl, N. and L.F. Rickelt. 1985. Iridoid glucosides from Pedicularis. Phytochemistry 24(3) : 491-494.
32. Bernhard, H., Ivan, K., Siegfried, J., Detlef, G., Manfred, H. and H.Schmid 1973. Chaenorhine, a macrocyclic spermine alkaloid ; 149. Report on alkaloids Helv.Chim.Acta 56(4) 1266-1303.
33. Bessey, C.E. 1915. The phylogenetic taxonomy of flowering plants. Ann.Missouri Bot.Gard. 2 : 109-164.
34. Bhaskar, A. and T.R.Seshadri. 1973. New Chemical Components of Didymocarpus pedicellata leaves, Indian J.Chem. 11(4) : 404-405.
35. Bianco, A., Guiso, M., Iavarone, C., Passacantilli, P. and C.Trogolo. 1980. 6-O- β -xylopyranosyl aucubin from Verbascum sinuatum Phytochemistry 19(4) : 571-574.
36. Bianco, A., Francesconi, A. and P.Passacantilli. 1981. Iridoid glucosides of Odontites verna ssp serotina Phytochemistry 20(6) : 1421-1422.
37. Bianco, A., Guiso, M., Iavarone, C., Passacantilli, P. and C.Trogolo. 1981a. Iridoids : 29, Sinuatol from Verbascum

- sinuatum. Planta Med. 41(1) : 75-79.
38. Bianco, A., Daniela, B. and P.Passacantilli. 1982.
6-O- β glucopyranosyl aucubin, a new iridoid from
Odontites verna ssp serotina. Planta Med. 44(2): 97-99.
39. Bianco, A., Passacantilli, P. and P.Giovanna. 1982a.
Epilogamic acid and 7- β -hydroxy-8-epi-iridoidal
glucoside : 2. New iridoid glucosides from Linaria
cymbalaria. Planta Med. 46(1): 38-41.
40. Boeshore, I. 1920. The morphological continuity of the
Scrophulariaceae and Orobanchaceae, Contrib.Bot.Lab. Univ.
Penn. 5 : 139-176.
41. Boivin, B. 1956. Les families de Tracheophytes. Bull.
Soc.Bot.France 103 : 490-505.
42. Borodin, L.I., Litvinenko, V.I. and N.V.Kurinnaya 1970.
Lignoside, A new flavonoid glycoside of Gratiola
officinalis. Farm Z.H. 25(2) : 62-67.
43. Borodin, L.I., Litvienenko, V.I. and N.V.Kurinnaya. 1970a.
New flavonoid c-glycosides of Gratiola officinalis
Khim.Prir.Soedin (Tashk.) 6(1): 19-24.
- *44. Bracke, M. 1924. Bull.Soc.Chim.Biol. 6, 665.
45. Bremekamp, C.E.B. 1953. The delimitation of the
Acanthaceae. K.Nederl.Akad.Wetenshap.Proc.Ser.C. 56 :
533-546.
46. Brew, E.J.C. and R.H. Thomson. 1971. Naturally occurring
quinones XX : Anthraquinones in Digitalis purpurea.
J.Chem.Soc.Sect.C.Org.Chem. 10 : 2007-2010.

47. Brooker, S.E. and K.J. Harkiss. 1974. Investigation of the occurrence of alkaloids in Antirrhinum molle, A.mollissimum and A.hispanicum. Planta Med. 26(4) : 305-310.
48. Bruhl, P. 1920. Notes on Lindenbergia urticifolia Lehm. and L.polyantha Royle. Ibid 17-30.
49. Burnett, A.R. and R.H. Thomson. 1968. Anthraquinones in two Digitalis species. Phytochemistry & 7(8) : 1423.
50. Burnett, G.T. 1835. Outlines of Botany etc. 2 Vols. London.
51. Burt, B.L. 1962. Studies on the Gesneriaceae of the Old World XXIV. Tentative keys to the tribes and genera. Notes Roy. Bot.Gard.Edinb. 24 : 205-220.
52. Campbell, D.H. 1930. The relationship of Paulownia Bull.Torr.Bot.Club. 57 : 47-50.
53. Caruel, T. 1881. System novum regni vegetabilis. Nuovo Giorn Bot. Ital. 13 : 217-228.
54. Chaubal, P.D. 1966. Palynological studies on the family Acanthaceae Ph.D. Thesis, Poona University.
55. Choisy, J.D. 1848. Selagineae. In de Candolle. A Prod. Syst.Nat.Regn. Veg. XII Victoris. Masson, Paris.
56. Crete ,P. 1955. L' application de certaines donnees embryologiques a la systematique des Orobanchacees et de quelques familles voisines. Phytomorphology : 5: 422-435.
57. Crete , P. 1959. Precis de botanique : Vol.2, Systematique des Angiosperms.

58. Cronquist, A. 1968. The evolution and classification of flowering plants. Houghtan Mifflin Company, Boston.
59. Cronquist, A. 1981. An integrated system of classification of flowering plants. Columbia University Press. New York.
60. Cronquist, A. 1987. A botanical critique of cladison Bot. Rev. 53 : 1-52.
- *61. Dahlgren and Rao 1971. In Systematic palynology of tribe Leucophylleae (Scrophulariaceae) and selected Myoporaceae by Neizgoda, C.J. and A.S. Tomb (1975). Pollen ET. Spores 17 (4): 495-513.
62. Dahlgren, R., Jensen, S.R. and B.E. Nielsen 1981. A revised classification of the Angiosperms with comments on correlation between chemical and other characters. Phytochemistry and Angiosperm Phylogeny, Young, D.A. and S. Seigler (Ed.) 149-204.
63. Daniel, M. and S.D. Sabnis, 1979. Chemotaxonomy of Loganiaceae Curr.Sci. 48 : 383-385.
64. Daniel, M. and S.D. Sabnis 1987. The chemical phylogeny of the order Gentianales. In Phytochemistry and Taxonomy (Bilgrami, K.S. and G.V.V. Dogra Ed.) Print House (India), Lucknow.
65. Daniel, M and S.D. Sabnis, 1987a Chemosystematics of some Indian members of the Acanthaceae. Proc. Indian Acad. Sci. (Plant Sci). 97 (4): 315-323.
66. Das, V.S.R., Rao K.N. and G.V.S. Rao 1966. Phenolic acid in some members of Pedaliaceae. Curr.Sci. 35 (6): 160.

67. De Eds, F. 1968. In Comprehensive Biochemistry (Florkin, M and E.H. Stotz, Ed.) Elsevier Publishing Co., Amsterdam.
68. Degot, A.V., Litvinenko, V.I and P.I. Kovalev 1971. Iridoids from Odontites serotina (Lam) Dum. Rast.Resur. 7 (3): 390-396.
69. Degot, A.V.; Litvinenko, V.I and N.V. Kurinnaya 1971a. Flavonoids from Ortantha lutea. Khim Priir.Soedin (Tashk.) 7 (1) : 119.
70. Den Ouder, R.W. and W.L.H. Van Veenendaal 1983. Wood anatomy of Uncarina leandrii (Pedaliaceae) and its relation to Bignoniaceae, Iawa Bull 4(1): 53-59.
71. De Pascual, T. Diaz, J.F. and M. Grande, 1980. Components of Verbascum thapsus L. 3. Contribution to the study of saponins. An. Quim Ser C Quim Org Bio Quim, 76 (2): 107-110.
72. Diaz, M.D. 1977. Flavonoids and morphological studies of Lindernia Allion. (Scrophulariaceae) in South America. Bot. Jour. Linn. Soc. 75 : 47-67.
73. Diffluvia, T. 1979. The endosperm and embryo in the system of Tubiflorae with special reference to Boraginaceae and Hydrophyllaceae. Kurlziana 12/13 (0): 101-112.
74. Dighe, S.S. and A.B. Kulkarni 1973. Chemical investigation of Aegnetia indica Linn. Indian J. Chem. 11(4): 404.
75. D'ohot, A.V., Lytvynenko, V.I., Chernykh, N.O. and I.H. Zoz, 1972. Iridoids in the Scrophulariaceae family, Farm.Z.H. 27 (1): 66-73.
76. Dominguez and A. Xorge 1970. Chemical components of Leucophyllum texanum, Planta Med. 17(4): 366-368.

77. Don, G. 1831-1838. A general history of the dichlamydeous plants etc. London.
78. Drude, O. 1887. Die naturliche systematische Anordnung der Blütenpflanzen. Sber.Abhandl.naturwiss Ges Isis in Dresden, Sber 4, 7 : 75-84.
79. Dutta, N.M. and S. Chanda. 1979. A contribution to the taxonomy and palynology of Gratioleae of eastern India Trans.Rose Res.Inst. (Calcutta) 42(1/2) : 1-34.
80. Eichenberger, W. and M. Wilhelm. 1966. Sterols in leaves and Chloroplasts. Z.Naturforsch 21b(9) : 859-867.
81. Elisens, W.J. 1986. Pollenmorphology and Systematic relationships among New World species in Tribe Antirrhineae (Scrophulariaceae) Amer.J.Bot. 73(9) : 1298-1311.
82. Elisens, W.J. and A.S. Tomb. 1983. Seed Morphology in New World Antirrhineae, Systematic and Phylogenetic implications. Plant Syst.Evol. 142(1/2): 23-48.
83. Endlicher, S.L. 1836-40. Genera Plantarum Secundum ordines naturales disposita. Vindobonae.
84. Endlicher, S.L. 1841. Enchiridion botanicum exhibens classes et ordines Plantarum. etc. Lipsiae.
85. Engler, A. and K.Prantl. 1897-1899. Die natürlichen Pflanzen familien Leipzig.
86. Engler, A. and L.Diels. 1914. Scrophulariaceae. Ibid Nachfrage 4 Zu 4(36) : 273-279 Leipzig.
87. Erdtman, G. 1952. Pollen morphology and Plant taxonomy Angiosperms, Almquist and Wiksell, Stockholm.

88. Euw, J.V. and T. Reichstein 1966. The biosynthesis of cardenolides and spirostanol glycoside in Digitalis lanata Ehrh. Comparison of the formation ratio : Helv.Chim.Acta. 49 (5): 1468-1474.
89. Evans, F.G. 1973. Steryl esters of Digitalis purpurea L. herb. J. Pharm.Pharmacol. 25 (2): 156-163.
90. Falsone, G., Laryeae, M.D., Crea A.E.G. and E. Finner, 1982. Iridoids from Verbascum sinuatum. Planta Med. 44(3):150-153.
91. Feitag, H., Sigrid S., Horst H.A.L and K. Meyer 1967. Glycosides in the leaves of Isoplexis isabelliana (Webb) Masf. Helv. Chim. Acta 50 (5) : 1336-1365.
92. Fernald, M.L., 1960 Gray's Manual of Botany 8th ed. American Book Co., New York.
93. Flyr. 1970. In Systematic palynology of tribe Leucophylleae (Scrophulariaceae) and selected Myoporaceae by Neizgoda, C.J. and A.S. Tomb 1975. Pollen ET. Spores 17(4), 495-513.
94. Freire, R., Gonzalez, A.G. and E. Suarez. 1970. New sources of steroid sapogenins VII. Sceptrumgenin and isoplexigenin A, B, C, D from Isoplexis sceptrum. Tetrahedron 26(13):3233-3244.
95. Fruchier, A., Jean-Pierre, R., Claude, A and G. Privat. 1981. A tropone derivative from Orobanche rapum genistae Phytochemistry 20 (4): 777-780.
96. Gates, F.C. 1940. Annotated List of the plants of Kansas: Ferns and Flowering plants (C.f. Gibbs, R.D. Vol.IV)
97. Gentry A.H. and A.S.Tomb 1979. Taxonomic implications of Bignoniaceae palynology. Ann. Missouri Bot. Gard. 66 : 756-777.

98. Gentry, A.H. 1980. Bignoniaceae Part I (Crescentieae and Tourrettieae) Fl. Neotropica Monogr. 25.
99. Ghosh, A., and L.B. Jack, 1979. Seed Lipid constituents of species of Proboscides, J. Nat. Prod. (Lloydia) 42 (3): 287-292.
100. Gibbs, R.D. 1974. Chemotaxonomy of flowering plants. McGill Queens Univ. Press, Montreal.
101. Giessman, T.A. (ed.) 1962. "Chemistry of the flavonoid compounds" Pergamon Press, Oxford.
102. Gilbert, R.I. 1973. Chalcone glycosides of Antirrhinum majus Phytochemistry 12(4): 809-810.
103. Gilbert, R.I. 1975. Aurone glycosides of Antirrhinum orontium Phytochemistry 14(5/6): 1461.
104. Gill, S., Jozef, K., and H. Mionskowski 1969. Identification of luteolin 7-glucoside of Digitalis ambigua Murray. Diss Pharm. Pharmacol 21 (5): 413-415.
105. Gleason, H.A. 1952. The New Britton and Brown Illustrated Flora of the North Eastern United States and Adjacent Canada. III Lancaster Press, Inc. Lancaster, Pa.
106. Glisic, L.M. 1929. "Uber die Endosperm Und Haustorien bildung bei Orobanchae hederae Duby und Orobanchae gracilis Sm. Bull. Inst. Jard. Bot. Univ. Beograd 1:106-141.
107. Gonzalez, A.G., Barrera, R.F., Rocio, J.A.S and E.S. Lopez 1972. New natural sources of sapogenins XII. Funchaligenin, new steroidal sapogenin from Isoplexis sceptrum (Linn.) Steudel. An. Quim 68(7/8):1021-1027.

108. Grisebach, A.H.R. 1854. Grundriss der systematischen Botanik für akademische Vorlesungen. Göttingen.
109. Gundersen, A. 1950. Families of Dicotyledons. Waltham, Mass.
110. Hallier, H. 1903. Über die Abgrenzung und Verwandtschaft der einzelnen Sippen bei den Scrophulariaceae. Bull. Herb. Boiss Ser 2, 3 : 181-207.
111. Hallier, H. 1905. Provisional scheme of the natural (Phylogenetic) system of flowering plants. New Phytol. 4 : 141-162.
112. Hallier, H. 1912. L' Origine et le système phylétique des angiospermes exposés à l'aide de leur arbre généalogique. Arch. Néerl. Sci. Exactes et Nat. Ser III B, 1 : 146-234.
113. Harborne, J.B. 1959. Chromat. Rev. 2: 108.
114. Harborne, J.B. 1966. The evolution of flavonoid pigments in plants. In Comparative phytochemistry (Swain T. Ed) Academic Press. London and New York.
115. Harborne, J.B. 1967. Comparative Biochemistry of Flavonoids. Academic Press, London.
116. Harborne, J.B. 1971. Identification of Scutellarein 4'-methylether in Linaria aeruginea. Phytochemistry. 10(11): 2850-2851.
117. Harborne, J.B. 1977. Flavonoids and the evolution of the angiosperms. Biochem. Syst. Ecol. 5 : 7-22.
118. Harborne, J.B. 1984. Phytochemical methods 2nd edn. Chapman and Hall, London.
119. Harborne, J.B. and C.A. Williams. 1971. 6-Hydroxy luteolin and scutellarein as phyletic markers in higher plants .

- Phytochemistry 10 : 367-368.
120. Harborne, J.B. and E. Hall. 1964. Plant Polyphenols-XII: The occurrence of tricin and glycoflavone in grasses. Phytochemistry 3 : 421.
121. Harborne, J.B. and T. Swain. 1979. Flavonoids of the Solanaceae, in The Biology and Taxonomy of the Solanaceae. (Hawkes, J.G., Lester, R.N. and A.D. Skelding, Ed.) Academic Press. London. 257-268.
122. Harkiss, K.J. 1971. Studies in the Scrophulariaceae. III. Investigation of the occurrence of alkaloids in Antirrhinum majus L. Planta Med. 20(2) : 108-113.
123. Harkiss, K.J. 1972. Studies in the Scrophulariaceae VI. Quantitative determination of choline in Antirrhinum majus L. A. orontium, L. and Linaria vulgaris Mill. Planta Med. 21(4) : 353-357.
124. Hariss, K.J. and P. Timmins. 1973. Phytochemical investigation of Euphrasia officinalis, Planta Med. 23(4) : 342-347.
125. Hartl, D. 1957. Die Stellung von Lindenbergia Lehmann in System der Scrophulariaceen. Beitr. Biol. Pflanzen 33 : 265-277.
126. Hartl, D. 1959. Das alveolierte Endosperm bei Scrophulariaceen, Seine Entstehung, Anatomie und taxonomische Bedeutung. Beitr. Biol. Pflanzen 35 : 95-110.
127. Hegnauer, R. 1964-73. Chemotaxonomic der Pflanzen. Birkhouser. Basel and Stuttgart.

128. Hegnauer, R. 1966. Aucubinartige Glucoside Uber die Verbreitung and Bedeulung als Systematisches Merkmal. Pharm.Acta Helv. 41 : 557-567.
129. Hegnauer, R. 1969. Chemical evidence for the classification of some plant taxa. In Perspectives in Phytochemistry (Swain, T. and J.B. Harborne, Eds.) 1; 121-138.
130. Hegnauer, R. 1971. Pflanzan steffe und Pflanzen Systematik. Naturwiss 58 : 585-598.
- *131. Hein, S. 1959. Planta Med. 7, 185.
132. Heinricher, E. 1897-1902. Die grunnen Halbuschmaretzer. Jahrbucher f. Wissenschaftliche Botanik.
133. Heirmann, A., Kartnig, T., Seligmann, O, and H.Wagner. 1977. Flavonoids in the leaves of Digitalis lanata Planta Med. 32(1) : 24-26.
134. Hooker, J.D. 1884. Scrophulariaceae. The flora of the British India (Reeve & Co: Kent) 4 : 279-286.
135. Hungund, B.L. and C.H. Pathak. 1971. U.S.D.A. Forest Service Research Paper NE - 201.
136. Hutchinson, J. 1926. The families of flowering plants I. Dicotyledons Macmillan and Co. Ltd. London.
137. Hutchinson, J. 1948. British Flowering Plants. Academic Press. London.374.
138. Hutchinson, J. 1959. The families of flowering plants. 2nd ed. 2 Vols. Oxford University Press. New York.
139. Hutchinson, J. 1969. Evolution and phylogeny of flowering plants. Academic Press. London.

- *140. Ibrahim, R.K. and G.H.N.Towers. 1960. Arch.Biochem. and Biophys. 87 : 125.
141. Imre, S. 1972. Flavone and anthraquinone pigments of Digitalis species III. Further anthraquinone and flavone compounds from the leaves of Digitalis viridiflora. Planta Med. 21(3) : 274-278.
142. Imre, S. and L.Ersoy, 1973. About flavone and anthraquinone pigments of Digitalis : V. The structure of digiterrol and digiferruginol. Z.Naturforsch.Teil.C.Biochem.Biophys.Biol. Virol. 28(7/8) : 471-472.
143. Imre, S., Sezen, S. and R.H.Thomsan. 1976. Anthraquinones in Digitalis species. Phytochemistry 15(2) : 317-320.
144. Imre, Z. and O.Ersoy. 1977. Cardiac glycosides from the leaves of Digitalis schischkinii (Ivan) Werner. Arch.Pharm. (weenheim) 310(2) : 142-151.
145. Imre, Z., Ersoy, O. and T.Yurdan. 1982. Cardiac glycoside composition in the leaves of Digitalis schischkimii. Planta 45(4) : 203-206.
146. Inamdar, J.A. 1969. Epidermal Structure and Stomatal Ontogeny in Vegetative and Floral organs of Martynia annua L., Pedaliium murex L. and Sesamum indicum L. Flora Abt. B, Bd. 158 : 526-537.
147. Inouye, H. and Y. Aoki. 1968. Constituents of Veronicastrum sibiricum Pennell var japonicum Hara. Phytochemistry 7(9) : 1709.
148. Inoue, K., Shinichi, U., Hidekazu, N. and H. Inouye 1983. Quinones of Streptocarpus dunii. Phytochemistry : 22 (3) : 737-742.

149. Jensen, S.R., Nielsen, B.J. and R.D. Dahlgren. 1975. Iridoid compounds, their occurrence and systematic importance in Angiosperms. Bot. Notiser. 128: 148-180.
- *150. Jirawongse, V.A. 1964. A chemotaxonomic study of the Scrophulariaceae. Thesis Purdue University.
151. Job, K., 1969. The Biology of Parasitic Flowering Plants. University of California Press. Berkeley and Los Angeles.
152. Johne, S. and D. Groeger 1968. Alkaloids in Linaria species. Pharmazie 23 (1) : 35-36.
153. Johri, B.M. and H. Singh 1959. The morphology, embryology and systematic position of Elytraria acaulis (Linn.f.) Lindau Bot. Nat. 112 : 227-251.
154. Junior, P. 1982. Barbatoside, a new iridoid glucoside from Penstemon barbatus Planta Med. 45 (2): 127-128.
155. Junior, P. 1983. Nemoroside and nemorososide, new iridoid glucoside from Penstemon nemorosus Planta Med. 47 (2): 67-70.
156. Junior, P. 1983a. 8-Epivalerosidate, a new iridoid glucoside from Penstemon serrulatus Menz. Planta Med. 47 (3) : 161-163.
157. Jussieu, A.L. de. 1791. Genera Plantarum. Paris. 132.
158. Kamienski, F. 1897. Lentibulariace. In Die Naturlichen Pflanzen families (Engler, A and K. Prantl, Ed.) 4, Abt. 3b: 108-123. Leipzig.
159. Karimova, S.G. 1974. Biologically active substances of some species of Pedicularis L. of north eastern Khangay Rastit Resur. 10 (3): 368-371.

160. Kawai Ken Ichi and S. Shibata 1978. Pseudojujubogenin, a new sapogenin from Bacopa monniera Phytochemistry 17 (2): 287-290.
161. Kemertelidze, E.P. 1973. Cardenolides of Digitalis ciliata Khim. Prir. Soedin (Tashk.) 9 (4): 563-564.
162. Khan, M. Israr and K.A. Zirvi 1971. Chemical constituents of Orobanche aegyptica. Pak. J. Sci.Ind.Res. 14 (5):345.
163. KhakimdzhanoV,S., Abdusamatov,A. and S.Yu. Yunusov.1973. The structure of pedicularidine. Khim. Prir. Soedin. (Tashk.) 9 (1): 132-133.
164. Kikuchi, T., Satoko, M., Yoko,K. and T.Namba 1983. New Iridoid glycosides from Harpagophytum procumbens DC. Chem. Pharm. Bull (Tokyo) 31 (7): 2296-2301.
165. Kitagawa,I., Toshiyuki, K., Yonetani, K. and M.Yoshihara 1980. Constituents of Linaria japonica. 4: Chemical correlation of cis-clerodane diterpene with trans-clerodane diterpene. Chem.Pharm.Bull (Tokyo) 28(8): 2403-2413.
166. Kobayashi, H and J. Komatsu 1983. The constituents of Cistanche herba : 1 Yakugaku Zasshi 103 (5):508-511.
167. Konishi Tenji and Junzo Shoji 1981. Constituents of Biochniakia rossica (Chem. et.Schltdl) Fedtsch.etc. Chem. Pharm.Bull (Tokyo) 29 (10) : 2807-2815.
168. Kooiman,P. 1970. The occurrence of iridoid glycosides in the Scrophulariaceae. Acta Bot. Neerl. 19 (3):329-340.
169. KostECKO M.O. and D. Janina 1968. The search for aucubin containing plants. Herba Pol. 14 (4): 270-274.

170. Krishnaswamy, N.R., Seshadri, T.R. and P.J. Tahir 1970. Flavone glucosides of Nepeta Hindustana and Sesamum indicum. Nepetrin and Pedalin and their aglucones. Indian J. Chem. 8 (12): 1074-1078.
- *171 Krolikowska, M. 1967. Rocz. Chem. 41, 529.
172. Krueger Dagmar, Peter Junior and M. Wichtl. 1983. New cardiac glycosides from Digitalis lanata Ehrh. Planta. Med. 49 (2): 74-78.
173. Kubitzki, R. 1969. Chemosystematische Betrachtungen zur Grossgliederung der Dicotylen. Taxon. 18 : 360-368.
174. Kulkarni, A.R. 1968. Contribution to the embryology of Pedaliaceae I. Embryo development in Sesamum lacinatum Klein. J. Shivaji Univ. (Sci) 1 (½) : 114-117.
175. Kulshreshtha, D.K. and R.P. Rastogi 1973. Bacogenin A₁: A novel dammarane triterpene sapogenin from Bacopa monniera Phytochemistry 12 (4): 887-892.
176. Kulshreshtha, D.K. and R.P. Rastogi. 1974. Bacogenin A₂: A novel sapogenin from bacosides. Phytochemistry 13 (7): 1205-1206.
177. Kumar, S. and G.S. Paliwal 1982. Foliar anatomy of Scrophulariaceae and Acanthaceae. Geophytology 12 (1): 22-29.
178. Kuptsova, L.P. and A.I. Bankovskii 1970. A new flavonoid from some Linaria species. Khim. Prir. Soedin (Tashk.) 6 (1): 128-129.
179. Lein Feliner, W. 1973. The gynoscium of Bignoniaceae: II The U shaped placenta of Schlegelia (Crescentieae) Oesterr Bot. Z. 121 (½) 13-22.

180. Li.H. 1947. Relationship and taxonomy of the genus Brandisia. J. Arn. Arb. 28 : 127-136.
181. Lincoln, D.E. 1980. Leaf resin flavonoids of Diplacus aurantiacum. Biochem. Syst. Ecol. 8 (4): 397-400.
182. Lindau, G. 1895. Acanthaceae in Die naturtichen pflanzen familien (Engler and K. Prantl.Eds.) IV 3b:274-354.
183. Lindley, J. 1836. A natural system of Botany etc. 2nd Ed. London.
184. Lindley, J. 1853. The Vegetable Kingdom 3rd ed. London.
185. Lowry, J.B. 1972. Anthocyanins of some Malaysian members of the Gesneriaceae. Phytochemistry 11 (11):3267-3269.
186. Lowry, J.B. 1973. Rhabdothamnus solandri, some phytochemical results. N.Z.J.Bot. 11(3) : 555-560.
187. Mabry, T.J., Markham, K.R. and M.B. Thomas. 1970. The systematic identification of flavonoids. Springer-Verlag, Berlin.
188. Mahato, S.B., Das, M.C. and N.P.Sahu, 1981. Triterpenoids of Scoparia dulcis. Phytochemistry 20(1) : 171-173.
189. Manchand P.S. and P.J. Blount. 1975. X-ray structure and absolute stereochemistry of stemarin, a diterpene with a new skeleton. J.Chem.Soc.Chem.Comm 21 : 894-895.
190. Manske, R.H.F. 1944. The alkaloids. Ann.Rev.Biochem. 13 : 533-548.
191. Marloth, R. 1932. The flora of the South Africa III. Darter Bros. and Co., Capetown.
192. Markham, K.R. 1982. Techniques of Flavonoid identification. Academic Press, London.

193. Martius, C.F.P.Von. 1835. Conspectus regni vegetabilis (etc. Nurnberg, Sehrag.
194. Mecoy Jeffy, W. and F.R.Stermitz. 1983. Alkaloids from Castilleja miniata and Penstemon whippleanus, 2 host species for the plume moth Amblypkilia pica. J.Nat.Prod. (Lloydia) 46(6) : 902-907.
195. Mecuse, A.D.J. 1970. The descent of the flowering plants in the light of new evidence from phytochemistry and from other sources I & II. Acta.Bot.Neerl. 19 : 61-72.
- *196. Meier, W. and A.Furst. 1962. Helv.Chim.Acta, 45, 232.
197. Melchior, H. 1964. In Engler's Syllabus der Pflanzen familien (Revised ed.) Vol.2. Berlin.
198. Metcalfe, C.R. and L.Chalk. 1950. Anatomy of the Dicotyledons. Clarendon Press. Oxford.
199. Meyers, F.H., Jawetz, E. and Goldfien. 1972. Review of Medical Pharmacology 3rd edn. Lange Medical Publications. Los Altos, California.
200. Minatsakanyan, V.A., Arutyanyan, L.S. and M.I. Eribekyan . 1983. Iridoid glycosides of Verbascum saccatum. Khim. Prir Soedin (Tashk.) 0(1) : 38-41.
201. Mohan Ram, H.Y. and M.Wadhi. 1965. Embryology and the delimitation of the Acanthaceae. Phytomorphology 15 : 201-205.
202. Mukherjee, J. 1975. Pollen morphology and plant taxonomy of Selagineae. (Scrophulariaceae). J.Palyn. XI : 1&2: 143-147.

203. Nagendran, C.R., Ramakrishna, T.M. and B.M. Narayana. 1980. Development of embryosac and endosperm haustoria in Striga densiflora. Indian J.Bot. 3(1) : 51-54.
204. Nair, A., Ramachandran, G., Ramesh, P., Sankara, S.S. and B.S.Joshi. 1976. Angeflorin; A new flavone glucoside from the leaves of Angelonia grandiflora. Indian J. Chem.(Sect. B) 14(6): 463-464.
205. Nakanishi, T., Ozaki, J., Inada, A., Murata, H., Nishi, M., Linuma, M. and K.Yoneda. 1985. Flavonoids of Striga asiatica. J.Nat.Pro.(Lloydia) 48(3): 491-493.
206. Niezgoda, C.J. and A.S.Tomb. 1975. Systematic Palynology of Tribe Leucophylleae (Scrophulariaceae) and selected Myoporaceae. Pollen E.Spores 17(4): 495-513.
207. Nonaka, G. and I.Nishioka. 1972. Bitter phenyl propanoid glycosides from Conandron ramoidioides. Phytochemistry 16(8) : 1265-1268.
- *208. Obdulio, F. and M.P. Lobeto. 1944. Chem.Zbl. I, 32.
209. Ozaki, Y., Johne, S. and M.Hesse. 1979. Iridoid glucosides from Leuocarpus perfoliatus. Helv.Chim.Acta 62(8): 2708-2711.
210. Padhye, P.M. 1982. Chemical systematics of Bignoniaceae and Verbenaceae Ph.D. Thesis, M.S.University, Baroda.
211. Paech, K. and M.V.Tracey 1965. Modern Methods of Plant Analysis Vol.II. Springer-Verlag.
212. Paris, R.R. and G.Bacidouin 1969. The presence of glucuronoxylsides of luteolin in the leaves of Digitalis purpurea L. C.R.Hebd. Seances Acad. Sci.Ser. D. Sci.Natur (Paris) 269 (17): 1675-1677.

213. Pawar, J.S. and A.R. Kulkarni. 1971. Contribution to the morphology of Pedaliaceae IV: Wood anatomy of Sesamum alatum Thonn. J. Shivaji Univ. 4 (B) : 117-120.
214. Peev Dimitar, R. 1982. Different substitution tendencies of leaf flavones in the Veronica hederifolia group. Plant Syst. Evol. 140 (2/3) : 235-242.
215. Pennell, F.W. 1920. Scrophulariaceae of Colombia. I. Proc. Acad. Nat. Sci. Phila. 72 : 136-188.
216. Pennell, F.W. 1921. Scrophulariaceae of the West gulf states. Proc. Acad. Nat. Sci (Phila.) 73: 459-536.
217. Pennell, F.W. 1935. The Scrophulariaceae, of eastern temperate North America. Acad. Nat. Sci.(Phila) Monograph 1 : 1-650.
218. Philcox, D. 1968. Revision of the Malesian species of Lindernia All. (Scrophulariaceae). Kew Bull. 22: 1-72.
219. Plouvier V., 1967. Research of flavone heterosides in some Fraxinus, Tilla and diverse plants. C.R.Hebd. Seances.Acad. Sci. Ser. D. Sci. Natur (Paris) 265 (21): 1647-1650.
220. Potter,S.D.L. 1897. Hand Book of Materia Medica, Pharmacy and Therapeutics. Manchester.
221. Price, J.P. 1963. The distribution of alkaloids in Rubiaceae. In Chemical Plant Taxonomy (Swain, T. ed.) Academic Press, London pp. 429-452.
222. Prijanto, B. 1969. The Asiatic species of Lindenbergia Lehm. Reinwardtia 7 (5): 543-560.

223. Prokosch, M., Prokosch, P., Weissenboeck, G. and R. Eloy
1982. Flavonoids from the leaf resin of Adenostoma
sparsifolium, Phytochemistry 21 (7) : 1835-1836.
224. Pulle, A. 1952. Compendium van de Terminologie, Homenclatuur
in Systematik der Zaadplanten 3rd ed. Utrecht.
225. Putudin, A.V., Smirnova, L.P., Glyzin, V.I. and A.I.
Ban'kovskii 1975. Search for pectolinarin and acetyl-
pectolinarin in plants of the genera Linaria Mill. and
Salvia L. Rastit Resur. 11 (2): 204-210.
226. Raj, B. 1961. Pollenmorphological studies in the Acanthaceae.
Grana Palynologica 3 : 1-108.
227. Reayat Khan, 1954. A contribution to the embryology of
Utricularia flexuosa Vahl. Phytomorphology 4: 70-117.
228. Reid, W.W. 1979. The diterpenes of Nicotiana species and
N. tabacum cultivars. In The Biology and Taxonomy of
the Solanaceae (Hawkes, T.G., Lester, R.N. and A.D. Skelding
Eds.) Academic Press, London, 273-278.
229. Rendle, A.B. 1925. The classification of flowering Plants
II Cambridge.
230. Rendle, A.B. 1938. The classification of flowering plants
2nd ed. Cambridge, Vol. 2.
231. Richard N. Lester 1979. The use of protein characters in
the taxonomy Solanum and other Solanaceae. In The Biology
and Taxonomy of the Solanaceae (Hawkes J.G. Lester, R.N.
and A.D. Skelding Eds.) Academic Press, London, 285-303.
232. Robertson, C. 1891. Flowers and insects. Asclepiadaceae to
Scrophulariaceae. Trans. Acad. Sc. St. Louis 5: 569-598.

233. Roby, M.R. and R.S. Frank 1984. Pyrrolizidine and pyridine monoterpene alkaloids from 2 (Castilleja Plant hosts of the plume moth Platyptilla pica. J.Nat.Prod.(Lloydia) 47 (5): 846-853.
234. Roby, M.R. and R.S.Frank, 1984a. Penstemonoside and other iridoids from Castilleja rhexifolia etc. J.Nat.Prod.(Lloydia) 47 (5): 854-857.
235. Robyns, W. 1931. L' organisation florals des Solanacees Zygomorphes. Mem. Acad. Roy. Belg. Coll. 8, Sc. 11 (8):1-96.
236. Rofle, R.A. 1883. On the Selagineae described by Linnaeus, Bergius, Linnaeus fil and Thunberg. Jour. Linn. Soc.Bot. 20: 338-358.
237. Rothmaler, W. 1954. Notes on Western Antirrhineae. Leafl. West. Bot. 7 : 113-117.
238. Rouy, G. 1909. Conspectus des tribus et des genres de la famille des Scrophulariacees. Rev. Gen. Bot. 21: 194-207.
239. Sahi, B. and S.N. Dixit 1969. Acanthaceae. A Phylogenetic and taxonomic consideration. Plant Sci. 1: 101-108.
240. Salama, O. and O. Sticher 1983. Iridoid glucosides from Euphrasia rostkoviana Planta Med. 47 (2): 90-94.
241. Salez, A.E., Mas, G.E., Bermejo, B.P. and P.P. Garcia 1984. Study of phenolic acids in two species of Digitalis An.R.Acad. Farm 50 (3) : 451-458.
242. Schmid, E. 1906. Beitrage zur Entwicklungsgeschichte der Scrophulariaceen. Beih. Bot. Centralb 20 (1): 175-299.
243. Seifert, K., Emmanuel, S., Seigfried, J. and H. Manfred 1982.

244. Seifert, K., Siefried, J. and H.Manfred 1982a.
Verbasconine, a macrocyclic spermine alkaloid isolated
from Verbascum, Helv. Chim.Acta 65 (8): 2540-2547.
245. Seikel, M.K. 1955. J. Am. Chem.Soc. 77, 5685.
246. Sen, A.B. and A.R. Chowdhury 1970. Celsianol. a novel sterol
from Celsia coromandeliana Vahl. J. Indian Chem.Soc. 47 (11):
1063-1967.
247. Shukla Y.N. and S.T. Reghunath, 1983. Hepta triacontan
4-one tetratriacontanyl octacosanoate and other constituents
from Pedaliium murex. Phytochemistry 22(4): 973-974.
248. Singh, B. and R.P. Rastogi 1972. Chemical examination of
Picrorhiza kurrooa VI. Reinvestigation of Kutkin. Indian J.
Chem. 10 (1): 29-31.
- *249. Skottsberg, C. 1940. Vaxhernas Liv.
Stockholm.
250. Soo, R. 1953. Die modernen Grundsätze der Phylogenie im
neuen System der Blütenpflanzen. Acta. Biol. Acad. Scient.
Hung. 4 : 257-306.
251. Srinivasan, A.R. 1942. Contribution to the morphology of
Pedaliium murex Linn. and Sesamum indicum D.C. Proc. Indian
Acad. Sci. 16 B : 155-162.
252. Srinivasan, Lucas, S.T., Burrowes, C.B., Wanderman, N.A.,
Redner, A., Bernstein, S. and P.N. Sawyen. 1971. Effects of
some flavonoids on the surface change characteristics of the
vascular system and their antithrombogenic characteristics
European Conf. Microcirculation 6, 394.

253. Steenis, C.G.G.V. 1949. Notes on the genus Wightia (Scrophulariaceae) Bull. Bot. Gard. Buitenzorg, III 18: 213-227.
254. Sticher, O., Meier, B., Lehmann, D. and L. Swiatek. 1980. Scrophularioside, a new iridoid glucoside from Scrophularia laterifolia. Planta Med. 38 (3): 246-254.
255. Sticher, O., Salama, O., Ratan, K. Chaudhuri and T. Winkler 1982. Structural analysis of eukovoside, a new phenyl propanoid glycoside from Euphrasia rostkoviana Hayne. Helv. Chim. Acta. 65 (5): 1538-1542.
256. Swaitek, L. 1969. Pharmacobotanic investigation of some species of family Scrophulariaceae III. Chemical constituents of herbs of Veronica spicata L. Diss. Pharm. Pharmacol. 21(6) : 545-551.
257. Switek, L. 1970. Pharmacobotanical investigation of some species of family Scrophulariaceae. IV. Chemical constituents of the herbs Scrophularia nodosa L. Diss. Pharm. Pharmacol. 22(5) : 323-328.
258. Swaitek, L. 1973. Phenolic acids of underground parts of Scrophularia nodosa L. Pol. J. Pharmacol. Pharm. 25(5) : 461-464.
259. Swaitek, L. and B. Broda. 1967. Pharmacobotanical investigation of some species of the Scrophulariaceae. Diss. Pharm. Pharmacol. 19(4) : 395-401.
260. Swaitek, L. and D. Elzbieta. 1976. Pharmacobotanical investigations on some species of Scrophulariaceae family

- V. Chemical constituents of Lathraea squamaria L.
Pol. J. Pharmacol. Pharm. 28 (1): 105-109.
261. Suryakanta, 1973. Pollenmorphological studies in the Bignoniaceae. J. Palyn (Lucknow) 9 : 45-82.
262. Takahashi, K., Yoshihisa, T. and S. Kishijara 1973. Studies on constituents of medicinal plants XII. The constituents of the flowers of Linaria japonica Miq. Yakugaku. Zasshi. 93 (6): 707-710.
263. Takeda, Y. and T. Fujika 1981. Iridoid glycosides of Melampyrum laxum Miq. Planta Med. 41 (2): 192-194.
264. Takhtajan, A. 1968. Classification and Phylogeny with special reference to the flowering plants. Proc. Linn. Soc. London 179 :221-227.
265. Takhtajan, A. 1969. Flowering plants: Origin and Dispersal. Edinburgh.
266. Takhtajan, A. 1980. Outline of the classification of flowering plants (Magnoliophyta) Bot. Rev. 46 (3): 225-359.
267. Thieret J.W. 1954. The tribes and genera of Central American Scrophulariaceae Ceiba 4 : 164-168.
268. Thieret, J.W. 1967. Supraspecific classification in the Scrophulariaceae, A review. Sida 3 (2): 87-106.
269. Thorne, R.F. 1968. Synopsis of a putatively phylogenetic classification of the flowering plants. Aliso 6 : 57-66.
270. Tiagi, B. 1956. A contribution to the embryology of Striga orobanchoides Benth and Striga euphrasioides Benth, Bull. Torr. Bot. Cl. 83 : 154-170.

271. Trim, A.R. and R. Hill 1951. The preparation and properties of Aucubin, Asperuloside, and some related glycosides. Biochem.J. 50: 130.
272. Tunmann, P and H.J. Bauersfeld 1975. Further components of the roots of Barbapogon procumbens DC Arch.Pharm (Weinheim) 308 (8): 665-667.
273. Ulubellen, A., Cetin, E.T. and A.Guran 1975. Steroidal compounds and hydrocarbons of Verbascum lasianthum Planta Med. 27 (1): 14-17.
274. Urban, I. 1926. Synopsis ilicifolia Repert. Spec.Nov.Regn. Veg. 22 : 369-371.
275. Valdes, B. 1970. Flavonoid pigments in flower and leaf of the genus Linaria. Phytochemistry 9 (6): 1253-1268.
276. Van Tieghem, P. 1891. Traite de Botanique 2 me pt.2me ed. F. Savy, Paris.
277. Van Tieghem, P. 1903. Structure de le'tamine Chez les Scrophulariacees. Ann.Sci.Nat.(Bot.) VIII,17 :363-371.
278. Varghese, T.M. 1967. Morphological studies in the family Scrophulariaceae. Agra Univ. J. Res.Sci. 16 (1): 153-157.
279. Varghese, T.M. 1971. Studies in the family Scrophulariaceae III Floral anatomy of some members of Gratiolaceae. Proc. India Acad. Sci. Sect. B 74 (1): 6-15.
280. Vij, S.P. and S.K.Kashyap 1976. Cytological studies in some North Indian Scrophulariaceae. Cytologia, 41: 685-695.
281. Wagner, W.H. Jr. 1980. Origin and Philosophy of ground plan-divergence methods of cladistics. Syst.Bot. 5:173-193.

282. Wang, M.H., Schermeister, L.J. and S.K.W. Khalil 1976. Thin-layer Chromatographic studies of the steroidal and cardenolide constituents of Penstemon species of Borth Dakota. Planta Med. 30 (2): 141-143.
283. Weinges, K. and K. Kuenstler 1977. Natural products from medicinal plants XXII Justus Liebigs Ann.Chem. (6): 1053-1057.
284. Wernham, H.F. 1911-1912. Floral evolution, with particular reference to the sympetalom dicotyledons New Phytologia 10: 73-83, 109-20; 145-59, 217-26, 293-305, 11 : 145-66, 217-35, 290-305, 373-97.
285. Wettstein, R. Von. 1891. Scrophulariaceae in Engler, A. and K. Prantl. Die natuerlichen Pflanzen familien 4 (3b): 39-123.
286. Wettstein, R. Von 1897. Scrophulariaceae. In Engler A. and K. Prantl. Die natuerlichen pflanzen familien. Nachtrage I Zu IV 3b; 293-299.
287. Wettstein, R. 1935. Handbuch der systematischen Botanik. 4th ed. Leipzig and Vienna.
288. Wieffering, J.R. 1966. Aucubinartige Glucoside (Pseudoindikane) und Verwandte Heteroside als systematische Merkmale. Phytochemistry 5 : 1053-1064.
289. Wink, M., Witte, L. and T. Hartmann 1981. Quinolizidine alkaloid composition of plants in Orobanche rapum genistae. Planta Med. 43 (4): 342-352.
290. Wollenweber, E., Cornelia, R. and V.H. Dietz 1981. The occurrence of aurentiancin and flavokawin B on Pityrogramma triangularis var. pallida and Didymocarpus species Phytochemistry 20 (5): 1167-1168.

291. Yang, R., Zhou, J., Zheng, Q. and J. Clardy 1983.
Melasmoside, a novel monoterpenoid glucoside from
Melasma arvense. Acta. Bot. yunnanica 5(2): 213-218.
292. Ziyaev, R., Abdusamatov, A and S.Yu. Yunusov 1971.
Alkaloids of Verbascum songoricum. Khim.Prir. Soedin.
(Tashk.) 7 (6) : 853-854.

* Not referred to the original article.