REFERENCES

Adjei, M. B., Quesenberry, K. H., & Chambliss, C. G. (2006). Nitrogen fixation and inoculation of forage legumes.

Afridi, R. A., Khan, M. A., Gul, H., & Daud, M. K. (2014). Allelopathic influence of rice extracts on phenology of various crops and weeds. *Pak. J. Bot*, *46*(4), 1211-1215.

Agrawal, A.R. ,Gehlot, A.,Verma, R and Rao ,P.B.(2002). Effect of weed extracts on seedling growth of same varieties of wheat. *Journal of environmental biology*.23:19-23.

Ahmad, A., Husain, A., Mujeeb, M., Khan, S. A., Alhadrami, H. A. A., & Bhandari, A. (2015). Quantification of total phenol, flavonoid content and pharmacognostical evaluation including HPTLC fingerprinting for the standardization of *Piper nigrum* Linn fruits. *Asian Pacific Journal of Tropical Biomedicine*, *5*(2), 101-107.

Aisha Ume, A., Yousaf, Z., Khan, F.,Hussain, U., Anjum, A.,Nayyab, Q. and Amina Younas, A.(2010). Evaluation of allelopathic potential of some selected medicinal species .African Journal of Biotechnology. **9(37):**6194-6206,

Akobundu, I.O. (1987). Weed science in the tropics. Principles and practices. 522pp. Chicester etc: Wiley

Allelopathy, an alternative tool to improve cropping systems. A review Manoel Bandeira de Albuquerque1*, Roseane Cavalcanti dos Santos2, Liziane Maria Lima2, Péricles de Albuquerque Melo Filho3, Rejane Jurema Mansur Custódio Nogueira4, Claudio Augusto Gomes da Camara ˆ 5, Alessandra de Rezende Ramos Agron. Sustain. Dev. (2011) 31: –379 395

Alstrom, S. (1996). Fundamentals of weed management in hot climate peasant agriculture. Vigyan Prakashan Jodhpur. pp. 5.

An, M. (2005). Mathematical modelling of dose-response relationship (Hormesis) in allelopathy and its application. *Nonlinearity in biology, toxicology, medicine*, *3*(2), nonlin-003.

Anjum, T. and Bajwa, R. (2005). Importance of Germination Indices in Interpretation of Allelochemical effects. *International journal of agriculture & biology*.**7:** 417-419.

Anon (1982) "In The Wealth of India: Raw Materials" Publication and Information Directorate, CSIR: New Delhi**, (10):** 78–81.

Anonymous (2004) "Wealth of India (Raw Materials)" Publications and Information Directorate, CSIR: New Delhi. **10:**77-81

Anonymous, (1976) "The Wealth of lndia"CSIR: New Delhi, pp.79-SO.

Ashraf, N., & Sen, D. N. (1978). Allelopathic potential of *Celosia argentea* in arid land crop fields. *Oecologia Plantarum*, *13*(4), 331-337.

Bais HP, Weir TL, Perry LG, Gilorys, Vivanco JM (2006). Annual Rev.Plant Biol.**57**: 233-266.

Bais, H. P., Park, S. W., Weir, T. L., Callaway, R. M., & Vivanco, J. M. (2004). How plants communicate using the underground information superhighway. *Trends in plant science*, *9*(1), 26-32.

Baličević,R., Ravlić, and Ivana Ravlić,M.(2015). Allelopathic effect of aromatic and medicinal plants on *Tripleurospermum inodorum* (l.) C.h. schultz. **Herbologia 15(2):** 41-53.

Baranwal A., Mazumder A., Chakraborthy, S. and Gupta, S. (2014). Phytopharmacological uses of *Tephrosia purpurea* - a review. *Pharmacophore*.**5 (4):** 658-665.

Barto, E.K. and Cipollini, D. (2009). Density-dependent phytotoxicity of *Impatiens pallida* plants exposed to extracts of *Alliaria petiolata*. *Journal of Chemical Ecology* **35**: 495-504.

Batish D.R., Kaur, S., Singh, H.P. and Kohli, R.K. (2009). Nature of interference potential of leaf debris of *Ageratum conyzoides. Plant growth regulation*.**57**:137-144.

Bell, D.T. and Koeppe, D.E.(1972). Noncompetatitve effect of giant foxtail on growth of corn. *Agronomy journal*.**64**:321-325.

Belz RG. **(2007).** Allelopathy in crop. Weed interactions. An update. *Pest Management Science*.**63**: 308.326.

Bhatt, A., Rawal, S., Dhar, U. (2006). Ecological features of a critically rare medicinal plant, *Swertia chirayita*, in Himalaya. *Plant Species Biol*. **21:** 49–52.

Bi, H.H., Zeng, R.S., Su, L.M., An, M., and Luo, S.M.(2007). Rice allelopathy induced by methyl jasmonate and methyl salicylate. *Journal of chemical ecology*.**33**:1089-1103.

Bold, H. C. 1957. *Morphology of plants*. Harper & Row, NY. 669 p.

Boyetchko, S. M., & Rosskopf, E. N. (2006). Strategies for developing bioherbicides for sustainable weed management. Handbook of sustainable weed management, 393-430.

Breen, J., & Ogasawara, M. (2011). A vision for Weed Science in the twenty-first century. *Weed Biology and Management*, 11(3), 113-117.

Caamal-Maidonado, J.A., Jimenez-Osorio, J.J., Torres - Barragan, A. and Anaya, A.L. (2001). The use of allelopathic legume cover and mulch species for weed control in cropping systems. *Agronomy journal.*93: 27-36.

Capinera, J.L. (2001). Handbook of Vegetable Pests. Academic Press, New York. 729 pp.

Cartwright, D.W., Langcake, P., Pryce, R.J., Leworthy, D.P. and Ride, J.P. (1977.) Chemical activation of host defence mechanisms as a basis for crop protection. *Nature*.**267**: 511-513

Chandler et al., 1984P.M. Chandler, J.A. Zucar, J.V. Jacobson, T.J.V. Higgins, A.S. Inglis(1984), The effect of gibberellic acid and abscisic acid on α-amylase mRNA levels in barley aleurone layers studies using an α-amylase c DNA clone Plant Mol. Biol., 3 pp. 407–408

Channappagoudar, B.B.,Jalageri, B.R.and Biradar, N.R.(2005). Allelopathic Effect of Aqueous Extracts of Weed Species on Germination and Seedling Growth of Some Crops.*Karnataka J.Agric.Sci.***18 (4):** 916-920.

Charudattan R.(2005). Use of plant pathogens as bioherbicides to manage weeds in horticultural crops. *Proceedings of florida state horticulture society.***118**:208-214.

Conrad,J.P.(1927).*American society of agronomy journal.***19**,1091.

Dangash, A. J., Pandya, N., Bharillya, A., Jhala, A., & Dharamchand, Jain (2014). Impact of Exogenous Elicitors on Artemisinin Production and Trichome Density in. *Notulae Scientia Biologicae,* 6(3), 349-353.

de Albuquerque, M. B., dos Santos, R. C., Lima, L. M., de Albuquerque Melo Filho, P., Nogueira, R. J. M. C., Da Câmara, C. A. G., & de Rezende Ramos, A. (2010). Allelopathy, an alternative tool to improve cropping systems. A review. *Agronomy for Sustainable Development*.

Devi, O.I. and Dutta, B.K.(2012). Allelopathic Effect of the Aqueous Extract of *Parthenium hysterophorus* and *Chromolaena Odorata* on the Seed Germination and Seedling Vigour of Zea mays L. In vitro.*Academic Journal of Plant Sciences.***5 (4):** 110-113.

Devi, Y. N., Dutta, B. K., Sagolshemcha, R., & Singh, N. I. (2014). Allelopathic effect of Parthenium hysterophorus L on growth and productivity of Zea mays L. and its phytochemical screening. *International Journal of Current Microbiogand and applied science.**3*** **(7):** 837-846.

Devine, M. D., S. O. Duke, and C. Fedtke. (1993). Physiology of the Herbicide Action. PTR Prentice Hall, Englewood Cliffs, New Jersey. 441 p.

Deyama,D.P.(1986). Allelopathic potential of *Parthenium hysterophorus* Linn.on the growth, nodulation and nitrogen content of *Leucaena leucocephala*. Leucaena Research Reports.**3 (7):** 36-37.

Dodd, A.P.(1940). The biological campaign against prickly pear.commonwealth prickly pear board, brisbane,Australia,pp.177.

Duke, S.O., and J. Lydon, (1987). Herbicides from natural compounds. *Weed Technology*. **1:** 122-128.

Duke,S.,Vaughn, K., EM. Croom E. and Elsholy, H.(1987). Weed Science.**35:**499-50.

Duke,S.O., Cedergreen,N., Vellini, E.D., and Belz, R.G. (2006). Hoemesis: is it an important factor in herbicides use in allelopathy? *Outlook in pests managment* **19**:29-33.

Einhelling,F.A. and Eckrich,D.C.(1984). Interaction of temperature and ferulic acid stress on grain sorghum and soyabean.*Journal of Chemical Ecology*.**10**:161-170.

Ercoli, L., Masoni, A., Pampana, S., & Arduini, I. (2007). Allelopathic effects of rye, brown mustard and hairy vetch on redroot pigweed, common lambsquarter and knotweed. *Allelopathy Journal*, 19(1), 249.

Fassoni, A.C. and Martins, M.L. (2013). Mathematical analysis of a model plant invasion mediated by allelopathy. Ecological complexity,artical in press.

Femina D Lakshmipriya P Subha S and Manonmani R. 2012, Allelopathic effects of weed (*Tridax procumbens* L.) extract on seed germination and seedling growth of some leguminous plants. International Research Journal of Pharmacy, 3(6): 90-95.

Flamini, G. (2012). Natural herbicides as a safer and more environmentally friendly approach to weed control: a review of the literature since 2000. Stud. *Nat. Prod. Chem*, 38, 353-96

Foy, C.L. (1999). How to make bioassays for allelopathy more relevant to field conditions with particular reference to cropland weeds. In: Inderjit, K.M.M. Dakshini & C.L. Foy (eds.), *Principals and Practices in Plant Ecology: Allelopathic Interactions*. CRC Press, Washington pp. 25–33.

Fuji,Y.,Parvez, S.,Parvez, M.,Ohmae, Y. And Iida O.(2003). Screening of 239 medicinal plant species for allelopathic activity using the sandwich method. *Weed biology and management*.**3**:233-241.

Fullaway, D.T. (1954). Biological control of cactus in Hawaii. *Four.econ.ent.*, **47**:696-700.

Gliessman, S. R. (1983). Allelopathic interactions in crop-weed mixtures. *Journal of Chemical Ecology*, *9*(8), 991-999.

Grodzinsky,A. M.(1982).evolutionary problems of the chemical interactions among plants. In: evolution and environment (eds., V.J.A.Nevak and J.mlikovsky). Pp.133-143.

Gulati, S. K., Garg, M. R., & Scott, T. W. (2005). Rumen protected protein and fat produced from oilseeds and/or meals by formaldehyde treatment; their role in ruminant production and product quality: a review. *Animal Production Science*, *45*(10), 1189-1203.

Hariprasad P, Ram Krishnan N (2003) Algae assay used for the determination of organic pollution level in freshwater body at Tiruvannamalai, *India. J Ecotoxicol Environ Monit* 13:241–248.

Hashimoto, T., Hayashi, A., Amano, Y., Kohno, J., Iwanari, H., Usuda, S., & Yamada, Y. (1991). Hyoscyamine 6 beta-hydroxylase, an enzyme involved in

tropane alkaloid biosynthesis, is localized at the pericycle of the root. *Journal of Biological Chemistry*, *266*(7), 4648-4653.

Hirai,N. (2003). Application of allelochemicals to agriculture. *Biological Science in Space*.**17**:4-5.

Hoagland, R. E., & Williams, R. D. (2003). Bioassays–Useful tools for the study of allelopathy. *Allelopathy: chemistry and mode of action of allelochemicals*, 315-451.

Hoagland, R. E., Weaver, M. A., & Boyette, C. D. (2007). *Myrothecium verrucaria* fungus: A bioherbicide and strategies to reduce its non-target risks. *Allelopathy Journal*, 19(1), 179.

Horton, David R. and Lewis, T. M. (2003). Numbers and types of arthropods overwintering on common mullein, *Verbascum thapsus* L. (Scrophulariaceae), in a central Washington fruit-growing region" . *Journal of the Entomological Society of British Columbia* **100**: 79–86.

Hou, J.Q. and J.T. Romo, (1998). Effects of chemicals stimulators on germination of Winterfat (*Ceratoideslanata* (Pursh) J. T. Howell). *Seed Science and Technology*, 26: 9-16.

http://www.fao.org/docrep/009/ag087e/AG087E05.htm
http://www.fao.org/docrep/009/ag087e/AG087E05.htm

Huffaker, C.B.(1970). Fundamental of Biological weed control In:Biological control of insect pests and weeds,Chapman and Hall ltd. Chapter 22,pp.631.

Ilori, O.J.,Otusanya,O.O.,Adelusi,A.A. and Sanni,R.O.(2010). Allelopathic activities of some weeds in the asteraceae family.*International journal of botany*.**6 (2):** 161-163.

Inderjit, K., & Keating, L.(1999). Allelopathy: principal and practice. John promises for biological control in: advance in agronomy, (eds). Sparks Dl. *Academic Press*, *67*, 141-231.

Inderjit, Seastedt, T.R., Callaway, R.M., Pollock, J.L. and Kaur, J. (2008). Allelopathy and plant invasions:Traditional, congeneric and bio-geographical approaches. *Biological Invasions***10:** 875-890.

Inderjit, W. R., and Callaway, RM (2011). The ecosystem and evolutionary contexts of allelopathy. *Trends Ecol. Evol*, *26*, 655-662.

J Wallace 2001. Organic field crop handbook. Canadian Organic Growers, 2nd ed., Ottawa, Ontario, Canada.

Jain A. And Jain A. (2012).*Tridax Procumbens (L.)*: A Weed With Immense Medicinal Importance: A Review. *International Journal of Pharma and Bio Sciences.3 (1):544-552.*

Jain, R., Fischer, S., Serra, S., & Chetty, R. (2010). The use of cytokeratin 19 (CK19) immunohistochemistry in lesions of the pancreas, gastrointestinal tract, and liver. *Applied Immunohistochemistry & Molecular Morphology*, *18*(1), 9-15.

Jamal, A. R. M. (2011). Ocimum sanctum L.: A review of phytochemical and pharmacological profile.

JG. Romangi J., Duck, S., and Dayan,E.(2000). Plant physiology. **123:725-**732.

Kader MA (2005) A comparison of seed germination calculation formulae and the associated interpretation of resulting data. Proc R Soc NSW 138:65–75

Kalinova, J. (2010). Allelopathy and organic farming. In Sociology, Organic Farming, Climate Change and Soil Science (pp. 379-418). Springer Netherlands.

Kato-Noguchi H, Ota K and Kujime H (2012). Absorption of momilactone A and B by Arabidopsis thaliana L. and the growth inhibitory effects.*Journal of Plant Physiology. 169(15):1471-1476.*

Kato-Noguchi, H., Kujime, H., and Ino, T. (2007). UV-induced momilactone Baccumulation in rice rhizosphere. *Journal of Plant Physiology.***164**: 1548–1551.

Khosravi, R., Sendi, J., & Ghadamyari, M. (2010). Effect of *Artemisia annua* L. on deterrence and nutritional efficiency of lesser mulberry pyralid (Glyphodes pylolais

Walker)(Lepidoptera: Pyralidae). *Journal of Plant Protection Research*, *50*(4), 423-428.

Kim,Y.O. and Lee,E.J.(2011). Comparison of phenolic compounds and the effect of invasive and native species in East Asia: support for the novel hypothesis. *Ecological research.***26**:87-94.

Kuldeep G. and Pathak, A (2013) Pahrmacognostic and Phytochemical evaluation of Tridax procumbens L. Journal of Pharmacognosy and phytochemistry. 1(5): 42-46.

Kumar, B., Vijayakumar, M., Govindarajan, R., & Pushpangadan, P. (2007). Ethnopharmacological approaches to wound healing—exploring medicinal plants of India. *Journal of Ethnopharmacology*, *114*(2), 103-113.

Kumar, G. Krishna, and T. Dennis Thomas.(2012): High frequency somatic embryogenesis and synthetic seed production in *Clitoria ternatea* Linn." *Plant Cell, Tissue and Organ Culture (PCTOC)* 110, no. 1 141-151.

Kumar, K. A., Sharvanee, S., Patel, J., & Choudhary, R. K. (2010). Chemical composition and antimicrobial activity of the essential oil of Desmostachya bipinnata linn. International Journal of Phytomedicine, 2(4).

Kumari A.K., RajaKumar K.N., Narasimha Rao, C.H. (2014) Adverse effects of chemical fertilizer and pesticides health and environment. *Journal of Chemical and Pharmaceutical sciences.*150-151.

Latham, P. 4.3. 3 Non-Native Invasive Species.

LEE, I.K., and MONSI, M. 1963. Ecological studies on*Pinus densiflora* forest. I. Effects of plant substances on the floristic composition of the undergrowth.*Bot. Mag (Tokyo)* 76:400–413.

Levin,D.A.(1976). *Annu.Rev.Ecolo.Syst.***7**,121-159.

Liang, G., Lin, S., & Liu, D. (2008). Quantitative determination of total saponins in polygalae by spectrophotometry. *Prog mod biomed*, *8*, 104-111.

Liebman, M., & Davis, A. S. (2009). Managing weeds in organic farming systems: an ecological approach. Organic farming: The ecological system. Francis C, editor. Madison: American Society of Agronomy, 173-196.

Liebman, M., & Gallandt, E. R. (1997). Many little hammers: ecological management of crop-weed interactions. Ecology in agriculture, 291-343.

Lodhi, S., & Singhai, A. K. (2013). Wound healing effect of flavonoid rich fraction and luteolin isolated from Martynia annua Linn. on streptozotocin induced diabetic rats. *Asian Pacific journal of tropical medicine*, *6*(4), 253-259.

Lorenzo, P., González, L. And Reigosa, M.J. (2010). The genus *Acacia* as invader: The characteristic case of *Acacia dealbata* Link in Europe. *Annals of Forest Science* **67**: 101.

Lowry, O. H., Rosebrough, N. J., Farr, A. L., & Randall, R. J. (1951). Protein measurement with the Folin phenol reagent. J biol Chem, 193(1), 265-275.

Lowry, O. H., Rosebrough, N. J., Farr, A. L., and Randall, R. J. (1951) Protein measurement with the Folin phenol reagent. *J. Biol. Chem.* **193,** 265–275.

Luis, M. D., Verdú, M., & Raventós, J. (2008). Early to rise makes a plant healthy, wealthy, and wise. *Ecology*, *89*(11), 3061-3071.

Maharjan,S., Shrestha,B. and Jha P.(2007).Allelopathic effects of aqueous extract of leaves of *Parthenium hysterophorus* l. On seed germination and seedling growth of some cultivated and wild herbaceous species.*Scientific World*.**5(5):**33-39.

Maisuria,K., And Patel, S. (2009). Seed germinability, root and shoot length and vigour index of soybean as influenced by rhizosphere fungi. Karnataka Journal of Agriculture Science. **22(5)** :1120-1122.

Mali A.A., Kanade M.B., **(2014)***Ann. Bio. Res*., 5 (3), 89-92.

Mandava, N.B. (1979). In plant growth substances (B.N.Mandava, ed.) P.135, ACS symposium series no.111, American Chemical Society , Washington DC.

Mandava, N.B. (1985).Chemistry and biology of allelopathic agents. In: *chemistry of allelopathy: biochemical interaction among plants* (Ed., A.C.Thompson). ACS Symposium Series No.268: 33-54. American Chemical Society, Washingtom DC.

Mani,V.S.,Gautam, K.C and Arora,R.K.(1975).Beware of the perdify of Parthnenium. *Indian farming*.**25(9):**15-16.

Marking, Syl (2002). Marestail Jumps Glyphosate Fence .Corn and Soybean Digest.

Mccalla, T.M. and Haskins, F.A.(1964). Phytotoxic substances from soil microorganisms and crop residues. *Bacteriological review*.**28**:181-207.

Mersie, W., & Singh, M. (1987). Allelopathic effect of parthenium (Parthenium hysterophorus L.) extract and residue on some agronomic crops and weeds. *Journal of Chemical Ecology*, *13*(7), 1739-1747.F

Miller, D.(1936). Biological control of noxious weeds.New zealand four.*Science and technology*.**18**: 581-584.

Mohsenzadeh S., Gholami M., Teixeira da Silva J.A. (2011). Allelopathic potential of *Ephedra*. *Med. Aromatic Plant Sci. Biotechnol*. 5: 160–162.

Molisch, H.(1937). Influence of one plant on another : allelopathy,Jena Germany: Gustave Fischer Verlag.

Moravcova, L.,Pysek, P.,Jarosik,V.and Zakravsky, P.(2011). Potential phytotoxic and shading effects of invasive *Fallopia* (Polygonaceae) taxa on the germination of dominant native species. *NeoBiota*.**9**:31-47.

Morvillo, C. M., de la Fuente, E. B., Gil, A., Martínez-Ghersa, M. A., & González-Andújar, J. L. (2011). Competitive and allelopathic interference between soybean crop and annual wormwood (*Artemisia annua* L.) under field conditions. *European Journal of Agronomy*, *34*(4), 211-221.

Mousavi, S. H., Alami-Saeid, K. H., & Moshatati, A. (2013). Effect of leaf, stem and root extract of alfalfa (Melilotus indicus) on seed germination and seedling

growth of wheat (*Triticum aestivum*). *International Journal of Agriculture and Crop Sciences*, *5*(1), 44.

Msafiri, C. J., Tarimo, M. T., & Ndakidemi, P. A. (2013). Allelopathic effects of Parthenium hysterophorus on seed germination, seedling growth, fresh and dry mass production of *Alysicurpus glumaceae* and *Chloris gayana*. *American Journal of Research Communication*. ***1*(11):**190-205.

Muller, C. H. (1969). Allelopathy as a factor in ecological process. *Plant Ecology*, *18*(1), 348-357.

Muzila, M., Setshogo, M. P., Moseki, B., & Morapedi, R. (2011). An assessment of *Prosopis* L. in the Bokspits area, south-western Botswana, based on morphology.

Nandagopal, S. and Ranjitha Kumari, B.D. (2007). Phytochemical and antibacterial studies of Chicory (*Cichorium intybus L.)*-A multipurpose medicinal plant. *Advan. Biol. Res.*,**1(1-2):** 17-21.

Narwal S.S.(2004). Allelopathy in crop production.Scientific Publisher.Jodhpur.pp-174-175.

Narwal, S. S., Palaniraj, R., Sati, S. C., Kadian, H. S., & Dahiya, D. S. (2003). Allelopathic plants: 8. *Parthenium hysterophorus*, 151-170.

Nasrine S., M-El-Dariera S.and Halilat M.El-Taher, H. (2011). Allelopathic effect from some medicinal plants and their potential uses as control of weed. International Conference on Biology, Environment and Chemistry.**24**:15-22.

Nekonam, M. S., Razmjoo, J., Kraimmojeni, H., Sharifnabi, B., Amini, H., & Bahrami, F. (2014). Assessment of some medicinal plants for their allelopathic potential against redroot pigweed (*Amaranthus retroflexus*). *Journal of Plant Protection Research*, *54*(1), 90-95.

Netsere, A. and Mendesil,E. (2011). Allelopathic effects of *Parthenium hysterophorus* L. aqueous extracts on soybean (*Glycine max* L.) and haricot bean

(*Phaseolus vulgaris* L.) seed germination, shoot and root growth and dry matter production. *Journal of Applied Botany and Food Quality*. **84**: 219 - 222

Nojiri ,H., Sugimora, M., Yamane, H., Nishimura, Y., Yamada, A., Shibuya, N., Kodama, O., Murofushi, N., and Omori, T.(1996). Involvement of jasmonic acid in elicitor-induced phytoalexin production in suspension-cultured rice cells. *Plant Physiology*. **110**: 387–392.

Pace, P. F., Cralle, H. T., El-Halawany, S. H., Cothren, J. T., & Senseman, S. A. (1999). Drought-induced changes in shoot and root growth of young cotton plants. *J. Cotton Sci*, *3*, 183-187.Wu, H., Pratley, J., Lemerle, D., Haig, T., An, M. (2001). Screening methods for the evaluation of crop

Pandey,D.K.(1994). Inhibition of Salvinia by Parthenium.ii. relative effect of flower,leaf,stem,and root residue on salvinia and paddy. *Journal of chemical ecology*.20:3123-3131.

Parihar,S.S. and Kandolia, K.C.(1987). Beware of the exotic weeds.*Indian farming*.**37(9):** 24-27.

Parker, K.F. (1972). An Illustrated Guideto Arizona Weeds. Univ. of Arizona Press. Tucson

Parmar, N. S., Tariq, M., & Ageel, A. M. (1988). Gastric anti-ulcer and cytoprotective effect of selenium in rats. *Toxicology and applied pharmacology*, *92*(1), 122-130.

Patade,Y., Maya, K., and Zakwan, A. (2011). Seed priming mediated germination improvement and tolerance to subsequent exposure to cold and salt stress in capsicum. Research. *Journal of. Seed Science***, 4 (3):** 125 -136.

Patel D.A. and Pandya N.R. (2013). Allelopathic effect of few medicinal plants on *Raphanus sativus* L. *International Journal of Research in Plant Science* 3(4) 88-93

Patel, D. D., & Kumbhar, B. A. (2016). Weed and its Management: A Major Threats to Crop Economy. *J Pharm Sci Bioscientific Res*, *6*(6), 453-758.

Pattanayak, P., Behera, P., Das, D. and Panda, S.(2010). An Overview.*Pharmacognsy* Rev. **4(7):** 95–105

Prasad, K. and V.C. Srivastava. 1991. Teletoxic effect of some weeds on germination and initial growth of groundnut (*Arachis hypogea*). *Ind. J. Agr. Sci.* 61:493–494.

Prati, D. And Bossdorf, O. (2004). Allelopathic inhibition of germination by *Alliaria petiolata* (Brassicaceae). *Americal Journal of Botany* 91: 285-288.

Pudełko, K., Majchrzak, L., & Narożna, D. (2014). Allelopathic effect of fibre hemp (*Cannabis sativa* L.) on monocot and dicot plant species. *Industrial Crops and Products*, *56*, 191-199.

Purohit,S. and Pandya, N.(2013). Allelopathic activity of *Ocimum sanctum* L. and *Tephrosia purpurea (L.) Pers.* Leaf extracts on few common legumes and weeds. *International Journal of Research in Plant Science.***3(1):** 5-9

Putnam,A.R.(1985). WeedAllelopathy.In:Weed Physiology,Reproduction and ecophysiology (Ed.,S.O.Duke) **1:** 131-155.CRC press,Boca Raton,Florida.USA.

Putnam,A.R.(1988). Allelopathy:problems and opportunity in weed management. In:Altirei,M.A. and liebman,M.(eds.), weeds management in agroecosystems: ecological approaches.pp.77-88. Florida.CRC.

Ragan, M. A. and J.S. Craigie. 1978. Phenolic compounds in brown and red algae. Pages 157–179in J.A. Hellebust and J.S. Craigie (eds.), Handbook of phycological methods: Physiological and biochemcial methods. Cambridge University Press, London.

Ram, M., Abdin, M. Z., Khan, M. A., & Jha, P. (2011). HPTLC fingerprint analysis: a quality Control for authentication of herbal phytochemicals. In *High-*

Performance Thin-Layer Chromatography (HPTLC) (pp. 105-116). Springer Berlin Heidelberg.

Ram, M., Khan, M. A., Jha, P., Khan, S., Kiran, U., Ahmad, M. M. & Abdin, M. Z. (2010). HMG-CoA reductase limits artemisinin biosynthesis and accumulation in Artemisia annua L. plants. *Acta physiologiae plantarum*, *32*(5), 859-866.

Ranal, M., Garcia de Santana, D. Ferreira,W. and Mendes-Rodrigues, C.(2009). Calculating germination measurements and organizing spread sheets. *Revista of Brasilian Botany*. **32(4):** 849-855**.**

Rastogi, R.P. and B.N. Mehrotra. 1993. Compendium of Indian Medicinal Plants, CDRI, Luknow and PID, New Delhi, India, Vol.1-3.

Rayalu, J.,Anuradha Laishram , A., Naik, J.,Reddy, S. and D. Jayasimha Rayalu.(2013). Phytochemical analysis, TLC profiling and antimicrobial activity of *Tephrosia purpurea*. *International Journal Of Pharmacy & Life Sciences*. 4**(2):**2375-2379.

Reigosa, M.J.R., Reigosa, M J., Pedrol, N.and González, L. (2006),*Allelopathy: a physiological process with ecological implications*, Springer, p. 1, ISBN 1-4020-4279-5

Ren, Y.Y. and West, C.A. (1992). Elicitation of diterpene biosynthesis in rice (*Oryza sativa* L.) By chitin. *Plant Physiology***. 99**: 1169–1178.

Rice, E.L.(1974).Allelopathy.Academic press,New york.

Rice, E.L.(1984). Allelopathy,2nd ed. Academic press,New york.

Ridenous,W.M. and Callaway,R.M.(2001). The relative importance of allelopathy in interference: the effects of an invasive weed on a native bunchgrass.*Oecologia*.**126**:444-450.

Rizvi, S. J. H., Haque, H., Singh, V. K., & Rizvi, V. (1992). A discipline called allelopathy. In *Allelopathy* (pp. 1-10). Springer Netherlands.

Robbins,W.W.,Crafts, A.S.,and Raynor, R.N.(1942). Weed control.mcgraw-Hill Book Co.New York and London,543 pp.

Robinson,T.(1967). The organic constituents of higher plants.5th edi.Cordus press.North Amherst, Massachusetts.

Rühl, A. T., Donath, T. W., Otte, A., & Eckstein, R. L. (2015). Impacts of short-term germination delay on fitness of the annual weed *Agrostemma githago* (L.). *Arable Weeds in the Face of Global Climate Change*, 67.

Saberi, M., Davari, A., Tarnian, F., Shahreki, M., & Shahreki, E. (2013). Allelopathic effects of *Eucalyptus camaldulensis* on seed germination and initial growth of four range species. *Annals of Biological Research*, ***4*(1):**152-159.

Sasidharan, S., Aravindran, S., Latha, L. Y., Vijenthi, R., Saravanan, D., & Amutha, S. (2010). In vitro antioxidant activity and hepatoprotective effects of *Lentinula edodes* against paracetamol-induced hepatotoxicity. *Molecules*, *15*(6), 4478-4489.

Scott,S.J., Jones R.A. and William W.A., (1984). Review of data analysis methods for seed germination. *Crop Science***(24):** 1192-1199.

Seyed Mohsen S.,Moussavi, N., Hossein, M., Keshavarzi, B. and Gharibdosti, A. (2011). Effect of aqueous extracts of allelopathic *Artemisia annua* on germination and early growth of Isabgol (*Plantago ovate*). *Annals of Biological Research*, **2 (6)**: 687-691.

Shikha R. and Jha A.K. (2016). Allelopathic Effect of Leaf Extract of *Parthenium hysterophorus L.* on Seed Germination and Growth of *Cicer aeritinum L.* **5(3):**652-655.

Shikha, R. and Jha. A.K. (2016). Evaluation of effect of leaf extract of *Parthenium hysterophorus* L. on seed germination, seedling growth and fresh weight of *Phaseolous mungo*. *American Journal of Research Communication*. **4(2):**86-103.

Siddiqui, S., Bhardwaj, S., Khan, S. S., & Meghvanshi, M. K. (2009). Allelopathic effect of different concentration of water extract of *Prosopsis juliflora* leaf on seed

germination and radicle length of wheat (*Triticum aestivum* Var-Lok-1). *American-Eurasian Journal of Scientific Research*, *4*(2), 81-84.

Singh, H.P; Batish , D.R ;Pandher,J.K. And Kohli,R.K. (2005). Phytotoxic effects of *Parthenium hysterophorus* residues on three *Brassica* species. *Weed Biology and Management.* **5(3):** 105-109.

Sisodia, S. And Siddiqui, M.B. (2010). Allelopathic effect by aquaous extracts of different parts of *Croton bonplandianum* Baill. On some crop and weed palnts.*J.Agric.Exten,Rural development.***2**:22-28.

Skrzypek, E., Repka, P., & Barabasz-Krasny, B. (2015). Allelopathic effect of aqueous extracts from the leaves of peppermint (*Mentha*× piperita L.) on selected physiological processes of common sunflower (*Helianthus annuus* L.). *Notulae Botanicae, Horti Agrobotanici, Cluj-Napoca.* ***43*(2):** 335-342.

Skrzypek, E., Repka, P., Stachurska-Swakon, A., Barabasz-Krasny, B., & Mozdzen, K. (2015). Allelopathic effect of aqueous extracts from the leaves of Peppermint. *Notulae Botanicae Horti Agrobotanici Cluj-Napoca*, *43*(2), 335.

Smith,A.M. and Secoy,D.M.(1977).Furrow.13,pp 27.

Smith,R.G. and Menalled, F.D.(2006). Integrated strategies for managing agricultural weeds: making cropping systems less susceptible to weed colonization and establishment.www.ipm.montana.edu.

Sobia T., Sidra M., Javeria H., Maryam H, and Bushra U., (2012). An overview of medicinal importance of *Swertia chirayita*. *International journal of applied science and technology.* **2(1):** 298-304.

Sodaeizadeh, H., & Hosseini, Z. (2012). Allelopathy an environmentally friendly method for weed control. INTECH Open Access Publisher.

Sohmer, S. H., & Gustafson, R. (1987). *Plants and Flowers of Hawai_i*. University of Hawaii Press.

Srivastava, J., Raghava, N., & Raghava, R. P. (2011). Allelopathic potential of *Parthenium* to reduce water absorption in germinating cowpea seeds. *Indian Journal of Scientific Research*, *2*(3), 59.

Steenhagen, D. A., & Zimdahl, R. L. (1979). Allelopathy of leafy spurge (*Euphorbia esula*). *Weed Science*, 1-3.

Sundaramoorthy, S., & Sen, D. N. (1990). Allelopathic effects of *Tephrosia purpurea* Pers. I: On germination and growth of some arid zone crops. *Journal of the Indian Botanical Society*, 69(3-4), 251-255.

Tabassum, S., Mahmood, S., Hanif, J., Hina, M., & Uzair, B. (2012). An overview of medicinal importance of *Swertia chirayita*. *International Journal of Applied*, *2*(1).

Tanveer, A., Safdar, M. E., Tariq, M. A., Yasin, M., & Noorka, I. R. (2014). Allelopathic Inhibition Of Germination And Seedling Vigor Of Some Selected Crops By Achyranthes Aspera L. *Herbologia*, *14*(2).

Tanveer, A.,Safdar , Muhammad,T., Muhammad, Y. and Noorka,I

Teasdale, John R., and Parthan Pillai. (2005) "Contribution of ammonium to stimulation of smooth pigweed (*Amaranthus hybridus* L.) germination by extracts of hairy vetch (*Vicia villosa* Roth) residue." *Weed Biology and Management* 5.1: 19-25.

Tefera, T.(2002). Allelopathic effects of *Parthenium hysterophorus* extracts on seed germination and seedling growth of eragrostis tef. *Journal of agronomy and crop science*.**188**:306-310.

Umer, A., Yousaf, Z., Khan, F., Hussain, U., Anjum, A., Nayyab, Q., & Younas, A. (2010). Evaluation of allelopathic potential of some selected medicinal species. *African Journal of Biotechnology*, 9(37), 6194-6206.

Urbano, B., Andres, F.G. and Ballesteros, A. (2006). Allelopathic potential of cover crops to control weeds in barley. *Allelopathy jornal*.**17 (1):** 53-64.

Vishwakarma, G. S., & Mittal, S. (2014). Bioherbicidal potential of essential oil from leaves of *Eucalyptus tereticornis* against *Echinochloa* crus-galli L. *Journal of Biopesticides*, 7, 47.

Vitonde, S., Thengane, R. J., & Ghole, V. S. (2014). Allelopathic effects of Cassia tora and *Cassia uniflora* on *Parthenium hysterophorous* L. *Journal of Medicinal Plants Research*, *8*(4), 194-196.

Wakjra,M.,Berecha,G. And Tulu,S.(2009).Allelopathic effects of an invasive alien weed *Parthenium hysterophorus* L. compost on lettuce germination and growth. *African Journal of Agricultural Research*.**4 (11):**1325-1330,

Waller,G.R. and Dermer,O.C.(1981).Enzymology of alkaloid metabolism in plants and microorganisms. *The biochemistry of plants*.7:317-402.

Weston, L.A.(1996). Utilization of allelopathy for weed managment in agroecosystems. *Agronomy Journal*.88:860-866.

Willis, R. J. (2007). *The history of allelopathy*. Springer Science & Business Media.

Willis,R.J.(1985). The historical basis of the concept of allelopathy. *Journal of history in biology*.**18**:71-102.

Wldmer, T.L., F. Guermache, My. Dolgovskaia &SY. Reznik. 2007. Enhanced growth and seed properties in introduced vs. native populations of yellow starthistle *(Centaurea solstitialis). Weed Science* 55: 465-473.

Wontermantel,W.M.,Hladky,L.L. and Cortez, A.A.(2009). A new expanded host range of cucurbit tellow stunting disorder virus includes three agriculture crops. *Plant disease*.**93(7):** 685-690.

Wu, H., J. Pratley, J., Lemerle,D., Haig,T. and Verbeek, B.(1998). Differential allelopathic potential among wheat accessions to annual ryegrass. Pp. 567–71. In: Proc. 9th Australian Agron. Conf., Wgga Wagga, NSW, Australia.

Wu, H., Pratley, J., Lemerle, D., Haig, T., An, M. (2001). Screening methods for the evaluation of crop allelopathic potential. *Botanical Review***67:** 403-415.

Xu,M., Galhano,R., Wiemann,P., Bueno,M., Tiernan,M., Wu,W., Chung,M., Gershenzon,J., Tudzynski,B., Sesma,A. And Peters,R.(2012).Genetic evidence for natural product mediated plant–plant allelopathy in rice (*Oryza sativa*). *New Phytologist* .**193**: 570–575.

Yang et al. — Allelopathic phenolics and chlorophyll accumulation Bot. Bull. Acad. Sin. (2004) 45: 119-125 119 Effects of three allelopathic phenolics on chlorophyll accumulation of rice (*Oryza sativa*) seedlings: II. Stimulation of consumption-orientation Chi-Ming Yang1,*, Ing-Feng Chang1 , Shu-Jin Lin1 , and Chang-Hung Chou2

Zhang, T. T., Zheng, C. Y., Hu, W., Xu, W. W., & Wang, H. F. (2010). The allelopathy and allelopathic mechanism of phenolic acids on toxic *Microcystis aeruginosa*. *Journal of applied phycology*, *22*(1), 71-77.

Zhou N. J., Geng C. A., Huang X. Y., Ma Y. B., Zhang X. M., Wang J. L.. (2015). Anti-hepatitis B virus active constituents from *Swertia chirayita*. Fitoterapia. **100:** 27–34.

zu Biesen, C. M. (2010). The rise to prominence of *Artemisia annua* L.–the transformation of a Chinese plant to a global pharmaceutical. *African Sociological Review/Revue Africaine de Sociologie*, *14*(2), 24-46.

Websites :

http://www.fao.org/docrep/009/ag087e/AG087E05.htm