

# Contents

|            | Content  | Page No       |
|------------|--|---------------|
|            | <b><u>Chapter 1</u></b><br><b>Introduction</b>   | <b>1-18</b>   |
| <b>1.1</b> | Liquid Crystal   | <b>1</b>      |
| <b>1.2</b> | History of Liquid Crystals   | <b>1-2</b>    |
| <b>1.3</b> | Classification of Liquid Crystals  | <b>2-11</b>   |
| <b>1.4</b> | Structure-property relation in thermotropic liquid crystal.  | <b>12-15</b>  |
| <b>1.5</b> | References   | <b>16-18</b>  |
|            | <b><u>Chapter 2</u></b><br><b>Schiff base of 4-n-alkoxy-2-hydroxy benzaldehyde with 4-amino acetophenone and their metal complexes: synthesis, characterization and mesomorphic behavior</b> | <b>19-63</b>  |
| <b>2.1</b> | Introduction   | <b>19-27</b>  |
| <b>2.2</b> | Experimental   | <b>28</b>     |
| <b>2.3</b> | Synthesis  | <b>29-34</b>  |
| <b>2.4</b> | Results and discussion   | <b>35-47</b>  |
| <b>2.5</b> | Structurally similar compounds   | <b>48-49</b>  |
| <b>2.6</b> | Conclusion   | <b>50</b>     |
| <b>2.7</b> | References   | <b>51-53</b>  |
|            | Supplementary data   | <b>54-63</b>  |
|            | <b><u>Chapter 3</u></b><br><b>Schiff base of 4-n-alkoxy-2-hydroxy benzaldehyde with 4-amino acetanilide and their metal complexes: synthesis, characterization and mesomorphic behavior</b>  | <b>64-103</b> |
| <b>3.1</b> | Introduction   | <b>64-65</b>  |
| <b>3.2</b> | Experimental   | <b>66</b>     |
| <b>3.3</b> | Synthesis  | <b>67-73</b>  |
| <b>3.4</b> | Results and discussion   | <b>74-83</b>  |
| <b>3.5</b> | Structurally similar compounds   | <b>84-85</b>  |
| <b>3.6</b> | Conclusion   | <b>86</b>     |
| <b>3.7</b> | References   | <b>87-89</b>  |
|            | Supplementary data   | <b>90-103</b> |
|            |  |               |

## Contents

|            |   |                |
|------------|---|----------------|
|            | <b><u>Chapter 4</u></b><br><b>Schiff base of 4-n-alkoxy-2-hydroxy benzaldehyde and 4-n-alkoxy benzaldehyde with thiophene derivatives: synthesis, characterization and mesomorphic behavior</b> | <b>104-194</b> |
| <b>4.1</b> | Introduction  | <b>104-115</b> |
| <b>4.2</b> | Experimental  | <b>116</b>     |
| <b>4.3</b> | Synthesis   | <b>117-129</b> |
| <b>4.4</b> | Results and discussion  | <b>130-148</b> |
| <b>4.5</b> | Comparison of mesomorphic behavior of homologous series-E and series-F  | <b>149-150</b> |
| <b>4.6</b> | Comparison of mesomorphic behavior of homologous series-E and series-A  | <b>151</b>     |
| <b>4.7</b> | Conclusion  | <b>152</b>     |
| <b>4.8</b> | References  | <b>153-159</b> |
|            | Supplementary data  | <b>160-194</b> |
|            | <b><u>Chapter 5</u></b><br><b>Schiff base of 4-n-alkoxy-2-hydroxy benzaldehyde with 1,3 phenylenediamine derivatives : synthesis, characterization and mesomorphic behaviour</b>                | <b>195-255</b> |
| <b>5.1</b> | Introduction  | <b>195-199</b> |
| <b>5.2</b> | Experimental  | <b>200-201</b> |
| <b>5.3</b> | Synthesis   | <b>202-209</b> |
| <b>5.4</b> | Results and discussion  | <b>210-227</b> |
| <b>5.5</b> | Structurally similar compounds  | <b>228-229</b> |
| <b>5.6</b> | Conclusion  | <b>230</b>     |
| <b>5.7</b> | References  | <b>231-237</b> |
|            | Supplementary data  | <b>238-256</b> |
|            | Publications  | <b>257</b>     |
|            | Conferences & Workshops   | <b>258-259</b> |