

CONTENTS

Abstract	(12)
Chapter 1 Introduction	(14)
1.1 Background and literature review	(15)
1.1.1 Drug Delivery Systems and Cancer Therapy	(16)
1.1.2 Understanding the Tumor microenvironment	(16)
1.1.3 Nanoparticles and cancer therapy	(20)
1.1.4 Stimuli responsive materials for cancer therapy	(22)
1.1.5 Carbohydrates as precursors for designing drug carriers	(30)
1.2 Thesis aim and hypothesis	(34)
1.3 References	(36)
Chapter 2 β-cyclodextrin based dual-responsive multifunctional nanotheranostics for cancer cell targeting and dual drug delivery	(45)
2.1 Introduction	(46)
2.2 Experimental Section	(48)
2.2.1 Materials	(48)
2.2.2 Synthesis of nanoconjugates	(48)
2.2.3 Characterization methods	(51)
2.2.4 Preparation of Drug loaded nanoconjugates	(52)
2.2.5 Cell Culture and Cell Viability assay	(53)
2.2.6 Hemolysis assay	(53)
2.2.7 Cellular Internalization	(54)
2.2.8 Cell mortality assay	(54)
2.2.9 In-vivo studies for anticancer potential	(54)
2.3.1 Synthesis of nanoconjugates	(55)
2.3.2 Characterization of nanoconjugates	(55)
2.3.3 Magnetic Hyperthermia Studies	(62)
2.3.4 Drug loading and release studies	(63)
2.3.5 Mathematical modelling of the drug release profiles	(66)
2.3.6 Cell Viability Assay	(76)
2.3.7 Hemolysis Assay	(78)
2.3.8 Internalization test	(80)
2.3.9 Cell mortality test	(80)
2.3.10 In-Vivo Studies	(82)
2.4 Conclusion	(85)
2.5 References	(86)
Chapter 3 Carbon nanotube (CNT) embedded cyclodextrin polymer derived injectable nanocarrier: A multiple faceted platform for stimulation of multi-drug resistance reversal	(89)
3.1 Introduction	(90)

3.2 Experimental Section	(93)
3.2.1 Materials	(93)
3.2.2 Synthesis of nanocarrier-multifunctional cyclodextrin embedded carbon nanotubes	(93)
3.2.3 Characterization	(95)
3.2.4 Determination of Photothermal Efficiency of CD-CNT	(96)
3.2.5 Preparation of Drug loaded CD-CNT	(96)
3.2.6 Drug Release Experiments for CD-CNT	(97)
3.2.7 <i>In-Vitro</i> Studies	(97)
3.2.8 <i>Ex-ovo</i> studies for assessment of antiangiogenic potential	(98)
3.2.9 <i>In-vivo</i> studies for anticancer potential	(99)
3.2.10 Statistical analysis	(99)
3.3 Results and discussion	(100)
3.3.1 Rational design of CD-CNT and its characterization	(100)
3.3.2 <i>In-vitro</i> drug loading and release	(105)
3.3.3 <i>In-Vitro</i> Studies	(108)
3.3.4 Assessment of <i>ex-ovo</i> anti-angiogenesis potential of the nanocarriers	(114)
3.3.5 <i>In-vivo</i> Studies	(117)
3.4 Conclusion	(119)
3.5 References	(120)
Chapter 4 Tumor homing dextran derived amphiphilic self-assembling polymer nanoarchitectures for stimuli responsive drug delivery	(125)
4.1 Introduction	(126)
4.2 Experimental Section	(128)
4.2.1 Materials and Methods	(128)
4.2.2 Synthesis of Dextran-Curcumin amphiphile	(128)
4.2.3 Synthesis of Dextran-Octylamine amphiphile	(130)
4.2.4 Assessment of self-assembly of the amphiphilic polymer	(131)
4.2.5 Encapsulation of anticancer drugs in the nanoarchitectures	(132)
4.2.6 <i>In-Vitro</i> Drug Release Studies for DOX.HCl and curcumin from nanocarriers	(133)
4.2.7 <i>In-vitro</i> studies	(134)
4.2.8 <i>In-vivo</i> studies	(134)
4.2.9 Statistical analysis	(135)
4.3 Result and Discussions	(135)
4.3.1 Designing of dextran derived amphiphilic polymers and their self-assembly into various nanoarchitectures	(135)
4.3.2 Analyzing the self-assembling phenomenon of the amphiphiles and drug loading capability	(142)
4.3.3 Detailed analytical characterization of the bare and drug loaded nanoarchitectures	(145)
4.3.4 <i>In-vitro</i> drug release from nanoarchitectures under conditions of mimicked tumor microenvironment (enzyme and pH responsive release)	(150)
4.3.5 Preclinical evaluation of the efficacy of nanoarchitectures for stimuli responsive anticancer therapy	(154)
4.4 Conclusion	(167)

4.5 References	(168)
Chapter 5 Targeting the tumor microenvironment via hierarchical disassembly of curcumin micelles: The pro-drug strategy	(172)
5.1 Introduction	(173)
5.2 Experimental Section	(175)
5.2.1 Materials	(175)
5.2.2 Characterization	(175)
5.2.3 Synthesis of curcumin derived amphiphilic polymers	(176)
5.2.4 Formation of micelles via self-assembly of amphiphiles and determination of its critical micelle concentration (CMC)	(179)
5.2.5 Study of pH-triggered hydrolysis of the acetal linkage in micelles	(179)
5.2.6 Assessment of pH responsive degradation of curcumin prodrug micelles	(179)
5.2.7 Monitoring size of the change of acetal linkage containing micelles in response to acidic pH	(180)
5.2.8 Drug Loading and Release Studies	(180)
5.2.9 <i>In-vitro</i> Studies	(181)
5.2.10 <i>In-vivo</i> Studies	(182)
5.2.11 Statistical analysis	(182)
5.3 Result and Discussion	(182)
5.3.1 Synthesis of various curcumin derived amphiphilic polymers	(188)
5.3.2 Self-assembly of amphiphiles into micelles	(191)
5.3.3 Hydrolysis of the acetal linkage containing polymers in response to pH variations	(196)
5.3.4 Loading and in Vitro Release of DOX in response to pH and enzyme	(198)
5.3.5 Targeting ability and antitumor activity of DOX loaded targeting micelles and a comparison of responsive disassembling micelles with non-responsive one	(206)
5.3.6 Combination Index Analysis	(209)
5.3.7 Assessment of the efficacy of self-therapeutic micelles as drug delivery system for anti-cancer effect in vivo	(213)
5.3.8 Mechanism of action of pro-drug micelles as devised from analytical and pre-clinical evaluations	(217)
5.4 Conclusion	(219)
5.5 References	(219)
Chapter 6: Summary and Future Outlook	(220)