

Contents

1	Introduction.....	3
1.1	Bioenergy and biofuels.....	3
1.2	Microalgae as the prominent source of biofuel and nutraceuticals.....	4
1.3	Factors influencing the biofuel production in microalgae	8
1.3.1	Environmental factors	9
1.3.2	Carbon source availability	12
1.3.3	Cultivation strategy	14
1.4	Metabolite synthesis pathways and their regulation in microalgae.....	18
1.4.1	Starch metabolism.....	18
1.4.2	Lipid metabolism	20
1.4.3	Biosynthesis of photosynthetic pigments.....	24
1.4.4	Role of carbon and nitrogen metabolism in regulating metabolite production	25
1.4.5	Role of signaling pathways in lipid metabolism.....	28
1.5	Techniques used to study the production of starch, chlorophyll, carotenoids, and lipids.....	31
1.5.1	UV-VIS Spectrophotometry	31
	High-performance liquid chromatography (HPLC).....	32
1.5.2	Fluorescence spectroscopy.....	34
1.5.3	Raman spectroscopy (RS).....	35
1.5.4	Microscopy	37
1.6	Single-cell analysis in microalgal biofuel production.....	38
1.7	The model microalga- <i>Chlamydomonas</i>	40
1.8	The foundation of the study	43
1.9	References	44
2	Effect of salt stress and varying cultivation strategies on bioenergy feedstock production in <i>Chlamydomonas reinhardtii</i>	65
2.1	Abstract	65
2.2	Introduction	66
2.3	Materials and Methods	68

2.3.1	Growth and maintenance of <i>Chlamydomonas reinhardtii</i> CC-125	68
2.3.2	Cultivation strategies of <i>C. reinhardtii</i> under salt stress.....	68
2.3.3	Cell count using a haematocytometer	70
2.3.4	Analysis of total photosynthetic pigments	70
2.3.5	Estimation of total starch content using Lugol's iodine	72
2.3.6	Estimation of neutral lipid content using Nile red dye	72
2.3.7	Epifluorescence live-cell imaging of lipid droplets	72
2.3.8	Total RNA extraction and quantitative real-time PCR	73
2.3.9	Statistical analysis	75
2.4	Results	75
2.4.1	Determination of cell density from OD ₆₈₀	75
2.4.2	Growth and cell density production in <i>C. reinhardtii</i> under salt stress ..	76
2.4.3	Chlorophyll and carotenoid production in <i>C. reinhardtii</i> under salt stress.....	79
2.4.4	Starch and lipid production in <i>C. reinhardtii</i> under salt stress	82
2.4.5	Expression analysis of genes involved in growth, autophagy, and <i>de novo</i> lipid synthesis in <i>C. reinhardtii</i>	86
2.4.6	Cell morphological changes in <i>C. reinhardtii</i> under salt stress	88
2.5	Discussion	106
2.6	Conclusion.....	113
2.7	References	114

3	Role of carbon concentration and mode of cultivation on growth and accumulation of lipid, starch, and photosynthetic pigments in <i>Chlamydomonas reinhardtii</i> under heterotrophy and mixotrophy	123
3.1	Abstract	123
3.2	Introduction	124
3.3	Materials and Methods	126
3.3.1	Microalgal growth conditions	126
3.3.2	Heterotrophic and mixotrophic cultivation strategies of <i>C. reinhardtii</i>	127
3.3.3	Measurement of cell density and growth parameters in <i>C. reinhardtii</i>	129
3.3.4	Analysis of total photosynthetic pigments.....	130
3.3.5	Estimation of total starch content using Lugol's iodine	130
3.3.6	Triacylglycerol estimation using Nile red fluorescence	130
3.3.7	Live-cell fluorescence imaging of lipid droplets	131

3.3.8	Total RNA extraction and quantitative real-time PCR	131
3.3.9	Statistical analysis	132
3.4	Results	132
3.4.1	Effect of sodium acetate concentration and mode of cultivation on hetero/mixotrophic growth of <i>C. reinhardtii</i>	132
3.4.2	Chlorophyll and carotenoid production in <i>C. reinhardtii</i> under hetero/mixotrophy	136
3.4.3	Starch and lipid production in <i>C. reinhardtii</i> under hetero/mixotrophy	139
3.4.4	Study of genes affecting growth, autophagy, and lipid production in <i>C. reinhardtii</i> under mixotrophy	143
3.4.5	Morphological changes and lipid droplet biogenesis in <i>C. reinhardtii</i> under mixotrophy	146
3.5	Discussion	150
3.6	Conclusion	157
3.7	References	158
4	Developing the non-invasive use of micro-Raman spectroscopy for the determination of biomolecular composition in <i>Chlamydomonas reinhardtii</i>	169
4.1	Abstract	169
4.2	Introduction	170
4.3	Materials and Methods	172
4.3.1	Microalgal strain, growth media, and culture conditions	172
4.3.2	Micro-Raman spectroscopy of <i>C. reinhardtii</i> cells	174
4.3.3	Signal processing of micro-Raman spectra	174
4.3.4	Micro-Raman spectral data analysis	175
4.3.5	Statistical analysis	176
4.4	Results	177
4.4.1	Assignment of Raman modes to cellular biomolecules	177
4.4.2	Validation of the Raman peak intensity with Nile red fluorescence representing the lipid production in <i>C. reinhardtii</i> cells under salt stress	185
4.4.3	Biomolecular correlation analysis by Raman spectroscopy in <i>C. reinhardtii</i> under salt stress	187
4.4.4	Chemical composition and population distribution heterogeneities of carotenoids and chlorophyll in <i>C. reinhardtii</i> under salt stress	190
4.4.5	Characterization and distribution heterogeneities of lipids produced by <i>C. reinhardtii</i> under salt stress	197

4.5	Discussion	208
4.6	Conclusion.....	215
4.7	References	216
5	Summary and Future Directions	225
5.1	Summary	225
5.2	Future Directions.....	229
Appendix		
Appendix.1.	Composition of Tris-Acetate Phosphate (TAP) medium.....	235
A1.1	Composition of Tris-Acetate Phosphate (TAP) medium.....	237
Appendix 2.	qRT-PCR: Primer specificity, amplification, and melting plots.....	241
A2.1.	Primer specificity.....	243
A2.2.	Amplification plots obtained in qRT-PCR.....	244
A2.3.	Melting plots obtained in qRT-PCR.....	245
Appendix 3.	Heterogeneities in the $\nu(\text{C-C})$ Raman mode cultivated under salt stress.....	247
A3.1.	Heterogeneities in the $\nu(\text{C-C})$ Raman mode in <i>C. reinhardtii</i> cultivated under salt stress.....	249
Appendix 4.	Pearson's correlation analysis for cellular components under salt stress.....	251
A4.1.	Pearson's correlation coefficients for cellular components in <i>C. reinhardtii</i> under salt stress obtained from Raman spectroscopy.....	253
Appendix 5.	Publications and Presentations.....	257
A5.1.	List of publications.....	259
A5.2.	List of presentations.....	264
Appendix 6.	Copyright Permissions.....	267
A6.1.	Copyright permissions for Figure 1.4.....	269
A6.2.	Copyright permission for Figure 1.7.....	270
A6.3.	Copyright permission for Figure 1.10.....	272
A6.4.	Copyright permission for Figure 1.11.....	273
A6.5.	Copyright permission for Publication 1.....	275
A6.6.	Copyright permission for Publication 2.....	277

List of Figures

Figure 1.1. Four generations of biofuel as based on the feedstock used for biofuel production.	3
Figure 1.2. Microscopic images of microalgae involved in biofuel production.	6
Figure 1.3. Factors governing biofuel production in microalgae and their impact on the environment and society.	8
Figure 1.4. A model representing morphological changes in <i>Chlamydomonas</i> sp. under saline stress.	11
Figure 1.5. Starch metabolism in microalgae	19
Figure 1.6. TAG biosynthesis pathway in microalgae.	22
Figure 1.7. Lipid catabolism and lipid droplet degradation in <i>Chlamydomonas</i> sp. ...	23
Figure 1.8. Interconnection between metabolic pathways of photosynthesis, lipid synthesis, protein metabolism, and starch metabolism in microalgae.	27
Figure 1.9. ROS-mediated regulation of autophagy and lipid production in microalgae.	29
Figure 1.10. Autophagosome formation in <i>Chlamydomonas</i> sp.	30
Figure 1.11. The loop between population-level and single-cell-level studies.	39
Figure 1.12. Diagrammatic representation of the structure of <i>Chlamydomonas reinhardtii</i> and the Cryo-EM FIB of frozen cells.	42
Figure 2.1. A schematic representation of salt stress addition in single-stage and two-stage methods of cultivation to achieve the final concentration of 100 mM NaCl.	Error! Bookmark not defined.
Figure 2.2. A schematic representation of single-stage, two-stage, and gradient strategy of cultivation in presence of 150 mM and 200 mM NaCl.	70
Figure 2.3. Linear plot of cell density measured using haematocytometer at the corresponding OD ₆₈₀	75
Figure 2.4. Effect of different modes of application of 100 mM NaCl on the growth of <i>C. reinhardtii</i> CC-125.	77
Figure 2.5. Effect of different modes of application of 150 mM and 200 mM NaCl on the growth of <i>C. reinhardtii</i> CC-125.	78
Figure 2.6. Effect of different modes of 100 mM NaCl application on the photosynthetic pigment production in <i>C. reinhardtii</i> CC-125.	80

Figure 2.7. Effect of different modes of 150 mM NaCl application on the photosynthetic pigment production in <i>C. reinhardtii</i> CC-125.....	80
Figure 2.8. Effect of different modes of 200 mM NaCl application on the photosynthetic pigment production in <i>C. reinhardtii</i> CC-125.....	81
Figure 2.9. Starch production in <i>C. reinhardtii</i> in presence of 100 mM NaCl.....	83
Figure 2.10. TAG production in <i>C. reinhardtii</i> under 100 mM NaCl as detected by Nile red fluorescence microscopy.....	84
Figure 2.11. Starch and TAG production in <i>C. reinhardtii</i> in presence of 150 mM NaCl.....	84
Figure 2.12. Starch and TAG production in <i>C. reinhardtii</i> in presence of 200 mM NaCl.....	85
Figure 2.13. Fold change in expression of genes affecting the growth, autophagy, and de novo lipid synthesis in <i>C. reinhardtii</i> under salt stress.....	87
Figure 2.14. Bright-field and Nile red stained epifluorescence microscopic images of <i>C. reinhardtii</i> observed at 100 X, grown in presence of 100 mM NaCl.....	90
Figure 2.15. Probability distribution of the size of <i>C. reinhardtii</i> cells when grown in presence of 100 mM NaCl.....	91
Figure 2.16. Probability distribution of the shape of <i>C. reinhardtii</i> cells when grown in presence of 100 mM NaCl.....	92
Figure 2.17. Probability distribution of the size of lipid droplets produced by <i>C. reinhardtii</i> in presence of 100 mM NaCl.....	93
Figure 2.18. Bright-field and Nile red epifluorescence microscopic images of <i>C. reinhardtii</i> observed at 100 X, grown in presence of 150 mM NaCl.....	95
Figure 2.19. Probability distribution of the size of <i>C. reinhardtii</i> cells when grown in presence of 150 mM NaCl.....	96
Figure 2.20. Probability distribution of the shape of <i>C. reinhardtii</i> cells when grown in presence of 150 mM NaCl.....	97
Figure 2.21. Probability distribution of the size of lipid droplets produced by <i>C. reinhardtii</i> in presence of 150 mM NaCl.....	98
Figure 2.22. Probability distribution of the number of lipid droplets distributed per area of the <i>C. reinhardtii</i> cell in presence of 150 mM NaCl.....	101
Figure 2.23. Probability distribution of the percentage area of the cell occupied by the lipid droplets produced by <i>C. reinhardtii</i> in presence of 150 mM NaCl.....	102

Figure 2.24. Fluctuations in the droplet growth dynamic properties in <i>C. reinhardtii</i> under the presence of 150 mM NaCl.	103
Figure 2.25. Spatial segregation of lipid droplets in the <i>C. reinhardtii</i> cells observed under the presence of 150 mM NaCl.	104
Figure 2.26. Liquid-liquid phase separation in lipid droplet biogenesis in <i>C. reinhardtii</i> under the presence of 150 mM NaCl.	105
Figure 2.27. Summary of the salt stress-induced effects on the growth, morphology, and metabolite state of the <i>C. reinhardtii</i> cell.	112
Figure 3.1. A schematic representation of hetero/mixotrophy cultivation strategies employed on <i>C. reinhardtii</i>	128
Figure 3.2. Growth curve fitting of the autotrophic control to determine lag time. ..	129
Figure 3.3. Cell density accumulation in <i>C. reinhardtii</i> culture under single-stage and gradient mode of hetero/mixotrophic cultivation.	135
Figure 3.4. Chlorophyll and carotenoid production in <i>C. reinhardtii</i> at different stages of the growth phase when grown heterotrophically in presence of sodium acetate.	137
Figure 3.5. Photosynthetic pigment accumulation in <i>C. reinhardtii</i> at different stages of the growth phase when grown mixotrophically in presence of sodium acetate.	138
Figure 3.6. Starch and triacylglycerol production in <i>C. reinhardtii</i> at different concentrations of sodium acetate and mode of heterotrophic cultivations.	141
Figure 3.7. Starch and triacylglycerol accumulation in <i>C. reinhardtii</i> at different stages of growth phase under mixotrophy in presence of sodium acetate.	142
Figure 3.8. Fold change in expression of genes affecting growth, autophagy, and lipid production in <i>C. reinhardtii</i> when grown mixotrophically in presence of sodium acetate.	145
Figure 3.9. Bright field and Nile red fluorescence images showing morphological features of mixotrophically grown <i>C. reinhardtii</i> cells.	148
Figure 3.10. Liquid-liquid phase separation in lipid droplet biogenesis in <i>C. reinhardtii</i> under mixotrophy.	149
Figure 3.11. Salient features of bioenergy feedstock production in <i>Chlamydomonas reinhardtii</i> under hetero/mixotrophy.	156
Figure 4.1. Tree diagram showing single-stage and two-stage strategies for salt stress.	173

Figure 4.2. Discrete wavelet transform (DWT) denoising of the Raman signal.	179
Figure 4.3. Deconvolution of the Raman spectra of <i>C. reinhardtii</i> cells yields distinct Raman peaks.	180
Figure 4.4. Representative peak centers of background-subtracted Raman signal before and after denoising.....	181
Figure 4.5. Deconvolution of the carotenoid peak at 1523 cm ⁻¹	182
Figure 4.6. Signal integrity and accuracy post-DWT treatment of the Raman spectra of <i>C. reinhardtii</i>	183
Figure 4.7. Probability distribution of lipid molecules in <i>C. reinhardtii</i> cells under salt stress measured by Raman spectroscopy and fluorescence microscopy.....	186
Figure 4.8. Correlation matrix of the cellular components of <i>C. reinhardtii</i> cultivated under different modes of salt stress.....	189
Figure 4.9. Heterogeneities in carotenoid chemical composition in <i>C. reinhardtii</i> under salinity stress as represented by the Raman peak positions.	191
Figure 4.10. Population-scale probability distribution of peak intensity obtained at peak center corresponding to carotenoids in <i>C. reinhardtii</i> under salinity stress.	194
Figure 4.11. Changes in Raman scattering intensity of chlorophyll mode in <i>C. reinhardtii</i> under salinity stress.	195
Figure 4.12. Pearson's correlation of chlorophyll and carotenoid as obtained from Raman modes in <i>C. reinhardtii</i> under salt stress.	196
Figure 4.13. Intensity fluctuations of Raman modes of chlorophyll and carotenoids obtained in <i>C. reinhardtii</i> under salt stress.	196
Figure 4.14. Biofuel properties of lipids produced by <i>C. reinhardtii</i> under different modes of salt stress.	199
Figure 4.15. Raman spectra of Triolein and microalgal oil after denoising using the DWT approach.....	200
Figure 4.16. Probability distribution of the ratio of I _α (CH ₂ /CH ₃) and I _{C=O} (I _β), as a function of incubation time and mode of application of salt stress in <i>C. reinhardtii</i>	202
Figure 4.17. Raman intensity at wavenumbers ~1440 and ~1656 cm ⁻¹ for peaks.	205
Figure 4.18. Population-scale heterogeneities in the lipid CH ₂ /CH ₃ scissoring mode and C=C stretching mode under salt stress in <i>C. reinhardtii</i>	206

Figure 4.19. Degree of unsaturation of lipids produced by <i>C. reinhardtii</i> under different modes of salt stress.....	207
Figure 4.20. The overall impact of the aging culture of <i>C. reinhardtii</i> and salt stress on lipid profile, carotenoid composition, and chlorophyll content detected by micro-Raman spectroscopy	214
Figure 5.1. Heat map representing the total yield of bioenergy feedstock obtained from <i>C. reinhardtii</i> grown under salt stress and hetero/mixotrophic cultivations.	228

Appendix

Figure A2.1. Gene amplification using mRNA-specific primers on <i>C. reinhardtii</i> genome	243
Figure A2.2. Representative amplification plots of qRT-PCR	244
Figure A2.3. Representative melting plots of qRT-PCR	245
Figure A3.1. Heterogeneities in the $\nu(\text{C-C})$ Raman mode in a population of 100 cells of <i>C. reinhardtii</i> cultivated under salt stress	249
Figure A4.1 Relationship among the cellular components as a function of culture age and salt stress in <i>C. reinhardtii</i>	254