

List of Tables

- Table 3.1** Calculated value of Adsorption energy (E_{ad}), LUMO energy (E_{LUMO}), Fermi level (E_F), HOMO energy (E_{HOMO}), HOMO-LUMO gap (E_G), relative change in E_F (ΔE_{FR}), relative change in E_G (ΔE_{GR}) and adsorption distance(d) of Adenine, Cytosine, Guanine, Thymine, and Uracil adsorbed over C_{24} fullerene, respectively (50)
- Table 3.2** The NBO second-order perturbation energy (E^2) measured in kcal/mol) signifies the extent of charge transfer occurring between C_{24} fullerene and each of the nucleobases: Adenine, Cytosine, Guanine, Thymine, and Uracil (52)
- Table 3.3** The computed variations in E_G (ΔE_G), recovery time (τ), and sensing response (S) for Adenine, Cytosine, Guanine, Thymine, and Uracil when they are adsorbed onto C_{24} fullerene, respectively (59)
- Table 3.4** Calculated value of Adsorption energy (E_{ad}), LUMO energy (E_{LUMO}), HOMO energy (E_{HOMO}), HOMO-LUMO gap (E_G), Recovery Time (τ) and Sensing Response (S) for Adenine, Cytosine, Guanine, Thymine and Uracil adsorbed C_{24} fullerene with solvent effect, respectively..... (61)
- Table 3.5** Dipole moments of Adenine, Cytosine, Guanine, Thymine, and Uracil when adsorbed onto C_{24} fullerene, both in the absence (μ_{gas}) and presence (μ_{water}) of solvent effect..... (62)
- Table 4.6** Calculated value of Adsorption energy (E_{ad}), LUMO energy (E_{LUMO}), Fermi level (E_F), HOMO energy (E_{HOMO}), HOMO-LUMO gap (E_G), relative change in E_F (ΔE_{FR}), relative change in E_G (ΔE_{GR}) and adsorption distance(d) of AMP, KET and MER adsorbed over C_{24} fullerene, respectively..... (73)
- Table 4.7** The NBO second-order perturbation energy (E^2 , kcal/mol) corresponds to the charge transfer between the C_{24} fullerene and AMP, KET, and MER, respectively..... (74)
- Table 4.8** Computed variations in E_G (ΔE_G), recovery time (τ), and sensing response (S) for Adenine, Cytosine, Guanine, Thymine, and Uracil when they are adsorbed onto C_{24} fullerene, respectively..... (79)
- Table 4.9:** Calculated value of Adsorption energy (E_{ad}), LUMO energy (E_{LUMO}), HOMO energy (E_{HOMO}), HOMO-LUMO gap (E_G), Recovery Time (τ) and Sensing Response (S) for AMP, KET, and MER adsorbed C_{24} fullerene with solvent effect, respectively..... (81)

Table 4.10	Comparison of the sensing property of C ₂₄ fullerene with other pristine nano-materials.....	(82)
Table 4.6	Detail of the simulation boxes which used in this work.....	(83)
Table 5.11	Calculated Values of HOMO-LUMO gap (E _G), cohesive energy (E _B), and Bond length of C ₂₄ , C ₃₆ , C ₅₀ and C ₇₀ fullerene.....	(95)
Table 5.2	Calculated value of Adsorption energy (E _{ad}), LUMO energy (E _{LUMO}), Fermi level (E _F), HOMO energy (E _{HOMO}), HOMO-LUMO gap (E _G), relative change in E _F (ΔE _{FR}), relative change in E _G (ΔE _{GR}) and adsorption distance(d) of LEU adsorbed over C ₂₄ , C ₃₆ , C ₅₀ and C ₇₀ fullerene, respectively.....	(100)
Table 5.3	The NBO second-order perturbation energy (E ² , kcal/mol) corresponds to the charge transfer between the LEU and C ₂₄ , C ₃₆ , C ₅₀ and C ₇₀ fullerene, respectively.....	(101)
Table 5.4	Calculated difference in E _G (ΔE _G), Recovery Time (τ) and Sensing Response(S) for LEU over C ₂₄ , C ₃₆ , C ₅₀ and C ₇₀ fullerene, respectively(T=310K)	(107)
Table 5.5	Global indices parameter of systems.....	(109)
Table 5.6	Calculated value of Adsorption energy (E _{ad}), LUMO energy (E _{LUMO}), HOMO energy (E _{HOMO}), HOMO-LUMO gap (E _G), Recovery Time (τ) and Sensing Response (S) for LEU adsorbed C ₂₄ , C ₃₆ , C ₅₀ and C ₇₀ fullerene with solvent effect, respectively.....	(110)
Table 5.7	Detail of the simulation boxes which used in this work.....	(111)