

8. Reprints of Published Articles

RESEARCH

Open Access



TNF- α differentially modulates subunit levels of respiratory electron transport complexes of ER/PR +ve/–ve breast cancer cells to regulate mitochondrial complex activity and tumorigenic potential

Anjali Shinde^{1†}, Hyeryeon Jung^{2†}, Hayun Lee^{2†}, Kritarth Singh^{3†}, Milton Roy¹, Dhruv Gohel¹, Han Byeol Kim², Minal Mane¹, Hitesh Vasiyani¹, Fatema Currim¹, Yu Ri Seo², Seojin Yang², Ara Cho², Eugene C. Yi^{2*} and Rajesh Singh^{1*} 

Abstract

Background: Tumor necrosis factor- α (TNF- α) is an immunostimulatory cytokine that is consistently high in the breast tumor microenvironment (TME); however, its differential role in mitochondrial functions and cell survival in ER/PR +ve and ER/PR –ve breast cancer cells is not well understood.

Methods: In the current study, we investigated TNF- α modulated mitochondrial proteome using high-resolution mass spectrometry and identified the differentially expressed proteins in two different breast cancer cell lines, ER/PR positive cell line; luminal, MCF-7 and ER/PR negative cell line; basal-like, MDA-MB-231 and explored its implication in regulating the tumorigenic potential of breast cancer cells. We also compared the activity of mitochondrial complexes, ATP, and ROS levels between MCF-7 and MDA-MB-231 in the presence of TNF- α . We used Tumor Immune Estimation Resource (TIMER) webserver to analyze the correlation between TNF- α and mitochondrial proteins in basal and luminal breast cancer patients. Kaplan-Meier method was used to analyze the correlation between mitochondrial protein expression and survival of breast cancer patients.

(Continued on next page)

* Correspondence: euyi@snu.ac.kr; singhraj1975@gmail.com

[†]Anjali Shinde, Hyeryeon Jung, Hayun Lee and Kritarth Singh contributed equally to this work.

²Department of Molecular Medicine and Biopharmaceutical Sciences, Graduate School of Convergence Science and Technology, Seoul National University, Seoul 03080, South Korea

¹Department of Bio-Chemistry, Faculty of Science, The Maharaja Sayajirao University of Baroda, Sayajigunj, Vadodara, Gujarat 390002, India
Full list of author information is available at the end of the article



© The Author(s). 2021 **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.



The analog of cGAMP, c-di-AMP, activates STING mediated cell death pathway in estrogen-receptor negative breast cancer cells

Hitesh Vasiyani¹ · Anjali Shinde¹ · Milton Roy¹ · Minal Mane¹ · Kritarth Singh² · Jyoti Singh¹ · Dhruv Gohel¹ · Fatema Currim¹ · Khushali Vaidya³ · Mahesh Chhabria³ · Rajesh Singh¹

Accepted: 30 March 2021 / Published online: 10 April 2021
© The Author(s), under exclusive licence to Springer Science+Business Media, LLC, part of Springer Nature 2021

Abstract

Immune adaptor protein like STING/MITA regulate innate immune response and plays a critical role in inflammation in the tumor microenvironment and regulation of metastasis including breast cancer. Chromosomal instability in highly metastatic cells releases fragmented chromosomal parts in the cytoplasm, hence the activation of STING via an increased level of cyclic dinucleotides (cDNs) synthesized by cGMP-AMP synthase (cGAS). Cyclic dinucleotides 2' 3'-cGAMP and its analog can potentially activate STING mediated pathways leading to nuclear translocation of p65 and IRF-3 and transcription of inflammatory genes. The differential modulation of STING pathway via 2' 3'-cGAMP and its analog and its implication in breast tumorigenesis is still not well explored. In the current study, we demonstrated that c-di-AMP can activate type-1 IFN response in ER negative breast cancer cell lines which correlate with STING expression. c-di-AMP binds to STING and activates downstream IFN pathways in STING positive metastatic MDA-MB-231/MX-1 cells. Prolonged treatment of c-di-AMP induces cell death in STING positive metastatic MDA-MB-231/MX-1 cells mediated by IRF-3. c-di-AMP induces IRF-3 translocation to mitochondria and initiates Caspase-9 mediated cell death and inhibits clonogenicity of triple-negative breast cancer cells. This study suggests that c-di-AMP can activate and modulates STING pathway to induce mitochondrial mediated apoptosis in estrogen-receptor negative breast cancer cells.

Keywords Stimulator of interferon gene (STING) · Cyclic GMP AMP synthase (cGAS) · Interferon regulatory factor3 (IRF-3) · Apoptosis · Cyclic dinucleotides (cDNs)

Introduction

The crosstalk between tumor cells, infiltrating immune cells and stroma in breast cancer tumor microenvironment (TME) provides an optimal niche for the growth and proliferation of cancer cells [1]. Hypoxic TME of solid tumors promotes the clonal evolution of the cancer cells which leads to the progression of the tumor [2]. Hypoxic TME can also induce necrotic cell death leading to the release of intrinsic danger-associated molecular patterns (DAMPs), which can activate

innate immune response [3]. The activation of the innate immune system and its regulation during tumorigenesis is emerging [4] however, its role in the acquisition of tumorigenic phenotype, its physiological and chemical modifiers are not well understood.

Our previous reports demonstrated that innate immune regulators are uniquely positioned at mitochondria which in turn links the inflammatory pathways and metabolism, hence playing an important role in the metabolic adaption of tumor cells [5]. STING (Stimulator of interferon gene) is also known as MITA, MPYS, TMEM 173 is localized at the ER/mitochondria contact site and is a major regulator of the type I immune response. Interestingly, STING is differentially expressed in ER/PR positive and negative breast cancer patients, therefore can differentially regulate inflammatory cell death [6]. The implication of increased level of STING in triple-negative breast cancer cells and association with metastasis and resistance to cell death is not well understood.

✉ Rajesh Singh
singhraj1975@gmail.com

¹ Department of Biochemistry, The M.S. University of Baroda, Vadodara, Gujarat 390002, India

² Department of Cell and Developmental Biology, University College London, Gower Street, London WC1E 6BT, UK


³ L. M. College of Pharmacy, Navrangpura, Ahmedabad, Gujarat 380 009, India



ELSEVIER

Contents lists available at ScienceDirect

Cellular Signalling

journal homepage: www.elsevier.com/locate/cellsig

TRIM32 regulates mitochondrial mediated ROS levels and sensitizes the oxidative stress induced cell death

Paresh Prajapati^{a,b,c}, Dhruv Gohel^a, Anjali Shinde^a, Milton Roy^a, Kritarth Singh^{a,d}, Rajesh Singh^{a,*}

^a Department of Biochemistry, Faculty of Science, The M.S. University of Baroda, Vadodara 390 002, Gujarat, India

^b Spinal Cord and Brain Injury Research Center, Department of Pathology and Laboratory Medicine, University of Kentucky, 800 S. Limestone, Lexington, KY 40536, USA

^c Neuroscience, University of Kentucky, Lexington, KY 40536, USA

^d Department of Cell and Developmental Biology, University College London, Gower Street, London WC1E 6BT, UK



ARTICLE INFO

Keywords:

Cell death
Mitochondria
Oxidative stress
ubiquitination
TRIM32
XIAP

ABSTRACT

Emerging evidence suggests that ubiquitin mediated post translational modification is a critical regulatory process involved in diverse cellular pathways including cell death. During ubiquitination, E3 ligases recognize target proteins and determine the topology of ubiquitin chains. Recruitment of E3 ligases to targets proteins under stress conditions including oxidative stress and their implication in cell death have not been systemically explored. In the present study, we characterized the role of TRIM32 as an E3 ligase in regulation of oxidative stress induced cell death. TRIM32 is ubiquitously expressed in cell lines of different origin and form cytoplasmic speckle like structures that transiently interact with mitochondria under oxidative stress conditions. The ectopic expression of TRIM32 sensitizes cell death induced by oxidative stress whereas TRIM32 knockdown shows a protective effect. The turnover of TRIM32 is enhanced during oxidative stress and its expression induces ROS generation, loss of mitochondrial transmembrane potential and decrease in complex-I activity. The pro-apoptotic effect was rescued by pan-caspase inhibitor or antioxidant treatment. E3 ligase activity of TRIM32 is essential for oxidative stress induced apoptotic cell death. Furthermore, TRIM32 decreases X-linked inhibitor of apoptosis (XIAP) level and overexpression of XIAP rescued cells from TRIM32 mediated oxidative stress and cell death. Overall, the results of this study provide the first evidence supporting the role of TRIM32 in regulating oxidative stress induced cell death, which has implications in numerous pathological conditions including cancer and neurodegeneration.

1. Introduction

Mitochondria are dynamic organelles and are implicated in various cellular functions including metabolism, cell death, inflammation, and immunity apart from its role in bioenergetics to maintain cellular homeostasis [1–3]. The equilibrium of the healthy mitochondrial network is maintained in the cells through the dynamic process of fusion and fission. The stressed or damaged mitochondria are labeled with ubiquitin (Ub) and selectively degraded through the process known as mitophagy in order to remove them from the healthy network [4]. Any defect in mitophagy or in the fusion and fission process leads to accumulation of defective mitochondria, resulting in the production of excessive ROS and initiates cell death [5]. This phenomenon has been observed in many pathological conditions including neurodegeneration and autoimmune diseases [6,7].

Ub is versatile molecule that can form different types of Ub chains of

different topology on the target proteins through seven conserved lysine residues [8,9]. The presence of atypical Ub chains on the target proteins regulate their stability and impart unique functional outcomes. The reported evidence suggests that ubiquitination during oxidative stress is initiated on the mitochondria which regulates mitophagy and cell death [10]. It was observed that K63 linked ubiquitination is initiated through Bre1 ubiquitin ligase during oxidative stress that determines cell survival [11]. The process of ubiquitination is achieved by the sequential action of three enzymes: E1 (Ub activating enzyme), E2 (Ub conjugating enzyme), E3 (Ub ligases). The E3 ligases are terminal proteins in ubiquitination process and provide pathway specificity as they recognize the substrate and initiate the transfer of Ub. Interestingly, the human genome contains more than 1000 E3 Ligases and their functions are not well understood. Moreover, the role of specific E3 Ligases, their recruitment to mitochondria and regulation of cell death pathways during oxidative environment have not been well studied [12].

* Corresponding author at: Department of Biochemistry, Faculty of Science, The Maharaja Sayajirao University of Baroda, Vadodara, Gujarat 390002, India.
E-mail address: rajesh.singh-biochem@msubaroda.ac.in (R. Singh).

<https://doi.org/10.1016/j.cellsig.2020.109777>

Received 22 June 2020; Received in revised form 6 September 2020; Accepted 7 September 2020

Available online 10 September 2020

0898-6568/ © 2020 Elsevier Inc. All rights reserved.



TNF- α -induced E3 ligase, TRIM15 inhibits TNF- α -regulated NF- κ B pathway by promoting turnover of K63 linked ubiquitination of TAK1

Milton Roy, Kritarth Singh¹, Anjali Shinde, Jyoti Singh, Minal Mane, Sawani Bedekar, Yamini Tailor, Dhruv Gohel, Hitesh Vasiyani, Fatema Currim, Rajesh Singh*

Department of Biochemistry, Faculty of Science, The MS University of Baroda, Vadodara, Gujarat 390002, India

ARTICLE INFO

Keywords:

TNF- α
NF- κ B
TRIM15
TRIM8
Ubiquitination
Functional antagonism

ABSTRACT

Ubiquitin E3-ligases are recruited at different steps of TNF- α -induced NF- κ B activation; however, their role in temporal regulation of the pathway remains elusive. The study systematically identified TRIMs as potential feedback regulators of the TNF- α -induced NF- κ B pathway. We further observed that TRIM15 is “late” response TNF- α -induced gene and inhibits the TNF- α -induced NF- κ B pathway in several human cell lines. TRIM15 promotes turnover of K63-linked ubiquitin chains in a PRY/SPRY domain-dependent manner. TRIM15 interacts with TAK1 and inhibits its K63-linked ubiquitination, thus NF- κ B activity. Further, TRIM15 interacts with TRIM8 and inhibits cytosolic translocation to antagonize TRIM8 modulated NF- κ B. TRIM8 and TRIM15 also show functionally inverse correlation in psoriasis condition. In conclusion, TRIM15 is TNF- α -induced late response gene and inhibits TNF- α induced NF- κ B pathway hence a feedback modulator to keep the proinflammatory NF- κ B pathway under control.

1. Introduction

The NF- κ B family of transcription factors are activated by a range of pathophysiological stimuli including viral and bacterial factors (LPS, dsRNA), antigen receptors, DNA damage, reactive oxygen species (ROS), and cytokines (TNF- α , IL-1), [1–3]. Activated NF- κ B promotes transcription of a several target genes that include growth factors, chemokines, cytokines, immune modulators, regulators of apoptosis, acute response genes, cell adhesion molecules [1,3]. NF- κ B pathway is tightly regulated by post-transcriptional and post-translational regulatory mechanisms, however, its dysregulation leads to prolonged activation of NF- κ B and chronic inflammatory conditions hence critical for organismal survival and fitness [4].

Pleiotropic cytokine TNF- α is an activator of the NF- κ B pathway and inflammation [1,5]. The increased level of TNF- α and persistent activation of proinflammatory NF- κ B is observed in many pathological conditions including cancer [6–9]. TNF- α activated NF- κ B target genes have been classified in ‘early’ ‘mid’ and ‘late’ response genes based on their temporal expression pattern [10]. Some of the early response genes

like NFKBIA (κ B α) and TNFIP3 (A20) act as negative feedback regulators of the pathway and dynamics of gene expression by controlling NF- κ B oscillations [10–12]. The modulation of the NF- κ B pathway by late responsive genes and its implication in feedback regulation to restrict inflammation had not yet been explored.

Ubiquitination plays a critical role in the regulation of TNF- α mediated NF- κ B pathway. cIAP1/2 mediated Lys-63-linked polyubiquitination of RIP1, recruits TAK1 [5,13]. After recruiting TAK1 to the complex, TAK1 is K(Lysine)-63-linked polyubiquitinated and activated to recruit κ B kinase (IKK) complex. Dual phosphorylation of κ B α by IKK complex leads its K48-linked polyubiquitination by SCF ^{β TRCP} ubiquitin ligase complex [5,13] which is degraded through ubiquitin proteasome system (UPS). Removal of K63-linked ubiquitin chains by deubiquitinase (DUB) like A20 and CYLD negatively regulates NF- κ B activation and plays a pivotal role in modulating NF- κ B pathway [4,5,13]. TNF- α stimulation also promotes the expression of several E3 ligases including various TRAFs, cIAPs, and XIAP and most of these modifiers positively regulate the pathway [1]. Surprisingly to date, no TNF- α -induced E3 ligase had been shown regulating the pathway in a

Abbreviations: TNF- α , Tumor necrosis facotr-alpha; TRIMs, Tripartite motif containing proteins; CCLE, Cancer Cell Line Encyclopedia; TRIM8, Tripartite motif containing protein 8; TRIM15, Tripartite motif containing protein 15; Baf-A1, Bafilomycin-A1; UPS, Ubiquitin proteasome system; GEO, Gene Expression Omnibus.

* Corresponding author.

E-mail address: singhraj1975@gmail.com (R. Singh).

¹ Current Affiliation: Department of Cell and Developmental Biology, University College London, Gower Street, London, UK, WC1E 6BT.

<https://doi.org/10.1016/j.cellsig.2021.110210>

Received 19 April 2021; Received in revised form 25 November 2021; Accepted 30 November 2021

Available online 3 December 2021

0898-6568/© 2021 Elsevier Inc. All rights reserved.



Exosome Release Is Modulated by the Mitochondrial-Lysosomal Crosstalk in Parkinson's Disease Stress Conditions

Fatema Currim¹ · Jyoti Singh¹ · Anjali Shinde¹ · Dhruv Gohel¹ · Milton Roy¹ · Kritarth Singh² · Shatakshi Shukla¹ · Minal Mane¹ · Hitesh Vasiyani¹ · Rajesh Singh¹

Received: 21 April 2020 / Accepted: 2 December 2020 / Published online: 6 January 2021
© The Author(s), under exclusive licence to Springer Science+Business Media, LLC part of Springer Nature 2021

Abstract

Parkinson's disease (PD) is a neurodegenerative disorder characterized by the progressive loss of dopaminergic neurons in the substantia nigra (SN) pars compacta region of the brain. The main pathological hallmark involves cytoplasmic inclusions of α -synuclein and mitochondrial dysfunction, which is observed in other part of the central nervous system other than SN suggesting the spread of pathogenesis to bystander neurons. The inter-neuronal communication through exosomes may play an important role in the spread of the disease; however, the mechanisms are not well elucidated. Mitochondria and its role in inter-organellar crosstalk with multivesicular body (MVB) and lysosome and its role in modulation of exosome release in PD is not well understood. In the current study, we investigated the mitochondria-lysosome crosstalk modulating the exosome release in neuronal and glial cells. We observed that PD stress showed enhanced release of exosomes in dopaminergic neurons and glial cells. The PD stress condition in these cells showed fragmented network and mitochondrial dysfunction which further leads to functional deficit of lysosomes and hence inhibition of autophagy flux. Neuronal and glial cells treated with rapamycin showed enhanced autophagy and inhibited the exosomal release. The results here suggest that maintenance of mitochondrial function is important for the lysosomal function and hence exosomal release which is important for the pathogenesis of PD.

Keywords Mitochondrial dysfunctions · Mitochondria-lysosome crosstalk · Exosome release · Parkinson's disease

Introduction

Parkinson's disease (PD) is a chronic, progressive, neurodegenerative movement disorder including motor as well as non-motor symptoms. It affects 1% of the population of over 60 years of age and 3% of people over 80 years of age, and an estimated seven to ten million people are affected worldwide [1, 2]. Neuronal loss in the substantia nigra leads to a decrease in the dopamine levels in the corpus striatum, which leads to the motor symptoms, namely, tremor, rigidity and bradykinesia. The cellular hallmark of PD is the presence of the intracytoplasmic Lewy bodies and Lewy neurites, composed of protein aggregates, fats and polysaccharides. The protein aggregates contain α -synuclein, neurofilaments,

ubiquitin, Parkin and Synphilin [3]. There are emerging evidences which show that misfolded proteins spread through the brain along anatomically connected networks to other neuronal regions thereby promoting progressive decline [4]. PD occurs sporadically as well as in a familial form. Mutations in genes like SNCA, LRRK2 and VPS35 are related to autosomal dominant forms of PD, while PARKIN, PINK1 and DJ1 are associated with autosomal recessive form of PD [5]. The key molecular pathways regulated by these genes involved in PD are emerging; however, their role in progression to different brain regions is not well understood.

The dopaminergic neurons are specifically vulnerable in PD; however, α -synuclein aggregation and neuronal degeneration are observed in non-dopaminergic parts of the brain as well, like neocortex, brain stem and olfactory bulb [6], suggesting that the pathology spreads to other types of brain cells including microglia and astrocytes and other parts of the brain. Emerging studies suggest that exosomes play a major role in inter-neuronal and neuron-glia communication in the brain [7]. Exosomes are a class of extracellular vesicles ranging in the size approximately 30–150 nm, which are released from almost all cell types including neuron, glial and astrocytes [8].

✉ Rajesh Singh
singhraj1975@gmail.com

¹ Department of Biochemistry, Faculty of Science, The MS University of Baroda, Vadodara, Gujarat 390002, India

² Department of Cell and Developmental Biology, University College London, Gower Street, London WC1E 6BT, UK



DNA damage induces STING mediated IL-6-STAT3 survival pathway in triple-negative breast cancer cells and decreased survival of breast cancer patients

Hitesh Vasiyani¹ · Minal Mane¹ · Khushboo Rana² · Anjali Shinde¹ · Milton Roy³ · Jyoti Singh¹ · Dhruv Gohel¹ · Fatema Currim¹ · Ratika Srivastava² · Rajesh Singh^{1,4}

Accepted: 10 August 2022 / Published online: 26 August 2022

© The Author(s), under exclusive licence to Springer Science+Business Media, LLC, part of Springer Nature 2022

Abstract

Triple-negative breast cancer is aggressive and metastatic breast cancer type and shows immune evasion, drug resistance, relapse and poor survival. Anti-cancer therapy like ionizing radiation and chemotherapeutic drug majorly induces DNA damage hence, alteration in DNA damage repair and downstream pathways may contribute to tumor cell survival. DNA damage during chemotherapy is sensed by cyclic GMP-AMP synthase(cGAS)-stimulator of interferon genes (STING), which determines the anti-tumor immune response by modulating the expression of programmed cell death ligand-1 (PD-L1), immune suppressor, in the tumor microenvironment. Triple-negative breast cancer cells are cGAS-STING positive and modulation of this pathway during DNA damage response for survival and immune escape mechanism is not well understood. Here we demonstrate that doxorubicin-mediated DNA damage induces STING mediated NF- κ B activation in triple-negative as compared to ER/PR positive breast cancer cells. STING-mediated NF- κ B induces the expression of IL-6 in triple-negative breast cancer cells and activates pSTAT3, which enhances cell survival and PD-L1 expression. Doxorubicin and STAT3 inhibitor act synergistically and inhibit cell survival and clonogenicity in triple-negative breast cancer cells. Knockdown of STING in triple-negative breast cancer cells enhances CD8 mediated immune cell death of breast cancer cells. The combinatorial treatment of triple-negative breast cells with doxorubicin and STAT3 inhibitor reduces PD-L1 expression and activates immune cell-mediated cancer cell death. Further STING and IL-6 levels show a positive correlation in breast cancer patients and poor survival outcomes. The study here strongly suggests that STING mediated activation of NF- κ B enhances IL-6 mediated STAT3 in triple-negative breast cancer cells which induces cell survival and immune-suppressive mechanism.

✉ Rajesh Singh
rajesh.singh-biochem@msubaroda.ac.in;
singhraj1975@gmail.com

Hitesh Vasiyani
hiteshvasiyani@yahoo.com

Minal Mane
meenalmane14cir@gmail.com

Khushboo Rana
khoshboorana612@gmail.com

Anjali Shinde
anjalisshinde4891@gmail.com

Milton Roy
roymilton1989@gmail.com

Jyoti Singh
singhjyoti9315@gmail.com

Dhruv Gohel
dhruvgohilbaroda@gmail.com

Fatema Currim
currimfatema53@gmail.com


Ratika Srivastava
srivastava.ratika@gmail.com

¹ Department of Biochemistry, Faculty of Science, The Maharaja Sayajirao University of Baroda, Vadodara, Gujarat 390002, India

² Department of Microbiology & Biotechnology Center, The M.S. University of Baroda, Vadodara, Gujarat 390002, India

³ Institute for Cell Engineering, John Hopkins University School of Medicine, 733 North Broadway, MRB 731, Baltimore, MD 21205, USA

⁴ Department of Mol and Human Genetics, Banaras Hindu University, Varanasi, UP 221005, India



Expression of expanded *FMR1*-CGG repeats alters mitochondrial miRNAs and modulates mitochondrial functions and cell death in cellular model of FXTAS

Dhruv Gohel^a, Lakshmi Sripada^a, Paresh Prajapati^b, Fatema Currim^a, Milton Roy^a, Kritarth Singh^c, Anjali Shinde^a, Minal Mane^a, Darshan Kotadia^a, Flora Tassone^d, Nicolas Charlet-Berguerand^e, Rajesh Singh^{a,*}

^a Department of Biochemistry, Faculty of Science, The M.S. University of Baroda, Vadodara, 390002, Gujarat, India

^b SCoBIRC Department of Neuroscience, University of Kentucky, 741S. Limestone, BBSRB, Lexington, KY, 40536, USA

^c Department of Cell and Developmental Biology, University College London, Gower Street, London, WC1E 6BT, UK

^d Medical Investigation of Neurodevelopmental Disorders (MIND) Institute, University of California Davis, Davis, CA, 95817, USA

^e Institut de Genetique et de Biologie Moleculaire et Cellulaire (IGBMC), INSERM U1258, CNRS UMR7104, Université of Strasbourg, 67400, Illkirch, France

ARTICLE INFO

Keywords:
FXTAS
Mito-miRs
miR-320a
Mitotranscripts
Ago2
OXPHOS
Mitochondrial dysfunctions
Cell death

ABSTRACT

Fragile X-associated tremor/ataxia syndrome (FXTAS) is a progressive neurodegenerative disorder caused by an expansion of 55 to 200 CGG repeats located within 5'UTR of *FMR1*. These CGG repeats are transcribed into RNAs, which sequester several RNA binding proteins and alter the processing of miRNAs. CGG repeats are also translated into a toxic polyglycine-containing protein, FMRpolyG, that affects mitochondrial and nuclear functions reported in cell and animal models and patient studies. Nuclear-encoded small non-coding RNAs, including miRNAs, are transported to mitochondria; however, the role of mitochondrial miRNAs in FXTAS pathogenesis is not understood. Here, we analyzed mitochondrial miRNAs from HEK293 cells expressing expanded CGG repeats and their implication in the regulation of mitochondrial functions. The analysis of next generation sequencing (NGS) data of small RNAs from HEK293 cells expressing CGG premutation showed decreased level of cellular miRNAs and an altered pattern of association of miRNAs with mitochondria (mito-miRs). Among such mito-miRs, miR-320a was highly enriched in mitoplast and RNA immunoprecipitation of Ago2 (Argonaute-2) followed by Droplet digital PCR (ddPCR) suggested that miR-320a may form a complex with Ago2 and mitotranscripts. Finally, transfection of miR-320a mimic in cells expressing CGG premutation recovers mitochondrial functions and rescues cell death. Overall, this work reveals an altered translocation of miRNAs to mitochondria and the role of miR-320a in FXTAS pathology.

1. Introduction

FXTAS is a late onset inherited neurodegenerative disorder characterized by progressive intention tremor, gait ataxia and cognitive decline [1,2]. Nearly, 1 in ~3000 male and 1 in ~5000 female can be affected by FXTAS and disease symptoms get more pronounced with the age [3]. FXTAS is caused by an expansion of 55 to 200 CGG repeats (known as premutation) at the 5'UTR of the *FMR1* gene located on the long arm of X chromosome [4]. The expanded CGG repeats are transcribed into RNAs that titrate specific RNA binding proteins such as the DROSHA/DGCR8 complex involved in regulation of the processing of microRNAs

(miRNAs) [5]. Consequently, expression of various miRNAs are altered in FXTAS [6,7]. CGG repeats embedded in the 5'UTR of *FMR1* are translated into a toxic polyglycine-containing protein, FMRpolyG, through initiation via a non-canonical ACG start codon located upstream of the repeats [8–10]. However, it is still not understood if CGG RNA and/or FMRpolyG protein contribute to mitochondrial alterations leading neuronal cell dysfunctions and death [11]. Importantly, recent findings suggest that mitochondrial dysfunctions including loss of mitochondrial membrane potential, ATP and mitochondrial transcripts and proteins are associated with FXTAS pathogenesis [11–14]. We have recently shown decreased expression levels of mitochondrial transcripts

* Corresponding author.

E-mail address: rajesh.singh-biochem@msubaroda.ac.in (R. Singh).

<https://doi.org/10.1016/j.freeradbiomed.2021.01.038>

Received 4 June 2020; Received in revised form 30 December 2020; Accepted 17 January 2021

Available online 23 January 2021

0891-5849/© 2021 Elsevier Inc. All rights reserved.



hsa-miR-320a mediated exosome release under PD stress conditions rescue mitochondrial ROS and cell death in the recipient neuronal and glial cells

Shatakshi Shukla^a, Fatema Currim^a, Jyoti Singh^a, Shanikumar Goyani^a, M.V. Saranga^a, Anjali Shinde^a, Minal Mane^a, Nisha Chandak^a, Shyam Kishore^b, Rajesh Singh^{a,b,*}

^a Department of Biochemistry, Faculty of Science, The M.S. University of Baroda, Vadodara 390002, Gujarat, India

^b Department of Molecular and Human Genetics, Institute of Science, Banaras Hindu University, Varanasi UP 221005, India

ARTICLE INFO

Keywords:

Parkinson's disease
exosomes
hsa-miR-320a
autophagy
lysosome
mitochondria
cell-death

ABSTRACT

Parkinson's disease (PD) is a progressive neurodegenerative disorder characterized by dopaminergic neuronal cell death. Emerging evidence suggest exosomes as a crucial player in the progression and pathogenesis of PD via intercellular communication between different cell types in brain. Exosome release is enhanced from dysfunctional neurons/glia (source cells) under PD stress and mediates the transfer of biomolecules between different cell types (recipient) in brain leading to unique functional outcomes. Exosome release is modulated by alterations in the autophagy and lysosomal pathways; however, the molecular factors regulating these pathways remain elusive. Micro-RNAs (miRNAs) are class of non-coding RNAs that regulate gene expression post-transcriptionally by binding target mRNA and modulate its turnover and translation; however their role in modulating exosome release is not understood. Here, we analyzed the miRNAs-mRNAs network which target cellular processes regulating exosome release. hsa-miR-320a showed the maximum mRNA targets of autophagy, lysosome, mitochondria and exosome release pathways. hsa-miR-320a regulate ATG5 levels and modulate exosome release under PD stress conditions in neuronal SH-SY5Y and glial U-87 MG cells. hsa-miR-320a modulates autophagic flux, lysosomal functions, and mitochondrial ROS in neuronal SH-SY5Y and glial U-87 MG cells. Exosomes derived from hsa-miR-320a expressing source cells under PD stress conditions were actively internalized in the recipient cells and rescued cell death and mitochondrial ROS. These results suggest that hsa-miR-320a regulates autophagy and lysosomal pathways and modulates exosome release in the source cells and derived exosomes under PD stress conditions rescue cell death and mitochondrial ROS in the recipient neuronal and glial cells.

1. Introduction

Parkinson's disease is one of the most prevalent neurodegenerative disorder affecting approximately ten million people worldwide (Maserejian et al., 2020). The accumulation of misfolded α -synuclein aggregates also known as Lewy bodies in the substantia nigra region lead to neuronal loss results in both motor and non-motor defects in PD patients (Mehra et al., 2019). Non-motor symptoms appear prior to the onset of motor symptoms including pain, depression, sleep trouble, olfactory dysfunctions and gastrointestinal (GIT) alterations (Schapira et al., 2017). Evidences now suggest that substantia nigra is not the sole area with alpha synuclein aggregation and neuronal loss, instead it spreads to different brain regions (Dagher and Zeighami, 2018; Henrich et al., 2020; Port, 2018; Ruppert et al., 2020), however the molecular mechanisms are still not well understood. Emerging reports suggest the

role of nanovesicles termed as exosome in intercellular communication and its implication in the progression of PD pathogenesis (Yu et al., 2020). Brain cells including neurons, astrocytes and microglia actively communicate with each other via exosomes (Bavisotto et al., 2019). Moreover, exosomes mediated α -synuclein transfer to different cell types lead to cell death in specific regions of the brain (Guo et al., 2020b; Jacquet et al., 2021; Xia et al., 2019). Recent reports from our lab and other groups have shown enhanced exosome release and progression under PD stress conditions (Currim et al., 2021; Zhu et al., 2021); however the release mechanism and its effect after uptake in the bystander cells leading to neuronal dysfunction is not well understood. Inter-organellar crosstalk between different membrane bound organelles like mitochondria and ER with endo-lysosomal system is important for exchange of different biomolecules, maintaining the equilibrium either for degradation or secretion via exosome (Currim et al., 2021;

* Corresponding author at: Department of Molecular and Human Genetics, Institute of Science, Banaras Hindu University, Varanasi UP 221005, India.
E-mail address: rajeshs@bhu.ac.in (R. Singh).

<https://doi.org/10.1016/j.biociel.2023.106439>

Received 17 March 2023; Received in revised form 10 June 2023; Accepted 14 June 2023

Available online 8 July 2023

1357-2725/© 2023 Elsevier Ltd. All rights reserved.