

LIST OF FIGURES

Chapter 1

- Figure 1.1. Interorganellar crosstalk of mitochondria and lysosome in PD stress conditions.
- Figure 1.2. An overview of endolysosomal pathway, autophagy, and exosome release pathway in Parkinson's Disease.
- Figure 1.3. ESCRT-dependent exosome biogenesis pathway.
- Figure 1.4. MicroRNA sorting into exosomes.

Chapter 2

- Figure 2.1. Diagrammatic representation of the hypothesis of the study.

Chapter 4

- Figure 4.1.1. Isolation and characterization of exosomes.
- Figure 4.1.2. Exosome release is enhanced in PD stress conditions.
- Figure 4.1.3. PD stress conditions lead to mitochondrial dysfunctions.
- Figure 4.1.4. The number of lysosomes decline under PD stress conditions.
- Figure 4.1.5. Lysosome functions are altered under PD stress conditions.
- Figure 4.1.6. Autophagy flux is altered under PD stress conditions.
- Figure 4.1.7. Altered autophagic flux modulates exosome release in PD stress conditions.
- Figure 4.1.8. Alteration of autophagic flux by wortmannin decreases exosome release in PD stress conditions.
- Figure 4.1.9. Enhancement of autophagy flux by Rapamycin decreases the release of exosomes in PD stress conditions.
- Figure 4.2.1. Neuronal and glial exosomes are internalized by recipient cells and localize to the mitochondria.
- Figure 4.2.2. Schematic flowchart showing isolation of PD-stressed exosomes from SH-SY5Y neuronal cells and treatment to bystander neuronal cells.
- Figure 4.2.3. PD-stressed neuronal exosomes modulate mitochondrial functions in recipient neuronal cells leading to cell death.
- Figure 4.2.4. PD-stressed neuronal exosomes modulate mitochondrial functions in recipient glial cells leading to cell death.
- Figure 4.2.5. Schematic flowchart showing NGS analysis of exosomal RNA of SK-N-SH neuronal cells treated with 6-OHDA.
- Figure 4.2.6. Read length of miRNAs in NGS and sorting according to read counts.
- Figure 4.2.7. 6-OHDA treatment leads to differential miRNA enrichment in SK-N-SH exosomes.
- Figure 4.2.8. Putative bioinformatic analysis of targets of upregulated exosomal miRNAs.
- Figure 4.2.9. Putative bioinformatic analysis of targets of downregulated exosomal miRNAs.
- Figure 4.2.10. Validation of miRNAs from NGS in SK-N-SH neuronal cells and exosomes.

- Figure 4.2.11. Validation of miRNAs from NGS in SH-SY5Y neuronal cells and exosomes.
- Figure 4.2.12. Validation of miRNAs in rat primary cortical and midbrain neurons and exosomes.
- Figure 4.2.13. Cellular targets of hsa-miR-30a-5p and hsa-miR-181c-5p involved in mitochondrial pathways, inflammation, and cell death.
- Figure 4.2.14. miRNA sorting in exosomes is specific.
- Figure 4.2.15. miR-30a-5p and miR-181c-5p are selectively sorted in neuronal exosomes and are internalized by bystander neuronal cells.
- Figure 4.2.16. Schematic flowchart showing mimic-loaded exosomes from SH-SY5Y neuroblastoma cells treated to bystander neuronal cells pre-treated with PD stress inducers.
- Figure 4.2.17. miRNA mimic-loaded exosomes rescue mitochondrial dysfunction and cell death in PD-stressed neuronal cells.
- Figure 4.2.18. miRNA mimic-loaded exosomes rescue mitochondrial dysfunction and cell death in PD-stressed glial cells.
- Figure 4.3.1. Dose-dependent toxicity of rotenone in primary neurons.
- Figure 4.3.2. Exosome release is enhanced in primary neurons on treatment of the PD-stress inducer rotenone.
- Figure 4.3.3. Body weights of 3-month-old male Sprague-Dawley rats recorded before and after the administration of rotenone.
- Figure 4.3.4. Exosome release is enhanced in biofluids of acute rotenone-treated rat models of PD.
- Figure 4.3.5. Exosomal miRNA levels are altered in primary neuronal cells and exosomes on treatment with rotenone.
- Figure 4.3.6. Exosomal miRNAs are differentially expressed in biofluids of rotenone-treated rat models of PD.
- Figure 4.3.7. Serum exosomes alter mitochondrial functions in primary neurons from rat embryos.
- Figure 4.3.8. Serum exosomes from acute rotenone-treated rat models of PD decreases the cell viability of primary neurons from rat embryos.
- Figure 4.3.9. R8 exo from serum of acute rotenone-treated rat models of PD exacerbates rotenone-induced cell death of primary midbrain neurons from rat embryos.

Chapter 5

- Figure 5.1. Schematic representation of interorganellar crosstalk between mitochondria and lysosome in PD regulating the autophagy pathway, hence, the exosome pathway.
- Figure 5.2. Schematic representation of neuronal functions modulated by exosomal miRNAs.