

## LIST OF TABLES

Table no.	Title	Page no.
2.1	<b>Prevalence of Constipation in India (last 10 years)</b>	<b>11</b>
4.1	<b>Experimental Design for the study</b>	<b>45</b>
5.1.1	<b>General Information of the respondents &amp; its association with presence of functional constipation (N= 364)</b>	<b>73</b>
5.1.2	<b>Number (%) of subjects suffering from medical illnesses and their association with presence of functional constipation (N= 364)</b>	<b>74</b>
5.1.3	<b>Number (%) of subjects with a family history of constipation and its association with presence of Functional Constipation (N= 364)</b>	<b>76</b>
5.1.4	<b>Number (%) of subjects with specific personal and its association with presence of functional constipation (N= 364)</b>	<b>77</b>
5.1.5	<b>Number (%) of subjects' physical activity profile its association with presence of functional constipation (N= 364)</b>	<b>78</b>
5.1.6.1	<b>Number (%) of subjects with perceptions related to constipation and their association with presence of functional constipation (N=364)</b>	<b>79</b>
5.1.6.2	<b>Number (%) of subjects with general practices related to constipation and their association with presence of functional constipation (N= 364)</b>	<b>80</b>
5.1.6.3	<b>Number (%) of subjects with dietary practices related to constipation and their association with presence of functional constipation (N= 364)</b>	<b>81</b>
5.1.7	<b>Number (%) of subjects suffering from constipation and its symptoms (N-364)</b>	<b>83-84</b>
5.1.8.1	<b>Number (%) of subjects' chrono nutrition profile and its association with the presence of constipation on a working day (N=364)</b>	<b>86</b>

<b>5.1.8.2</b>	<b>Number (%) of subjects' chrono nutrition profile and its association with the presence of constipation on a free day (N=364)</b>	<b>87</b>
<b>5.1.9.1</b>	<b>Number (%) of subjects' type of diet consumed and its association with the presence of constipation (N=364)</b>	<b>89</b>
<b>5.1.9.2</b>	<b>Percent calorie and nutrient intake of RDA &amp; EAR by constipated subjects before supplementation (N=40)</b>	<b>89</b>
<b>5.1.10</b>	<b>Number (%) of subjects' food frequency of various foodstuffs and its association with constipation status (N=364)</b>	<b>90-91</b>
<b>5.2.1.1</b>	<b>Mean values for sensory qualities of gummies standardized with varying levels of water</b>	<b>94</b>
<b>5.2.1.2</b>	<b>Mean values for sensory qualities of gummies standardized with varying levels of sugar</b>	<b>95</b>
<b>5.2.1.3</b>	<b>Mean values for sensory qualities of gummies standardized with varying levels of agar</b>	<b>95</b>
<b>5.2.1.4</b>	<b>Mean values for sensory qualities of gummies standardized with varying levels of citric acid</b>	<b>96</b>
<b>5.2.2.1</b>	<b>Mean values for sensory qualities of gummies standardized with varying levels of GOS</b>	<b>97</b>
<b>5.2.2.2</b>	<b>Mean values for difference test for sucralose addition in 100% GOS gummies</b>	<b>97</b>
<b>5.2.3</b>	<b>Mean values of physico chemical characteristics of the 100% GOS gummies</b>	<b>99</b>
<b>5.2.5.1</b>	<b>Mean values of microbial count (Log10CFU/g of sample) of GOS gummies stored for a period of 6 months (37°C)</b>	<b>101</b>
<b>5.2.5.2</b>	<b>Mean values for sensory parameters of GOS Gummies stored for a period of 6 months (37°C)</b>	<b>102</b>
<b>5.3.1</b>	<b>General information of the respondents for the clinical trial (N= 35)</b>	<b>106</b>

<b>5.3.2</b>	<b>Mean scores of parameters of constipation profile before and after supplementation in experimental and control group (N=35)</b>	<b>108</b>
<b>5.3.3</b>	<b>Mean values of selected gut microbes analyzed in the constipated subjects (N=35)</b>	<b>110</b>
<b>5.3.4</b>	<b>Mean values of fecal short chain fatty acids analyzed in the constipated subjects (N=15)</b>	<b>112</b>
<b>5.3.5</b>	<b>Mean scores of depression profile before and after supplementation in experimental and control group (N=35)</b>	<b>113</b>
<b>5.3.6</b>	<b>Mean scores of quality of life before and after supplementation in experimental and control group (N=35)</b>	<b>114</b>
<b>5.3.7</b>	<b>Associations between various factors such as gut flora, SCFA, depression, QOL with presence of functional constipation at baseline (N=15)</b>	<b>117</b>