

BIBLIOGRAPHY

- Abowei, J. F. N. (2010). Salinity, dissolved oxygen, pH and surface water Temperature conditions in Nkoro River, Niger Delta, Nigeria. *Advanced Journal of Food Science and Technology*, 2(1), 16–21.
- Acharya, K. V., Dadhaniya, P., Shendage, A., Badne, A., & Chavan, A. (2019). Estuarine Fisheries resource of India. *International Journal of Sciences and Applied Research*, 6(1), 1–12.
- Afolabi, O. A., Arawomo, O. A., & Oke, O. L. (1984). Quality changes of Nigerian traditionally processed freshwater fish species. I. Nutritive and organoleptic changes. *International Journal of Food Science & Technology*, 19(3), 333–340.
- Aguirre, A. L. I., & Gallardo-Cabello, M. (2004). Reproduction of *Mugil cephalus* and *M. curema* (Pisces: Mugilidae) from a coastal lagoon in the Gulf of Mexico. *Bulletin of Marine Science*, 75(1), 37–49.
- Ahammad, A. K. S., Hasan, N. A., Haque, M. M., Bashar, A., Ahmed, Md. B. U., Alam, M. A., Asaduzzaman, Md., Bashar, Md. A., & Mahmud, Y. (2021). Environmental Factors and Genetic Diversity as Drivers of Early Gonadal Maturation: A Gonadosomatic Index Based Investigation on Indian Shad, *Tenualosa ilisha* Population of Bangladesh. *Frontiers in Marine Science*, 8. <https://doi.org/10.3389/fmars.2021.758868>
- Ajah, P. O., & Udoh, S. A. (2012). Food and feeding habits, condition factor and length-weight relationships of *Mugil cephalus* and *Pseudotolithus elongatus* in Cross River estuary, Cross River State, Nigeria. *Tropical Freshwater Biology*, 21(2), 59–70.
- Akin, S., Buhan, E., Winemiller, K. O., & Yilmaz, H. (2005). Fish assemblage structure of Koycegiz Lagoon–Estuary, Turkey: Spatial and temporal distribution patterns in relation to environmental variation. *Estuarine, Coastal and Shelf Science*, 64(4), 671–684.
- Akombo, P. M., Atile, J. I., Adikwu, I. A., & Araoye, P. A. (2011). Morphometric measurements and growth patterns of four species of the genus *Synodontis* (Cuvier, 1816) from Lower Benue River, Makurdi, Nigeria. *International Journal of Fisheries and Aquaculture*, 3, 263–270.

- Albieri, R. J., & Araújo, F. G. (2010). Reproductive biology of the mullet *Mugil liza* (Teleostei: Mugilidae) in a tropical Brazilian bay. *Zoologia (Curitiba)*, 27(3), 331–340. <https://doi.org/10.1590/s1984-46702010000300003>
- Aleleye-Wokoma, I. P., Hart, S. A., & Hart, A. I. (2001). Age and growth of *Mugil cephalus* (Linnaeus, 1758) (Perciformes: Mugilidae) in Bonny Estuary. In *14th Annual Conference of Fisheries Society of Nigeria (FISON)*, 19(23), 119–129.
- Ali, M., Salam, A., & Anas, M. (2002). Weight-length relationship and relative condition of *Oreochromis mossambicus*, an exotic fish species, in river Indus, Pakistan. *Pakistan Journal of Zoology*, 34, 125–128.
- Ali, S. S. R., Ramachandran, M., Chakma, S. K., & Sheriff, M. A. (2017). Proximate composition of commercially important marine fishes and shrimps from the Chennai coast, India. *International Journal of Fisheries and Aquatic Studies*, 5(5), 113–119.
- Allen, G. R. (1997). Marine Fishes of Tropical Australia and South-East Asia. *Western Australia Museum, Perth, Australia*, pp. 292.
- Anene, A. (2005). Condition factors of four cichlid species of a man-made lake in Imo state, Southeast, Nigeria. *Turkish Journal of Fisheries and Aquatic Sciences*, 5, 43–47.
- Anukima, A.M. (1963). Biological Abstract, 46, 23348.
- AOAC International (1995) Official methods of analysis of AOAC International, 16th edition. Arlington, VA, USA, Association of Analytical Communities, 2.
- Aragão, C., Costas, B., Vargas-Chacoff, L., Ruiz-Jarabo, I., Dinis, M. T., Mancera, J. M., & Conceição, L. E. (2010). Changes in plasma amino acid levels in a euryhalinefish exposed to different environmental salinities. *Amino Acids*, 38, 311–317.
- Azis, P. M. A., & Nair, N. B. (1983). Meiofauna of the EdavaNadayara Paravur backwater system of south west coast of India. *Mahasagar-Bulletin of the National Institute of Oceanography and Fisheries*, 16, 55–66.
- Azpurua, J., De La Cruz, D., Valderama, A., & Windsor, D. (2010). Lutzomyia sand fly diversity and rates of infection by Wolbachia and an exotic *Leishmania* species on Barro Colorado Island, Panama. *PLoS Neglected Tropical Diseases*, 4(3), e627. <https://doi.org/10.1371/journal.pntd.0000627>

Bibliography & Webliography

- Bacheler, N. M., Wong, R. A., Buckel, J. A. (2005). Movements and mortality rates of striped mullet in North Carolina. *North American Journal of Fisheries Management*, 25, 361–373.
- Bailey, B. E. (1952). Marine oils with particular reference to those of Canada. *Bulletin - Fisheries Research Board of Canada*, 89, 1–413.
- Baker, C. S., Lento, G. M., Cipriano, F., & Palumbi, S. R. (2000). Predicted decline of protected whales based on molecular genetic monitoring of Japanese and Korean markets. Proceedings. *Biological sciences*, 267(1449), 1191–1199.
- Baldwin, C. C., Mounts, J. H., Smith, D. G., & Weigt, L. A. (2009). Genetic identification and color descriptions of early life-history stages of Belizean *Phaeoptyx* and *Astrapogon* (Teleostei: Apogonidae) with comments on identification of adult *Phaeoptyx*. *Zootaxa*, 1–22.
- Bandarra, N. M., Batista, I., Nunes, M. L., & Empis, J. M. (2001). Seasonal variation in the chemical composition of horse-mackerel (*Trachurus trachurus*). *European Food Research and Technology*, 212, 535–539.
- Barletta, M., Lima, A. R., Costa, M. F., & Dantas, D. V. (2017). Estuarine ecoclines and the associated fauna: ecological information as the basis for ecosystem conservation. *Coastal wetlands: alteration and remediation*, 479–512. https://doi.org/10.1007/978-3-319-56179-0_16
- Baskaran, P. R. (1993). On the nutritional and reproductive cycles of mullets *Liza dussumieri* VAL_ and *Mugil cephalus* L of Muthupet saline swamp Tamil Nadu in the context of aquaculture. Ph.D. Thesis. <https://shodhganga.inflibnet.ac.in/handle/10603/116555>
- Bell, P. A., O'Neill, M. F., Leigh, G. M., Courtney, A. J., & Peel, S. L. (2005). Stock assessment of the Queensland-New South Wales Sea mullet fishery (*Mugil cephalus*). Queensland Department of Primary Industries and Fisheries, Queensland, Australia.
- Benakappa, S., Reddy, M. P. M., & Hariharan, V. (1980). Hydrographic conditions in the fishing grounds off Mukka-Kaup, South Kanara. *Mahasagar-Bulletin of the National Institute of Oceanography and Fisheries*, 13(1), 1–7.
- Ben-Yami, M. (1981). Handling, transportation and stocking of fry. In: O.H. Oren (ed.). *Aquaculture of Grey Mullet*, IBP 26. Cambridge, Cambridge University Press, International Biological Programme, 26. pp. 335–359.

- Berg, L. S. (1940). Classification of fishes both recent and fossil. Travaux de l'Institut Zoologique de l'Academie des Sciences de l'URSS. Leningrad: Izdatel'stvo Akademii Nauk SSSR [in Russian].
- Berg, L. S. (1947). Classification of fishes both recent and fossil. JW Edwards, (2).
- Bhadja, P., & Kundu, R. (2012). Status of the seawater quality at few industrially important coasts of Gujarat (India) off Arabian Sea. *Indian Journal of Geo-Marine Sciences*, 41(1), 90-97.
- Bhakta, D., Meetei, W. A., Vaisakh, G., Kamble, S. P., Das, S., & Das, B. K. (2018). Finfish Diversity of Narmada Estuary in Gujarat of India. *Proceedings of the Zoological Society*, 72(3), 257-262. <https://doi.org/10.1007/s12595-018-0263-1>
- Bhakta, D., Meetei, W. A., Vaisakh, G., Kamble, S., Chanu, T. N., & Das, S. K. (2017). An account of indigenous fishing gears of Narmada Estuary, Gujarat, India. *Journal of the Indian Society of Coastal Agricultural Research*, 35(1), 76-81.
- Bhatt, D. M., & Mankodi, P. C. (2020). Diversity record of family mugilidae (cuvier, 1829) from the estuary of Sabarmati river, Gujarat, India. *Journal of Biological Studies*, 3(2), 58-68.
- Bhatt, D. M., Sarma, K. J., Thakkar, N. J., & Mankodi, P. C. (2022). Growth Index, Gonadosomatic Index and Gut-Content Analysis of *Mugil Cephalus* Linnaeus, 1758 Found in Three Major Estuaries of Gujarat, India. In: *Proceedings of National Conference on Present Day Biology: Recent Advancements in Biological Sciences 10-11 December, 2021, Ahmedabad, Gujarat, India*. ISBN No: 978-93-9996-06-3
- Blaber, S. J. (1997). Fish and fisheries in tropical estuaries. Springer Science & Business Media, (22).
- Blaber, S. J. M. (1976). The food and feeding ecology of Mugilidae in the St. Lucia lake system. *Biological Journal of the Linnean Society*, 8(3), 267-277. <https://doi.org/10.1111/j.1095-8312.1976.tb00249.x>
- Blaber, S. J. M. (1977). The feeding ecology and relative abundance of mullet (Mugilidae) in Natal and Pondoland estuaries. *Biological Journal of the Linnean Society*, 9(3), 259-275.

Bibliography & Webliography

- Blaber, S. J. M. (1987). Factors affecting recruitment and survival of mugilids in estuaries and coastal waters of southeastern Africa. *In American Fisheries Society Symposium*, 1, 507–518.
- Blaber, S. J. M. (2000). *Tropical Estuarine Fishes: Ecology, Exploitation and Conservation*. Blackwell, Oxford, 372 pp.
- Blaber, S. J. M., & Whitfield, A. K. (1977). The feeding ecology of juvenile mullet (Mugilidae) in south-east African estuaries. *Biological Journal of the Linnean Society*, 9(3), 277–284. <https://doi.org/10.1111/j.1095-8312.1977.tb00270.x>
- Blanc, M., & Hureau, J. (1971). Catalogue critique des types de poissons du Muséum national d'Histoire naturelle. (Suite) (Mugiliformes et Polynémiformes). *Bulletin du Muséum National d'Histoire Naturelle*, 3, 673–735. [In French].
- Bleeker, P. (1853). Nalezingen op de ichthyologische fauna van Bengalen en Hindostan. *Bataviaasch Genootschap van Kunsten en Wetenschappen*, 1-70.
- Bœuf, G., & Payan, P. (2001). How should salinity influence fish growth? *Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology*, 130(4), 411–423.
- Boisneau, P. (1998). Les effets de l'artificialisation des estuaires sur les baisses de productions halieutiques et la necessite d'approches globales des impacts. *Les estuaires francais: evolution naturelle et artificielle*, 91-99.
- Bolger, T., & Connolly, P. L. (1989). The selection of suitable indices for the measurement and analysis of fish condition. *Journal of Fish Biology*, 34(2), 171–182.
- Boran, G., & Karaçam, H. (2011). Seasonal changes in proximate composition of some fish species from the Black Sea. *Turkish Journal of Fisheries and Aquatic Sciences*, 11(1), 1–5.
- Boyd, P. W., Watson, A. J., Law, C. S., Abraham, E. R., Trull, T., Murdoch, R., Bakker, D. C. E., Bowie, A. R., Buesseler, K. O., Chang, H., Charette, M., Croot, P., Downing, K., Frew, R., Gall, M., Hadfield, M., Hall, J., Harvey, M., Jameson, G., & LaRoche, J. (2000). A mesoscale phytoplankton bloom in the polar Southern Ocean stimulated by iron fertilization. *Nature*, 407(6805), 695–702. <https://doi.org/10.1038/35037500>

- Bradford, M. M. (1976). A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Analytical biochemistry*, 72(1-2), 248–254.
- Brahmane, V. T., Temkar, G. S., Metar, S. Y., Sikotaria, K. M., & Desai, A. Y. (2014). Ichthyofaunal diversity in the vicinity of marine protected areas, Jamnagar, Gulf of Kachchh, India. *Asian Journal of Advanced Basic Sciences*, 3(1), 78–88.
- Breder, C. M. (1940). The spawning of *Mugil cephalus* on the Florida west coast. *Copeia*, 1, 138–139.
- Briggs, J. C. (1960). Fishes of worldwide (circumtropical) distribution. *Copeia*, 171–180.
- Broadhead, G. C. (1953). Investigations of the Black Mullet, *Mugil cephalus* L., in Northwest Florida. *Florida Board of Conservation, Marine Research Lab, Technical Series No. 7. Florida Department of Natural Resources, Division of Marine Resources, USA*, 1-33.
- Bromhall, J. D. (1954). Notes on the reproduction of the grey mullet, *Mugil cephalus* L. *Hong Kong University Fishries Journal*, (1), 19-34.
- Brusle, J. (1981). Sexuality and biology of reproduction in grey mullets, in "Aquaculture of grey mullets" (ed. by O. H. Oren), *Cambridge University Press, Cambridge*, 99–154.
- Cameron, J. N. (1970). The influence of environmental variables on the hematology of pinfish (*Lagodon rhomboides*) and striped mullet (*Mugil cephalus*). *Comparative Biochemistry and Physiology*, 32(2), 175–192.
- Cardona, L. (2000). Effects of salinity on the habitat selection and growth performance of Mediterranean flathead grey mullet *Mugil cephalus* (Osteichthyes, Mugilidae). *Estuarine, Coastal and Shelf Science*, 50(5), 727–737.
- Cardona, L. (2006). Habitat selection by grey mullets (Osteichthyes, Mugilidae) in Mediterranean estuaries: the role of salinity. *Scientia marina*, 70, 443–455.
- Cardona, L. (2016). Food and feeding of Mugilidae. *Biology, ecology and culture of grey mullets (Mugilidae)*, 165-195.
- Cardona, L., Royo, P., & Torras, X. (2001). Effects of leaping grey mullet *Liza saliens* (Osteichthyes, Mugilidae) in the macrophyte beds of oligohaline

- Mediterranean coastal lagoons. *Hydrobiologia*, 462(1/3), 233–240.
<https://doi.org/10.1023/a:1013106905602>
- Carpentier, A., Como, S., Dupuy, C., Lefrançois, C., & Feunteun, E. (2014). Feeding ecology of *Liza* spp. in a tidal flat: Evidence of the importance of primary production (biofilm) and associated meiofauna. *Journal of Sea Research*, 92, 86–91. DOI: 10.1016/j.seares.2013.10.007.
- Cassifour, P. (1975). Contribution a l'étude de la biologie des mugilidés des réservoirs a poissons du bassin d'Arcachon. Universidad de Burdeos, Francia, 1, 104.
- Cech Jr, J. J., & Wohlschlag, D. E. (1973). Respiratory responses of the striped mullet, *Mugil cephalus* (L.) to hypoxic conditions. *Journal of Fish Biology*, 5(4), 421–428.
- Cervigón, F. (1985). The ichthyofauna of the Orinoco estuarine water delta in the west Atlantic coast, Caribbean. In a Yanez-Arancibia (Ed.), *Fish Community Ecology in Estuaries and Coastal lagoons: Towards an Ecosystem Integration*, UNAM Press, Mexico, 57–78.
- Chacko, P. I., & Ganapati, S. V. (1949). Some observations on the Adyar river with special reference to its hydrobiological conditions. *The Indian Geographical Journal*, 24(3), 35–49.
- Chandran, M. R. (1969). Reproductive and nutritional cycles in a marine perch, *Psammoperca waigiensis* (Cuvier). Ph.D. Thesis, University of Madurai, Madurai.
- Chandran, R., & Ramamoorthy, K. (1984). Hydrobiological studies in the gradient zone of the Vellar Estuary: 1. Physico-chemical parameters. *Mahasagar-Bulletin of the National Institute of Oceanography and Fisheries*, 17(2) 69–77.
- Chang, C. W., & Iizuka, Y. (2012). Estuarine use and movement patterns of seven sympatric Mugilidae fishes: The Tatu Creek estuary, central western Taiwan. *Estuarine, Coastal and Shelf Science*, 106(1), 121–126.
<https://doi.org/10.1016/j.ecss.2012.04.023>
- Chang, C. W., & Tzeng, W. N. (2000). Species composition and seasonal occurrence of mullets (Pisces, Mugilidae) in the Tanshui estuary northwest Taiwan. *Journal of the Fisheries Society of Taiwan*, 27(4), 253–262.

- Chang, C. W., Iizuka, Y., & Tzeng, W. N. (2004). Migratory environmental history of the grey mullet *Mugil cephalus* as revealed by otolith Sr:Ca ratios. *Marine Ecology Progress Series*, 269, 277–288. <https://doi.org/10.3354/meps269277>
- Chang, C. W., Tzeng, W. N., & Lee, Y. C. (2000). Recruitment and hatching dates of grey mullet (*Mugil cephalus* L.) juveniles in the Tanshui Estuary of Northwest Taiwan. *Zoological Studies*, 39, 99–106.
- Chaoui, L., Kara, M. H., Faure, E., & Quignard, J. P. (2006). The fish fauna of Mellah Lagoon (north-east Algeria): diversity, production and commercial catches analysis. *Cybium*, 30, 123–132.
- Chapman, L. J., & McKenzie, D. J. (2009). Behavioral responses and ecological consequences. In *Fish Physiology: Hypoxia*, 27, 25–77.
- Chaturvedi, L. D., Joshi, B. D., & Gupta, D. K. (1976). Biochemical composition of some tissues in *Heteropneustes fossilis* during pre-spawning period. *Matsya*, 2, 16–18.
- Chaudhuri, B. L. (1917). Fauna of the Chilika Lake: fish part III. *Memoirs of Indian Museum*, 5(6), 491–508.
- Chidambaram, K., Krishnamurthy, C. G., Venkataraman, R., & Chari, S. T. (1952). Studies on mackerel: Fat variations—and certain biological aspects. In *Proceedings/Indian Academy of Sciences, New Delhi-Springer*, 35(2), 43–68.
- Chubb, C. F., Potter, I. C., Grant, C. J., Lenanton, R. C. J., & Wallace, J. (1981). Age, structure, growth rates and movements of sea mullet, *Mugil cephalus* L., and Yellow-eye Mullet, *Aldrichetta forsteri* (Valenciennes), in the Swan-Avon river system, Western Australia. *Australian Journal of Marine and Freshwater Research*, 32(4), 605–628.
- Clark, B. M., Hutchings, K., & Lamberth, S. J. (2009). Long-term variations in composition and abundance of fish in the Berg Estuary, South Africa. *Transactions of the Royal Society of South Africa*, 64(2), 238–258.
- Crosetti, D., & Blaber, S. J. (2016). Biology, ecology and culture of grey mullets (Mugilidae). Boca Raton: CRC Press.
- Crosetti, D., & Blaber, S. J. M. (2016). Biology, ecology and culture of grey mullet (Mugilidae). *CRC Press*, 538.

Bibliography & Webliography

- Cyrus, D. P., & Blaber, S. J. M. (1987). The influence of turbidity on juvenile marine fishes in estuaries. Part 1. Field studies at Lake St. Lucia on the southeastern coast of Africa. *Journal of Experimental Marine Biology and Ecology*, 109(1), 53–70.
- Cyrus, D. P., & Blaber, S. J. M. (1987). The influence of turbidity on juvenile marine fishes in estuaries. Part 2. Laboratory studies, comparisons with field data and conclusions. *Journal of Experimental Marine Biology and Ecology*, 109(1), 71–91.
- Cywinska, A., Hunter, F., & Hebert, P. (2007). Identifying Canadian mosquito species through DNA Barcodes. *Medical and Veterinary Entomology*, 20, 413–424. <https://doi.org/10.1111/j.1365-2915.2006.00653.x>
- Das, H. P. (1978). Maturity and spawning in *Mugil cephalus* Linnaeus in Goa waters. *Mahasagar-Bulletin of the National Institute of Oceanography*, 11(1-2), 63–71.
- Dawnay, N., Ogden, R., McEwing, R., Carvalho, G. R., & Thorpe, R. S. (2007). Validation of the barcoding gene COI for use in forensic genetic species identification. *Forensic Science International*, 173(1), 1–6. <https://doi.org/10.1016/j.forsciint.2006.09.013>
- Day, F. (1888). Observations on the Fishes of India. *Proceedings of the Zoological Society of London, Oxford: Blackwell Publishing Ltd*, 56(1), 258–267.
- Day, F. (1889a). Fauna of British India, including Ceylon and Burma. Taylor and Francis, London, (2).
- Day, F. (1889b). Fauna of British India Fishes. Taylor and Francis Ltd., London, 1, 1-548.
- Day, J. H. (1951). The ecology of South African estuaries. Part 1. General considerations. *Transactions of the Royal Society of South Africa*, 33(1), 53–91.
- Day, J. H. (1981). The nature, origin and classification of estuaries. In J. H. Day (Ed.), *Estuarine Ecology with Particular Reference to Southern Africa*, A.A. Balkema, Cape Town, South Africa, 1–6.
- Denizci, R. (1958). Some thoughts about the biology of common grey mullet (*Mugil cephalus* L.) in the waters of Istanbul and its surroundings. *Rapports et*

Proces-Verbaux des Reunions - Commission Internationale pour l' Exploration Scientifique de la Mer Mediterranee, 14, 359–368.

Deshkar, S., Lakhmapurkar, J., & Gavali, D. (2012). State of three estuaries of Gulf of Khambhat. *Indian Journal of Geo-Marine Sciences, 41, 70–75.*

Deshkar, S., Lakhmapurkar, J., & Gavali, D. (2014). Temporal variation in the nutrient fluxes in Narmada estuary. *Indian Journal of Geo-Marine Sciences, 43(02), 284–288.*

Devasundaram, M. P. (1952). Scale study of *Mugil cephalus*, L., of Chilka Lake. *Madras Univ. Jour, 22(1), 147–163.*

Devasundaram, M. P. (1954). A report on the fisheries of the Chilika lake from 1948 to 1950. *Orissa Government Publication Cuttack: Govt Press, 1–34.*

Devlaming, V., Grossman, G., & Chapman, F. (1982). On the use of the gonosomatic index. *Comparative Biochemistry and Physiology Part A: Physiology, 73(1), 31–39.*

Digby, M. J., Saenger, P., Whelan, M. B., McCdonchie, D., Eyre, B. N. H., & Bucher, D. (1999). A Physical Classification of Australian Estuaries. Land and Water Resource Research and Development Corporation, Centre for Coastal Management, Southern Cross University, Lismore, NSW, 1–47.

Duncker, G. (1923). Korrelation zwischen Laänge und Gewicht bei Fischen. *Mitteilungen Aus Dem Zoologischen Staatsinstitut Und Zoologischen Museum Zu Hamburg, 4, 1–50.*

Durairaja, R., Jayakumar, N., Jawahar, P., & Ahilan, B. (2020). Length-weight relationship and condition factor of grey mullet, *Mugil cephalus* Linnaeus, 1758 from Pulicat lake, Tiruvallur (dt), Tamil Nadu. *Journal of Entomology and Zoology Studies, 8(6), 228–234.*

Durand, J. D., Chen, W. J., Shen, K. N., Fu, C., & Borsa, P. (2012). Genus-level taxonomic changes implied by the mitochondrial phylogeny of grey mullets (Teleostei: Mugilidae). *Comptes Rendus Biologies, 335(10-11), 687–697.*

Durand, J. D., Hubert, N., Shen, K. N., & Borsa, P. (2016). DNA barcoding grey mullets. *Reviews in Fish Biology and Fisheries, 27(1), 233–243.*
<https://doi.org/10.1007/s11160-016-9457-7>

Dyer, K. R. (1973). *Estuaries: a physical introduction.* John Wiley & Sons, London.

Bibliography & Webliography

- Ege, R., & Krogh, A. (1914). On the relation between the temperature and the respiratory exchange in fishes. *Internationale Revue der gesamten Hydrobiologie und Hydrographie*, 7(1), 48–55.
- Elaine, E. B., Gallardo-Cabello, M., Puente-Gomez, M., & Garcia-Boa, A. (2016). Reproduction of *Mugil cephalus* (Percoidei: Mugilidae) off the Central Mexican Pacific Coast. *Fisheries and Aquaculture Journal*, 7(4), 1–9.
- EMECS, (2007). Water quality standards for coastal water bodies. International Centre for the Environmental Management of Enclosed Coastal Seas. <https://www.emecs.or.jp/>
- Erman, F. (1961). On the biology of thick-lipped grey mullet (*Mugil chelo* Cuv.). *Commission Internationale pour l'Exploration Scientifique de la Méditerranée*, 16(2), 277–285.
- Erman, P. (1959). Observations on the biology of the grey mullet (*Mugil cephalus*). *General Fisheries Council for the Mediterranean*, 5, 157–169.
- Eschmeyer, W.N., & Fong, J.D., (2014). Species by Family/Subfamily. <http://research.calacademy.org/research/ichthyology/catalog/SpeciesByFamily.asp>
- Espino-Barr, E., Gallardo-Cabello, M., Puente-Gómez, M., & Garcia-Boa, A. (2016). Reproduction of *Mugil cephalus* (Percoidei: Mugilidae) off the Central Mexican Pacific Coast. *Fisheries and Aquaculture Journal*, 7, 180. doi:10.4172/2150-3508.1000180
- Evangeline, G., & Subbiah, K. (1969). Hydrobiology and fishery of the Ennore estuary from June, 1965 to May, 1967. *Madras J. Fisheries*, 5, 139–159.
- Ezzat, A. (1965). Contribution à l'étude de la biologie de quelques Mugilidae de la région de l'étang de Berre et de Port de Bouc. Ph.D. Thesis.
- Farrugio, H. (1975). Les muges (Poisons téléostéens) de Tunisie. Répartition et pêche, contribution à leur étude systématique et biologique. Ph.D. Thesis. <https://archimer.ifremer.fr/doc/00105/21585/>
- Farrugio, H., & Quignard, J. P. (1974). Biologie de *Mugil* (Liza) *ramada* Risso, 1826. et de *Mugil* (Chelon) *labrosus* Risso, 1826 (Poissons, Teleostreens, Mugilidae) du lac de Tunis. Age et Croissance. *Bulletin de L'institut National Scientifique et Technique D'oceanographie et de Peche de Salamambo*, 3, 1–4.

- Fazio, F., Marafioti, S., Arfuso, F., Piccione, G., & Faggio, C. (2013). Influence of different salinity on haematological and biochemical parameters of the widely cultured mullet, *Mugil cephalus*. *Marine and Freshwater Behaviour and Physiology*, 46(4), 211–218.
- Ferreira, M., Antunes, P., Gil, O., Vale, C., & Reis-Henriques, M. A. (2004). Organochlorine contaminants in flounder (*Platichthys flesus*) and mullet (*Mugil cephalus*) from Douro estuary, and their use as sentinel species for environmental monitoring. *Aquatic toxicology*, 69(4), 347–357.
- Folch, J., Lees, M., & Stanley, G. S. (1957). A simple method for the isolation and purification of total lipids from animal tissues. *Journal of biological chemistry*, 226(1), 497–509.
- Folmer, O., Black, M., Hoeh, W., Lutz, R., Vrijenhoek, R. (1994). DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. *Molecular Marine Biology and Biotechnology*, 3, 294–299.
- Fontaine, M., & Hatey, J. (1953). Contribution to the study of glucose metabolism in the salmon (*Salmo salar*) at different stages of its development and of its migrations. *Physiologia comparata et oecologia*, 3, 37–52.
- French, C. J., Hochachka, P. W., & Mommsen, T. P. (1983). Metabolic organization of liver during spawning migration of sockeye salmon. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology*, 245(6), R827–R830.
- Fry, F. E. J. (1957). The aquatic respiration of fish. *The physiology of fishes*, Academic Press, New York, (1), 1–64.
- Fry, F. E. J. (1964). Animals in aquatic environments: fishes. *Handbook of physiology*, 4(44), 715–728.
- Fulton, T. W. (1904). The rate of growth of fishes. Twenty-Second Annual Report, Part III. *Fisheries Board of Scotland, Edinburgh*, 141–241.
- Fulton, T. W. (1904). The rate of growth of fishes. Twenty-second Annual Report, Part III. *Fisheries Board of Scotland, Edinburgh*, 141–241.
- Fulton, T. W. (1911). The sovereignty of the sea: an historical account of the claims of England to the dominion of the British seas, and of the evolution of the territorial waters. *W. Blackwood, Edinburgh, Londo*, 799.

- Gajbhiye, S. N., Ram, J., Nair, V. R., & Desai, B. N. (1981). Plankton of the Narmada estuary and adjacent creek. *Mahasagar-Bulletin of the National Institute of Oceanography and Fisheries*, 14, 23–31.
- Ganias, K., Somarakis, S., Koutsikopoulos, C., & Machias, A. (2007). Factors affecting the spawning period of sardine in two highly oligotrophic Seas. *Marine Biology*, 151, 1559–1569. <https://doi.org/10.1007/s00227-006-0601-0>
- Gautier, D., & Hussenot, J. (2005). Les mulets des mers d'Europe. Synthèse des connaissances sur les bases biologiques et les techniques d'aquaculture. Ifremer, Plouzané
- Gayanilo, F. C., & Pauly, D. (1997). FAO ICLARM stock assessment tools (FISAT): References manual, FAO Computerized information series: 262.
- Gholampoor, T. E., Imanpoor, M. R., Shabanpoor, B., & Hosseini, S. A. (2011). The Study of Growth Performance, Body Composition and Some Blood Parameters of *Rutilus frisii kutum* (Kamenskii, 1901) Fingerlings at Different Salinities. *Journal of Agriculture, Science and Technology*, 13, 869–876.
- Ghosh, A. R., & Guha, B. C. (1934) Investigations on the nutritive value of Indian food stuffs - 2. *Indian Journal of Medical Research*, 21, 761–764.
- Girling, P., Purse, J., & Nowak, B. (2003). Effects of acute salinity and water quality changes on juveniles of green black flounder *Rhombosolea tapirina*. *Acta Ichthyologica et Piscatoria*, 33, 1–16.
- Glamuzina, B., Dulčić, J., Conides, A., Bartulović, V., Matić-Skoko, S., & Papaconstantinou, C. (2007). Some Biological Parameters of the Thin-Lippes Mullet *Liza Ramada* (Pisces, Mugilidae) in the Neretva River Delta (Eastern Adriatic, Croatian Coast). *Vie et milieu*, 57(3), 131–136.
- Gohil, M. N., & Mankodi, P. C. (2013). Diversity of Fish Fauna from Downstream Zone of River Mahisagar, Gujarat State, India. *Journal of Environmental Sciences*, 3, 23–26.
- Gokhale, K. S., & Patil, N. N. (1989). Relationship of metals in different organs of *Myatus gullo* (Ham.) with environmental parameters and biology of fish. *Journal of Environmental Biology*, 10(3), 265–271.

- Gomiero, L. M., Junior, G. A. V., & Naous, F. (2008). Relação peso-comprimento e fator de condição de *Cichla kelberi* (Perciformes, Cichlidae) introduzidos em um lago artificial no Sudeste brasileiro. *Acta Scientiarum. Biological Sciences*, 30(2), 173–178.
- Gopi, K. C., & Mishra, S. S. (2015) Diversity of marine fish of India. In Marine faunal diversity in India, Academic Press, 171–193.
- Gothandaraman. N. (1993). Studies on Micro Zooplankton, Ph.D. Thesis, Annamalai University, India, 180.
- Gray, C. A., Johnson, D. D., Young, D. J., & Broadhurst, M. K. (2004). Discards from the commercial gillnet fishery for dusky flathead, *Platycephalus fuscus*, in New South Wales, Australia: spatial variability and initial effects of change in minimum legal length of target species. *Fisheries Management and Ecology*, 11, 323–333.
- Green, J. (1968). The Biology of Estuarine Animals. Sidgwick & Jackson, London.
- Greenwood, P. H., Rosen, D. E., Weitzman, S. H., & Myers, G. S. (1966). Phyletic studies of teleostean fishes, with a provisional classification of living forms. *Bulletin of the American Museum of Natural History*, 131, 339–456.
- Grigorakis, K., Alexis, M. N., Taylor, K. A., & Hole, M. (2002). Comparison of wild and cultured gilthead sea bream (*Sparus aurata*); composition, appearance and seasonal variations. *International Journal of Food Science & Technology*, 37(5), 477–484.
- Gunter, G. (1967). Some relationships of estuaries to the fisheries of the Gulf of Mexico. In G. H. Lauff (Ed.), *Estuaries*. American Association for the Advancement of Science, Washington, 621–638.
- Gupta, R., Tatu, K., Christian, L., Joshi, D., & Kamboj, R. D. (2018). Seasonal Assessment of Some Water Quality Parameters for Estuarine and Riverine Zones of Narmada River, Gujarat. *International Journal of Scientific Research in Biological Sciences*, 5(5), 42–51.
- Hajibabaei, M., Shokralla, S., Zhou, X., Singer, G. A., & Baird, D. J. (2011). Environmental barcoding: a next-generation sequencing approach for biomonitoring applications using river benthos. *PLoS one*, 6(4), e17497. <https://doi.org/10.1371/journal.pone.0017497>

Bibliography & Webliography

- Hajibabaei, M., Singer, G. A., Hebert, P. D., & Hickey, D. A. (2007). DNA barcoding: How it complements taxonomy, molecular phylogenetics and population genetics. *Trends in Genetics*, 23(4), 167–172.
- Hakimelahi, M., Kamrani, E., Taghavi, M. S., Ghodrati, S. M., & Vahabnezhad, A. (2010). Growth parameters and mortality rates of *Liza klunzingeri* in the Iranian waters of the Persian Gulf and Oman Sea, using Length Frequency Data. *Iranian Journal of Fisheries Sciences*, 9(1), 87–96.
- Hall, G. M., & Ahmad, N. H. (1997). Surimi and fish-mince products. In *Fish Processing Technology* (G.M. Hall, ed.), Chapman and Hall, London, UK, pp. 74–92.
- Hamilton, F. (1822). An account of the fishes found in the river Ganges and its branches (Vol. 1). Archibald Constable.
- Hamilton, F. (1822). An account of the fishes found in the river Ganges and its branches. *Archibald Constable*, 1- 405.
- Handbook on Fisheries Statistics, India (2018). Department of Animal Husbandry, Dairying and Fisheries, Government of India. <http://dof.gov.in/sites/default/files/Handbook%20on%20FS%202018.pdf>
- Harrison, I. J., & Howes, G. J. (1991). The pharyngobranchial organ of mugilid fishes; its structure, variability, ontogeny, possible function and taxonomic utility. *Bulletin of the British Museum Natural History*, 57, 111–132.
- Harrison, I. J., & Senou, H. (2002). Order Mugiliformes, Mugilidae. *The living marine resources of the western central Atlantic*, 2, 1071–1085.
- Hasan, R., & Jafri, A. K. (1964). A physico-chemical analysis of ripe, unspawned eggs of some freshwater teleosts. In *Proceedings/Indian Academy of Sciences, New Delhi-Springer*, 60, 1–11.
- Hassan, H. U., Ali, Q. M., Ahmad, N., Masood, Z., Hossain, M. Y., Gabol, K., ... & Kamal, M. (2021). Assessment of growth characteristics, the survival rate and body composition of Asian Sea bass *Lates calcarifer* (Bloch, 1790) under different feeding rates in closed aquaculture system. *Saudi Journal of Biological Sciences*, 28(2), 1324–1330.
- Hebert, P. D. N., Ratnasingham, S., Zakharov, E. V., Telfer, A. C., Levesque-Beaudin, V., Milton, M. A., Pedersen, S., Jannetta, P., & Dewaard, J. R. (2016). Counting

- animal species with DNA barcodes: Canadian insects. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 371(1702). 20150333. <https://doi.org/10.1098/rstb.2015.0333>
- Hebert, P. D., Cywinska, A., & Ball, S. L. (2003). Biological identifications through DNA barcodes. *Proceedings of the Royal Society of London B: Biological Sciences*, 270(1512), 313–321.
- Hedge, J. E., Hofreiter, B. T., & Whistler, R. L. (1962). Carbohydrate chemistry. Academic Press, New York, 17, 371–80.
- Heino, & Kaitala, (1999). Evolution of resource allocation between growth and reproduction in animals with indeterminate growth. *Journal of Evolutionary Biology*, 12(3), 423–429.
- Heinrichs, J., Kreier, H. P., Feldberg, K., Schmidt, A. R., Zhu, R. L., Shaw, B., Shaw, A. J., & Wissemann, V. (2011). Formalizing morphologically cryptic biological entities: new insights from DNA taxonomy, hybridization, and biogeography in the leafy liverwort *Porella platyphylla* (Jungermanniopsida, Porellales). *American Journal of Botany*, 98(8), 1252–1262. <https://doi.org/10.3732/ajb.1100115>
- Henderson, R. J., Sargent, J. R., & Hopkins, C. C. E. (1984). Changes in the content and fatty acid composition of lipid in an isolated population of the capelin during sexual maturation and spawning. *Marine Biology*, 78, 255–263.
- Hendricks, L. J. (1961). The striped mullet, *Mugil cephalus* Linnaeus. The Ecology of the Salton Sea, California in Relation to the Sportfishery. *California Fish and Game, Fishery Bulletin*, 113, 1–204.
- Herre, W. A. (1939). On a collection of littoral and freshwater fishes from the Andaman Islands. *Records of the Zoological Survey of India*, 327–372.
- Herre, W. A. (1941). List of the fishes known from the Andaman Islands. *Memoirs of Indian Museum*, 13(3), 331–403.
- Hickling, C. F. (1970). A Contribution to the Natural History of the English Grey Mulletts (Pisces, Mugilidae). *Journal of the Marine Biological Association of the United Kingdom*, 50, 609–633.
- Hinton, M., Taylor, R.G., & Murphy, M., (1997). Use of gonad indices to estimate the status of reproductive activity of female swordfish, *Xiphias gladius*: a validated classification method. *Fishery Bulletin*, 95, 80–84.

Bibliography & Webliography

- Hoar, W. S. (1953). Control and timing of fish migration. *Biological Reviews*, 28(4), 437–452.
- Hoar, W. S. (1957). Endocrine organs chapter VI. *The physiology of fishes, edited by Margaret E. Brown. Academic Press, Inc. New York*, 245–277.
- Hoese, H. D. (1985). Jumping mullet – the internal diving bell hypothesis. *Environmental Biology of Fishes*, 13, 309–314.
- Hofmann, G. E., & Somero, G. N. (1996). Interspecific variation in thermal denaturation of proteins in the congeneric mussels *Mytilus trossulus* and *M. galloprovincialis*: evidence from the heat-shock response and protein ubiquitination. *Marine Biology*, 126, 65–75.
- Hossain, M. Y. (2010). Morphometric relationships of length-weight and length of four Cyprinid small indigenous fish species from the Padma River (NW Bangladesh). *Turkish Journal of Fisheries and Aquatic Sciences*, 10, 131–134.
- Hung, C. M., & Shaw, D. (2006). The impact of upstream catch and global warming on the grey mullet fishery in Taiwan: a non-cooperative game analysis. *Marine Resource Economics*, 21(3), 285–300. <https://doi.org/10.1086/mre.21.3.42629512>
- Huss, H. H. (1995). Quality and quality changes in fresh fish. FAO Fisheries Technical. Food and Agriculture Organization of the United Nations, pp. 348 <http://www.fao.org/3/V7180E/V7180E00.htm>
- Hwang, S. Y., Kuo, C. L., & Tanaka, S. (1990). Stock assessment of grey mullet *Mugil cephalus* in Taiwan by cohort analysis. *Nippon Suisan Gakkaishi*, 56(12), 1955–1963. <https://doi.org/10.2331/suisan.56.1955>
- Ivanova, N. V., Zemlak, T. S., Hanner, R. H., & Hebert, P. D. (2007). Universal primer cocktails for fish DNA barcoding. *Molecular Ecology Notes*, 7(4), 544–548.
- Jacob, P. K., & Krishnamurthi, B. (1948). Breeding and feeding habits of mullets (*Mugil*) in Ennore Creek. *Journal of the Bombay Natural History Society*, 47, 663–668.
- Jacquot, R. (1961). Organic constituents of fish and other aquatic animal foods. In *Fish as food*. Academic Press, New York, pp. 145–209.
- Jayaraman, R. (1954). Seasonal variations in salinity, dissolved oxygen and nutrient salts in the inshore waters of the Gulf of Mannar and Palk Bay near Mandapam (S. India). *Indian Journal of Fisheries*, 1(1&2), 345–364.

- Jayram, K. C. (1981). The freshwater fishes of India, Pakistan, Bangladesh, Burma and Srilanka. *A handbook of Zoological Survey of India*, 2, 475.
- Jeyaseelan, M. (1981). Role of mangrove forest of Pichavaram as fish nursery. *Proceedings of the Indian National Science Academy*, 46, 548–553.
- Jhingran, V. G. (1958). Observations on the seaward migration of *Mugil cephalus* Linnaeus from the Chilika Lake for breeding. *Current Science*, 27, 181–182.
- Jhingran, V. G. (1959). On the breeding migration of *Mugil cephalus* Linnaeus from the Chilika Lake to the sea. *Proceedings of the Indian Science Congress*, 46, 46.
- Jhingran, V. G. (1963). Report on the fisheries of the Chilka Lake, 1957-1960. *Bulletin*, 1, 1–12.
- Jhingran, V. G. (1982). Fishes and Fisheries of India. *Hindustan Publishing corporation (India) Delhi*, 645.
- Jhingran, V. G., & Mishra, K. N. (1962). Further fish tagging experiments in the Chilika Lake (1959) with special reference to *Mugil cephalus* Linnaeus. *Indian Journal of Fisheries Section-A*, 9(2), 477–498.
- Jhingran, V. G., & Natarajan, A. V. (1966). Final report on the fisheries of the Chilka Lake (1957-65). *Bulletin*, Central Inland Fisheries Research Institute, Barrackpore, India (8).
- Jhingran, V. G., & Natarajan, A. V. (1969). A study of the fisheries and fish populations of the Chilka Lake during the period 1957-65. *Journal of Inland Fisheries Society of India*, 1(1), 49–126.
- Jhingran, V. G., & Natarajan, A. V. (1973). Fishery resources of the Chilka Lake and its bearing on fisheries in adjacent areas of Bay of Bengal. In *Proceedings of the symposium on living resources of the seas around India. Special publication, CMFRI, Kerala*, 365-372.
- Jin, B., Fu, C., Zhong, J., Li, B., Chen, J., & Wu, J. (2007). Fish utilization of a salt marsh intertidal creek in the Yangtze River estuary, China. *Estuarine, Coastal and Shelf Science*, 73(3-4), 844–852.
- Jisr, N., Younes, G., Sukhn, C., & El-Dakdouki, M. H. (2018). Length-weight relationships and relative condition factor of fish inhabiting the marine area of the Eastern Mediterranean city, Tripoli-Lebanon. *The Egyptian Journal of Aquatic Research*, 44(4), 299–305.

- Jobling, M. (2012). R. J. Roberts (ed): Fish pathology Vol. 20. *Aquaculture International*, 811-812. <http://dx.doi.org/10.1007/s10499-012-9552-2>
- John, C. M. (1955). The grey mullets of Kayamkulam Lake, India, and their fishery. *Copeia*, 3, 225-230.
- Jones, S., & Sujansingani, K. H. (1954). Fish and fisheries of the Chilka Lake with statistics of fish catches for the years 1948-1950. *Indian Journal of Fisheries*, 1(1-2), 256-344.
- Joseph, N. I. (1976). Feeding habits and rate of growth of mullet fingerlings in the Kayamkulam lake, Kerala, India. *Journal of the Marine Biological Association of India*, 18(3), 488-492.
- Joshi, A., Parmar, E. A. R., Temkar, G. S., Desai, A. Y., & Bhatt, A. J. (2018). Ichthyofaunal Biodiversity of Kharakuva Fish Market, Veraval, Gujarat, India. *International Journal of Bio-Resource and Stress Management*, 9(5), 596-605. <https://doi.org/10.23910/ijbsm/2018.9.5.1902>
- Joshi, K.K., Varsha, M.S. & Sethulakshmi, M. (2018) Ichthyofaunal Diversity of India- Challenges Ahead for a Mega Biodiversity Country. In: ICAR Sponsored Winter School on Recent Advances in Fishery Biology Techniques for Biodiversity Evaluation and Conservation, Kochi, India. Teaching Resource, Retrieved from <http://eprints.cmfri.org.in/>, pp. 1-20.
- Kaliyaperumal, C. (1992). Studies on the interrelationship between phytoplankton and zooplankton in the waterways of the pitchavaram Mangroves. Ph.D Thesis, Annamalai University, India, 215.
- Kannappan, T., & Karthikeyan, M. M. (2013). Diversity of fishes in relation to physico-chemical properties of Manakudy estuary, Southwest coast of India. *International Journal of Biodiversity and Conservation*, 5(7), 396-407.
- Karuppasamy, P. K. (1997). Studies on zooplankton in the Pitchavaram mangroves a laboratory culture of rotifer *Brachionus plicatilis*. M.Phil. Thesis, Annamalai University, India, 85.
- Kastooribai, R. S. (1991). A comparative study of two tropical lentic systems in the context of aquaculture. Ph. D. Thesis, University of Madras, India.
- Kathiresan, K., Rajendran, N., & Thangadurai, G. (1996). Growth of mangrove seedlings in the intertidal area of Vellar estuary, southeast coast of India. *Indian Journal of Marine Sciences*, 25(3), 240-243.

- Katselis, G., Koutsikopoulos, C., Dimitriou, E., & Rogdakis, Y. (2003). Spatial patterns and temporal trends in the fishery landing of the Messolonghi-Etoliko lagoon system (western Greek coast). *Scientia marina*, 67, 501–511.
- Kennedy, M., & Fitzmaurice, P. (1969). Age and growth of thick-lipped grey mullet *Crenimugil labrosus* in Irish waters. *Journal of the Marine Biological Association of the United Kingdom*, 49(3), 683–699.
- Kesteven, G. L. (1942). Account of the fishery and preliminary statement of the biology of *Mugil dobula* Gunther. *Division of Fisheries Council for Scientific and Industrial Research-Melbourne*, 92–98.
- Kesteven, G. L. (1953). Further results of tagging sea mullet, *Mugil cephalus* Linnaeus, on the eastern Australia coast. *Marine and Freshwater Research*, 4(2), 251–306.
- Khériji, S., El Cafsi, M., Masmoudi, W., Castell, J. D., & Romdhane, M. S. (2003). Salinity and temperature effects on the lipid composition of mullet sea fry (*Mugil cephalus*, Linne, 1758). *Aquaculture international*, 11, 571–582.
- Kim, S. J., Lee, Y. D., Yeo, I. K., Baek, H. J., Kim, H. B., Nagaе, M., Soyano, K., & Hara, A. (2004). Reproductive cycle of the female grey mullet, *Mugil cephalus*, on the coast of Jeju Island, Korea. *Journal of Environmental Toxicology*, 19, 73–80.
- Klein, B. P., & Bardy, P. L. (1984) Experimental foods. Department of food nutrition, University of Illinois, U.S.A.
- Koutrakis, E. T. (1999). Review of the history and the systematics of the Mediterranean grey mullets (Mugilidae). *Geosciences Journal*, 10, 365–374.
- Koutrakis, E. T., & Tsikliras, A. C. (2003). Length-weight relationships of fishes from three northern Aegean estuarine systems (Greece). *Journal of Applied Ichthyology*, 19(4), 258–260. <https://doi.org/10.1046/j.1439-0426.2003.00456.x>
- Krishnakumar, K. (1992). Biology of the mullets of the south west coast of India. Ph.D. Thesis, 481.
- Kumar, C. S., & Prabu, V. A. (2014). Physico-chemical parameters in Parangipettai coastal waters and Vellar estuary, Southeast coast of India. *International Journal of Current Microbiology and Applied Sciences*, 3(9), 85–93.
- Kumar, J. I. N., Kumar, R. N., Patel, Y., & Viyol, S. (2010). Assessment of assessment of hydro-geochemical properties in relation to phytoplankton occurrence in

- tropical estuary - Narmada of Gujarat, India. *Indian Journal of Environmental Sciences*, 14(1), 19–28.
- Kumar, P., Arasu, A. R. T., Kailasam, M., Sukumarran Krishna, Subburj, R., Tyagraj, G., & Natarajan, M. (2015). Gonadal development and steroid hormone profile of wild caught grey mullet (*Mugil cephalus*). *Biological rhythm research*, 4, 601–610
- Kumar, P., Behera, P., Biswas, G., & Ghoshal, T. K. (2022). Oocyte growth, gonadosomatic index, hepatosomatic index and levels of reproductive hormones in goldspot mullet *Planiliza parsia* (Hamilton, 1822) reared in captivity. *Indian Journal Fisheries*, 69(1), 84–96.
- Kumaran, R. S., Ravi, V., Gunalan, B., Murugan, S., & Sundramanickam, A. (2012). Estimation of proximate, amino acids, fatty acids and mineral composition of mullet (*Mugil cephalus*) of Parangipettai, Southeast Coast of India. *Advances in Applied Science Research*, 3(4), 2015–2019.
- Kurma Rao, R., & Babu, K. R. (2016). Reproductive biology of the flathead grey mullet, *Mugil cephalus* (Linnaeus, 1758) from Krishna Estuarine region, East coast of Andhra Pradesh, India. *International Journal of Fisheries and Aquatic Studies*, 4, 483–488.
- Kutty, M. N., Sukumaran, N., & Kasim, H. M. (1980). Influence of temperature and salinity on survival of the freshwater mullet, *Rhinomugil corsula* (Hamilton). *Aquaculture*, 20, 261–274.
- Laffaille, P., Feunteun, E., & Lefeuvre, J. C. (2000). Composition of fish communities in a European macrotidal salt marsh (the Mont Saint-Michel Bay, France). *Estuarine, Coastal and Shelf Science*, 51(4), 429–438.
- Lawson, E. O., Akintola, S. O., & Olatunde, O. A. (2010). Aspects of the Biology of Sickie fin mullet, *Liza falcipinnis* (Valenciennes, 1836) from Badagry creek, Lagos, Nigeria. *Nature and Science*, 8, 168–182.
- Le Cren, E. D. (1951). The Length-Weight Relationship and Seasonal Cycle in Gonad Weight and Condition in the Perch (*Perca fluviatilis*). *The Journal of Animal Ecology*, 20(2), 201–219. <https://doi.org/10.2307/1540>
- Lemos, V. M., Varela Jr., A. S., Schiwingel, P. R., Muelbert, J. H., & Vieira, J. P. (2014). Migration and reproductive biology of *Mugil liza* (Teleostei: Mugilidae) in south Brazil. *Journal of Fish Biology*, 85, 671–687.

- Lester, W. F. (1975). Polluted river: River Trent, England. In River Ecology, Whitton BA (ed.). *Blackwell Scientific Publications-Oxford*, 489–513.
- Levinton, J. S. (2017). Marine biology: function, biodiversity, ecology. Oxford University Press, 1–530.
- Levinton, J. S. (2017). Marine biology: function, biodiversity, ecology. Oxford University Press, 1–530.
- Liao, I. C. (1981). Cultivation methods. In OH Oren, ed. Aquaculture of grey mullets. IBP 26. Cambridge, UK: Cambridge Univ. Press, pp. 361–390.
- Linder, D. R., Strawn, K., & Luebke, R. W. (1975). The culture of striped mullet (*Mugil cephalus* Linnaeus) in ponds receiving heated effluent from a power plant. *Aquaculture*, 5(2), 151–161.
- Lloyd, R. (1992). Pollution and Freshwater Fish (1st edition). Fishing News Books Ltd.: London, UK.
- Luther, G. (1963). Some observations on the biology of *Liza macrolepis* (Smith) and *Mugil cephalus* Linnaeus (Mugilidae) with notes on the fishery of grey mullets near Mandapam. *Indian Journal of Fisheries*, 10(2), 642–666.
- Luther, G. (1967). The grey mullets. In: CMFRI, Mandapam, editor. Souvenir 20th Anniversary Central Marine Fisheries Research Institute. Ernakulam: City Printers, pp. 70–74.
- Luther, G. (1973). The Grey mullet fishery resources of India. In: CMFRI, Kochi, editor. *Proceeding of living resources of the sea around India*. Cochin: Central Marine Fisheries Research Institute, pp. 455–460.
- Luther, G. (1974). On the little know mullet, *Valamugil buchanani* (Bleeker) (Mugilidae: Pisces) in the Indian waters. *Journal of the Marine Biological Association of India*, 16(1), 1–290.
- Luther, G. (1977). New characters for consideration in the taxonomic appraisal of grey mullets. *Journal of the Marine Biological Association of India*, 19(1), 1–9.
- Luther, G. (1967). The grey mullets. In: Souvenir 20th Anniversary Central Marine Fisheries Research Institute, Mandapam, pp 70–74.
- Luther, J. (1973). The Grey mullet fishery resources of India. In *Proceeding of the symposium on living resources of the sea around India*, Special Publication, Central Marine Fisheries Research Institute, Cochin, India. 455–460.

- Maiti, S. (2011). Handbook of methods in environmental studies, Volume 1: Water and Wastewater analysis. Oxford Book Company, Jaipur
- Maiti, S. K. (2003). Handbook of methods in environmental studies. ABD publishers, Jaipur, (2), 110-121.
- Maitland, P.S. & Herdson, D. (2009). Key to the marine and fresh- water fishes of Britain and Ireland. Environment Agency, Bristol. ISBN 978-1-84911-049-5.
- Mallo, S., Vallespinós, F., Ferrer, S. G., & Vaqué, D. (1993). Microbial activities in estuarine sediments (Ebro Delta, Spain) influenced by organic matter influx. *Scientia Marina*, 57(1), 31–40.
- Mani, P. (1989). Phytoplankton ecology of mangrove ecosystem. Ph.D. Thesis, Annamalai University, Indian, 72.
- Marais, J. F. K., & Erasmus, T. (1977). Body composition of *Mugil cephalus*, *Liza dumerili*, *Liza richardsoni* and *Liza tricuspidens* (Teleostei: Mugilidae) caught in the Swartkops estuary. *Aquaculture*, 10(1), 75–86.
- Mariani, S. (2001). Can spatial distribution of ichthyofauna describe marine influence on coastal lagoons? A central Mediterranean case study. *Estuarine, Coastal and Shelf Science*, 52(2), 261–267.
- McDonough, C. J., Roumillat, W. A., & Wenner, C. (2003). Fecundity and spawning season of striped mullet (*Mugil cephalus* L.) in South Carolina estuaries. *Fishery Bulletin*, 101, 822–834.
- McDonough, C. J., Roumillat, W. A., Wenner, C. (2005). Sexual differentiation and gonad development in striped mullet (*Mugil cephalus*) from South Carolina estuaries. *Fishery Bulletin*, 103, 601–619.
- McPherson, L. R., Ganas, K., & Marshall, C. T. (2011). Inaccuracies in routinely collected Atlantic herring (*Clupea harengus*) maturity data and correction using a gonadosomatic index model. *Journal of the Marine Biological Association of the United Kingdom*, 91(7), 1477–1487. Doi:10.1017/S002531541100018X
- McQuinn, I. H. (1989). Identification of spring-and autumn-spawning herring (*Clupea harengus harengus*) using maturity stages assigned from a gonadosomatic index model. *Canadian Journal of Fisheries and Aquatic Sciences*, 46(6), 969–980.

- Meien, V. A. (1927). Observations on the yearly variations of the ovaries of the perch (*Perca fluviatilis* L.). *Russkii Zoologicheskii Zhurnal*, 7(4).
- Miranda, M. T. P., Sunitha, S., & Jeena Mol, J. M. (2011). Biochemical composition of certain demersal fishes of Kollam (Quilon) in Kerala, southwest coast of India, with special reference to *Mugil cephalus* (Linnaeus) and *Johnius carutta* (Bloch). *Journal of Applied Zoological Researches*, 22(2), 113–118.
- Mishra, R. R., Rath, B., & Thatoi, H. (2008). Water Quality Assessment of Aquaculture Ponds Located in Bhitarkanika Mangrove Ecosystem, Orissa, India. *Turkish Journal of Fisheries and Aquatic Sciences*, 8, 71–77.
- Mishra, S., Panda, D., & Panigrahy, R. C. (1993). Physico-chemical characteristics of the Bahuda estuary (Orissa), east coast of India. *Indian Journal of Marine Sciences*, 22, 75–77.
- Misra, K. S. (1959). An aid to the identification of the common commercial fishes of India and Pakistan. *Record of the Indian Museum*, 57, 1–320.
- Mohamed, A. M., & Mohamed, M. A. (2019). Organoleptic evaluation of wet-salted marine fishes *Mugil cephalus*, *Chanos chanos* and *Gerres oyena* from Port Sudan coast. *Environmental Research*, 5(5), 684–698.
- Mohan Raj, V., Padmavathy, S., & Sivakumar, S. (2013). Water quality Parameters and it influences in the Ennore estuary and near Coastal Environment with respect to Industrial and Domestic sewage. *International Research Journal of Environment Sciences*, 2(7), 20–25.
- Mohanty, S. S., Dash, B. P., & Pramanik, D. S. (2015). Proximate composition of three marine fishes of Chandipur, Bay of Bengal, India. *International Journal of Fisheries and Aquatic Studies*, 2, 354–358.
- Mommsen, T. P., Danulat, E., & Walsh, P. J. (1992). Metabolic actions of glucagon and dexamethasone in liver of the ureogenic teleost *Opsanus beta*. *General and comparative endocrinology*, 85(2), 316–326.
[https://doi.org/10.1016/0016-6480\(92\)90016-D](https://doi.org/10.1016/0016-6480(92)90016-D)
- Montero, D., Blazer, V. S., Socorro, J., Izqwendo, M. S., & Turt, L. (1999). Dietary and culture influences on macrophage aggregate parameters in gilthead sea bream (*Sparus aurata*) juveniles. *Aquaculture*, 179, 523–534.

Bibliography & Webliography

- Moon, H. N., Park, J. H., Park, C., Namgung, J., Kim, K. H., & Yeo, I. K. (2017). Physiological responses of Gray mullet *Mugil cephalus* to low-pH water. *Korean Journal of Fisheries and Aquatic Sciences*, 50(2), 153–159.
- Moore, R. H. (1974). General ecology, distribution and relative abundance of *Mugil cephalus* and *Mugil curema* on the south Texas coast. *Contributions in Marine Science*, 18, 241–245.
- Morey, G., Moranta, J., Massuti, E., Grau, A., Linde, M., Riera, F., & Morales-Nin, B. (2003). Weight–length relationships of littoral to lower slope fishes from the western Mediterranean. *Fisheries Research*, 62(1), 89–96.
- Morovic, D. (1957). Les muges de l'Adriatique avec la bibliographie des muges. Inst. Zaslakvodno Ribar. Zagreb.
- Moslen, M., & Miebaka, C. A. (2017). Length-weight relationship and condition factor of *Mugil cephalus* and *Oreochromis niloticus* from a Tidal creek in the Niger Delta, Nigeria. *Archives of Agriculture and Environmental Science*, 2(4), 287–292.
- Muellner-Riehl, A., Schaefer, H., & Lahaye, R. (2011). Evaluation of candidate DNA barcoding loci for economically important timber species of the mahogany family (Meliaceae). *Molecular Ecology Resources*, 11, 450–460. <https://doi.org/10.1111/j.1755-0998.2011.02984.x>
- Murugan, S., Ashiq UR Rahman, M., Ajmal Khan, S., & Lyla, P. S. (2014). Growth and population dynamics of flathead grey mullet, *Mugil cephalus* (Linnaeus, 1758) from Parangipettai water (Southeast Coast of India). *Thalassa*, 30(2), 47–56.
- Murugan, S., Khan, S. A., Lyla, P. S., Kumar, C. P., Jalal, K. C., Kamaruzzaman, B. Y., & John, B. A. (2012). Length-weight relationship of *Mugil cephalus* (Linnaeus 1758) in Vellar estuary, southeast coast of India. *Pakistan Journal of Biological Sciences*, 15(12), 595–599.
- Mustafa, S., & Jafri, A. K. (1978). Some biochemical constituents and the calorific values of different regions of the body musculature of pond murrel *Channa punctatus* Bloch. *Fish. Techno.*, 15, 57–59.
- Nair, N. B., Abdul Azis, P. K., Krishna Kumar, K., Dharmaraj, K., & Arunachalam, M. (1984). Ecology of Indian estuaries. VI: Physico-chemical conditions in

- Kadinamkulam backwater, SW Coast of India. *Indian journal of marine sciences*, 13(2), 69–74.
- Nair, V. M., & Radhakrishnan, K. V. (1988). Regional variation in lipid and moisture content of *Tor khudree* muscle. In: The First Indian Fisheries Forum, *Proceedings: Asian Fisheries Society, Indian Branch, Manga1ore*, pp.89–90.
- Nelson, J. S. (1984) *Fishes of the World* (2nd edition). John Wiley & Sons, New York, 523.
- Nelson, J. S. (1984). *Fishes of the World*. 2nd ed. New York: John Wiley & Sons.
- Nelson, J. S. (1994). *Fishes of the World*. 3rd ed. New York: John Wiley & Sons.
- Njinkoue, J. M., Gouado, I., Tchoumboungang, F., Ngueguim, J. H. Y., Ndinteh, D. T., Fomogne-Fodjo, C. Y., & Schweigert, F. J. (2016). Proximate composition, mineral content and fatty acid profile of two marine fishes from Cameroonian coast: *Pseudotolithus typus* (Bleeker, 1863) and *Pseudotolithus elongatus* (Bowdich, 1825). *Journal of Nutrition & Food Science*, 4, 27–31. <https://doi.org/10.1016/j.nfs.2016.07.002>
- Nordlie, F. G. (1985). Osmotic regulation in the sheepshead minnow *Cyprinodon variegatus* Lacepede. *Journal of Fish Biology*, 26(2), 161–170.
- Nordlie, F. G. (2015) Adaptation to salinity and osmoregulation in Mugilidae. In *Biology, Ecology and Culture of Grey Mulletts (Mugilidae)*, CRC Press, 303–333.
- Nordlie, F. G., & Haney, D. C. (1998). Adaptations in salt marsh teleosts to life in waters of varying salinity. *Italian Journal of Zoology*, 65, 405–409.
- Nordlie, F. G., Szelistowski, W. A., & Nordlie, W. C. (1982). Ontogenesis of osmotic regulation in the striped mullet, *Mugil cephalus* L. *Journal of Fish Biology*, 20(1), 79–86.
- Odum, W. E. (1970). Utilization of the direct grazing and plant detritus food chains by the striped mullet *Mugil cephalus*. In J. H. Steele (Ed.), *Marine Food Chains*, Oliver & Boyd, London, 220–240.
- Odum, W. E. (1988). Comparative Ecology of Tidal Freshwater and Salt Marshes. *Annual Review of Ecology and Systematics*, 19(1), 147–176. <https://doi.org/10.1146/annurev.es.19.110188.001051>

Bibliography & Webliography

- Offem, B. O., Samsons, Y. A., & Omoniyi, I. T. (2009). Length-weight relationship, condition factor and sex ratio of forty-six important fishes in a tropical flood river. *Research Journal of Fisheries and Hydrobiology*, 4, 65–72.
- Orban, E., Nevigato, T., Masci, M., Di Lena, G., Casini, I., Caproni, R., Gambelli, L., De Angelis, P. & Rampacci, M. (2007). Nutritional quality and safety of European perch (*Perca fluviatilis*) from three lakes of Central Italy. *Food chemistry*, 100, 482–490.
- Oren, O. H. (1971). International Biological Programme: Coordinated studies on grey mullets. *Marine Biology*, 10(1), 30–33.
- Oren, O. H. (1981). Aquaculture of grey mullets. Cambridge University Press, London, pp. 409.
- Oren, O. H. (Ed.). (1981). Aquaculture of grey mullets. Cambridge University Press, London, (26), 409.
- Özyurt, C. E. (2003). The Determination of mesh size for some commercially important demersal fish species captured by deep trawl net at Babadillimanı Bight (Silifke-Mersin). Ph.D. Thesis, Cukurova University, Adana, 124.
- Özyurt, G., & Polat, A. (2006). Amino acid and fatty acid composition of wild sea bass (*Dicentrarchus labrax*): a seasonal differentiation. *European Food Research and Technology*, 222, 316–320.
- Özyurt, G., Polat, A., & Özkütük, S. (2005). Seasonal changes in the fatty acids of gilthead sea bream (*Sparus aurata*) and white sea bream (*Diplodus sargus*) captured in Iskenderun Bay, eastern Mediterranean coast of Turkey. *European Food Research and Technology*, 220, 120–124.
- Packer, L., Gibbs, J. C., & Jason, H. (2009). Barcoding methodology and applications DNA barcoding and the mediocrity of morphology. *Molecular Ecology Resources*, 9(1), 42–50.
- Panchakshari, V., Prabhavathi, K., Krishna, P. V., & Srikanth, B. (2015). The physico-chemical parameters of water from Nizampatnam bay, east coast of India. *International Journal of Advanced Research*, 3(9), 1259–1264.
- Panda, D., Mohanty, S. K., Pattnaik, A. K., Das, S., & Karna, S. K. (2018). Growth, mortality and stock status of mullets (Mugilidae) in Chilika Lake, India. *Lakes & Reservoirs: Research & Management*, 23(1), 4–16.

- Pandey, B. N., Datta Munshi, J. S., Choubey, B. J., & Pandey, P. K. (1976). Seasonal variation in body composition in relation to breeding cycle of an air-breathing fish, *Heteropneustes fossilis* (Bloch). *Journal of the Inland Fisheries Society of India*, 8, 91–95.
- Pandey, V., Jha, D. K., Ratnam, K., Venkatnarayanan, S., Sathish Kumar, P., Verma, P., & Dharani, G. (2022). Seasonal variability and its impact on the sub-tidal macrobenthic fauna of Diu Island, north-eastern Arabian Sea: A multivariate approach. *Frontiers in Marine Science*, 9, 1006689.
- Pandian, T. J. (1966). Feeding and reproductive cycles of the fish *Mystus gulio* in the Cooum backwaters, Madras. *Indian Journal of Fisheries*, 13(1 & 2), 322–333.
- Panfili, J., Thior, D., Ecoutin, J. M., Ndiaye, P., & Albaret, J. J. (2006). Influence of salinity on the size at maturity for fish species reproducing in contrasting West African estuaries. *Journal of Fish Biology*, 69(1), 95–113.
- Parmar, H., Barad, D., & Parasharya, D. (2015). Reef dependent ichthyofauna of the Gulf of Kachchh, Gujarat, Western India. *International Journal of Fisheries and Aquatic Studies*, 2(6), 33–37.
- Patale, V. V., & Tank, J. G. (2022). Evaluation of the edaphic and water properties of Diu coast (Saurashtra, Gujarat, India) in relation to the population density of *Avicennia marina*. *Applied Water Science*, 12(4), 78.
- Paterson, A. W., & Whitfield, A. K. (2003). The fishes associated with three intertidal salt marsh creeks in a temperate southern African estuary. *Wetlands Ecology and Management*, 11, 305–315.
- Patnaik, D. A. (1962). On the biology of *Mugil cephalus* Linnaeus of the Chilika Lake. *Proceedings of the Second All India Congress of Zoology Varanasi, (Pt. II)*, 457–464.
- Pauly, D. (1985). Ecology of coastal and estuarine fishes in Southeast Asia: a Philippine case study. In A. Yafiez-Arancibia (Ed.), *Fish Community Ecology in Estuaries and Coastal Lagoons: towards an ecosystem integration*, UNAM Press, Mexico, 499–514.
- Pentinsaari, M., Hebert, P. D., & Mutanen, M. (2014). Barcoding beetles: a regional survey of 1872 species reveals high identification success and unusually deep interspecific divergences. *PLoS One*, 9(9), e108651.

- Perera, P. A. B., & De Silva, S. S. (1978). Studies on the chemical biology of young grey mullet, *Mugil cephalus* L. *Journal of Fish Biology*, 13(3), 297–304.
- Perkins, E. J. (1976). The biology of estuaries and coastal waters. Academic Press, London.
- Petrakis, G., & Stergiou, K. I. (1995). Weight-length relationships for 33 fish species in Greek waters. *Fisheries Research*, 21(3-4), 465–469. [https://doi.org/10.1016/0165-7836\(94\)00294-7](https://doi.org/10.1016/0165-7836(94)00294-7)
- Pigott, G. M., & Tucker, B. (2017). Seafood: effects of technology on nutrition. CRC press.
- Pillay, S. R. (1962a). A Revision of the Indian Mugilidae. Part I. *Journal of the Bombay Natural History Society*, 59(1), 254–270.
- Pillay, S. R. (1962b). A Revision of the Indian Mugilidae. Part II. *Journal of the Bombay Natural History Society*, 59(2), 547–576.
- Pillay, T. V. R. (1953) Studies on the food, feeding habits and alimentary tract of the grey mullet, *Mugil tade* Forsskål, 1775. *Ibid*, 19, 777–827.
- Pillay, T. V. R. (1954). The biology of the grey mullet *Mugil tade* Forskal, with notes on its fishery in Bengal. In: *Proceedings of the Indian National Science Academy*, 20(2), 187–217.
- Polat, A., Kuzu, S., Oezyurt, G., & Tokur, B. (2009). Fatty acid composition of red mullet (*Mullus barbatus*): a seasonal differentiation. *Journal of Muscle Foods*, 20(1), 70–78.
- Potter, I. C., Tweedley, J. R., Elliott, M., & Whitfield, A. K. (2015). The ways in which fish use estuaries: a refinement and expansion of the guild approach. *Fish and Fisheries*, 16(2), 230–239.
- Prabhavathi, K., Krishna, P. V., Panchakshari, V., & Swapna, C. H. (2015). Fish resources in Nizampatnam coast, Andhra Pradesh. *International Journal of Advanced Research*, 3, 999–1004.
- Prakoso, V. A., Subagja, J., & Chang, Y. J. (2015). Low water temperature and its effects on stress response of grey mullets *Mugil cephalus* acclimated in freshwater. *Indonesian Aquaculture Journal*, 10(1), 41–45.
- Pramanik, D. S., & Mohanty, S. S. (2016). Length-weight relationship and biology of some common edible fish species at Chandipur, Bay of Bengal, Odisha. *International Journal of Fisheries and Aquatic Studies*, 4(1), 335–340.

- Pritchard, D. W. (1968). What is an estuary, physical viewpoint. *In Estuaries*, American Association for the Advancement of Science, Washington, 3–5.
- Qasim, S. Z. (1957). The biology of *Blennius oholis* L. (Teleostei). *The Proceedings of the Zoological Society of London*, 12, 161–208.
- Rabeh, I., Telahigue, K., Chetoui, I., Masmoudi, W., & El Cafsi, M. (2010). The effects of low salinity on lipid composition in the gills of grey mullet " *Liza ramada*". *INSTM Bulletin: Marine and Freshwater Sciences*, 37, 65–73.
- Raghunathan, C., Gupta, R. S., Wangikar, U., & Lakhmapurkar, J. (2004). A record of live corals along the Saurashtra coast of Gujarat, Arabian Sea. *Current Science*, 1131–1138.
- Ramamirtham, C. P., & Jayaraman, R. (1963). Some aspects of the hydrographical conditions of the backwaters around Willingdon Island (Cochin). *Journal of the Marine Biological Association of India*, 5(2), 170–177.
- Raman, K., Kaliyamurthy, M., & Joseph, K. O. (1977). Observations on the ecology and fisheries of the Pulicat lake during drought and normal periods. *Journal of the Marine Biological Association of India*, 19(1), 16–20.
- Ramos, S., Cowen, R. K., Paris, C., Ré, P., & Bordalo, A. A. (2006). Environmental forcing and larval fish assemblage dynamics in the Lima River estuary (northwest Portugal). *Journal of Plankton Research*, 28(3), 275–286.
- Rana, M. M., Chakraborty, S. C., & Saeid, A. (2020). Comparative studies of nutritional, microbial and organoleptic properties of different indigenous dried fish from local market in Bangladesh. *Advanced Journal of Chemistry-Section A*, 3(3), 318–327.
- Rangaswamy, C. (1975). Maturity and spawning of *Mugil cephalus* Linnaeus of Lake Pulicat. *Recent research in estuarine biology*, 47–60.
- Rangaswamy, C. (1976). Length-weight relationship in grey mullet *Mugil cephalus* Linnaeus. *Matsya*, 2, 19–22.
- Rao, K. S. (1967). Reproductive cycles and lipid levels of *Leiognathus splendens* (Cuv.). *Journal of the Marine Biological Association of India*, 9(2), 303–322.
- Rao, R. K., & Babu, K. R. (2013). Studies on food and feeding habits of *Mugil cephalus* (Linnaeus 1758) east coast off Andhra Pradesh India. *Canadian Journal of Pure and Applied Sciences*, 7(3), 2499–2504.

- Rao, S. N. Y., & Sastry, D. R. K. (2005). Fauna of Marine National Park, Gulf of Kachchh (Gujarat): An Overview. Conservation Area Series. Zoological Survey of India, Kolkata, 23, 1–79, Plates I-VIII.
- Ravagnan, G. (1992). Vallicoltura integrata. Edizioni Edagricole, Bologna, pp 502.
- Raval, I. H., Das, K. C., & Haldar, S. (2017). Collection of mullet fish (*Mugil cephalus*) from west coast of India: evaluation of its quality with relation to food safety. *Environmental Science and Pollution Research*, 24(11), 10833–10845. <https://doi.org/10.1007/s11356-017-8555-y>
- Rekha, M. U., Tomy, S., Sukumaran, K., Vidya, R., Kailasam, M., Balasubramanian, C. P., & Vijayan, K. K. (2021). Comparison of the reproductive biology of two stocks of Indian subcontinental *Mugil cephalus* Linnaeus, 1758 with special reference to reproductive isolation and philopatry. *Indian Journal of Geo-Marine Science*, 50(2), 130–140.
- Render, J. H., Thompson, B. A., & Allen, R. L. (1995). Reproductive development of striped mullet in Louisiana estuarine waters with notes on the applicability of reproductive assessment methods for isochronal species. *Transactions of the American Fisheries Society*, 124(1), 26–36.
- Rindi, F., Guiry, M., & Lopez-Bautista, J. (2008). Distribution, morphology, and phylogeny of *Klebsormidium* (Klebsormidiales, Charophyceae) in urban environment in Europe. *Journal of Phycology*, 44, 1529–1540. <https://doi.org/10.1111/j.1529-8817.2008.00593.x>
- Rollo, F., Ubaldi, M., Marota, I., Luciani, S., & Ermini, L. (2002). DNA Diagenesis: Effect of Environment and Time on Human Bone. *Ancient Biomolecules*, 4, 1–7. <https://doi.org/10.1080/13586120290018473>
- Rowshan-Ali, M., Roni, M. A. H., Haque, M. A., & Uddin, M. H. (2014). A study on chemical characterization and proximate composition of flathead mullet fish (*Mugil cephalus*) of estuarine region of the Bay of Bengal. *International Letters of Natural Sciences*, 12(2), 132–138.
- Sahoo, D. K., Karna, S. K., & Panda, S. (2012). Length weight relationship (lwr) and growth estimation of *Mugil cephalus* (Linnaeus, 1758) in the Chilika Lagoon, Orissa, India. *Asian Journal of Experimental Biological Sciences*, 3(4), 700–705.

- Saitou, N., & Nei, M. (1987). The neighbor-joining method: A new method for reconstructing phylogenetic trees. *Molecular Biology and Evolution*, 4, 406–425.
- Sankaranarayanan, V. N., & Qasim, S. Z. (1968). The influence of some hydrographical factors on the fisheries of Cochin area. *Proceedings of the National Institute of Sciences of India. Part B, Biological sciences*, 38, 148–163.
- Sankaranarayanan, V. N., & Qasim, S. Z. (1969). Nutrients of the Cochin Backwater in relation to environmental characteristics. *Marine Biology*, 2, 236–247.
- Saravanakumar, A., Rajkumar, M., Sesh Serebiah, J., & Thivakaran, G. A. (2009). Fishery resources in arid zone mangroves in gulf of Kachchh, Gujarat, northwest coast of India. *Journal of Ocean University of China*, 8(3), 233–240. <https://doi.org/10.1007/s11802-009-0233-3>
- Sargent, J. R. (1995). Origins and function of eggs lipids: Nutritional implication. *Broodstock Management and Egg and Larval Quality*, 353–372.
- Sarkar, S. (1977). Further studies on seasonal and spatial variations of salinity in Chilika lake. *Journal of the Inland Fisheries Society of India*, 9, 1–9.
- Sarkar, S. (1979). Further studies on seasonal and spatial variations of salinity in Chilka lake. *Journal of the Inland Fisheries Society of India*, 11(2), 1–9.
- Sarkar, U. K., Jena, J. K., Singh, S. P., Singh, A. K. & Rebello, S. C. (2012). Documenting coastal fish biodiversity of India: status, issues and challenges. *In: International Day for Biological Diversity Marine Biodiversity Uttar Pradesh State Biodiversity Board*, 22–28.
- Sarma, K. J., & Mankodi, P. C. (2017). Deciphering Identification of Inland Fishes of Gujarat Using DNA Barcoding. *Turkish Journal of Fisheries and Aquatic Sciences*, 17(5), 1055–1060.
- Sarojini, K. K. (1951) The fishery of the Indian grey mullets—A review. *Zoological Society of India*, 3, 159–79.
- Sarojini, K. K. (1957) Biology and fisheries of the grey mullets of Bengal I. Biology of *Mugil parsia* Hamilton with Notes on its Fishery in Bengal. *Indian Journal of Fisheries*, 4(1), 160–207.
- Sarojini, K. K. (1958) Biology and fisheries of the grey mullets of Bengal. II. Biology of *Mugil cunnesius* Valenciennes. *Indian Journal of Fisheries*, 5(1), 56–76.

Bibliography & Webliography

- Schultz, E. T., & McCormick, S. D. (2013). Euryhalinity in an evolutionary context. In S. D. McCormick, A. P. Farrell, & C. J. Brauner (Eds.), *Euryhalinity in an evolutionary context*, Elsevier Inc., New York, 477–533.
- Schultz, L. P. (1953). Family Mugilidae. In: Schultz LP, Herald ES, Lachner EA, et al., editors. *Fishes of the Marshall and Marianas Islands, Vol. I. Families from Asymmetriontidae Through Siganidae. Bulletin: United States National Museum, 202*, 310–322.
- Schultz, L. P. (1946). A revision of the genera of mullets, fishes of the family Mugilidae, with descriptions of three new genera. *Proceedings of the United States Natural Museum, 96*, 377–395.
- Shehadeh, Z. H., & Ellis, J. N. (1970). Induced spawning of the striped mullet (*Mugil cephalus* L.). *Journal of fish biology, 2*, 355–360.
- Shreni, K. D., & Jafri, A. K. (1978). Distribution pattern of some biochemical constituents in muscle of the common cat fish, *Heteropneustes fossilis* (Bloch). *Fish. Technol, 15*, 121–123.
- Skjæraasen, J. E., Kennedy, J., Thorsen, A., Fonn, M., Strand, B. N., Mayer, I., & Kjesbu, O. S. (2009). Mechanisms regulating oocyte recruitment and skipped spawning in Northeast Arctic cod (*Gadus morhua*). *Canadian Journal of Fisheries and Aquatic Sciences, 66*(9), 1582–1596.
- Smith, K., & Deguara, K. (2002). Review of the Biological Information and Stock Assessment for the NSW Sea Mullet Resource. *NSW Fisheries, Cronulla, 60–69*.
- Smith, M. A., Fisher, B. L., & Hebert, P. D. N. (2005). DNA barcoding for effective biodiversity assessment of a hyperdiverse arthropod group: the ants of Madagascar. *Philosophical Transactions of the Royal Society B: Biological Sciences, 360*(1462), 1825–1834.
- Sobhana, B. (1976). Studies on certain cyprinid fishes of Kerala. Ph. D. Thesis, University of Kerala, Trivandrum, India, 453.
- Solai, A., Gandhi, M. S., & Sriram, E. (2010). Implications of physical parameters and trace elements in surface water off Pondicherry, Bay of Bengal, South East Coast of India. *International Journal of Environmental Sciences, 1*(4), 529–542.

- Soundarapandian, P., Premkumar, T., & Dinakaran, G. K. (2009). Studies on the Physico-chemical characteristics and Nutrients in the Uppanar estuary of Cuddalore, South east coast of India. *Current Research Journal of Biological Sciences*, 1(3), 102–105.
- Sreenivasan, A., & Natarajan, M. V. (1963) Variations in the chemical composition of three fresh water fishes of Bhavanisagar Reservoir. *Indian Journal of Fisheries*, 8(2), 436–439.
- Stansby, M. E. (1954). Composition of certain species of fresh-water fish. 1. Introduction: the determination of the variation of composition of fish. *Food Research*, 19, 231–234.
- Stearns, S. C. (1992). The evolution of life histories. Oxford University Press, Oxford. 249.
- Su, W. C., & Kawasaki, T. (1995). Characteristics of the life history of grey mullet from Taiwanese waters. *Fisheries science*, 61(3), 377–381.
- Subrahmanyam, C. (1963). Studies on the reproductive cycle of *Penaeus indicus* and development of a few fishes of Madras. *Proceedings of the Indian National Science Academy*, 22, 279–295.
- Sunil, M. S. (2000). Length-weight relationship in *Rasbora daniconius* (Ham.) from Achancoil River, Pathanamthitta, Kerala, India. *Indian Journal of Fisheries*, 47(3), 271–274.
- Sweeney, B. W., Battle, J. M., Jackson, J. K., & Dapkey, T. (2011). Can DNA barcodes of stream macroinvertebrates improve descriptions of community structure and water quality? *Journal of the North American Benthological Society*, 30(1), 195–216.
- Tacon, A.G. J. (1996). Trends in aquaculture production with particular reference to low income food deficit countries 1984-83. *FAO Aquaculture, Aquaculture Newsletter*, 12, 6–9.
- Talwar, P. K., & Jhingran, A. G. (1991). Inland Fishes of India and Adjacent Countries. CRC Press, (2).
- Tampi, P. R. S. (1959). Ecological and fisheries characteristics of a salt water lagoon near Mandapam. *Journal of the Marine Biological Association of India*, 1(2), 113–130.

Bibliography & Webliography

- Tamura, K., Nei, M., & Kumar, S. (2004). Prospects for inferring very large phylogenies by using the neighbor-joining method. *Proceedings of the National Academy of Sciences (USA)*, 101, 11030–11035.
- Tamura, K., Stecher, G., & Kumar, S. (2021). MEGA11: molecular evolutionary genetics analysis version 11. *Molecular biology and evolution*, 38(7), 3022–3027.
- Tang, Y. A. (1975). Collection handling and distribution of grey mullet fingerlings in Taiwan. *Aquaculture*, 5, 81–84.
- Thakur, N. K. (1970). Observations on the mullet fishery in the Hooghly and Mahanadi estuarine systems. *Indian Journal of Fisheries*, 17(1 & 2), 1–12.
- Thangaraj, G.S., (1985). Ecobiology of the marine zone of the veiiar estuary. Ph.D. Thesis, Annamalai University, India, 194.
- Thompson, D. W. (1947). A glossary of Greek fishes. Oxford University Press, Geoffrey Cumberlege, pp. 297.
- Thomson, D. W. (1954). The Mugilidae of Australia and adjacent Seas. *Marine and Freshwater Research*, 5, 70–131.
- Thomson, D. W. (1997). The Mugilidae of the world. *Memoirs of the Queensland Museum*, 41, 457–562.
- Thomson, J. M. (1951). Growth and Habits of the Sea Mullet, *Mugil dobula* Gnnther, in Western Australia. *Australian journal of Marine and Freshwater Research*, 2(2), 193–225.
- Thomson, J. M. (1954). The Mugilidae of Australia and adjacent seas. *Marine and Freshwater Research*, 5(1), 70–131.
- Thomson, J. M. (1955). The movements and migrations of mullet (*Mugil cephalus* L.). *Australian Journal of Marine and Freshwater Research*, 6, 469–485.
- Thomson, J. M. (1963). Synopsis of biological data on the grey mullet *Mugil cephalus* Linnaeus 1758. *Australia: CSIRO Division of Fisheries and Oceanography, Fisheries Synopsis*, 1, 1–75.
- Thomson, J. M. (1966). The grey mullets. *Oceanography and Marine Biology an Annual Review*, 4(1), 301–335.
- Thomson, J. M. (1981). The taxonomy of grey mullets. *Aquaculture of grey mullets*, 1–16.

- Titouah, H., Rahmani, A., Mouloud, M. N., Aberkane, B., Ichalal, K., Debbou-Iouknane, N., Inouri-Iskounen, A., Khellouf, A., & Iguer-Ouada, M. (2022). Reproduction Activity of Flathead Mullet *Mugil cephalus* (Linnaeus, 1758) in Wild Condition in the Northeast of Algeria. *Russian Journal of Marine Biology*, 48(6), 504–512. <https://doi.org/10.1134/s1063074022060128>
- Tsogas, A., Vatavali, K., Dimitriou, E., Badeka, A., Kontakos, S., & Kontominas, M. G. (2019). Combined effect of light salting and vacuum packaging on the microbiological, chemical, and sensory attributes of mullet fillets (*Mugil cephalus*) during refrigerated and frozen/refrigerated storage. *Journal of food processing and preservation*, 43(8), e14009.
- Tulgar, A., & Berik, N. (2012). Effect of seasonal changes on proximate composition of red mullet (*Mullus barbatus*) and hake (*Merluccius merluccius*) were caught from Saroz Bay. *Research journal of biological sciences*, 2(2), 45–50.
- Vaitheeswaran, T., Malathi, S., & Felix, S. (2016). Studies on length-weight relationship of *Mugil cephalus* (Linnaeus, 1758), (Family: Mugilidae) Chennai coast of Tamil Nadu. *International Journal of Fisheries and Aquatic Studies*, 4(4), 4–17.
- Valentini, A., Pompanon, F., & Taberlet, P. (2009). DNA barcoding for ecologists. *Trends in Ecology & Evolution*, 24(2), 110–117.
- Vasanth, K., (1989). Studies on hydrobiology and decomposition of macrophytes in portonovo marine environment southeast coast of India. Ph.D. Thesis, Annamalai University, India, 252.
- Venkataraman, K., & Raghunathan, C. (2015). Coastal and Marine Biodiversity of India. In K. Venkataraman & C. Sivaperuman (Eds.), *Marine Faunal Diversity in India*, 303–348. <https://doi.org/10.1016/b978-0-12-801948-1.00019-7>
- Vitale, F., Svedäng, H., & Cardinale, M. (2006). Histological analysis invalidates macroscopically determined maturity ogives of the Kattegat cod (*Gadus morhua*) and suggests new proxies for estimating maturity status of individual fish. *ICES Journal of Marine Science*, 63(3), 485–492.
- Vivier, L., Cyrus, D. P., & Jerling, H. L. (2010). Fish community structure of the St Lucia estuarine system under prolonged drought conditions and its potential for recovery after mouth breaching. *Estuarine, Coastal and Shelf Science*, 86(4), 568–579.

Bibliography & Webliography

- Wafar, M., Venkataraman, K., Ingole, B., Ajmal Khan, S., & LokaBharathi, P. (2011). State of knowledge of coastal and marine biodiversity of Indian Ocean countries. *PLoS one*, 6(1), e14613.
- Wallace, J. H. (1975). The estuarine fishes of the east coast of South Africa. I. Species composition and length distribution in the estuarine and marine environments. II. Seasonal abundance and migrations. *Investigational Report, Oceanographic Research Institute, Durban*, 40(1), 1–72.
- Ward, R. D., Zemplak, T. S., Innes, B. H., Last, P. R., & Hebert, P. D. (2005). DNA barcoding Australia's fish species. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 360(1462), 1847–1857. <https://doi.org/10.1098/rstb.2005.1716>
- Weber-Scan, P. K., & Duffy, L. K., (2007). Effects of total dissolved solids on aquatic organisms: a review of literature and recommendation for salmonid species. *American Journal of Environmental Sciences*, 3(1), 1–6. <https://doi.org/10.3844/ajessp.2007.1.6>
- West, G. (1990). Methods of assessing ovarian development in fishes: a review. *Marine and Freshwater Research*, 41(2), 199–222.
- Whitehouse, R. H. (1922). The grey mullets of Tuticorin. *Madras fisheries bulletin*, 15, 71–98.
- Whitfield, A. K. (1996). a review of estuarine Ichthyology in south Africa over the past 50 years. *Transactions of the Royal Society of South Africa*, 51(1), 79–89. <https://doi.org/10.1080/00359199609520601>
- Whitfield, A. (1980). Factors Influencing the Recruitment of Juvenile Fishes into the Mhlanga Estuary. *South African Journal of Zoology*, 15(3), 166–169. <https://doi.org/10.1080/02541858.1980.11447705>
- Whitfield, A. K., Panfili, J., & Durand, J. D. (2012). A global review of the cosmopolitan flathead mullet *Mugil cephalus* Linnaeus 1758 (Teleostei: Mugilidae), with emphasis on the biology, genetics, ecology and fisheries aspects of this apparent species complex. *Reviews in Fish Biology and Fisheries*, 22(3), 641–681. <https://doi.org/10.1007/s11160-012-9263-9>
- Wiegand, M. D. (1996). Composition, accumulation, and utilization of yolk lipids in teleost fish. *Reviews in Fish Biology and Fisheries*, 6, 259–286.

- Wiens, J., Chippindale, P., & Hillis, D. (2003). When are phylogenetic analyses misled by convergence? A case study in Texas cave salamanders. *Systematic Biology*, 52(4), 501–514
- Winberg, G. G. (1956). Rate of metabolism and food requirements of fishes. *Fisheries Research Board of Canada Translation Series no 433*, 1–251.
- Winemiller, K. O., & Leslie, M. A. (1992). Fish assemblages across a complex, tropical freshwater/marine ecotone. *Environmental Biology of Fishes*, 34(1), 29–50. <https://doi.org/10.1007/bf00004783>
- Winkler, L. (1888). Determination of dissolved oxygen in water. *Reports of the German Chemical Society*, 21(2), 2843–2854.
- Wootton, R. J. (1990). Ecology of teleost fishes. Chapman and Hall. Upper Saddle River, New Jersey, USA.
- Wootton, R. J. (1998). Ecology of teleost fishes. Kluwer, Dordrecht, the Netherlands, 2, 386.
- Yang, W. T., & Kim, U. B. (1962). A preliminary report on the artificial culture of grey mullet in Korea. *Proceedings: Indo-pacific fisheries council*, 9(2-3), 62–70.
- Yashouv, A., & Berner-Samsonov, E. (1970). Contribution to the knowledge of eggs and early larval stages of mullets (Mugilidae) along the Israeli coast. *Bamidgeh*, 22(3), 72–89.
- Yehia, H. M., Alkhuriji, A. F., Al-Masoud, A. H., Tsiraki, M., & Mosilhey, S. H. (2022). The quality of handling and extended the shelf life and preservation of lagoon mullet fish (*Mugil cephalus*). *Food Science and Technology*, 42, e53722.
- Zubia, M., Rehana, Y., Muhammad, S. H., & Omer, M. T. Lakht-e-Zehra., & Adeyemi, S. O. (2014). Length-Weight relationship, condition and relative condition factor of four Mugilid species (Family mugilidae) from the Karachi Coast of Pakistan. *Journal of Coastal Development*, 17(385), 1–6.

WEBLIOGRAPHY

- Bleeker, P. (1853). Nalezingen op de ichthyologische fauna van Bengalen en Hindostan. Verhandelingen van het Bataviaasch Genootschap van Kunsten en Wetenschappen 25,1–164. Pls. 1-6. [in French].
- Bru, E. (2022). Chelon subviridis. The IUCN Red List of Threatened Species 2022: e.T197050A2478575. (<https://dx.doi.org/10.2305/IUCN.UK.2022-1.RLTS.T197050A2478575.en>). Accessed on 14th April, 2024.
- Camara, K.D., Carpenter, K.E., Djiman, R., Nunoo, F., Sagna, A., Sidibé, A., Sylla, M., de Morais, L., Jelks, H., Tolan, J., Vega-Cendejas, M., Espinosa-Perez, H., Chao, L., Moretzsohn, F. & Sparks, J.S. (2019). Mugil cephalus. The IUCN Red List of Threatened Species 2019: e.T135567A127923853. (<https://dx.doi.org/10.2305/IUCN.UK.2019-2.RLTS.T135567A127923853.en>). Accessed on 13th April, 2024.
- Dahanukar, N. (2010). Rhinomugil corsula. The IUCN Red List of Threatened Species 2010: e.T166478A6217637. (<https://dx.doi.org/10.2305/IUCN.UK.2010-4.RLTS.T166478A6217637.en>). Accessed on 13th April, 2024.
- Dahanukar, N. (2018). *Minimugil cascasia* (amended version of 2010 assessment). The IUCN Red List of Threatened Species 2018: e.T166422A128744264. (<https://dx.doi.org/10.2305/IUCN.UK.2010-4.RLTS.T166422A128744264.en>). Accessed on 13th April, 2024.
- Eschmeyer, W. N. (ed.). (2014). Catalogue of Fishes: Genera, Species, References. (<http://research.calacademy.org/research/ichthyology/catalog/fishcatman.asp>), Electronic version accessed on 1st September, 2014.
- FAO (2020). FishStat Aquaculture production 1950–2013 and Capture fisheries production 1950–2013. (<http://www.fao.org/fishery/topic/16140/en>), Accessed 10 December, 2023.
- Fricke, R., Eschmeyer, W. N., & Van der Laan, R. (2018). Catalog of fishes: genera, species, references. California Academy of Sciences, San Francisco, CA, USA. <http://researcharchive.calacademy.org/research/ichthyology/catalog/fishcatmain.asp>.
- Fricke, R., Eschmeyer, W.N. & Van der Laan R. (eds) (2023). Eschmeyer's Catalog of Fishes: Genera, Species, References.

- (<http://researcharchive.calacademy.org/research/ichthyology/catalog/fishcatmain.asp>). Electronic version accessed 12 December, 2023.
- Froese, R. (2006). Cube law, condition factor and weight-length relationships: history, meta-analysis and recommendations. *Journal of Applied Ichthyology*, 22(4), 241–253. <https://doi.org/10.1111/j.1439-0426.2006.00805.x>
- Froese, R., & Pauly, D. Editors. (2024). FishBase. *Chelon ramada* (Risso, 1827). Accessed through: World Register of Marine Species at: <https://www.marinespecies.org/aphia.php?p=taxdetails&id=1042830> on 2024-04-06.
- Froese, R., & Pauly, D. Editors. (2024). FishBase. *Planiliza haematocheilus* (Temminck & Schlegel, 1845). Accessed through: World Register of Marine Species at: <https://www.marinespecies.org/aphia.php?p=taxdetails&id=1509599> on 2024-04-06.
- Hoese, D. (2018). *Planiliza tade* (amended version of 2012 assessment). The IUCN Red List of Threatened Species: e.T166470A128738324. (<https://dx.doi.org/10.2305/IUCN.UK.2012-2.RLTS.T166470A128738324.en>). Accessed on 14th April, 2024.
- Palmer-Newton, A. (2020). *Crenimugil seheli*. The IUCN Red List of Threatened Species 2020 e.T186086A1809877. (<https://dx.doi.org/10.2305/IUCN.UK.2020-3.RLTS.T186086A1809877.en>) Accessed on 13th April, 2024.