

## References

- Acharya, K. P., Paudel, P. K., Jnawali, S. R., Neupane, P. R., & Koehl, M. (2017). Can forest fragmentation and configuration work as indicators of human–wildlife conflict? Evidences from human death and injury by wildlife attacks in Nepal. *Ecological indicators*, 80:74-83.
- Adriaensen, F., Chardon, J.P., De Blust, G., Swinnen, E., Villalba, S., Gulinck, H., & Matthysen, E. (2003). The application of ‘least-cost’ modelling as a functional landscape model. *Landscape Urban Plan*, 64(4): 233-247.
- Ahmadi, M., Kaboli, M., Nourani, E., Alizadeh Shabani, A., & Ashrafi, S. (2013). A predictive spatial model for gray wolf (*Canis lupus*) denning sites in a human-dominated landscape in western Iran. *Ecological Research*, 28:513–521.
- Akhtar, N., Bargali, H.S., & Chauhan, N.P.S. (2007). Characteristics of sloth bear day dens and use in disturbed and unprotected habitat of North Bilaspur Forest Division, Chhattisgarh, Central India. *Ursus*, 18 (2):203-208.
- Akhtar, N., Singh Bargali, H., & Chauhan, N. P. S. (2004). Sloth bear habitat use in disturbed and unprotected areas of Madhya Pradesh, India. *Ursus*, 15(2): 203–211.
- Araújo, M. B., Cabeza, M., Thuiller, W., Hannah, L., & Williams, P. H. (2004). Would climate change drive species out of reserves? An assessment of existing reserve-selection methods. *Global change biology*, 10(9):1618-1626.
- Athreya, V., Srivathsa, A., Puri, M., Karanth, K. K., Kumar, N. S., & Karanth, K. U. (2015). Spotted in the news: using media reports to examine leopard distribution, depredation, and management practices

outside protected areas in Southern India. *PLoS ONE* 10:e0142647. doi: 10.1371/journal.pone.0142647

- Babu, S., Karthik, T., Srinivas, G. & Kumara, H. N. (2015). Linking critical patches of sloth bear *Melursus ursinus* for their conservation in Meghamalai Hills, Western Ghats, India. *Current Science*, 109(8):1492–1498.
- Baldwin, R. A. (2009). Use of maximum entropy modeling in wildlife research. *Entropy* 11(4):854-866. <https://doi.org/10.3390/e11040854>.
- Ball, L. C., Doherty Jr, P. F., & McDonald, M. W. (2005). An occupancy modelling approach to evaluating a Palm Springs ground squirrel habitat model. *The Journal of Wildlife Management*, 69(3):894-904.
- Bandhyopadhyay, N., Bhuiyan, C. & Saha, A. K. (2016). Heat waves, temperature extremes and their impacts on monsoon rainfall and meteorological drought in Gujarat, India. *Natural Hazards*, 82: 367–388.
- Bargali, H. S., Akhtar, N., & Chauhan, N. P. S. (2005). Characteristics of sloth bear attacks and human casualties in North Bilaspur Forest Division, Chhattisgarh, India. *Ursus*, 16 (2): 263-267.
- Bargali, H.S., Akhtar, N., & Chauhan, N.P.S. (2012). The sloth bear activity and movement in highly fragmented and disturbed habitat in Central India. *World Journal of Zoology*, 7 (4):312–319.
- Bargali, H.S., Akhtar, N., Chauhan, N.P.S. (2012). Sloth bear (*Melursus ursinus*) habitat in the forests of North Bilaspur Forest Division, Chhattisgarh. *Indian Forester*, 138(10): 876.
- Baruch-Mordo, S., Wilson, K. R., Lewis, D. L., Broderick, J., Mao, J. S., & Breck, S. W. (2014). Stochasticity in natural forage production affects use of urban areas by black bears: implications to management of human-bear conflicts. *PLoS ONE* 9:e85122. doi: 10.1371/journal.pone.0085122
- Basak, K., Suraj, M., Ahmed, M., Kumar, U., & Bhattacharjee, S. (2023). The first attempt of assessing the population status of large predators and

their prey in an unexplored central Indian Protected Area. *Wildlife Letters*, 1(2): 47-58.

- Bashir, T., Bhattacharya, T., Poudyal, K., Qureshi, Q., & Sathyakumar, S. (2018). Understanding patterns of distribution and space-use by *Ursus thibetanus* in Khangchendzonga, India: Initiative towards conservation. *Mammalian Biology*, 92:11-20.
- Baskaran, N., Venkatesh, S., Srivasthava, S.K., & Desai, A.A. (2015). On the behavioural ecology of sloth bear (*Melursus ursinus* Shaw 1791) in Mudumalai Wildlife sanctuary, Western Ghats, India. *Animal Diversity, Natural History and Conservation*, 5:313-333.
- Beier, P., Majka, D.R., & Spencer, W.D. (2008). Forks in the road: choices in procedures for designing wildland linkages. *Conservation Biology* 22(4): 836-851.
- Benson, J. F., & Chamberlain, M. J. (2007). Space use and habitat selection by female Louisiana black bears in the Tensas River Basin of Louisiana. *The Journal of Wildlife Management*, 71(1):117-126.
- Bhandari, S., Crego, R. D., & Stabach, J. A. (2022). Spatial segregation between wild ungulates and livestock outside protected areas in the lowlands of Nepal. *Plos one*, 17(1), e0263122.
- Bhuvan, Indian Geo Platform for ISRO (<https://bhuvan-appl.nrsc.gov.in>).
- Bombieri, G., Penteriani, V., Almasieh, K., Ambarlı, H., Ashrafzadeh, M. R., Das, C. S., ... & del Mar Delgado, M. (2023). A worldwide perspective on large carnivore attacks on humans. *PLoS biology*, 21(1), e3001946.
- Boroushaki, S., & Malczewski, J. (2008). Implementing an extension of the analytical hierarchy process using ordered weighted averaging operators with fuzzy quantifiers in ArcGIS. *Computers & Geosciences*, 34(4):399-410.

- Brodie, J. F., Paxton, M., Nagulendran, K., Balamurugan, G., Clements, G. R., Reynolds, G., ... & Hon, J. (2016). Connecting science, policy, and implementation for landscape-scale habitat connectivity. *Conservation Biology*, 30(5):950-961.
- Buchholtz, E.K., Stronza, A., Songhurst, A., McCulloch, G., Fitzgerald, L.A. (2020). Using landscape connectivity to predict human-wildlife conflict. *Biology Conservation*, 248:108677.
- Can, Ö. E., Cruze, N. D., Garshelis, D. L., Beecham, J. J. & Macdonald, D. W. (2014). Resolving human–bear conflict: a global survey of countries, experts, and key factors. *Conservation Letters*, 7:501–513.
- Carricondo-Sanchez, D., Odden, M., Kulkarni, A., & Vanak, A. T. (2019). Scale-dependent strategies for coexistence of mesocarnivores in human-dominated landscapes. *Biotropica*, 51(5):781–791.
- Carroll, C., McRae, B., & Brookes, A. (2012). Use of linkage mapping and centrality analysis across habitat gradients to conserve connectivity of gray wolf populations in western North America. *Conservation Biology*, 26:78–87. doi:10.1111/j.1523-1739.2011. 01753.x
- Champion, H. G., & Seth, S. K. (1968). A revised survey of the forest types of India. Manager of publications. Nataraj Publisher, Dehradun.
- Choudhary, K., & Singh Nama, K. (2016). Human-Sloth Bear Conflict: Causes and Mitigation Measures in Kota, Rajasthan, India. *Pelagia Research Library Advances in Applied Science Research*, 7(6):1-7.
- Compton, B.W., McGarigal, K., Cushman, S.A., & Gamble, L.R. (2007). A resistant-kernel model of connectivity for amphibians that breed in vernal pools. *Conservation Biology*, 21(3):788-799.
- Conesa, F. C., Devanthery, N., Balbo, A. L., Madella, M. & Monserrat, O. (2014). Use of satellite SAR for understanding long-term human occupation dynamics in the monsoonal semi-arid plains of North Gujarat, India. *Remote Sensing*, 6(11):11420–11443.

- Cozzi, G., Chynoweth, M., Kusak, J., Coban, E., Coban, A., Ozgul, A. & Skercioglu, C. H. (2016). Anthropogenic food resources foster the coexistence of distinct life history strategies: year-round sedentary and migratory brown bears. *Journal of Zoology*, 300(2):142–150.
- Crutzen, P. J. (2006). The “Anthropocene.” In *Earth System Science in the Anthropocene* (pp. 13–18). Berlin/Heidelberg: Springer-Verlag. [https://doi.org/10.1007/3-540-26590-2\\_3](https://doi.org/10.1007/3-540-26590-2_3)
- Cushman, S. A., Landguth, E. L., & Flather, C. H. (2012). Evaluating the sufficiency of protected lands for maintaining wildlife population connectivity in the US northern Rocky Mountains. *Diversity and Distributions*, 18(9): 873-884.
- Debata, S., Swain, K. K., Sahu, H. K., et al. (2017). Human–sloth bear conflict in a human-dominated landscape of northern Odisha, India. *Ursus*, 27(2):90–98.
- Desai, V. (2020). *Assessment of habitat and feeding niche of sloth bear in Balaram Ambaji wildlife sanctuary, Gujarat*. MSc thesis. Hemchandracharya North Gujarat University, Gujarat, India.
- Devarajan, K., Morelli, T. L., & Tenan, S. (2020). Multi-species occupancy models: Review, roadmap, and recommendations. *Ecography*, 43(11):1612–1624.
- Devkar, R., Patel, P., & Upadhyay, K. (2013). Present scenario and possible threats to the Microchiropteran Bats near Jambughoda Wildlife Sanctuary, Gujarat. *Small Mammal Mail*, 5(1):28-31.
- Dhamorikar, A. H., Mehta, P., Bargali, H. & Gore, K. (2017). Characteristics of human–sloth bear (*Melursus ursinus*) encounters and the resulting human casualties in the Kanha–Pench corridor, Madhya Pradesh, India. *PLoS ONE* 12(4), e0176612.
- Dharaiya, A., Shah, V., Gadhavi, D., & Dharaiya, N. (2021). Assessment of wildlife habitat and natural resources with special reference to water

management in dry deciduous forest ecosystem of Gujarat state, India. *Journal of Geomatics*, 15(2).

- Dharaiya, N. & Gajera, N. (2008). Occurrence, Distribution and Status of Small and Certain Rare Species of Mammals in North Gujarat Region. 10.13140/RG.2.2.14497.04962.
- Dharaiya, N. (2008). Study on the status, distribution and occurrence of certain rare and small mammals in North Gujarat region. Final Report Submitted to Gujarat Forest Research Institute, 80.
- Dharaiya, N. (2009). Evaluating habitat & human–bear conflicts in North Gujarat, India, to seek solutions for human–bear coexistence. Research Project Report I-submitted to the Small Grants Division, Rufford Foundation, London, England, UK.
- Dharaiya, N., & Ratnayeke, S. (2009). Escalating Human-Sloth Bear Conflicts in North Gujarat: a tough time to encourage support for bear conservation. *International Bear News*, 18(3).
- Dharaiya, N., Bargali, H.S. & Sharp, T. (2020). *Melursus ursinus* (amended version of 2016 assessment). The IUCN Red List of Threatened Species 2020: e.T13143A166519315. <https://dx.doi.org/10.2305/IUCN.UK.20201.RLTS.T13143A166519315.en>. Accessed on 10 April 2024.
- Dharaiya, N., Bargali, H.S., & Sharp, T. (2016). *Melursus ursinus* The IUCN red list of threatened species 2016: e. T13143A45033815.
- Dharaiya, N., & Singh, C.P. (2017). Habitat Suitability & Corridor analysis for Sloth Bear in Gujarat using Remote Sensing and Ecological Modelling. Submitted to Space Application Centre Indian Space Research Organization, Ahmedabad
- Dickman A.J., Macdonald E.A., & Macdonald D.W. (2011). A review of financial instruments to pay for predator conservation and encourage

human-carnivore coexistence *Proceedings of the National Academy of Sciences*, 108:13937–13944.

- Dutta, T., Sharma, S., & DeFries, R. (2018). Targeting restoration sites to improve connectivity in a tiger conservation landscape in India. *Peer J*, 6, e5587.
- Dutta, T., Sharma, S., Maldonado, J. E., Panwar, H. S. & Seidensticker, J. (2015). Genetic Variation, Structure, and Gene Flow in a Sloth Bear (*Melursus ursinus*) Meta-Population in the Satpura-Maikal Landscape of Central India. *PLoS ONE*, 10(5), e0123384.
- Dutta, T., Sharma, S., McRae, B. H., Roy, P. S., & DeFries, R. (2016). Connecting the dots: mapping habitat connectivity for tigers in central India. *Regional Environmental Change*, 16:53-67.
- Elith, J., & Leathwick, J. R. (2009). Species distribution models: ecological explanation and prediction across space and time. *Annual Review of Ecology, Evolution, and Systematics*, 40:677-697.
- Elith, J., H. Graham, C., P. Anderson, R., Dudík, M., Ferrier, S., Guisan, A., ... & E. Zimmermann, N. (2006). Novel methods improve prediction of species' distributions from occurrence data. *Ecography*, 29(2):129-151.
- Elith, J., Phillips, S. J., Hastie, T., Dudík, M., Chee, Y. E., & Yates, C. J. (2011). A statistical explanation of MaxEnt for ecologists. *Diversity and Distributions*, 17(1):43–57. [https://doi.org/10.1111/J.1472-4642.2010.00725.X@10.1111/\(ISSN\)1472-4642.SPECIES-DISTRIBUTION-MODELS-INCONSERVATION-BIOGEOGRAPHY](https://doi.org/10.1111/J.1472-4642.2010.00725.X@10.1111/(ISSN)1472-4642.SPECIES-DISTRIBUTION-MODELS-INCONSERVATION-BIOGEOGRAPHY)
- Ferraz, G., Nichols, J. D., Hines, J. E., Stouffer, P. C., Bierregaard Jr, R. O., & Lovejoy, T. E. (2007). A large-scale deforestation experiment: effects of patch area and isolation on Amazon birds. *Science*, 315(5809): 238-241.

- Fiske, I. & Chandler, R. (2011) Unmarked: an R package for fitting hierarchical models of wildlife occurrence and abundance. *Journal of Statistical Software*,43:1–23
- Fragoso, J. M., Levi, T., Oliveira, L. F., Luzar, J. B., Overman, H., Read, J. M., & Silvius, K. M. (2016). Line transect surveys underdetect terrestrial mammals: implications for the sustainability of subsistence hunting. *PloS one*, 11(4), e0152659.
- Frank, B. (2016). Human–wildlife conflicts and the need to include tolerance and coexistence: an introductory comment. *Society and Natural Resources*,29(6):738-743.<https://doi.org/10.1080/08941920.2015.1103388>
- Frank, B., & J. A. Glikman. (2019). Human–wildlife conflicts and the need to include coexistence. *Human–wildlife interactions: Turning conflict into coexistence*:1-19. <https://doi.org/10.1017/9781108235730.004>
- Franklin, J. (2010). Moving beyond static species distribution models in support of conservation biogeography. *Diversity and Distributions*, 16(3): 321-330.
- FSI (2021). Forest Survey of India. Indian state of forest report. Forest Survey of India, Ministry of Environment, Forest and Climate change, Dehradun.
- Garcia, K.C., Joshi, H.M., & Dharaiya, N. (2016). Assessment of human–sloth bear conflicts in North Gujarat, India. *Ursus*, 27(1): 5–10. <https://doi.org/10.2192/URSUS-D-15-00012.1>
- Garshelis, D.L., Joshi, A.R., Smith, J.L. (1999a). Estimating density and relative abundance of sloth bears. *Ursus*, 87-98.
- Garshelis, D. L., Joshi, A. R., Smith, J. L. D., & Rice, C. G. (1999b). Sloth Bear Conservation Action Plan. In *Bears: status survey and conservation*

action plan. (pp. 225–240). International Union for the Conservation of Nature and Natural Resources. Gland, Switzerland.

- Gehr, B., Hofer, E. J., Muff, S., Ryser, A., Vimercati, E., Vogt, K., & Keller, L. F. (2017). A landscape of coexistence for a large predator in a human dominated landscape. *Oikos*, 126(10):1389–1399.
- Ghimire, D. & Thapa, T. B. (2014) Distribution and Habitat Preference of Sloth Bear in Chitwan National Park, Nepal. *Journal of Natural History Museum*, 28, 9–17.
- Goswami, V. R., Medhi, K., Nichols, J. D., & Oli, M. K. (2015). Mechanistic understanding of human-wildlife conflict through a novel application of dynamic occupancy models. *Conservation Biology*, 29, 1100–1110. doi: 10.1111/cobi.12475
- Guan, T., Wang, F., Li, S., & McShea, W. J. (2015). Nature reserve requirements for landscape-dependent ungulates: The case of endangered takin (*Budorcas taxicolor*) in Southwestern China. *Biological Conservation*, 182:63-71.
- Guisan, A., & Thuiller, W. (2005). Predicting species distribution: offering more than simple habitat models. *Ecology letters*, 8(9):993-1009.
- Gujarat State Gazetteers (1972). Panchmahals and Vadodara District. Government of Gujarat Publication, Ahmedabad.
- GujaratForestDepartment.<https://forests.gujarat.gov.in/writereaddata/images/pdf/Gujarat-Forest-Statistic-2021-22.pdf> Accessed on 10 December 2022.
- Gupta, R., & Sharma, L. K. (2020). Efficacy of Spatial Land Change Modeler as a forecasting indicator for anthropogenic change dynamics over five decades: A case study of Shoolpaneshwar Wildlife Sanctuary, Gujarat, India. *Ecological Indicators*, 112:106171.

- Harris, L.D., & Silva-Lopez, G. (1992). Forest fragmentation and the conservation of biological diversity. In: Fiedler PL, Jain SK, editors. *Conservation Biology*. Boston, MA: Springer US; 197–237.
- Hernandez, P. A., Graham, C. H., Master, L. L., & Albert, D. L. (2006). The effect of sample size and species characteristics on performance of different species distribution modelling methods. *Ecography*, 29(5):773-785.
- Hines, J.E. (2006). PRESENCE3 –Software to estimate patch occupancy and related parameters. USGS–PWRC. URL:<http://www.mbrpwr.usgs.gov/software/presence.html>.
- Inskip, C., & Zimmermann, A. (2009). Human–felid conflict: A review of patterns and priorities worldwide. *Oryx*, 43:18–34.
- Jain, P., Ahmed, R., Sajjad, H., Sahana, M., Jaafari, A., Dou, J., & Hong, H. (2021). Habitat suitability mapping of sloth bear (*Melursus ursinus*) in the Sariska Tiger Reserve (India) using a GIS-based Fuzzy analytical hierarchy process. In *Remote sensing and GIScience* (pp. 205-227). Springer, Cham.
- Jangid, A., Sahu, A., & Dharaiya, N. (2016). Record of Sloth Bear (*Melursus ursinus ursinus*, Shaw, 1791) from Aravalli District, Gujarat. *Journal of New Biological Reports*, 5(3):129-132.
- Jayadevan, A., Nayak, R., Karanth, K. K., Krishnaswamy, J., DeFries, R., Karanth, K. U., & Vaidyanathan, S. (2020). Navigating paved paradise: Evaluating landscape permeability to movement for large mammals in two conservation priority landscapes in India. *Biological Conservation*, 247:108613.
- Jena, J., Borah, J., Dave, C., & Vattakaven, J. (2021). Lifeline for Tigers: Status and Conservation of the KanhaPench Corridor. New Delhi (India): WWF-India. Report. [http://awsassets.wwfindia.org/downloads/kanha\\_pench\\_corridor\\_report.pdf](http://awsassets.wwfindia.org/downloads/kanha_pench_corridor_report.pdf)

- Jennings, S. B, Brown, N. D., & Sheil, D. (1999). Assessing forest canopies and understory illumination: canopy closure, canopy cover and other measures. *Forestry*, 22(1) :59-73.
- Jhala, Y.V., Qureshi, Q., & Sinha, P.R. (2011). Status of tigers, co-predators and prey in India. National Tiger Conservation Authority, Govt of India and the Wildlife Institute of India, New Delhi and Dehra Dun, India.
- Johnson, A., Wang, X., Kong, F., & Xue, M. (2013). Object-based evaluation of the impact of horizontal grid spacing on convection-allowing forecasts. *Monthly Weather Review*, 141(10):3413-3425.
- Joshi, A., Vaidyanathan, S., Mondol, S., Edgaonkar, A., & Ramakrishnan, U. (2013). Connectivity of tiger (*Panthera tigris*) populations in the human-influenced forest mosaic of central India. *PloS one*, 8(11), e77980.
- Joshi, H., Dharaiya, N. A., & Singh, C. P. (2015). Habitat Modeling to Detect Corridors for Sloth Bears in Gujarat, India. *International Bear News*, 24(3):28–19.
- Kaminski, D. J., Comer, C. E., Garner, N. P., Hung, I., & Calkins, G. E. (2013). Using GIS-based, regional extent habitat suitability modelling to identify conservation priority areas: A case study of the Louisiana black bear in East Texas. *The Journal of Wildlife Management*, 77(8):1639–1649.
- Kanagaraj, R., Wiegand, T., Kramer-Schadt, S., & Goyal, S. P. (2013). Using individual-based movement models to assess inter-patch connectivity for large carnivores in fragmented landscapes. *Biological Conservation*, 167:298-309.
- Karanth, K. K., & Kudalkar, S. (2017). History, location, and species matter: insights for human–wildlife conflict mitigation from India. *Human Dimensions of Wildlife*, 22(4): 331-346.

- Karanth, K. U., Gopaldaswamy, A. M., Kumar, N. S., Vaidyanathan, S., Nichols, J. D., & MacKenzie, D. I. (2011). Monitoring carnivore populations at the landscape scale: occupancy modelling of tigers from sign surveys. *Journal of Applied Ecology*, 48(4):1048-1056.
- Kéry, M., Guillera-Arroita, G., & Lahoz-Monfort, J. J. (2013). Analysing and mapping species range dynamics using occupancy models. *Journal of Biogeography*, 40(8):1463-1474.
- Ketting, J. (2020). *Social Factors Drive Sloth Bear Conflict In Gujarat: An Integrated Interdisciplinary Approach To Human-Wildlife Conflict And Coexistence*. MSc thesis. Utrecht University, Netherlands.
- Khattak, R. H., L. Teng, T. Mehmood, S. Ahmad, F. Bari, E. U. Rehman, & Z. Liu. (2021). Understanding the dynamics of human–wildlife conflicts in north-western Pakistan: implications for sustainable conservation. *Sustainability* 13(19): 10793. <https://doi.org/10.3390/su131910793>
- Khezri S., H. Shahabi & Ahmad B. (2013). Application of GIS and Remote sensing Techniques in Assessment of Natural Hazards in the Central Zab Basin, Northwest of Iran. 3(March 2013), 765–773.
- Koen, E.L., Bowman, J., Sadowski, C., & Walpole, A.A. (2014). Landscape connectivity for wildlife: development and validation of multispecies linkage maps. *Methods in Ecology & Evolution*, 5(7):626-633.
- Koirala, S., Baral, S., Garber, P. A., Basnet, H., Katuwal, H. B., Gurung, S., ... & Li, M. (2022). Identifying the environmental and anthropogenic causes, distribution, and intensity of human rhesus macaque conflict in Nepal. *Journal of Environmental Management*, 316:115276.
- Krausman, P. R., Rosentstock, S. S. & Cain III, J. W. (2006). Developed water for wildlife: Science, Perception, Values, and Controversy. *Wildlife Society Bulletin*, 34(3):563–569.

- Krofel, M., Elfström, M., Ambarlı, H., et al. (2020). Human–Bear Conflicts at the Beginning of the Twenty-First Century: Patterns, Determinants, and Mitigation Measures. In: Penteriani V, Melletti M, eds. *Bears of the World: Ecology, Conservation and Management*. Cambridge: Cambridge University Press;213-226. doi:10.1017/9781108692571.016.
- Kumar, G., Paul, D. K., & Kumari, D (2022). Human-Sloth Bear (*Melursus ursinus*) Conflicts in Nawada Forest Division, Nawada, Bihar, India. bioRxiv; 2022. DOI: 10.1101/2022.10.14.512280.
- Kumar, Nirmal J. I., Soni, H., & Kumar, R. N. (2006). Assessment of Anthropogenic Interventions in Jambughoda Wildlife Sanctuary, Gujarat, India. *Ecology Environment & Conservation*, 12(2): 235.
- Kumar, P., Singh, B. K., & Rani, M. (2013). An efficient hybrid classification approach for land use/land cover analysis in a semi-desert area using  $\{\rm ETM\}^{+}$  and LISS-III sensor. *IEEE Sensors Journal*, 13(6):2161-2165.
- Kumar, S., & Ramana, M. G. (2017). Management Plan for Shoolpaneshwar Wildlife Sanctuary. Narmada Forest Division, Gujarat Forest Department, Gujarat, India.
- Larsen, R. T., Bissonette, J. A., Flinders, J. T. & Whiting, J. C. (2012). Framework for understanding the influences of wildlife water developments in the western United States. *California Fish and Game*, 98(3):148–163.
- Larson, M. A. (2003). Landscape-level habitat suitability models for twelve wildlife species in southern Missouri (U.S Forest Service General Technical Report NC- 233). St. Paul: North Central Research Station.
- Lehman, J. (2021). All Men are Created Equal: Originalism, the Second Amendment, and Mental Disability. *Journal of Undergraduate Social Work Research*, 5(1) :31-40.

- Levine, J. M., Bascompte, J., Adler, P. B., & Allesina, S. (2017). Beyond pairwise mechanisms of species coexistence in complex communities. *Nature*, 546(7656):56–64.
- Li, Z., Wang, T., Smith, J. L., Feng, R., Feng, L., Mou, P., & Ge, J. (2019). Coexistence of two sympatric flagship carnivores in the human-dominated forest landscapes of Northeast Asia. *Landscape Ecology*, 34: 291–305.
- Lischka, S. A., Teel, T. L., Johnson, H. E., Reed, S. E., Breck, S., Carlos, A. D., & Crooks, K. R. (2018). A conceptual model for the integration of social and ecological information to understand human-wildlife interactions. *Biological Conservation*, 225:80-87.
- Long, H., D. Mojo, C. Fu, G. Wang, E. Kanga, A. M. Oduor, & L. Zhang. (2020). Patterns of human–wildlife conflict and management implications in Kenya: a national perspective. *Human Dimensions of Wildlife* 25(2):121-135. <https://doi.org/10.1080/10871209.2019.1695984>
- Lunn, D., Spiegelhalter, D., Thomas, A. & Best, N. (2009). The BUGS project: evolution, critique, and future directions. *Statistics in Medicine*, 28:3049–3067.
- MacKenzie, D. I., Nichols, J. D., Lachman, G. B., Droege, S., Andrew Royle, J., & Langtimm, C. A. (2002). Estimating site occupancy rates when detection probabilities are less than one. *Ecology*, 83(8), 2248–2255.
- MacKenzie, D. I., Nichols, J. D., Royle, J. A., Pollock, K. H., Bailey, L., & Hines, J. E. (2017). Occupancy estimation and modeling: Inferring patterns and dynamics of species occurrence. *Elsevier*.
- Malik, A. Dharaiya, N. & Proctor, M. (2018). Use of Geospatial techniques to target water sources for sloth bears aimed at alleviating conflicts with people. *International Bear News*, 27(3):44–46.

- Malik, A., Singh, C. P., Dharaiya, N., & Padate, G. (2023). Potential connectivity for sloth bear habitats in Western India's landscape. *European Journal of Wildlife Research*, 69(6):118.
- Manral, U., Sengupta, S., Hussain, S. A., Rana, S. A. K. S. H. I., & Badola, R. (2016). Human wildlife conflict in India: a review of economic implication of loss and preventive measures. *Indian Forester*, 142(10):928-940.
- Mark, D. M. (1988). "Network Models in Geomorphology." *Modelling Geomorphological Systems*, ed. M. G. Anderson. New York: John Wiley. 73–97.
- McRae, B. (2012). Centrality Mapper connectivity analysis software. The Nature Conservancy, Fort Collins.
- McRae, B. H., Shah, V., & Edelman, A. (2016). Circuitscape: modeling landscape connectivity to promote conservation and human health. *The Nature Conservancy*, 14:1-14.
- McRae, B., & Kavanagh, D.M. (2011). Linkage Mapper connectivity analysis software. The Nature Conservancy, Fort Collins.
- McRae, B.H., Dickson, B.G., Keitt, T.H., & Shah, V.B. (2008). Using circuit theory to model connectivity in ecology, evolution, and conservation. *Ecology*, 89(10):2712-2724.
- McRae, B.H., Shirk, A., Platt, J. (2013). Gnarly landscape utilities: resistance and habitat calculator user guide. *The Nature Conservancy*, Fort Collins.
- Mewada, T. P. (2015) Index of Relative Importance of the Dietary Proportions of Sloth Bear (*Melursus ursinus*) in Semi-Arid Region. *Notulae Scientia Biologicae*, 7(3):281–288.
- Mewada, T., & Dharaiya, N. (2010). Seasonal dietary composition of sloth bear (*Melursus ursinus*) in the reserve forest of Vijaynagar, North Gujarat, India. Regional office for Asia and the Pacific.

- Mewada, T.P., Tiwari, U. L., & Kotia, A. (2019). Fruiting species influence the seasonal use of the habitat by sloth bear in and around Balaram Ambaji Wildlife Sanctuary, Gujarat, India. *Notulae Scientia Biologicae*, 11(2):183-190.
- Miller, J. R. B. (2015). Mapping attack hotspots to mitigate human–carnivore conflict: approaches and applications of spatial predation risk modeling. *Biodiversity & Conservation*. 24:2887–2911. doi: 10.1007/s10531-015-0993-6.
- Milleret, C., Ordiz, A., Chapron, G., Andreassen, H. P., Kindberg, J., Månsson, J., Tallian, A., Wabakken, P., Wikenros, C., Zimmermann, B., Swenson, J. E., & Sand, H. (2018). Habitat segregation between brown bears and gray wolves in a human-dominated landscape. *Ecology and Evolution*, 8(23):11450–11466.
- Morell, N., Appleton, R. D. & Arcese, P. (2021). Roads, forest cover, and topography as factors affecting the occurrence of large carnivores: The case of the Andean bear (*Tremarctos ornatus*). *Global Ecology and Conservation*, 26, e01473.
- Morelle, K., and Lejeune, P. (2015). Seasonal variations of wild boar *Sus scrofa* distribution in agricultural landscapes: a species distribution modelling approach. *European Journal of Wildlife Research*, 61:45–56. doi: 10.1007/s10344-014-0872-6
- Mukherjee, S., Duckwoth, J. W., Silwa, A., Appel, A., & Kittle, A. (2017). *Prionailurus rubiginosus*. The IUCN Red List of Threatened Species 2016: e. T18149A50662471.
- Naim Akhtar, N. A., & Chauhan, N. P. S. (2008). Status of human-wildlife conflict and mitigation strategies in Marwahi Forest Division, Bilaspur Chhattisgarh, 1349-1358.

- Nichols, J. D., Hines, J. E., Mackenzie, D. I., Seamans, M. E., & Gutierrez, R. J. (2007). Occupancy estimation and modeling with multiple states and state uncertainty. *Ecology*, 88(6),1395–1400.
- NWA (National Wetland Atlas): India, SAC/EPISA/ABHG/NWIA/ATLAS/34/2011, Space Applications Centre, ISRO, Ahmedabad.
- Odden, M., Athreya, V., Rattan, S., & Linnell, J. D. (2014). Adaptable neighbours: movement patterns of GPS-collared leopards in human dominated landscapes in India. *PLoS One*, 9(11), e112044.
- Open Street Map. [www.openstreetmap.org](http://www.openstreetmap.org) Accessed on 9<sup>th</sup> July 2023.
- Ordiz, A., Kindberg, J., Sæbø, S., Swenson, J. E. & Støen, O. (2014). Brown bear circadian behavior reveals human-environmental encroachment. *Biological Conservation*, 173:1–9.
- Oriol-Cotterill, A., Valeix, M., Frank, L. G., Riginos, C., & Macdonald, D. W. (2015). Landscapes of coexistence for terrestrial carnivores: The ecological consequences of being downgraded from ultimate to penultimate predator by humans. *Oikos*, 124(10):1263–1273.
- Padalia, H., V. Srivastava, & Kushwaha, S. (2014). Modeling potential invasion range of alien invasive species, *Hyptis suaveolens* (L.) Poit. in India: comparison of MaxEnt and GARP. *Ecological Informatics*, 22:36-43. <https://doi.org/10.1016/j.ecoinf.2014.04.002>
- Palita, S. K., Kar, T., & Debata, S. (2014). Human-Sloth Bear Interactions: Preliminary survey from Semiliguda Range of Koraput Forest Division, Southern Odisha, India. *Journal of Zoological Society of Orissa*, 26:13-21.
- Pandya, T.M. and Oza, G.M. (1998). Bioregion Common Property Resource Management Studies. *International Society of Naturalists*, Vadodara, India. 197pp.

- Pant, B., Sharma, H. P., Dahal, B. R., Regmi, S., & Belant, J. L. (2023). Spatio-temporal patterns of human-wildlife conflicts and effectiveness of mitigation in Shuklaphanta National Park, Nepal. *PLoS One*, 18(4), e0282654.
- Pariwakam, M., Joshi, A., Navgire, S. & Vaidyanathan, S. (2018): A policy framework for connectivity conservation and smart green linear infrastructure development in the Central Indian and Eastern Ghats tiger landscape. Wildlife Conservation Trust (WCT), Mumbai.
- Patel, K. (2019). *Assessment of food resource sharing between human and sloth bear in Jessore wildlife sanctuary: An approach to understand human-sloth bear interaction*. MSc thesis. Hemchandracharya North Gujarat University, Gujarat, India.
- Patel, P. M., Gami, B. & Patel, B. (2015). Seasonal impact on physical-chemical properties of soil in North and South Gujarat. *IOSR Journal of Agriculture and Veterinary Science*, 8(6):26–30.
- Paudel, R.P., Kadariya, R., Lamichhane, B.R., Subedi, N., Sashika, M., Shimosuru, M., & Tsubota, T. (2022). Habitat occupancy of sloth bear *Melursus ursinus* in Chitwan National Park, Nepal. *Ecology & Evolution*, 12(3):e8699.
- Pearson, R. G., Raxworthy, C. J., Nakamura, M., & Peterson, A. T. (2007). Predicting species distributions from small numbers of occurrence records: A test case using cryptic geckos in Madagascar. *Journal of Biogeography*, 34:102–117.
- Penteriani, V., & M. Melletti, eds. (2020). *Bears of the world: ecology, conservation and management*. Cambridge University Press, Cambridge.
- Penteriani, V., Bombieri, G., del Mar Delgado M, et al. (2020). Patterns of Bear Attacks on Humans, Factors Triggering Risky Scenarios, and How to Reduce Them. In: Penteriani V, Melletti M, eds. *Bears of the*

*World: Ecology, Conservation and Management*. Cambridge: Cambridge University Press, pp. 239-249. doi:10.1017/9781108692571.018

- Peterson, A. T., & Robins, C. R. (2003). Using Ecological-Niche Modeling to Predict Barred Owl Invasions with Implications for Spotted Owl Conservation. *Conservation Biology*, 17:1161-1165.
- Phillips, S. J. (2005). A brief tutorial on Maxent. AT&T Research 190:231-259.  
[https://biodiversityinformatics.amnh.org/open\\_source/maxent/Maxent\\_tutorial2017.pdf](https://biodiversityinformatics.amnh.org/open_source/maxent/Maxent_tutorial2017.pdf)
- Phillips, S. J., & Dudík, M. (2008). Modeling of species distributions with Maxent: new extensions and a comprehensive evaluation. *Ecography*, 31(2):161–175. <https://doi.org/10.1111/j.0906-7590.2008.5203.x>
- Phillips, S. J., Anderson, R. P., Dudík, M., Schapire, R. E., & Blair, M. E. (2017). Opening the black box: an open-source release of Maxent. *Ecography*, 40(7):887–893. <https://doi.org/10.1111/ecog.03049>
- Phillips, S. J., Anderson, R. P., and Schapire, R. E. (2006). Maximum entropy modeling of species geographic distributions. *Ecological Modelling* 190(3-4):231-259. <https://doi.org/10.1016/j.ecolmodel.2005.03.026>
- Plumptre, A. J. (2000). Monitoring mammal populations with line transect techniques in African forests. *Journal of Applied Ecology*, 37:356–368.
- Pokharel, M., Subba, A., Rai, D., Bhandari, S., & Ghimirey, Y. (2022). Fine-scale ecological and anthropogenic variables predict the habitat use and detectability of sloth bears in the Churia habitat of east Nepal. *Ecology & Evolution*, 12(1):e8512.
- Pradhan, P. M., Ghose, M. K. & Kharke, Y. R. (2012). Automatic association of Strahler's order and attributes with the drainage system.

*International Journal of Advanced Computer Science and Applications*, 3(8):30–33.

- Proosdij, A. J., Sosef, M., Wieringa, J., & Raes, N. (2016). Minimum required number of specimen records to develop accurate species distribution models. *Ecography*, 39 :542:552.
- Puri, M., Srivathsa, A., Karanth, K. K., Kumar, N. S., & Karanth, K. U. (2015). Multiscale distribution models for conserving widespread species: the case of sloth bear *Melursus ursinus* in India. *Diversity and Distributions*, 21(9):1087–1100. <https://doi.org/10.1111/ddi.12335>
- Puri, M., Srivathsa, A., Karanth, K. K., Patel, I., & Kumar, N. S. (2023). Safe space in the woods: Mechanistic spatial models for predicting risks of human–bear conflicts in India. *Biotropica*, 55(2):504-516.
- Quigley, H., & Herrero, S. (2005). Characteristics and prevention of attacks on humans. In: Woodroffe R, Thirgood S, Rabinowitz A (eds) *People and wildlife: conflict or coexistence?* Cambridge University Press, Cambridge, pp 27–48.
- Rajpurohit, K. S. & Chauhan, N. P. S. (1996). Study of animal damage problems in and around protected areas and managed forest in India. Phase I: Madhya Pradesh, Bihar and Orissa. Wildlife Institute of India, Dehradun.
- Rajpurohit, K. S. & Krausman, P. R. (2000). Human–sloth-bear conflicts in Madhya Pradesh, India. *Wildlife Society Bulletin*, 28: 393–399.
- Ramesh, T., Kalle, R., Sankar, K., & Qureshi, Q. (2012). Factors affecting habitat patch use by sloth bears in Mudumalai Tiger Reserve, Western Ghats, India. *Ursus*, 23(1):78–85. <https://doi.org/10.2192/ursus-d-11-00006.1>
- Ratnayeke, S., van Manen, F. T. & Padmalal, U. K. G. K. (2007)a. Home ranges and habitat use of sloth bears *Melurus Ursinus Inornatus* in Wasgomuwa National Park, Sri Lanka. *Wildlife Biology*, 13(3):272–284.

- Ratnayeke, S., van Manen, F. T., Pieris, R., & Pragash, V. S. J. (2007)b. Landscape characteristics of sloth bear range in Sri Lanka. *Ursus*, 18(2):189-202.
- Ratnayeke, S., van Manen, F. T., Pieris, R., & Pragash, V. S. (2014). Challenges of large carnivore conservation: sloth bear attacks in Sri Lanka. *Human Ecology*, 42(3):467-479.
- Reddy, P. A., Puyravaud, J. P., Cushman, S. A., & Segu, H. (2019). Spatial variation in the response of tiger gene flow to landscape features and limiting factors. *Animal Conservation*, 22(5):472-480.
- Regmi, S., Belant, J. L., Pant, B., & Sharma, H. P. (2023). Factors influencing mammalian community occupancy in Dhorpatan Hunting Reserve, Nepal. *Ecology and Evolution*, 13(4): e9980.
- Rodgers, W. A., Panwar, H. S., & Mathur, V. B. (2002). Wildlife protected area network in India: A Review (Executive Summary). Wildlife Institute of India, Dehradun, India. 44 pp.
- Rot, J., Jangid, A. K., Singh, C. P., & Dharaiya, N. A. (2023). Escaping Neobiota: Habitat use and avoidance by sloth bears in Jessore Sloth bear Sanctuary India. *Trees, Forests and People*, 100400.be
- Rudnick, D. A., Sadie, J. R., Beier, P., Cushman, S. A., Dieffenbach, F., Epps, C. W., Gerber, L. et al. (2012). The role of landscape connectivity in planning and implementing conservation and restoration priorities. *Issues in ecology*, 16: 1-23.
- Saaty, T. L. (2004). Decision making—the analytic hierarchy and network processes (AHP/ANP). *J Syst Sci Syst.*, 13(1):1-35.
- Sahlén, E., Støen, O.-G., & Swenson, J. E. (2011). Brown bear den site concealment in relation to human activity in Sweden. *Ursus*. International Association for Bear Research and Management. <https://doi.org/10.2307/41304067>

- Sathyakumar, S., Kaul, R., Ashraf, N. V. K., Mookerjee, A., & Menon, V. (2012). National bear conservation and welfare action plan. Ministry of Environment and Forests, Wildlife Institute of India and Wildlife Trust of India, Delhi, India.
- Sawyer, S. C., Epps, C. W., & Brashares, J. S. (2011). Placing linkages among fragmented habitats: do least-cost models reflect how animals use landscapes? *Journal of Applied Ecology*, 48(3):668-678.
- Schwartz, M. K., Cushman, S. A., Mckelvey, K. S., Hayden, J. & Engkjer, C. (2006). Detecting genotyping errors and describing American black bear movement in northern Idaho. *Ursus*, 17(2):138–148.
- Seidensticker, J., Yoganand, K., & Johnsingh, A. J. T. (2011). Sloth bears living in seasonally dry tropical and moist broadleaf forests and their conservation. *The Ecology and Conservation of Seasonally Dry Forests in Asia; McShea, WJ, Davies, S., Bhumpakphan, N., Eds*, pp. 217-236.
- Sharma, P., N. Chettri, K. Uddin, K. Wangchuk, R. Joshi, T. Tandin, A. Pandey, K. S. Gaira, K. Basnet, & Wangdi, S. (2020). Mapping human–wildlife conflict hotspots in a transboundary landscape, Eastern Himalaya. *Global Ecology and Conservation* 24: e01284. <https://doi.org/10.1016/j.gecco.2020.e01284>.
- Sharma, S., Dutta, T., Maldonado, J. E., Wood, T. C., Panwar, H. S., & Seidensticker, J. (2013). Forest corridors maintain historical gene flow in a tiger metapopulation in the highlands of central India. *Proceedings of the Royal Society*; 280 (1767): 20131506. <http://dx.doi.org/10.1098/rspb.2013.1506>
- Sharp, T. R., Smith, T. S., Swaminathan, S., & Arun, A. S. (2022). Sloth bear attacks: regional differences and safety messaging. *Scientific Reports*, 12(1): 3943.

- Sharp, T. R., Swaminathan, S., Arun, A. S., Smith, T., Satyanarayan, K., & Seshamani, G. (2020). Sloth bear attacks on the Deccan Plateau of Karnataka, India. *Ursus*, 2020(31e8):1-11.
- Sillero-Zubiri, C., & Laurenson, M. K. (2001). Interactions between carnivores and local communities: Conflict or co-existence? *Conservation Biology Series*, 5:282–312.
- Simpsons, N. O., Stewart, K. M. & Bleich, V. C. (2011). What have we learned about water developments for wildlife? Not enough! *California Fish and Game*, 97(4):190–209.
- Singh, H. (2008). Biological Diversity in Gujarat. *Gujarat, Perspectives of the Future*, 1(16):247.
- Singh, H., & Kariyappa, A. (2020). Records of Rusty-Spotted Cat *Prionailurus rubiginosus* (I. Geoffroy Saint-Hilaire, 1831) (Mammalia: Carnivora: Felidae) in Mount Abu Wildlife Sanctuary, Rajasthan, India. *Journal of Threatened Taxa*, 12(16):17258-17262.
- Singh, N., Sonone, S. & Dharaiya, N. (2018). Sloth bear attacks on humans in central India: Implications for species conservation. *Human Wildlife Interactions*, 12:338–347.
- Smith P. M. Q., & Ruxton, G. D. (2020). Effective use of the McNemar test. *Behavioral Ecology and Sociobiology*, 74:1-9.
- Socioeconomic Data and Application Centre, SEDAC (<https://sedac.ciesin.columbia.edu/>)
- Soga, M., & Gaston, K. J. (2020). The ecology of human–nature interactions. *Proceedings of the Royal Society B: Biological Sciences*, 287(1918):20191882.
- Soliku, O., & Schraml, U. (2020). Protected areas management: a comparison of perceived outcomes associated with different co-management types. *Forest Policy and Economics*, 118:102258.

- Srivastava, A., Raypa, S., & Singhvi, V. K. (2022). Rainfall variability over Gujarat state 2022. *Published by Gujarat Institute of Disaster Management, Gandhinagar, Gujarat, ISBN 978-81-957623-5-4.*
- Srivathsa, A., Puri, M., Kumar, N. S., Jathanna, D., & Karanth, K. U. (2018). Substituting space for time: Empirical evaluation of spatial replication as a surrogate for temporal replication in occupancy modelling. *Journal of Applied Ecology*, 55(2):754–765.
- Sukhadiya, D., Joshi, J. U., & Dharaiya, N. (2013). Feeding ecology and habitat use of sloth bear (*Melursus ursinus*) in Jassore Wildlife Sanctuary, Gujarat, India. *Indian Journal of Ecology*, 40(1):14-18.
- Sultana, F., Khan, S. & Nabi, G. (2015). Occupancy and habitat use of sloth bear (*Melursus ursinus*) in Mukundara Hills Tiger Reserve, Rajasthan, India. *Flora and Fauna*, 21:203–208.
- Sundjaja, J. H., Shrestha, R., & Krishan, K. (2023). McNemar, & Mann-Whitney U Tests. [Updated 2023 Jul 17]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing;2024Jan-Availablefrom: <https://www.ncbi.nlm.nih.gov/books/NBK560699/>
- Suraci, J. P., Nickel, B. A., & Wilmers, C. C. (2020). Fine-scale movement decisions by a large carnivore inform conservation planning in human-dominated landscapes. *Landscape Ecology*, 35:1635–1649.
- Tarjuelo, R., Morales, M. B., Traba, J., & Delgado, M. P. (2014). Are species coexistence areas a good option for conservation management? Applications from fine scale modelling in two steppe birds. *Plos one*, 9(1), e87847.
- Thatte, P., Chandramouli, A., Tyagi, A., Patel, K., Baro, P., Chhattani, H., & Ramakrishnan, U. (2020). Human footprint differentially impacts genetic connectivity of four wide-ranging mammals in a fragmented landscape. *Diversity and Distributions*, 26(3):299-314.

- Thatte, P., Tyagi, A., Neelakantan, A., Natesh, M., Sen, M., & Thekaekara, T. (2021). Trends in wildlife connectivity science from the biodiverse and human-dominated South Asia. *Journal of the Indian Institute of Science*, 101(2):177-193.
- Tischendorf, L., & Fahrig, L. (2000). On the usage and measurement of landscape connectivity. *Oikos*, 90(1): 7-19.
- Treves, A., & Karanth, K. U. (2003). Human-carnivore conflict and perspectives on carnivore management worldwide. *Conservation Biology*, 17(6):1491-1499.
- Tucker, M. A., Böhning-Gaese, K., Fagan, W. F., Fryxell, J. M., Van Moorter, B., Alberts, S. C., ... & Mueller, T. (2018). Moving in the Anthropocene: Global reductions in terrestrial mammalian movements. *Science*, 359(6374), 466-469.
- Vyas, R., & Upadhyay, K. (2014). Sighting and distribution of Rusty spotted Cat in Gujarat State, India. *Cat News*, 61: 26-29.
- Wang, R. L., Li, Q., Feng, C. H., & Shi, Z. P. (2017). Predicting potential ecological distribution of *Locusta migratoria tibetensis* in China using MaxEnt ecological niche modeling. *Acta Ecologica Sinica*, 37(24):8556–856
- Wang, S., Wu, M., Hu, M., Fan, C., Wang, T., & Xia, B. (2021). Promoting landscape connectivity of highly urbanized area: An ecological network approach. *Ecological Indicators*, 125:107487.
- Warriar, R., Noon, B. R., & Bailey, L. L. (2021). A framework for estimating human-wildlife conflict probabilities conditional on species occupancy. *Frontiers in Conservation Science*, 2:679028.
- White, G. C., & Burnham, K. P. (1999). Program MARK: survival estimation from populations of marked animals. *Bird study*, 46(sup1): S120-S139.

- Wisz, M. S., Pottier, J., Kissling, W. D., Pellissier, L., Lenoir, J., Damgaard, C. F., ... & Svenning, J. C. (2013). The role of biotic interactions in shaping distributions and realised assemblages of species: implications for species distribution modelling. *Biological Reviews*, 88(1): 15-30.
- Yoganand, K., Rice, C. G., & Johnsingh, A. J. T. (2005). Evaluating Panna National Park with special reference to ecology of sloth bear (*Melursus ursinus*). Final project report. Wildlife Institute of India, Dehradun, India.
- Yoganand, K., Rice, C. G., Johnsingh, A. J. T., & Seidensticker, J. (2006). Is the sloth bear in India secure? A preliminary report on distribution, threats and conservation requirements. *The Journal of the Bombay Natural History Society*: 103 (2-3).
- Young, J. C., Marzano, M., White, R. M., McCracken, D. I., Redpath, S. M., Carss, D. N., ... Watt, A. D. (2010). The emergence of biodiversity conflicts from biodiversity impacts: Characteristics and management strategies. *Biodiversity and Conservation*, 19(14):3973–3990. <https://doi.org/10.1007/s10531-010-9941-7>
- Zvidzai, M., Mawere, K. K., N'andu, R. J., Ndaimani, H., Zanamwe, C., & Zengeya, F. M. (2023). Application of maximum entropy (MaxEnt) to understand the spatial dimension of human–wildlife conflict (HWC) risk in areas adjacent to Gonarezhou National Park of Zimbabwe. *Ecology and Society*, 28(3). <https://doi.org/10.5751/ES-14420-280318>