

# Bibliography

## 8. Bibliography

- Abdi, H., & Williams, L. J. (2010). Principal component analysis. *Wiley Interdisciplinary Reviews: Computational Statistics*, 2(4), 433–459. <https://doi.org/10.1002/WICS.101>
- Adams, D. C., Rohlf, F. J., & Slice, D. E. (2004). Geometric morphometrics: Ten years of progress following the ‘revolution’. *Italian Journal of Zoology*, 71(1), 5–16. <https://doi.org/10.1080/11250000409356545>
- Adebowale, A., Naidoo, Y., Lamb, J., & Nicholas, A. (2014). Comparative foliar epidermal micromorphology of Southern African *Strychnos* L. (Loganiaceae): Taxonomic, ecological and cytological considerations. *Plant Systematics and Evolution*, 300(1), 127–138. <https://doi.org/10.1007/S00606-013-0865-Z>
- Alagumariappan, P., Dewan, N. J., Muthukrishnan, G. N., Raju, B. K. B., Bilal, R. A. A., & Sankaran, V. (2020). Intelligent Plant Disease Identification System Using Machine Learning. *Engineering Proceedings*, 2(1). <https://doi.org/10.3390/ecsa-7-08160>
- Andrade, I. M., Mayo, S. J., Kirkup, D., & Van Den Berg, C. (2008). Comparative morphology of populations of *Monstera* Adans. (Araceae) from natural forest fragments in Northeast Brazil using elliptic Fourier Analysis of leaf outlines. *Kew Bulletin*, 63(2), 193–211. <https://doi.org/10.1007/S12225-008-9032-Z>
- APG – IV. Angiosperm Phylogeny Group. Chase, M. W., Christenhusz, M. J., Fay, M. F., Byng, J. W., Judd, W. S., ... & Stevens, P. F. (2016). An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants: APG IV. *Botanical journal of the Linnean Society*, 181(1), 1-20. <https://doi.org/10.1111/boj.12385>
- Austin, D. F. (1975a). Convolvulaceae. In R. E. J. Woodson & R. W. Schery (Eds.), *Flora of Panama* (Vol. 9, pp. 157–224). *Annals of the Missouri Botanical Garden*.
- Austin, D. F. (1975b). Typification of the New World Subdivisions Of *Ipomoea* L. (Convolvulaceae). *Taxon*, 24(1), 107–110. <https://doi.org/10.2307/1219005>
- Austin, D. F. (1979). An Infrageneric Classification for *Ipomoea* (Convolvulaceae). In *Source: Taxon*, 28 (4): 359-961.
- Austin, D. F. (1980). Convolvulaceae. In M. D. Dassanayake & F. R. Fosberg (Eds.), *A revised handbook to the flora of Ceylon*, Vol. 1, pp. 288–363. Amerind Publication Co.
- Austin, D. F. (1982). Convolvulaceae. In G. Harling & B. Sparre (Eds.), *Flora of Ecuador*, Vol. 15, pp. 3–99. University of Goteborg.
- Austin, D. F. (2007). Water Spinach (*Ipomoea aquatica*, Convolvulaceae): A Food Gone Wild. In *Ethnobotany Research & Applications* (Vol. 5, pp. 123–146). University of Hawaii at Manoa.
- Austin, D. F., & Ghazanfar, S. (1979). Convolvulaceae. In E. Nasir & S. I. Ali (Eds.), *Flora of West Pakistan*, pp. 1–64. Agricultural Research Council.

- Babu, A. K., Kumaresan, G., Raj, V. A. A., & Velraj, R. (2018). Review of leaf drying: Mechanism and influencing parameters, drying methods, nutrient preservation, and mathematical models. *Renewable and Sustainable Energy Reviews*, 90: 536–556. <https://doi.org/10.1016/J.RSER.2018.04.002>
- Belhumeur, P. N., Chen, D., Feiner, S., Jacobs, D. W., Kress, W. J., Ling, H., Lopez, I., Ramamoorthi, R., Sheorey, S., White, S., & Zhang, L. (2008). Searching the World's Herbaria: A System for Visual Identification of Plant Species. In: D. Forsyth, P. Torr, and A. Zisserman (Eds.): ECCV 2008, Part IV, LNCS 5305, pp. 116–129. Springer-Verlag Berlin Heidelberg.
- Belo, D. P., Louzada, R. B., & Buriel, M. T. (2024). Alpha-taxonomy underestimates the diversity in *Jacquemontia* (Convolvulaceae): a new hypothesis for *Jacquemontia evolvuloides* complex based on morphological, anatomical, and morphometric analyses. *Plant Biosystems*, 1–19. <https://doi.org/10.1080/11263504.2024.2347853>
- Benítez, H. A., Lemic, D., Villalobos-Leiva, A., Bažok, R., Órdenes-Claveria, R., Živković, I. P., & Mikac, K. M. (2020). Breaking symmetry: Fluctuating asymmetry and geometric morphometrics as tools for evaluating developmental instability under diverse agroecosystems. *Symmetry*, 12(11): 1–13. <https://doi.org/10.3390/sym12111789>
- Bentham, G., & Hooker, J. D. (1876). *Genera Plantarum* (Vol. 2). Reeves & Co.
- Biju, C., & Dhar, T. P. (2020). Palynological studies in species of *Ipomoea* linn. *The Journal of Indian Botanical Society*, 99(1–2): 86–91. <https://doi.org/10.5958/2455-7218.2020.00010.8>
- Bisen, D. (2021). Deep convolutional neural network-based plant species recognition through features of leaf. *Multimedia Tools and Applications*, 80(4): 6443–6456. <https://doi.org/10.1007/s11042-020-10038-w>
- Blackith, R. E., & Reyment, R. A. (1971). *Multivariate Morphometrics*. Academic Press Inc.
- Bookstein, F. L. (1982). Foundations of morphometrics. *Annual Review of Ecology and Systematics*, 13: 451–470. <https://doi.org/10.1146/ANNUREV.ES.13.110182.002315>
- Bookstein, F. L. (1991). Morphometric Tools for Landmark Data: Geometry and Biology. In *Morphometric Tools for Landmark Data*. Cambridge University Press. <https://doi.org/10.1017/CBO9780511573064>
- Bookstein, F. L. (1996). Combining the Tools of Geometric Morphometrics. In F. M. Leslie, M. Corti, A. Loy, G. J. P. Naylor, & D. E. Slice (Eds.), *Advances in Morphometrics* (pp. 131–151). Springer, Boston, MA. [https://doi.org/10.1007/978-1-4757-9083-2\\_12](https://doi.org/10.1007/978-1-4757-9083-2_12)
- Bookstein, F. L. (2021). A New Method for Landmark-Based Studies of the Dynamic Stability of Growth, with Implications for Evolutionary Analyses. *Evolutionary Biology*, 48(4): 428–457. <https://doi.org/10.1007/S11692-021-09548-8>

- Bookstein, F. L., & Mitteroecker, P. (2014). Comparing Covariance Matrices by Relative Eigenanalysis, with Applications to Organismal Biology. *Evolutionary Biology*, 41(2): 336–350. <https://doi.org/10.1007/S11692-013-9260-5>
- Budd, G. E. (2006). On the origin and evolution of major morphological characters. *Biological Reviews*, 81(4): 609–628. <https://doi.org/10.1017/S1464793106007135>
- Buril, M. T., Oliveira, P. P., Rodrigues, R., Santos, F. de A. R. Dos, & Alves, M. (2015). Pollen morphology and taxonomic implications in *Jacquemontia Choisy* (Convolvulaceae). *Grana*, 54(1): 1–11. <https://doi.org/10.1080/00173134.2014.946961>
- Cáez-Ramírez, G., Alamilla-Beltrán, L., & Gutiérrez-López, G. F. (2018). Morphometric analysis and tissue structural continuity evaluation of senescence progression in fresh cut papaya (*Carica papaya* L.). *Journal of Food Engineering*, 216: 107–119. <https://doi.org/10.1016/J.JFOODENG.2017.08.004>
- Carlquist, S., & Hanson, M. A. (1991). Wood and Stem Anatomy of Convolvulaceae. *Aliso: A Journal of Systematic and Floristic Botany*, 13(1): 51–94. <https://doi.org/10.5642/aliso.19911301.03>
- Chen, Y., Jabbour, F., Novikov, A., Wang, W., & Gerber, S. (2018). A study of floral shape variation in Delphinieae (Ranunculaceae) using geometric morphometrics on herbarium specimens. *Botany Letters*, 165(3–4): 368–376. <https://doi.org/10.1080/23818107.2018.1427145>
- Chitwood, D. H., & Otoni, W. C. (2017). Morphometric analysis of *Passiflora* leaves: the relationship between landmarks of the vasculature and elliptical Fourier descriptors of the blade. *GigaScience*, 6(1). <https://doi.org/10.1093/GIGASCIENCE/GIW008>
- Chitwood, D. H., & Sinha, N. R. (2016). Evolutionary and Environmental Forces Sculpting Leaf Development. *Current Biology*, 26: 297–306. <https://doi.org/10.1016/j.cub.2016.02.033>
- Choisy, J. D. (1833). *Convolvulaceis orientalis*. *Mem. Soc. Phys. Geneve*.
- Cooke, Theodore. (1908). The flora of the presidency of Bombay. In *The flora of the presidency of Bombay / By Theodore Cooke. (Vol. 2)*. Taylor and Francis. <https://doi.org/https://doi.org/10.5962/bhl.title.10884>
- Cope, J. S., Corney, D., Clark, J. Y., Remagnino, P., & Wilkin, P. (2012a). Plant species identification using digital morphometrics: A review. *Expert Systems with Applications*, 39(8), 7562–7573. <https://doi.org/10.1016/J.ESWA.2012.01.073>
- Cronquist, A. (1968). *The evolution and classification of flowering plants*. New York Botanical Garden.
- Dahlgren, G. (1989). An updated angiosperm classification. *Botanical Journal of the Linnean Society*, 100(3): 197–203. <https://doi.org/10.1111/J.1095-8339.1989.TB01717.X>

- Du, J. X., Wang, X. F., & Zhang, G. J. (2007). Leaf shape based plant species recognition. *Applied Mathematics and Computation*, 185(2): 883–893. <https://doi.org/10.1016/j.amc.2006.07.072>
- Ferguson, I. K., Verdcourt, B., & Poole, M. M. (1977). Pollen Morphology in the Genera *Merremia* and *Operculina* (Convolvulaceae) and Its Taxonomic Significance. *Kew Bulletin*, 31(4): 763. <https://doi.org/10.2307/4109548>
- Fisher, R. (1954). The Analysis of Variance with Various Binomial Transformations. *Biometrics*, 10(1): 130. <https://doi.org/10.2307/3001667>
- Fleming, A. J. (2005). The control of leaf development. *New Phytologist*, 166(1): 9–20. <https://doi.org/10.1111/J.1469-8137.2004.01292.X>
- Flückiger, F. A., & Hanbury, D. (1874). *Pharmacographia: A History of the Principal Drugs of Vegetable Origin*. MacMillan and Co.
- Furuta, N., Ninomiya, S., Takahashi, N., Ohmori, H., & Ukai, Y. (1995). Quantitative Evaluation of Soybean (*Glycine max* L. Merr.) Leaflet Shape by Principal Component Scores Based on Elliptic Fourier Descriptor. *Japanese Journal of Breeding*, 45(3): 315–320. <https://doi.org/10.1270/JSBBS1951.45.315>
- Galton, F. (1889). Co-relations and their measurement, chiefly from anthropometric data. *Proceedings of the Royal Society of London*, 45(273–279): 135–145. <https://doi.org/10.1098/rspl.1888.0082>
- Galton, F. (1907). *Inquiries into Human Faculty and Its Development* (2nd ed.). MacMillan and Co.
- Gamble, J. (1923). *Flora of Presidency of Madras* (Vol. 5).
- George, W. (1893). *A Dictionary of the Economic Products of India* (Vol. 6). Cambridge University Press.
- Goodall, C. (1991). Procrustes Methods in the Statistical Analysis of Shape. *Journal of the Royal Statistical Society: Series B (Methodological)*, 53(2): 285–321. <https://doi.org/10.1111/J.2517-6161.1991.TB01825.X>
- Gornall, R. J., & Bohm, B. A. (1978). Angiosperm Flavonoid Evolution: A Reappraisal. *Systematic Botany*, 3(4): 353. <https://doi.org/10.2307/2418744>
- Govil, C. M. (1972). Morphological studies in the family convolvulaceae - IV. Vascular anatomy of the flower. *Proceedings of the Indian Academy of Sciences - Section B*, 75(6): 271–282. <https://doi.org/10.1007/BF03045721>
- Griffin, W. J., & Lin, G. D. (2000). Chemotaxonomy and geographical distribution of tropane alkaloids. *Phytochemistry*, 53(6): 623–637. [https://doi.org/10.1016/S0031-9422\(99\)00475-6](https://doi.org/10.1016/S0031-9422(99)00475-6)
- Gurgel-Gonçalves, R., Ferreira, J. B. C., Rosa, A. F., Bar, M. E., & Galvão, C. (2011). Geometric morphometrics and ecological niche modelling for delimitation of near-sibling triatomine

- species. *Medical and Veterinary Entomology*, 25(1): 84–93. <https://doi.org/10.1111/J.1365-2915.2010.00920.X>
- Hallier, H. (1893). *Convolvulaceae africanae*.
- Hammer, O., Harper, D., & Ryan, P. (2001). PAST: Paleontological Statistics software package for education and data analysis (4.16c; pp. 1–9). *Palaeontologia Electronica*.
- Harborne, J. B. (2000). Arsenal for survival: secondary plant products. *TAXON*, 49(3): 435–449. <https://doi.org/10.2307/1224343>
- Hâruta, O. (2011). Elliptic Fourier analysis of crown shapes in *Quercus petraea* trees. *Annals of Forest Research*, 54(1), 99–117.
- Heniger, J. (1986). *Hendrik Adriaan van Reede tot Drakenstein (1636-1691) and Hortus Malabaricus*. Rotterdam & Boston.
- Hoogland, R. D. (1953). A review of the genus *Erycibe* Roxb. *Blumea: Biodiversity, Evolution and Biogeography of Plants*, 7(2), 342–361.
- Hooker, J. D. (1885). The flora of British India. In *The flora of British India* /By J. D. Hooker assisted by various botanists. Published under the authority of the secretary of state for India in council. (Vol. 4). L. Reeve.
- Hotelling, H. (1933). Analysis of a complex of statistical variables into principal components. *Journal of Educational Psychology*, 24(6), 417–441. <https://doi.org/10.1037/H0071325>
- Hutchinson, J. (1926). *The families of flowering plants* (Vol. 1). Clarendon Press.
- Huxley, J. S. (1950). Relative growth and form transformation. *Proceedings of the Royal Society of London. Series B - Biological Sciences*, 137(889), 465–469. <https://doi.org/10.1098/RSPB.1950.0055>
- Idrees, M., Pathak, M. L., Memon, N. H., Khan, S., Zhang, Z. Y., & Gao, X. F. (2021). Morphological And Morphometric Analysis of Genus *Eriobotrya* Lindl. (Rosaceae). *The Journal of Animal & Plant Sciences*, 31(4): 1087–1100. <https://doi.org/10.36899/JAPS.2021.4.0306>
- Igual, L., Perez-Sala, X., Escalera, S., Angulo, C., & De La Torre, F. (2014). Continuous Generalized Procrustes analysis. *Pattern Recognition*, 47(2): 659–671. <https://doi.org/10.1016/J.PATCOG.2013.08.006>
- Iwata, H., & Ukai, Y. (2002). SHAPE: A Computer Program Package for Quantitative Evaluation of Biological Shapes Based on Elliptic Fourier Descriptors. *Journal of Heredity*, 93(5): 384–385. <https://doi.org/10.1093/JHERED/93.5.384>
- Iwata, H., Niikura, S., Matsuura, S., Takano, Y., & Ukai, Y. (1998). Evaluation of variation of root shape of Japanese radish (*Raphanus sativus* L.) based on image analysis using elliptic Fourier

- descriptors. *Euphytica* 1998 102:2, 102(2): 143–149.  
<https://doi.org/10.1023/A:1018392531226>
- Jain, S. K., & Rao, R. R. (1976). *A Handbook of Field and Herbarium Methods* (1st ed.). Today and Tomorrow's publishers.
- Jensen, J. K., Rasmussen, L. H., Raulund-Rasmussen, K., & Borggaard, O. K. (2008). Influence of soil properties on the growth of sycamore (*Acer pseudoplatanus* L.) in Denmark. *European Journal of Forest Research*, 127(4): 263–274. <https://doi.org/10.1007/S10342-008-0202-1/FIGURES/5>
- Jensen, R. J. (2003). The Conundrum of Morphometrics. In *Source: Taxon*, 52(4), 663–671.
- Jumawan, J. H., & Buot, I. E. (2021). Leaf Outline Evaluation in Selected Philippine Hoya R. Br. Species (Apocynaceae) Using Elliptic Fourier Analysis. *Natural History Museum Journal*, 15(1): 21–30. <https://doi.org/10.14456/thnhmj.2021.2>
- Jungers, W. L., Falsetti, A. B., & Wall, C. E. (1995). Shape, relative size, and size-adjustments in morphometrics. *American Journal of Physical Anthropology*, 38(21) 137–161. <https://doi.org/10.1002/AJPA.1330380608>
- Jussieu, A. L. de. (1789). *Genera Plantarum Secundum Ordines Naturales*. Apud Viduam Herissant et Theophilum Barrois.
- Kattee, A. (2020). Taxonomic revision of the genus *Ipomoea* L. (Convolvulaceae) for India [Ph.D. Thesis]. Shivaji University.
- Kendall, D. G. (1989). A Survey of the Statistical Theory of Shape. *Statistical Science*, 4(2): 87–99. <https://doi.org/10.1214/SS/1177012582>
- Klein, L. L., Caito, M., Chapnick, C., Kitchen, C., O'Hanlon, R., Chitwood, D. H., & Miller, A. J. (2017). Digital morphometrics of two north American grapevines (*Vitis*: Vitaceae) quantifies leaf variation between species, within species, and among individuals. *Frontiers in Plant Science*, 8. 373: 1-10. <https://doi.org/10.3389/fpls.2017.00373>
- Klein, L., & Svoboda, H. (2017). Comprehensive Methods for Leaf Geometric Morphometric Analyses. *BIO-PROTOCOL*, 7(9): 1-21. <https://doi.org/10.21769/bioprotoc.2269>
- Klingenberg, C. P. (2011). MorphoJ: An integrated software package for geometric morphometrics. *Molecular Ecology Resources*, 11(2): 353–357. <https://doi.org/10.1111/j.1755-0998.2010.02924.x>
- Klingenberg, C. P., Duttke, S., Whelan, S., & Kim, M. (2012). Developmental plasticity, morphological variation and evolvability: a multilevel analysis of morphometric integration in the shape of compound leaves. *Journal of Evolutionary Biology*, 25(1): 115–129. <https://doi.org/10.1111/J.1420-9101.2011.02410.X>

- Koch, M. A., & German, D. A. (2013). Taxonomy and systematics are key to biological information: *Arabidopsis*, *Eutrema* (*Thellungiella*), *Noccaea* and *Schrenkiella* (*Brassicaceae*) as examples. *Frontiers in Plant Science*, 4, 54338. <https://doi.org/10.3389/FPLS.2013.00267>
- Kuhl, F. P., & Giardina, C. R. (1982). Elliptic Fourier features of a closed contour. *Computer Graphics and Image Processing*, 18(3): 236–258. [https://doi.org/10.1016/0146-664X\(82\)90034-X](https://doi.org/10.1016/0146-664X(82)90034-X)
- Kupe, M., Sayıncı, B., Demir, B., Ercisli, S., Baron, M., & Sochor, J. (2021). Morphological Characteristics of Grapevine Cultivars and Closed Contour Analysis with Elliptic Fourier Descriptors. *Plants*, 10(7): 1350. <https://doi.org/10.3390/PLANTS10071350>
- Lauder, G. V. (1990). Functional Morphology and Systematics: Studying Functional Patterns in an Historical Context. *Source: Annual Review of Ecology and Systematics*, 21: 317–340.
- Lawand, P. R. (2019). Taxonomic Revision of *Argyreia* Lour *Convolvulaceae* from India [Shivaji University].
- Lawand, P. R., Kapadane, K. K., Gondaliya, A. D., Shimpale, V. B., & Rajput, K. S. (2023). Formation of inter-and intraxylary phloem in some species of *Argyreia* Lour. (*Convolvulaceae*). *Flora*, 304: 152302. <https://doi.org/10.1016/J.FLORA.2023.152302>
- Lawing, A. M., Polly, P. D., David Polly, C. P., & Le Comber, S. (2010). Geometric morphometrics: recent applications to the study of evolution and development. *Journal of Zoology*, 280(1): 1–7. <https://doi.org/10.1111/J.1469-7998.2009.00620.X>
- Linnaeus, C. (1753). *Species Plantarum*, Vol. 2.
- Liu, Y., Li, Y., Song, J., Zhang, R., Yan, Y., Wang, Y., & Du, F. K. (2018). Geometric morphometric analyses of leaf shapes in two sympatric Chinese oaks: *Quercus dentata* Thunberg and *Quercus aliena* Blume (*Fagaceae*). *Annals of Forest Science*, 75(4): 1–12. <https://doi.org/10.1007/S13595-018-0770-2>
- Lo Bianco, R., & Mirabella, F. (2018). Use of Leaf and Fruit Morphometric Analysis to Identify and Classify White Mulberry (*Morus alba* L.) Genotypes. *Agriculture*, 8(10): 157. <https://doi.org/10.3390/AGRICULTURE8100157>
- Mabberley, D. (2017). *Mabberley's Plant-Book: A Portable Dictionary of Plants, Their Classification and Uses* (4th ed.). Cambridge University Press.
- Manos, P. S., Miller, R. E., & Wilkin, P. (2001). Phylogenetic Analysis of *Ipomoea*, *Argyreia*, *Stictocardia*, and *Turbina* Suggests a Generalized Model of Morphological Evolution in Morning Glories. 26(3): 585–602. <https://doi.org/10.1043/0363-6445-26.3.585>
- Marcus, L. (1990). Traditional Morphometrics. In F. J. Rohlf & F. L. Bookstein (Eds.), *Proceedings of the Michigan Morphometrics Workshop* (pp. 77–122). Museum of Zoology, The University of Michigan.

- Mccooy, M. W., Benjamin, A. E., Bolker, M., Ae, C. W. O., Miner, B. G., & Vonesh, J. R. (2006). Size correction: comparing morphological traits among populations and environments. *Oecologia*, 148: 547-554. <https://doi.org/10.1007/s00442-006-0403-6>
- McLellan, T., & Endler, J. (1998). The relative success of some methods for measuring and describing the shape of complex objects. *Systematic Biology*, 47(2): 264–281. <https://academic.oup.com/sysbio/article-abstract/47/2/264/1655073>
- Migicovsky, Z., Li, M., Chitwood, D. H., & Myles, S. (2018). Morphometrics Reveals Complex and Heritable Apple Leaf Shapes. *Frontiers in Plant Science*, 8. <https://doi.org/10.3389/FPLS.2017.02185>
- Mitteröcker, P. (2021). Morphometrics in Evolutionary Developmental Biology. *Evolutionary Developmental Biology*, 941–951. [https://doi.org/10.1007/978-3-319-32979-6\\_119](https://doi.org/10.1007/978-3-319-32979-6_119)
- Mitteroecker, P., & Bookstein, F. (2011). Linear Discrimination, Ordination, and the Visualization of Selection Gradients in Modern Morphometrics. *Evolutionary Biology*, 38(1): 100–114. <https://doi.org/10.1007/S11692-011-9109-8>
- Mitteroecker, P., & Schaefer, K. (2022). Thirty years of geometric morphometrics: Achievements, challenges, and the ongoing quest for biological meaningfulness. *American Journal of Biological Anthropology. Yearbook Biological Anthropology*, 178(Suppl. 74): 181–210. <https://doi.org/10.1002/ajpa.24531>
- Moles, A. T. (2018). Being John Harper: Using evolutionary ideas to improve understanding of global patterns in plant traits. *Journal of Ecology*, 106(1): 1–18. <https://doi.org/10.1111/1365-2745.12887>
- Moles, A. T., Perkins, S. E., Laffan, S. W., Flores-Moreno, H., Awasthy, M., Tindall, M. L., Sack, L., Pitman, A., Kattge, J., Aarssen, L. W., Anand, M., Bahn, M., Blonder, B., Cavender-Bares, J., Cornelissen, J. H. C., Cornwell, W. K., Díaz, S., Dickie, J. B., Freschet, G. T., ... Bonser, S. P. (2014). Which is a better predictor of plant traits: temperature or precipitation? *Journal of Vegetation Science*, 25(5): 1167–1180. <https://doi.org/10.1111/JVS.12190>
- Myint, T. (1966). Revision of the genus *Stylisma* (Convolvulaceae). *Brittonia*, 18(2): 97–117. <https://doi.org/10.2307/2805192>
- Myint, T., & Ward, D. B. (1968). A taxonomic revision of the genus *Bonamia* (Convolvulaceae). *Phytologia*, 17: 121–239.
- Neto, J. C., Meyer, G. E., Jones, D. D., & Samal, A. K. (2006). Plant species identification using Elliptic Fourier leaf shape analysis. *Computers and Electronics in Agriculture*, 50(2): 121–134. <https://doi.org/10.1016/J.COMPAG.2005.09.004>
- O’HIGGINS, P. (2000). The study of morphological variation in the hominid fossil record: biology, landmarks and geometry. *The Journal of Anatomy*, 197(1): 103–120. <https://doi.org/10.1046/J.1469-7580.2000.19710103.X>

- Pearson, K. (1897). Mathematical contributions to the theory of evolution. On a form of spurious correlation which may arise when indices are used in the measurement of organs. *Proceedings of the Royal Society of London*, 60(359–367), 489–498. <https://doi.org/10.1098/RSPL.1896.0076>
- Pessoa, E. M., Cordeiro, J. M. P., Felix, L. P., Almeida, E. M., Costa, L., Nepomuceno, Á., Souza, G., Chase, M. W., Alves, M., & Van Den Berg, C. (2020). Too many species: morphometrics, molecular phylogenetics and genome structure of a Brazilian species complex in *Epidendrum* (Laeliinae; Orchidaceae) reveal fewer species than previously thought. *Botanical Journal of the Linnean Society*, 195(2): 161–188. <https://doi.org/10.1093/botlinnean/boaa071>
- Peter, A. (1891). Convolvulaceae. In A. Engler & K. Prantl (Eds.), *Die Natürlichen Pflanzenfamilien* (3rd ed., Vol. 4, pp. 1–40). Wilhelm Engelmann.
- Pietrusewsky, M. (2018). Traditional Morphometrics and Biological Distance. *Biological Anthropology of the Human Skeleton*, 545–591. <https://doi.org/10.1002/9781119151647.CH17>
- Pires, N. D., & Dolan, L. (2012). Morphological evolution in land plants: new designs with old genes. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 367(1588): 508–518. <https://doi.org/10.1098/RSTB.2011.0252>
- Pisuttimarn, P., Simões, A. R. G., Petrongari, F. S., Simão-Bianchini, R., Barbosa, J. C. J., de Man, I., Fonseca, L. H. M., Janssens, S. B., Patil, S. B., Shimpale, V. B., Pornpongrungrueng, P., Leliaert, F., & Chatrou, L. W. (2023). *Distimake vitifolius* (Convolvulaceae): reclassification of a widespread species in view of phylogenetics and convergent pollen evolution. *Botanical Journal of the Linnean Society*, 202(3): 363–388. <https://doi.org/10.1093/BOTLINNEAN/BOAC077>
- Polly, P. D., Michelle, A. L., Anne-Claire, F., & Goswami, A. (2013). Phylogenetic Principal Components Analysis and Geometric Morphometrics. *Hystrix the Italian Journal of Mammalogy*, 24(1): 33–41. <https://doi.org/10.4404/hystrix-24.1-6383>
- POWO. (2024). *Plants of the World Online*. Facilitated by the Royal Botanic Gardens, Kew.
- Prain, D. (1906). Convolvulaceae. In K. George & J. S. Gamble (Eds.), *Materials for a Flora of the Malayan Peninsula*. *The Journal of Asiatic Society of Bengal*, 74: 284–327.
- Proietti, E., Filesi, L., Di Marzio, P., Di Pietro, R., Masin, R., Conte, A. L., & Fortini, P. (2021). Morphology, geometric morphometrics, and taxonomy in relict deciduous oaks woods in northern Italy. *Rendiconti Lincei*, 32(3): 549–564. <https://doi.org/10.1007/s12210-021-01001-4>
- Pryer, K. M., Tomasi, C., Wang, X., Meineke, E. K., & Windham, M. D. (2020). Using computer vision on herbarium specimen images to discriminate among closely related horsetails

- (Equisetum). Applications in Plant Sciences, 8(6): e11372. <https://doi.org/10.1002/aps3.11372>
- R Core Team. (2024). R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing.
- Rajput, K. S., Kapadane, K. K., Ramoliya, D. G., Thacker, K. D., & Gondaliya, A. D. (2022). Inter- and Intraxylary Phloem in Vascular Plants: A Review of Subtypes, Occurrences, and Development. *Forests* 2022, 3(12): 2174. <https://doi.org/10.3390/F13122174>
- Rajput, K. S., Ramoliya, D. G., Gondaliya, A. D., Shimpale, V. B., Patil, S. B., & Raole, V. M. (2024). Ontogeny of the multiple cambial variants and secondary growth in some species of *Merremia* and related genera (Convolvulaceae). *Plant Biosystems*, 158(3): 572–588. <https://doi.org/10.1080/11263504.2024.2347848>
- Ramoliya, D. G., Gondaliya, A. D., Raole, V. M., & Rajput, K. S. (2023). Structure and development of phloem in some species of *Merremia* Dennst. ex Endl. sensu lato (Convolvulaceae). *Nordic Journal of Botany*, 2023(2): e03776. <https://doi.org/10.1111/NJB.03776>
- Reich, P. B., & Oleksyn, J. (2004). Global patterns of plant leaf N and P in relation to temperature and latitude. *Proceedings of the National Academy of Sciences of the United States of America*, 101(30): 11001–11006. <https://doi.org/10.1073/PNAS.0403588101>
- Ren, T., He, N., Liu, Z., Li, M., Zhang, J., Li, A., Wei, C., Lü, X., & Han, X. (2021). Environmental filtering rather than phylogeny determines plant leaf size in three floristically distinctive plateaus. *Ecological Indicators*, 130: 108049. <https://doi.org/10.1016/J.ECOLIND.2021.108049>
- Revell, L. J. (2009). Size-Correction and Principal Components For Interspecific Comparative Studies. *Evolution*, 63(12): 3258–3268. <https://doi.org/10.1111/J.1558-5646.2009.00804.X>
- Reyment, R. (1990). Reification of Classical Multivariate Analysis in Morphometry. In F. J. Rohlf & F. L. Bookstein (Eds.), *Proceedings of Michigan Morphometrics Workshop* (pp. 123–144). Museum of Zoology, University of Michigan.
- Robertson, K. R. (1971). A revision of the genus *Jacquemontia* (Convolvulaceae) in North and Central America and the West Indies. Washington University in St. Louis.
- Rohlf, F. J. (1986). Relationships among eigen shape analysis, Fourier analysis, and analysis of coordinates. *Mathematical Geology*, 18(8): 845–854. <https://doi.org/10.1007/BF00899747>
- Rohlf, F. J. (1990a). An overview of Image Processing and Analysis Techniques for Morphometrics. In F. J. Rohlf & F. L. Bookstein (Eds.), *Proceedings of the Michigan Morphometrics Workshop*, pp. 37–60. Museum of Zoology, University of Michigan.

- Rohlf, F. J. (1990b). Fitting Curves to Outlines. In Rohlf F.J. & F. L. Bookstein (Eds.), *Proceeding of the Michigan Morphometrics Workshop*, pp. 167–178. Museum of Zoology, University of Michigan.
- Rohlf, F. J. (1999). Shape statistics: Procrustes superimpositions and tangent spaces. *Journal of Classification*, 16(2): 197–223. <https://doi.org/10.1007/S003579900054>
- Rohlf, F. J., & Archie, J. W. (1984). A Comparison of Fourier Methods for the Description of Wing Shape in Mosquitoes (Diptera: Culicidae). *Systematic Biology*, 33(3): 302–317. <https://doi.org/10.2307/2413076>
- Rohlf, F. J., & Bookstein, F. L. (1990). *Proceedings of the Michigan Morphometrics Workshop*.
- Rohlf, F. J., & Marcus, L. F. (1993). A Revolution in Morphometrics. *Trends in Ecology and Evolution*, 8(4): 1–4.
- Roxburgh, W. (1832). *Flora Indica Or Descriptions of Indian Plants*. Printed for W. Thacker and Co., Calcutta and Parbury Allen and Co., London. Serampore
- Royer, D. F., Lockwood, C. A., Scott, J. E., & Grine, F. E. (2009). Size variation in early human mandibles and molars from Klasies River, South Africa: Comparison with other middle and late Pleistocene assemblages and with modern humans. *American Journal of Physical Anthropology*, 140(2): 312–323. <https://doi.org/10.1002/AJPA.21071>
- RStudio Team. (2023). *RStudio: Integrated Development for R (2023.06.1+524)*. Posit.
- Sack, L., & Scoffoni, C. (2013). Leaf venation: structure, function, development, evolution, ecology and applications in the past, present and future. *New Phytologist*, 198(4): 983–1000. <https://doi.org/10.1111/NPH.12253>
- Sampathkumar, R. (1982). Studies on the Cotyledonary Leaves of Some Convolvulaceae. *Taxon*, 31(1): 53-56.
- Sattler, R., & Rutishauser, R. (2022). Fundamentals of Plant Morphology and Plant Evo-Devo (Evolutionary Developmental Morphology). *Plants*, 12(1):118. <https://doi.org/10.3390/PLANTS12010118>
- Savriama, Y., & Klingenberg, C. P. (2011). Beyond bilateral symmetry: geometric morphometric methods for any type of symmetry. *BMC Evolutionary Biology*, 11(1): 1–24. <https://doi.org/10.1186/1471-2148-11-280>
- Savriama, Y., Gómez, J. M., Perfectti, F., & Klingenberg, C. P. (2012). Geometric morphometrics of corolla shape: dissecting components of symmetric and asymmetric variation in *Erysimum mediohispanicum* (Brassicaceae). *New Phytologist*, 196(3): 945–954. <https://doi.org/10.1111/J.1469-8137.2012.04312.X>

- Sengupta, S. (1972). On the pollen morphology of convolvulaceae with special reference to taxonomy. *Review of Palaeobotany and Palynology*, 13(3–4): 157–212. [https://doi.org/10.1016/0034-6667\(72\)90030-9](https://doi.org/10.1016/0034-6667(72)90030-9)
- Shah, G. (1978). *Flora Of Gujarat State* (1st ed., Vol. 1). Sardar Patel University.
- Shanmukha Rao, S. R., & Leela, M. (1993). Seed morphology (LM and SEM) in some Ipomoea L. (Convolvulaceae). *Feddes Repertorium*, 104(3–4): 209–213.
- Simões, A. R. G., Eserman, L. A., Zuntini, A. R., Chatrou, L. W., Utteridge, T. M. A., Maurin, O., Rokni, S., Roy, S., Forest, F., Baker, W. J., & Stefanović, S. (2022). A Bird's Eye View of the Systematics of Convolvulaceae: Novel Insights from Nuclear Genomic Data. *Frontiers in Plant Science*, 13: 889988. <https://doi.org/10.3389/FPLS.2022.889988>
- Simões, A. R. G., Furness, C. A., & Da Luz, C. F. P. (2019a). The systematic value of pollen morphology in Operculina (Convolvulaceae). *Grana*, 58(1): 13. <https://doi.org/10.1080/00173134.2018.1511750>
- Simões, A. R. G., Pisuttimarn, P., da Luz, C. F. P., Furness, C., Pornpongrungrueng, P., & Chatrou, L. W. (2021a). Palynological characterization of the Southeast Asian woody climbers Decalobanthus Ooststr. (Convolvulaceae). *Grana*, 60(5): 356–369. <https://doi.org/10.1080/00173134.2021.1874512>
- Simões, A. R., & Staples, G. (2017). Dissolution of Convolvulaceae tribe Merremieae and a new classification of the constituent genera. In *Botanical Journal of the Linnean Society*, 183(4): 561–586. <https://doi.org/10.1093/botlinnean/box007>
- Simoies, A. R., Culham, A., & Carine, M. (2015). Resolving the unresolved tribe: a molecular phylogenetic framework for the Merremieae (Convolvulaceae). *Botanical Journal of the Linnean Society*, 179: 374–387. <https://doi.org/10.1111/boj.12339>
- Slice, D. E. (2005). Modern Morphometrics. In D. E. Slice (Ed.), *Modern Morphometrics in Physical Anthropology. Developments in Primatology: Progress and Prospects*, pp. 1–45. Springer, Boston, MA. [https://doi.org/10.1007/0-387-27614-9\\_1](https://doi.org/10.1007/0-387-27614-9_1)
- Srivastava, G., Mehrotra, R. C., & Dilcher, D. L. (2018). Paleocene ipomoea (convolvulaceae) from India with implications for an East gondwana origin of convolvulaceae. *Proceedings of the National Academy of Sciences of the United States of America*, 115(23): 6028–6033. <https://doi.org/10.1073/PNAS.1800626115>
- Staples, G. W. (1990). Preliminary taxonomic consideration of the Poraneae (Convolvulaceae). *Journal of the Arnold Arboretum*, 71(2): 251–258. <https://doi.org/10.5962/bhl.part.24932>
- Staples, G. W., & Traiperm, P. (2017). A nomenclatural review of Argyreia (Convolvulaceae). *Taxon*, 66(2): 445–477. <https://doi.org/10.12705/662.12>

- Staples, G. W., Simões, A. R., & Austin†, D. F. (2020). A Monograph of Operculina (Convolvulaceae). *Annals of the Missouri Botanical Garden*, 105(1): 64–138. <https://doi.org/10.3417/2020435>
- Stefanović, S., Krueger, L., & Olmstead, R. G. (2002). Monophyly of the Convolvulaceae and circumscription of their major lineages based on DNA sequences of multiple chloroplast loci. *American Journal of Botany*, 89(9): 1510–1522. <https://doi.org/10.3732/AJB.89.9.1510>
- Stegmann, M. B., & Delgado Gomez, D. (2002). A Brief Introduction to Statistical Shape Analysis.
- Sundberg, P. (1989). Shape and Size-Constrained Principal Components Analysis. *Systematic Zoology*, 38(2), 166–168.
- Takhtajan, A. L. (1980). Outline of the classification of flowering plants (magnoliophyta). *The Botanical Review*, 46(3): 225–359. <https://doi.org/10.1007/BF02861558>
- Terral, J. F., Newton, C., Ivorra, S., Gros-Balthazard, M., de Morais, C.T., Picq, S., Tengberg, M. and Pintaud, J.C. (2012), Insights into the historical biogeography of the date palm (*Phoenix dactylifera* L.) using geometric morphometry of modern and ancient seeds. *Journal of Biogeography*, 39: 929-941. <https://doi.org/10.1111/j.1365-2699.2011.02649.x>
- Thompson, D. W. (1945). *On Growth and Form* (1st ed.). Cambridge University Press.
- Thorne, R. F. (1992). Classification and geography of the flowering plants. *The Botanical Review*, 58(3): 225–327. <https://doi.org/10.1007/BF02858611>
- van der Niet, T., Zollikofer, C. P. E., León, M. S. P. de, Johnson, S. D., & Linder, H. P. (2010). Three-dimensional geometric morphometrics for studying floral shape variation. *Trends in Plant Science*, 15(8): 423–426. <https://doi.org/10.1016/j.tplants.2010.05.005>
- van Ooststroom, S. J. (1938). The Convolvulaceae of Malaysia - I. Blumea: Biodiversity, Evolution and Biogeography of Plants, 3(1): 62–94.
- van Ooststroom, S. J. (1939). The Convolvulaceae of Malaysia, II. Blumea: Biodiversity, Evolution and Biogeography of Plants, 3(2): 267–371.
- van Ooststroom, S. J. (1940). The Convolvulaceae of Malaysia - III. Blumea: Biodiversity, Evolution and Biogeography of Plants, 3(3): 481–582.
- van Ooststroom, S. J., & Hoogland, R. D. (1953). Convolvulaceae. In S. van Steenis (Ed.), *Flora Malesiana - Series 1, Vol. 4, Issue 1*, pp. 388–512.
- Venable, D. L., Dyreson, E., Pinero, D., & Becerra, J. X. (1998). Seed Morphometrics and Adaptive Geographic Differentiation. *Evolution*, 52(2): 344–354. <https://doi.org/10.1111/j.1558-5646.1998.tb01636.x>
- Verdcourt, B. (1963). Convolvulaceae. In C. E. Hubbard & E. Milne-Redhead (Eds.), *Flora of Tropical East Africa*. Crown Agents for Oversea Governments and Administrations.

- Viscosi, V., & Cardini, A. (2011). Leaf morphology, taxonomy and geometric morphometrics: A simplified protocol for beginners. *PLoS ONE*, 6(10). <https://doi.org/10.1371/journal.pone.0025630>
- Viscosi, V., & Fortini, P. (2011). Leaf shape variation and differentiation in three sympatric white oak species revealed by elliptic Fourier analysis. *Nordic Journal of Botany*, 29(5): 632–640. <https://doi.org/10.1111/J.1756-1051.2011.01098.X>
- Viscosi, V., Antonecchia, G., Lepais, O., Fortini, P., Gerber, S., & Loy, A. (2012). Leaf Shape and Size Differentiation in White Oaks: Assessment of Allometric Relationships among Three Sympatric Species and Their Hybrids. *International Journal of Plant Sciences*, 173(8), 875–884. <https://doi.org/10.1086/667234>
- Viscosi, V., Lepais, O., Gerber, S., & Fortini, P. (2009). Leaf morphological analyses in four European oak species (*Quercus*) and their hybrids: A comparison of traditional and geometric morphometric methods. *Plant Biosystems*, 143(3): 564–574. <https://doi.org/10.1080/11263500902723129>
- Wäldchen, J., & Mäder, P. (2018). Plant Species Identification Using Computer Vision Techniques: A Systematic Literature Review. *Archives of Computational Methods in Engineering*, 25(2): 507–543. <https://doi.org/10.1007/s11831-016-9206-z>
- Webster, M., & Sheets, H. D. (2010). A Practical Introduction to Landmark-Based Geometric Morphometrics. *The Paleontological Society Papers*, 16, 163–188. <https://doi.org/10.1017/s1089332600001868>
- Wight, R. 1837. *Clavis analytica of the Convolvulaceae of the Peninsula of India*. *Madras Journal of literature science*, 5: 15–23.
- Wood, J. R. I., Muñoz Rodríguez, P., & Scotland, R. W. (2022). Priorities for taxonomic studies of Indian Convolvulaceae. *Journal of Economic and Taxonomic Botany*, 46(1 & 2): 5–14.
- Wood, J. R. I., Muñoz-Rodríguez, P., Williams, B. R. M., & Scotland, R. W. (2020). A foundation monograph of *Ipomoea* (Convolvulaceae) in the New World. *PhytoKeys*, 143: 1-831. <https://doi.org/10.3897/PHYTOKEYS.143.32821>
- Wood, J. R. I., Williams, B. R. M., Mitchell, T. C., Carine, M. A., Harris, D. J., & Scotland, R. W. (2015). A foundation monograph of *Convolvulus* L. (Convolvulaceae). *PhytoKeys*, 51(1): 1–282. <https://doi.org/10.3897/phytokeys.51.7104>
- Woodson, R. E., Schery, R. W., & Austin, D. F. (1975). Flora of Panama. Part IX. Family 164. Convolvulaceae. *Annals of the Missouri Botanical Garden*, 62(1): 157. <https://doi.org/10.2307/2395053>
- Yoshioka, Y., Iwata, H., Ohsawa, R., & Ninomiya, S. (2004). Analysis of Petal Shape Variation of *Primula sieboldii* by Elliptic Fourier Descriptors and Principal Component Analysis. *Annals of Botany*, 94(5): 657–664. <https://doi.org/10.1093/AOB/MCH190>