

LIST OF FIGURES

Figure No.	Title	Page No.
Fig. 1.1	Paleogeographic map shows the position of the different continents. a. Bajocian, M. Jurassic, shows the splitting of east and west Gondwana, b. L. Cretaceous-Albian, shows south-eastward movement of Indian Plate, c. Paleocene, Thanetian-Danian, shows the northward drifting of Indian plate lying in southern hemisphere, and d. Indian plate lying in northern hemisphere during Bartonian-Lutetian, M. Eocene (Smith et al., 1994).	4
Fig. 1.2	A part of Saurashtra Peninsula map showing the location of the study area.	5
Fig. 1.3	Drainage map of Saurashtra Peninsula showing the radial pattern originating from the central Jasdan plateau.	10
Fig. 2.1	a. General geological and structural map of the Saurashtra Peninsula. It is bounded by major faults (Western Cambay Margin Fault (WCBMF), Western Extension of Narmada Son Fault (NSF), Extension of Western Kathiawar Fault (WKF), Northern Kathiawar Fault (NKF)), b. Northern part of Saurashtra showing the Mesozoic units, Deccan Traps, and Intertrappean sites (modified after Biswas, Aslam, Bhatt 2000).	16
Fig. 3.1	Flowchart showing methodology that adopted for analysing the samples for palynofossils.	36
Fig. 3.2	Palynofossil extraction technique chart (after Grey (1965), Green (2001)).	41
Fig. 4.1	Diagrammatically illustrated varying positions of the sedimentary successions in relation with the different phases of lava flow, (a) Infratrappean, (b) Intertrappean, (c) Intratrappean, and (d) Supratrappean.	49
Fig. 4.2	Composite litholog of representing lithofacies of Ninama Basin.	55

Fig. 4.3	Composite litholog of representing lithofacies of Chotila Basin.	61
Fig. 5.1	Measured lithologs of the Ninama Basin intertrappean sequence show the lithofacies distribution. a. Composite litholog of the Ninama section, Ninama village. b. Shekhdod hill section, Shekhdod village. c. Moti Lakhavad section, Moti Lakhavad village. d. Motamatra hill section, Motamatra village and e. Motamatra well section, Motamatra village.	71
Fig. 5.2	Measured lithologs of the Chotila intertrappean sequence show the lithofacies distribution. a. Chanpa Hill Section East. b. Chanpa Hill Section West. c. Actionware Industry Section. d. Bhalgam Quarry Section. e. Bamanbor Road section. f. Rangpar GIDC Section (RGIDC). g. Bamanbor Navagam section and h. Garida Section.	82
Fig. 5.3	Plot of the geochemical proxies for paleosalinity, a. $Ca/(Ca+Fe)$, b. Sr/Ba ; paleoredox c. $V/V+Ni$, d. V/Cr ; paleodepth e. $Fe/(Ca+Mg)$, of the Ninama Basin.	95
Fig. 5.4	Plot of the geochemical proxies for paleoweathering, volcanic vs detrital input and paleoclimate, a. Ti/Al , b. Na/K , c. $K/Fe+Mg$, d. Al/Ti , e. CIA-K, and f. MAP, of the Ninama Basin.	96
Fig. 5.5	Plot of the geochemical proxies for paleosalinity, a. $Ca/(Ca+Fe)$, a. Sr/Ba ; paleoredox c. $V/V+Ni$, d. V/Cr ; paleodepth and, e. $Fe/(Ca+Mg)$, of the Chotila Basin.	103
Fig. 5.6	Plot of the geochemical proxies for paleoweathering, volcanic vs detrital input and paleoclimate, a. Ti/Al , b. Na/K , c. $K/Fe+Mg$, d. Al/Ti , e. CIA-K, and f. MAP, of the Chotila Basin.	104
Fig. 8.1	Generalized litholog shows lithostratigraphy, lithofacies and systematic collected of the samples of the Ninama Intertrappeans. Palynofossils distribution shows their relative abundance in the systematically collected samples with their palynostratigraphic zonation. Ninama Limestone contains abundant and diverse palynofossils as compared to the lower Sukhbhadar Formation.	186

Fig. 8.2 Generalized litholog shows sample positions, lithostratigraphy and lithofacies of the Chotila Intertrappeans. Palynofossils distribution shows their relative abundance in the systematically collected samples with their palynostratigraphic zonation. Chotila Chert and Bamanbor Formation contain abundant and most diverse palynofossils as compared to lower Rangpar Formation. 194

Fig. 9.1 Three-dimensional model of Ninama paleolake, depicting 3 phases of deposition. Phase 1- Deposition of LGW facies, reflects high clastic input, brackish to saline, oxic-dysoxic moderately deep lake and absence of palynofossils; Phase 2- Deposition of GSH and SM facies with moderate to high terrestrial input, freshwater to brackish water, anoxic deep lake with abundant fungal spores and moderate pollens; Phase 3- Deposition of ML, CL and GBL facies with low to moderate terrestrial input, saline, oxic-anoxic shallow lake with abundant of fungal spores and pollens. 207

Fig. 9.2 Three-dimensional model of Chotila paleolake, depicting 3 phases of deposition. Phase 1- Deposition of CS facies of Rangpar formation, showing high rate of fine argillaceous clastic input, in brackish, anoxic, shallow to moderately deep lake with fungal spores; Phase 2- Deposition of LC, MC and MS facies, of Chotila Chert with variable terrestrial input, in brackish water, anoxic deep lake and abundant fungal spores and moderate pollens; Phase 3- Deposition of FSS and CS facies of Bamanbor Formation, in shallow to deep lake, high terrestrial input, variable salinity, and oxic-anoxic condition, with abundance of fungal spores and pollens. 210

Fig. 9.3 Paleogeographic position of Insular India during its northward drift in the Tethys Ocean during Late Cretaceous, Paleocene and Early Eocene. 213