

LIST OF PLATES

Plate No.	Title	Page No.
Plate 3.1	a. Geochemistry lab showing fume chamber and working space for palynological processing of samples, b. Hotplate, c. Oven, d. Centrifuge, e. Mortar and Pestle and f. Weighing scale.	42
Plate 3.2.	a and b. Sample demineralization in fume hood (HCl, HF and HNO ₃ treatments), c. Sample washing, d and e. Sample settling, f. KOH treatment and washing by centrifuge, g and h. sieving using 10–15-micron mesh, i. Ultrasonicator, j. Slide preparation, smearing, staining, k. Cooked, permanent slides with coverslip, l. Leica DM EP microscope.	44
Plate 4.1	a. Sukhbhadar river section showing the shales of Sukhbhadar Formation and Limestone of the Ninama Limestone. b. Close-up of the Sukhbhadar Formation shows the thinly bedded mudstone. c. Ninama well section shows the thick shale with intercalated lithic sandstone/greywacke. d. Contact between the shale of the Sukhbhadar Formation and Ninama Limestone near Motamatra, limestone (mudstone) is thinly bedded to flaggy in nature.	56
Plate 4.2	a. Thick limestone of the Linam Limestone overlain the shales of the Sukhbhadar Formation, Sukhbhadar river section near Ninama. b. The well section near Motamatra village shows Sukhbhadar Formation and Ninama Limestone nonconformably overlain the Deccan lava flow. c. Thick limestone bed grade to flaggy limestone in Shekhdod hill. d. Thick limestone bed in the Shekhdod hills. e. Flaggy limestone band exposed as superficial surface near Motamatra village.	58
Plate 4.3	Exposures of the Ninama Limestone around Ninama village and Ninama hill sections. a. Gradational contact between the marlite and calcareous shale and cherty limestone. b. Black limestone. c.	59

d. & e. The thick sequence of Ninama Limestone in the quarried section shows the bands of the marlite, calcareous shale and cherty limestone.

Plate 4.4 a. A freshly quarried section exposes the thick Chotila Chert with dark shale of the Rangpar Formation near Garida village. b. A freshly quarried section near Garida village shows an intratrappean sequence comprising a thick Chotila Chert and Rangpar shale sandwich between the younger lava flow, bending in the chert and shale beds is due to differential stress exerted from the different directions. c. Rangpar Formation and Chotila chert show the gradational contact near Garida village. d. Emplacement of the dyke in the Rangpar Formation and Chotila Chert has displaced the formations on either side. 62

Plate 4.5 Photographs of the freshly quarried section near the Rangpar industry area. a. All three formations of the Chotila Basin are exposed on the northern side of the industry area. It is capped by the younger lava flow. b. Well-developed Rangpar Formation and Chotila Chert are capped by the lava flow. c. Discordant relationship between the shale of Rangpar Formation and dyke. d. Gradational contact between the Rangpar Formation, Chotila Chert and Bamanbor Formation. 63

Plate 4.6 a. and b. Chanpa Hill Section, exposing the intratrappean consisting of mudshale and banded chert of Chotila Chert, sandwiched between the Deccan Traps. c. Thick banded chert and mudstone are exposed at the Chanpa Hill Section. d. The eastern part of the Chanpa Hill Section exposes the Chotila Chert, overlain and underlain by Deccan Traps. 64

Plate 4.7 a. Photograph of chert belonging to Chotila Chert, exposed at the Bamanbor-Navagam Section, which is highly disturbed due to the intruded Deccan traps. b. and c. Thick 1-1.5m Chotila Chert exposed at Bhalgam Quarry section, and is exposed as intratrappeans. d. Pseudo ripples due to the Deccan lava flows 65

preserved in the banded chert of Chotila Chert. e. and f. Highly deformed chert exposed at Bamanbor-Navagam Section. g. Thick unit of chert at Bamanbor-Navagam Section forming a yardang.

Plate 4.8 a. b. and c. Exposed Chotila Chert in the Vasundhara Hill Section. 66
 d. Chotila Chert and Bamanbor formation exposed near the Actionwear Industry Section.

Plate 4.9 a. Bamanbor Road Section exposing the thick sequence of 67
 Bamanbor Formation, overlain by Deccan Traps. b. Physa containing sandstone of Bamanbor Formation. c. Thick shale unit of Bamanbor Formation. d and e. Sandstone unit of Bamanbor Formation consisting of gastropods and bivalves. f. Moderately thick bed of the Bamanbor Formation exposed at Bamanbor Road Section.

Plate 5.1 a. The exposure of grey shale facies (GSH) on the right bank of the 73
 Sukhbhadar River near Ninama village. b. Laminated SH and Marlite (ML) facies exposed above the shale facies. c and d. Exposures of the calcareous shale (CSH) and cherty limestone (CL) facies. e. Thinly bedded silty mudstone (SM) facies shows gradational contact with shale facies. Photomicrographs of SM. f. Silt-sized quartz grains in a muddy matrix with iron oxide. g. Subordinate amount of quartz grains in the mud with iron oxide. h. Bands of lithic greywacke (LGW) facies are exposed in the dug well near Ninama village. Photomicrograph of lithic greywacke showing, i. Volcanic fragment (basalt), quartz grains with cleavable calcite and j. Texturally immature shows the cleavable calcite, iron oxide, broken quartz grains, and volcanic fragments, (Cc- Cleavable calcite, Qz- Quartz, Vf- Volcanic fragment, Io- Iron oxide, M- Mud; Scale- person height- 175cm, ruler length- 30cm).

Plate 5.2 a. Grey black limestone (GBL) facies from the Moti Lakhavad 76
 village showing jointed nature, exposed above the Deccan traps. b. Relics of a thin bed of highly jointed and weathered GBL at Sukhbhadar river, Ninama village. c. Large bouldery GBL at the

top of Ninama hill, Ninama village. Photomicrographs of mudstone microfacies. d. Sparitization and chertification in the groundmass. e. Organic matter and f. Chert, Silt-sized quartz grains and iron oxide, (Ms- Microspar, Om- Organic matter, Qz- Quartz, Ch-Chert, Io- Iron oxide; Scale- hammer length- 33cm, ruler length- 30cm).

Plate 5.3 a. Cherty limestone (CL) facies exposed at the Ninama hill, intercalated with marlite facies (ML). b. Thinly bedded CL near Shekhdod village. c. Close-up of CL facies at the top of Sukhbhadar River. Photomicrographs of peloidal cherty wackestone microfacies. d. Calcareous sphere of floating calcareous algae, with replacement by chert and sparite e. Elongated dark algal filaments, partially replaced by framboidal pyrite and chert, as well as groundmass is also recrystallized to microspar and replaced by chert. f. Framboidal pyrite replacing the algae, pellet centres recrystallized into microspar. Bioclastic bindstone microfacies, g. Cladoceran shell fragments and pellets bounded together by microbial mats. Ostracod shell fragment. H. Algae with sit-size quartz grains. i. Micritic pellets modified to microsars. j. PPL view of bioclastic bindstone with ostracod shell fragments and algae. Cone in cone calcitic microfacies and k. Cone in cone structures shows the growth of fibrous calcite which appeared as wavy laminae with organic matter, (Cs- Calcareous sphere, Ch- chert, Al- Algae, Fp- Framboidal pyrite, Pel- Pellet, Cl- Cladoceran, Os- Ostracod fragments, Fc- Fibrous calcite, La- Laminae Re - Recrystallized micrite, Ms- Microspar, ML- Microbial laminations; Scale- hammer length- 33cm, ruler length- 30cm).

Plate 5.4 a. Exposures of the marlite facies (ML) near Ninama village and b. Thick bedded ML facies. Photomicrographs of cladoceran bindstone microfacies. c. Algae, ostracod shell fragments and microspar. d. Elongated diatoms in the laminated organic matrix e. Cladoceran shell fragments in the algal mats. f. Prefer oriented

cladoceran appendages and their fragments. g. Algal filaments, cladoceran and ostracod fragments in well-laminated algal mats. h. Wavy-oriented organic matter and microspar bound with pellets and quartz, (Os- Ostracod fragments, Ms- Microspar, Al- Algae, Di- Diatoms, Cl- Cladoceran, Qz- Quartz, Om- Organic matter, Ml- Microbial laminations; Scale- person height- 175cm, ruler length- 30cm).

Plate 5.5 Fossiliferous shaly sandstone (FSS) lithofacies exposed at a. Bamanbor Road Section, overlain by silty shale facies (SS) and clay shale facies (CS) with b. Gastropod fossil, c. horizontal laminations in thin to thick beds of FSS. d. Photomicrographs of FSS d. and e. showing quartz, chert and muddy matrix embedded in a calcareous and ferruginous cement, (Gp- Gastropod, Qtz- Quartz, Ch- Chert, Sh rfx- Shale/mudstone lithic fragments, Mtx- Matrix). 83

Plate 5.6 Clay shale facies (CS) of Rangpar Formation exposed at, a. RGIDC section, with intercalated mudstone (MS) facies, b. Garida Road Section where it is overlain by thick MS. CS facies of Bamanbor Formation, exposed at c. RGIDC section, showing characteristic fissility, d. Bamanbor Road Section and e. Red CS facies at Actionware Industry section and f. CS facies overlying laminated chert (LC) facies. 84

Fig. 5.7 a. Silty shale facies (SS) and fossiliferous silty mudstone facies (FSM) intercalated between CS facies exposed at Rangpar GIDC section. Mudstone Lithofacies (MS) exposed at b. Rangpar GIDC Section, c. Bhalgam Quarry Section d. Garida Road Section, showing d. massive, thickly bedded nature and intercalation with CS facies and e. CS, MS and LC facies of Rangpar Formation, Chotila Chert and Bamanbor Formation. 86

Plate 5.8 a. Fossiliferous silty mudstone facies (FSM) exposed at Rangapar GIDC Section and b. *Valvata* and various gastropods embedded in FSM. 87

Plate 5.9 Laminated Chert Lithofacies (LC) exposed at a. Chanpa Hill Section East, with a closeup of LC (inset). b. Tilted and folded LC facies at Bamanbor Navagam Section. c. Chanpa Section, d. Vertebrate fossil (Fish Fin) in LC facies. e. Pseudo ripples in LC facies. f. Bhalgam Quarry Section showing LC and black chert facies. Massive Chert Lithofacies (MC) exposed. g. Bamanbor Navagam Section, h. Chanpa Hill Section East and i. Black Chert lithofacies (BC) exposed at Vasundhara Hill Section (close-up in inset). 88

Plate 5.10 Photomicrographs of a. Mudstone facies. b, c and d. Laminated Chert Lithofacies showing alternate bands of microcrystalline and cryptocrystalline silica with dark bands of organic matter and iron oxide. e. thickly laminated chert facies with discontinuous laminae and dominated by microcrystalline quartz. f. Black chert facies with microfolds. 89

Plate 6.1 *Deccanoida conrugis* sp. nov. Fig. a, External view of holotype. Fig. b, Internal view of holotype. Fig. c-i, Paratypes of *D. conrugis*. Fig. j, X-ray image of the holotype showing umbonal angle and hinge plate (hp). Fig. k, Line drawing of the holotype shows internal morphology (ct-cardinal teeth, aa- anterior adductor muscle scar, lt-ligament tooth, lg-ligament, pl-pallial line, pa-posterior adductor muscle scar). All scale bars represent 1cm 122

Plate 6.2 *Deccanoida aleta* sp. nov. Fig. a-b, holotype: Fig. a, External view; Fig. b, Internal view. Fig. c-j, Paratypes: Fig. c, External view and internal shell shows the dentition and ligament (Fig. d). Fig. e, Internal view shows the dentition and ligament. Fig. f-j, External view. Fig. k. X-ray of holotype with umbonal angle and hinge plate (hp), Fig. l, Internal shell morphology illustrated of holotype. (ct-cardinal teeth, aa-anterior adductor muscle scar, lt-ligament tooth, lg- ligament, pl-pallial line, pa-posterior adductor muscle scar). All scale bars represent 1cm. 125

- Plate 6.3** *Deccanoida costaria* sp. nov. Fig. a-b holotype: Fig. a, left external view; Fig. b, right external view. Fig. c-d Paratype: Fig. c, left external view; Fig. d, right external view. Fig. e. X-ray of holotype (left valve) with umbonal angle and hinge plate. Fig. f Line drawing of holotype illustrate the internal shell morphology. (ct-cardinal teeth, aa-anterior adductor muscle scar, lt-ligament tooth, lg-ligament, pl-pallial line, pa-posterior adductor muscle scar). All scale bars represent 1cm. 127
- Plate 6.4** *Bamanboria oblongis* sp. nov. Fig. a-c, holotype: Fig. a, external view of right valve; Fig. b, internal of right view with umbonal angle; Fig. c, ventral view with highly corrugated commissure and tubercles. Fig. d-i, paratype: Fig. d-f, internal molds; Fig. g-h, external and internal view with cardinal teeth. Fig. i, external view. Fig. j, X-ray of holotype. Fig. k, internal shell morphology illustrated from the holotype. (ct-cardinal teeth, aa-anterior adductor muscle scar, lt-ligament tooth, lg-ligament, pl-pallial line, pa-posterior adductor muscle scar). All scale bars represent 1cm. 129
- Plate 6.5** *Chotilia trapeansis* sp. nov. Fig. a-b, holotype: Fig. a, external view of right valve. Fig. b, internal view of right valve. Fig. c-d, paratype: Fig. c, left valve view; Fig. d, right valve view and also shows prominent cardinal teeth fused together. Fig. e, X-ray of holotype with umbonal angle and hinge plate. Fig. f, illustrated internal shell morphology of holotype. (ct-cardinal teeth, aa-anterior adductor muscle scar, lt-ligament tooth, lg-ligament, pl-pallial line, pa-posterior adductor muscle scar). All scale bars represent 1cm. 131
- Plate 6.6** *Chotilia tuberculata* sp. nov. Fig. a-b, holotype: Fig. a, left valve external view showing prominent tubercles and cardinal teeth of right valve; Fig. b, external view of right valve. Fig. c. X-ray of holotype with hinge plate and umbonal angle. Fig. d. line drawing of the holotype shows the various internal morphological features. (ct-cardinal teeth, aa-anterior adductor muscle scar, lt-ligament 133

tooth, lg-ligament, pl-pallial line, pa-posterior adductor muscle scar, tb-tubercle). All scale bars represent 1 cm.

Plate 6.7 *Chotilia deccansis* sp. nov. Fig. a-e, holotype: Fig. a, external view of right valve; Fig. b, internal view of right valve with prominent cardinal teeth and opisthodontic parivincular ligament; Fig. c, ventral view with distinct coarsely corrugated commissure. Fig. d, X-ray image with umbonal angle and hinge plate Fig. e. Line drawing of the shell shows internal morphological features of holotype (ct-cardinal teeth, aa-anterior adductor muscle scar, lt-ligament tooth, lg-ligament, pl-pallial line, pa-posterior adductor muscle scar, cm-commissure). All scale bars represent 1 cm. 135

Plate 6.8 a. *Physa* from shaly sandstone, Bamanbor Formation, exposed at Bamanbor Roadcut section (Scale - 5₹ coin is 2.4 cm in diameter) 136
b. *Valvata* from yellow mudstone, Bamanbor Formation, exposed at Rangpar GIDC section.

Plate. 7.1 a. *Cythidites australis*, NL-NB, b. *Hammenisporis susannae*, NL-NB, c. *Arecipites bellus*, NL-NB, d. *Palmaepollenites communis*, NL-NB, e. *Palmidites* sp. NL-NB, f. *Longapertites* sp., NL-NB, g. *Longapertites retipilatus*, NL-NB, h. *Longapertites triangulates*, NL-NB i. *Matanomadhiasulcites maximus*, NL-NB, j-k. *Proxapertites cersus*, NL-NB, l. *Proxapertites crassimurus*, NL-NB, (NL- Ninama Limestone, SF- Sukhbhadar Formation, BF- Bamanbor Formation, CC- Chotila Chert, RF- Rangpar Formation, NB- Ninama Basin, CB- Chotila Basin). 145

Plate 7.2 a. *Dermatobrevicolporites* sp., NL-NB, b. *Dermatobrevicolporites dermatus*, CC-CB, c. *Lakiapollis ovatus*, NL-NB, d. *Meliapollis ramanujamii*, BF-CB, e. *Rhoipites kuctchensis*, BF-CB, f-g. *Tricolporopollenites* spp, SF-NB, h. *Aquilapollenites ovatus*, NL-NB, i. *Araliaceoipollenites psilatus*, NL-NB, j. *Araliaceoipollenites reticulatus*, NL-NB. 151

Plate 7.3 a. *Barringtoniapollenites retibaculatus*, NL-NB, b. 154
Cupuliferoipollenites ovatus, NL-NB, SF-NB, c.

Cupuliferoipollenites psillus, NL-NB, d. *Incrotonipollis neyvelii*, NL-NB e. *Cupanieidites flaccidiformis*, NL-NB, f. *Ephedripites* sp. SF-NB, g. *Malvacearumpollis baconyensis*, NL-NB, h. *Monocolpopollenites* sp. NL-NB, i. *Monocolpopollenites plicatus*, BF-CB.

Plate 7.4 a. *Florschuetzia rajpardiensis*, BF-CB, b. *Mulleripollis bolporensis*, BF-CB, c-e. *Rhombipollis geniculatus*, NL-NB, f-g. *Yeguapollis* sp. NL-NB. 157

Plate 7.5 a-c. *Perisporiacites varians*, a. NL-NB, b. CC-CB, RF- CB, d. *Meliolinites spinksii* RF-CB. 158

Plate 7.6 a. *Ceratohirudispora triradiata*, SF-NB, b. *Plureicellaesporites globatus*, BF-CB, c. *Plureicellaesporites planus*, BF-CB, d. *Polycellaesporonites bellus*, NL-NB, e-f. *Polycellaesporonites psilatus* CC-CB, g. *Polycellaesporonites saxenae*, BF-CB, h. *Staphlosporonites settyi* CC-CB. 161

Plate 7.7 a. *Frasnacritetrus indicus*, CC-CB, b. *Frasnacritetrus taugourdeai*, CC-CB, c. *Fusiformisporites* sp., NL-NB, d-e. *Papulosporonites multicellatus*, d, NL-NB, e. BF-CB, and f. *Dicellaesporites minutus*, BF-CB. 164

Plate 7.8 a-b. *Inapertisporites kedvessi*, a, SF-NB, b, NL-NB, c-d, *Inapertisporites maximus*, NL-NB, e. *Inapertisporites novus*, BF-CB, f-g. *Hypoxylonites subrotundus*, NL-NB, h. *Hypoxylonites minimus*, BF-CB, i. *Monoporisporites circularis*, NL-NB, j. *Diporisporites sirmaurensis*, CC-CB, k-l *Helicominites salvinites*, CC-CB. 166

Plate 7.9 a-g, *Polycellaesporonites* spp., a-c, NL-NB, d-f, CC-CB, g, BF-CB, h-l. *Diporisporites sirmaurensis*, CC-CB, m-q, *Pluricellaesporites* spp., m-o, CC-CB, p-q, BF-CB, r. *Ceratohirudispora* sp. BF-CB, s. *Helicominites* sp. CC-CB. 168

Plate 7.10	a. <i>Palaeomycites</i> spp., NL-NB, b. <i>Palaeomycites acinus</i> , NL-NB, c. <i>Udaria saxenae</i> , SF-NB, d. <i>Protocolletotrichum deccanensis</i> .	170
Plate 7.11	a-j <i>Palaeomycites</i> sp., a-c, SB-NB, d-f, NL-NB, g-j- RF-CB.	171
Plate 7.12	a-f, <i>Centropyxis aculeata</i> , a-c- NL-NB, d-f, RF-CB, g-m, <i>Arcella</i> <i>arenaria</i> , g-j- RF-CB, k-m, BF-CB.	174
Plate 7.13	a. <i>Botryococcus braunii</i> , BF-CB, b. NL-NB, c-e <i>Pediastrum</i> sp. NL-NB.	175
Plate 7.14	a. Rotifer, CC-CB, b-d. Cladocera, NL-NB, e-h- Phytoclast e-f. NL-NB, g. RF-CB, h. BF-CB, i- AOM.	177
Plate 7.15	a-f, Lepidopteran Scales, a-d, NL-NB, e-f, CC-CB.	178