

## TABLE OF CONTENT

Sr. No.	CONTENT	Page No.
<b>Chapter 1</b>	Introduction	1
<b>1.1</b>	Candida albicans	2
<b>1.2</b>	Candidiasis	2
<b>1.2.1</b>	Vaginal Candidiasis	3
<b>1.2.2</b>	Invasive Candidiasis	3
<b>1.2.3</b>	Oral Candidiasis	3
<b>1.3</b>	Head and neck radiotherapy effect in immunocompromised human patients	4
<b>1.4</b>	Factors that influence the virulence of <i>C. albicans</i>	4
<b>1.5</b>	Hyphal Stage Virulence Switch and its blocker	6
<b>1.6</b>	Cell Surface Composition	8
<b>1.7</b>	<i>C. albicans</i> Cell cycle	9
<b>1.8</b>	Cyclins/ CDK levels in Yeast vs hyphal morphogenesis	10
<b>1.9</b>	Ubiquitin	11
<b>1.10</b>	Ubiquitin Structure	11
<b>1.11</b>	Proximal and Distal ubiquitin structural diversity	12
<b>1.12</b>	UPS key functions as a central player in the cell cycle and filamentation of <i>C. albicans</i>	14
<b>1.13</b>	Transcriptional factors (Activators/ Inhibitors)	16
<b>1.13.1</b>	Activators	16
<b>1.13.2</b>	Inhibitors	16
<b>1.14</b>	Rationale	17
<b>Chapter 2</b>	<b>To study the cell cycle progression and morphology switching in <i>C. albicans</i> expressing ubiquitin mutations.</b>	22
<b>2.1</b>	Introduction	23
<b>2.2</b>	Materials and methods	25
<b>2.2.1</b>	Strains and media	25
<b>2.2.2</b>	Cloning of ubiquitin mutant genes	27

2.2.3	Transformation and selection of integrants	29
2.2.4	Influence of variants of ubiquitin on different antibiotic stress	29
2.2.5	Confocal microscopy & Fluorescence microscopy for morphological switching analysis	29
2.2.6	Cell cycle analysis	30
2.2.7	Detection of protein levels of Cdc28 by western blot analysis	30
2.2.8	Incorporation of variants in Lys63 and Lys48 linked chain formation by western blot analysis	31
2.2.9	SEM analysis	31
2.2.10	Calcofluor white staining	32
2.2.11	Congo red staining	32
2.2.12	Protease Extraction and Precipitation	32
2.2.13	Gelatin Zymography	33
2.2.14	Statistical analysis	34
2.3	Results	35
2.3.1	Site-directed mutagenesis of mutant forms of ubiquitin gene & cloning in pRC2312 vector	35
2.3.2	Confirmation of pRC2312 vector & Site-directed mutagenesis PCR products confirmation by gel electrophoresis	35
2.3.3	Site-Directed mutagenesis CTG to TTG ubiquitin mutants transformed in DH5 $\alpha$ cells	37
2.3.4	Cloning of mutant forms of ubiquitin gene in pTET25M-NC vector and expression in BWP17 strain of <i>C. albicans</i> for functional analysis	38
2.3.5	Ubiquitin overexpression dosage-dependent lethality & the survival capacity of the BWP17 strain of <i>C. albicans</i> .	40
2.3.6	UbEP42, UbL50P, and UbI61T show slow growth phenotype reduced survival and thermotolerance, and impaired heat stress-resisting capacity	41
2.3.7	UbEP42, UbL50P and UbI61T reduce resistance toward antibiotic stress	43
2.3.8	Yeast to hyphal transition was affected by mutants of ubiquitin UbEP42, UbL50P and UbI61T confirmed by using N-terminal GFP fusions	44
2.3.9	UbEP42, UbL50P and UbI61T disrupt cell cycle progression in <i>C. albicans</i>	45

2.3.10	Cdc28 protein kinase expression decrease in cells expressing UbEP42, UbL50P and Ubi61T	47
2.3.11	Expression of UbEP42, UbL50P and Ubi61T results in impairment of K63 and K48 linked polyubiquitinations	48
2.3.12	Cell surface analysis by Scanning Electron Microscope (SEM)	49
2.3.13	Slightly irregular deposition of chitin on <i>C. albicans</i> cell wall under the effect of mutant ubiquitin proteins observed	50
2.3.14	Unmasking the effect of mutant ubiquitin proteins over $\beta$ -glucan exposure on the cell surface of <i>C. albicans</i>	51
2.3.15	The effect of ubiquitin mutations over secretory aspartyl protease activity at different pH values	53
2.4	Discussion	57
<b>Chapter 3</b>	<b>Status of cyclins in the various stages of cell cycle in <i>C. albicans</i> expressing ubiquitin mutations.</b>	59
3.1	Introduction	60
3.2	Material and methods	60
3.2.1	Strains and media	60
3.2.2	Analysis of the expression levels of cyclins	61
3.3	Results	61
3.3.1	Status of cyclins in the various stages of cell cycle in <i>C. albicans</i> expressing ubiquitin mutations	61
3.4	Discussion	62
<b>Chapter 4</b>	<b>To Study the regulation transcription factors involved in cell cycle and morphogenesis in <i>C. albicans</i> expressing ubiquitin mutations</b>	64
4.1	Introduction	65
4.2	Material and Method	66
4.2.1	Strain and Media	66
4.2.2	Analysis of the expression levels of virulence-associated transcription factors	66
4.2.3	Sample preparation for proteome analysis	66

4.2.4	Data Analysis	67
4.3	Result and Discussion	69
4.3.1	Dysregulation of hyphal and bud-associated transcription factors in <i>C. albicans</i> expressing ubiquitin mutations	69
4.3.2	Characterization of Differentially Expressed Proteins in Ubiquitin Mutant Conditions	71
4.3.3	Common downregulation of some proteins in <i>Candida albicans</i> due to Overexpression of Ubiquitin mutants UbL50P, Ubi61T, and UbEP42	73
4.3.4	Metabolic Pathways and Stress Response	75
4.3.5	Cell Wall Synthesis and Morphogenesis	75
4.3.6	Protein Folding and Ubiquitin-Mediated Proteolysis	76
4.3.7	Secretion of Aspartyl Protease	76
4.3.8	Pathways Affected by Downregulated Proteins	76
4.3.9	Up and downregulated protein-protein functional interactions as revealed by String node interaction	77
4.3.10	Identification of common upregulated proteins in <i>Candida albicans</i> expressing UbEP42, UbL50P, and Ubi61T ubiquitin mutants	84
4.4	Conclusion	88
Chapter 5	<b>Generating di-glycine deletions of ubiquitin mutants to understand the mechanism of polyubiquitin chain formation in <i>C. albicans</i> expressing ubiquitin mutations.</b>	89
5.1	Introduction	90
5.1.1	Di-glycine motif significance in K48, K63 polyubiquitination chain formation	90
5.2	Material and Methods	93
5.2.1	Strain and growth conditions	93
5.2.2	Cloning and Transformation	93
5.2.3	Sequence analysis	96
5.2.4	Western Analysis	96
5.2.5	TrRosseta Structural Bioinformatics	97
5.3	Results	97
5.3.1	Plasmid PTET25M-CH isolation and Pcr for amplification of mutants	97

<b>5.3.2</b>	Plasmid PTET25M-CH Restriction Digestion and Di glycine Deletion Sequencing	98
<b>5.3.3</b>	Polyubiquitination of substrate protein under Di- glycine deletion mutants condition in <i>C. albicans</i>	99
<b>5.3.4</b>	3D & 2D structure prediction over +/- Di-glycine ubiquitin mutants	101
<b>5.4</b>	Discussion	103
<b>Summary</b>		105
<b>Bibliography</b>		109

## LIST OF FIGURES

<b>Figure No.</b>	<b>TITLE</b>	<b>Page No.</b>
<b>Figure 1.1</b>	Types of Candidiasis	2
<b>Figure 1.2</b>	Morphology Switch relation with Exogenous and Endogenous quorum sensing molecule	7
<b>Figure 1.3</b>	<i>C. albicans</i> Cell surface composition	8
<b>Figure 1.4</b>	Cell cycle stage cyclins/cdk	9
<b>Figure 1.5</b>	Tertiary structure of ubiquitin	12
<b>Figure 1.6</b>	Proximal and Distal ubiquitin structural diversity under K48, K63, and linear ubiquitin	14
<b>Figure 1.7</b>	Ubiquitin proteasome regulates cyclin and CDK inhibitors through ubiquitin ligase	15
<b>Figure 2.1</b>	Vector map of P-TET25MNC for ubiquitin mutants cloning and screening	26
<b>Figure 2.2</b>	Illustration of Cloning Strategy for site-directed mutagenesis of CTG codon in ubiquitin mutants and functional characterization	27
<b>Figure 2.3</b>	Protease extraction and Precipitation method	32
<b>Figure 2.4</b>	Illustration of gelatin zymography method	34
<b>Figure 2.5</b>	Transformation (A) Transformation of pRC2312 vector containing wild type and mutated ubiquitin into <i>DH5<math>\alpha</math></i> . (B) Requested Other inducible Vectors for further study such as: PTET25M-NC, PTET25-MCDC10-CH.	35

<b>Figure 2.6</b>	<p>Vector Confirmation (A) (a). 1% agarose gel showing 8.3 kb size pRC2312 vector. (b) Restriction digestion of pRC2312 plasmid with <i>Hind III</i> and <i>Bam HI</i>. The Lanes L1 to L4 contain: L(1):1kb ladder L(2) pRC2312 undigested plasmid, L(3) pRC2312 plasmid digested with <i>HindIII</i> and L(4) pRC2312 plasmid digested with <i>BamHI</i> for cloning purpose. (B) Site-directed mutagenesis PCR products run on 3% Agarose gel. Here, FR is denoted as = Forward and Reverse non mutagenic primer in combination, F*R= Forward mutagenic &amp; reverse non mutagenic, FR*=Forward non mutagenic and Reverse mutagenic</p>	36
<b>Figure 2.7</b>	<p>Site Directed mutagenesis (A) Transformants of wild type and mutated CTG codon ubiquitin mutants. (B) (a) 1% agarose gel showing results of restriction digestion of pRC2312 plasmid containing ubiquitin mutant genes is double digested with <i>Hind III</i> and <i>Bam HI</i>. The Lanes 1 contains: 100 bp ladder; Lanes 2 to 8 <i>Hind III</i> and <i>Bam HI</i> digestion of cloned Prc2312, PrcUbWT TTG, PrcS20F TTG, PrcUbEP42 TTG, PrcA46S TTG, PrcL50P TTG, PrcI61T TTG respectively. (b) 1% agarose gel showing results of PCR amplification for further clarification. (C) Ubiquitin CTG to TTG mutant confirmation by DNA sequencing</p>	37
<b>Figure 2.8</b>	<p>Cloning (A) Prc2312 plasmid carrying ubiquitin mutant gene isolated from transformed DH5<math>\alpha</math> cells and run on 1% agarose gel. The Lane 1 contain 100 bp ladder, Lane 2 to 7 showing pRC2312 UbWT TTG, UbS20F TTG, UbEP42 TTG, UbA46STTG, UbL50P TTG, UbI61T TTG respectively. (B) 2% agarose gel showing PCR products of ubiquitin mutant genes. (C) 1% agarose gel is showing result of restriction digestion of cloned vector pTET25M-NC with TTG ubiquitin mutant genes. The Lanes L (1):100 bp ladder, Lanes 2 showing undigested vector. Lane 3-8 contains <i>AatII</i> and <i>StuI</i> digestion of cloned pTET25M-NC UbWT, UbS20F, UbEP42, UbA46S, UbL50P, UbI61T respectively. (D) Transformation in BWP 17 Strain.</p>	39
<b>Figure 2.9</b>	(A) Impact of overexpression of UbEP42 and its single mutant	40

derivatives on *C. albicans*. BWP17 transformants with wild type and mutant forms of ubiquitin whose expression was driven by doxycycline-inducible TET1 promoter were grown in the presence of doxycycline inducer concentration 45ug/ml. BWP17 and it's transformed with the UbWt used as a positive control. (B) Growth was checked on SD minimal media through the streak plate method.

**Figure 2.10** The impact of ubiquitin protein mutants on the growth profile and thermotolerance of *C. albicans* cells. (A) BWP17 cells were genetically modified to express wild type and mutant forms of ubiquitin under the control of the doxycycline-inducible TET1 promoter. These cells were cultivated in the presence of a sublethal concentration of inducer, and their growth profile was monitored by measuring OD at 600nm at 2-hour intervals. Results revealed that strains expressing UbEP42, UbL50P, and Ubi61T exhibited slower growth compared to wild type. (B) The transformants were standardized and plated in the presence of the inducer (45 µg/ml doxycycline). Plates were then incubated at 30°C for varying durations (2, 4, 8, 12, and 16 hours), followed by 3-4 days of incubation. % Survival was determined by counting colonies formed. (C) Similarly, the transformants were plated in the presence of the inducer and exposed to 40°C for 2, 4, and 8 hours, followed by return to 30°C and incubation for 3-4 days. Percent % survival was calculated by counting colonies formed. (D) Additionally, BWP17 strain containing mutant ubiquitin was cultured to log phase and subjected to heat stress at 55°C for 0.5, 1.0, and 1.5 hours. This assay aimed to assess the tolerance of the strains to elevated temperatures.

**Figure 2.11** Impact of mutant ubiquitin proteins on the survival of *C. albicans* under antibiotic stress. Overall, the findings indicate that strains expressing UbEP42, UbL50P, and Ubi61T were unable to withstand antibiotic stress. The concentrations of antibiotics

	<p>utilized in this study were cycloheximide (75 <math>\mu\text{g/ml}</math>), tunicamycin (1 <math>\mu\text{g/ml}</math>), and canavanine (1 <math>\mu\text{g/ml}</math>). Plates were then incubated for 5-6 days at 30°C. Notably, UbS20F showed a slight decrease in viability, while tunicamycin exhibited a potent inhibitory effect on survival. The results are presented as Mean <math>\pm</math> SE, with n = 3 replicates. Statistical significance was denoted as P*** = &lt;0.001, while NS indicates non-significance.</p>	
<p><b>Figure 2.12</b></p>	<p>Effect of expression of UbWT, UbEP42 derived single mutants of ubiquitin on the yeast to hyphal transition monitored using fluorescence microscopy. (A) BWP17 cells and BWP17 UbWT as a control and the cells expressing wildtype and derived single mutant forms of ubiquitin were observed under GFP filter. Transformants were plated on YPD pates 10% serum condition at 37° C to study filamentous growth. (B) Confocal image of UBI61T resolved at higher magnification for better confirmation of morphology switching.</p>	<p>45</p>
<p><b>Figure 2.13</b></p>	<p>Impact of ubiquitin mutant expression on cell cycle progression. (A) FACS analysis was conducted on the BWP17 strain and its transformants. Cells expressing UbEP42, UbL50P, and Ubi61T exhibited a delayed G0/G1 phase, while those expressing UbS20F and UbA46S progressed through the G0/G1 phase similar to cells expressing wild type ubiquitin. The results are presented as Mean <math>\pm</math> SE, with n = 3 replicates. Statistical significance was denoted as P*** = &lt;0.001, while NS indicates non-significance</p>	<p>46</p>
<p><b>Figure 2.14</b></p>	<p>Examines the influence of ubiquitin mutant expression on cell cycle progression following hyphal induction. FACS analysis revealed a higher percentage of cells in the G1 phase specifically in BWP17/UbEP42, whereas BWP17/UbL50P and BWP17/Ubi61T expressing cells did not exhibit significant changes compared to the wild type. The results are presented as Mean <math>\pm</math> SE, with n = 3 replicates. Statistical significance was indicated as P*** = &lt;0.001, while NS denotes non-significance.</p>	<p>47</p>
<p><b>Figure 2.15</b></p>	<p>Illustrates the levels of Cdc28 protein kinase in BWP17 cells</p>	<p>48</p>

	expressing both wild type and mutant forms of ubiquitin. The results indicate a significant reduction in Cdc28 protein kinase levels in cells expressing UbEP42, UbL50P, and UbI61T.	
<b>Figure 2.16</b>	Western blot analysis reveals the polyubiquitination profile of K63 and K48 linkages in <i>C. albicans</i> cells expressing mutant forms of ubiquitin. It was observed that UbEP42, UbL50P, and UbI61T mutants exhibited impaired polyubiquitination with both K63 and K48 linkages. Conversely, the polyubiquitination profile of UbS20F and UbA46S mutants did not exhibit significant changes with respect to K63 and K48 linkages.	49
<b>Figure 2.17</b>	SEM image of <i>C. albicans</i> cells under ubiquitin mutant overexpression in the presence of doxycycline inducer concentration 45ug/ml. The minimum scale bar set 10µm and max 30 µm	50
<b>Figure 2.18</b>	Chitin deposition accumulation and aggregates of <i>C. albicans</i> cells under ubiquitin mutant's overexpression condition (A control B-F ubiquitin mutants) after the staining with Calcofluor white staining. The scale bar represents 20 µm	51
<b>Figure 2.19</b>	<i>C. albicans</i> cell surface β-glucan exposure under ubiquitin mutant overexpression condition after the Congo red staining (A control, (B-F mutants expressing cells). The scale bar represents 20 µm. yellow arrow indicates the higher intensity of dye fluorescence due to binding to β-glucan	52
<b>Figure 2.20</b>	(A) Extraction of protease visualize and folin lowry method standard curve obtain. (B) The secretion of aspartyl protease concentration acetone precipitated samples under different morphology form of candida albicans at different pH under influence of ubiquitin mutants' expression measured by using Folin Lowry method. Results represented in Mean ± SE. (n = 3), (Y= yeast, Y+D= logphase yeast incubated with doxycycline, H+D = After hyphal induction doxycycline inducer added	54
<b>Figure 2.21</b>	The secretion of aspartyl protease concentration acetone precipitated samples represented in chrt form.Under different	55

	<p>morphology forms of candida albicans at different pH under the influence of ubiquitin mutants' expression was measured by using the Folin Lowry method. Results represented in Mean <math>\pm</math> SE. (n = 3), (Y= yeast, Y+D= logphase yeast incubated with doxycycline, H+D = After hyphal induction doxycycline inducer added</p>	
<b>Figure 2.22</b>	<p>12% Zymography gel using gelatin (0.1%) as a substrate. Results showing the difference in secretory aspartyl protease secretion under pH5.9 and pH6.8 in presence of overexpression of Ubiquitin mutants.</p>	56
<b>Figure 2.23</b>	<p>Morphology of cells under different pH of YPD media with 10% bovine calf serum for 0 – 6hr at 37°C exhibited defects in hyphal growth. produced both pseudohyphae and true hyphal filaments. (A, C) Yeast to hyphal transition in presence of 10% serum without inducer conditions in respect to pH5.9, pH6.8. (B, D) Yeast to hyphal transition status in presence of 10% serum with inducer conditions in respect to pH 5.9, 6.8.</p>	57
<b>Figure 3.1</b>	<p>Effect of ubiquitin mutant on cyclins level in <i>Candida albicans</i>. N=3 All values are presented as Mean+ SEM *P&lt;0.05; **P&lt;0.01 and***P&lt;0.001 as compared to control.</p>	62
<b>Figure 4.1</b>	<p>Transcription factor m-RNA levels in <i>C. albicans</i> under ubiquitin mutants expression. N=3 All values are presented as Mean+ SEM *P&lt;0.05; **P&lt;0.01 and***P&lt;0.001 as compared to control.</p>	70
<b>Figure 4.2</b>	<p>Results from DIA quantitation of ubiquitin mutants. Plots (a), (b) and (c) represent the protein profile of upregulated vs downregulated in UbEP42, UbL50P, and Ubi61T respectively. Proteins showing an increase which is greater than 1.3-fold (log<sub>2</sub> fold change) and a decrease less than 1.3-fold (log<sub>2</sub> fold change) were considered and were represented by circular blue dots. Proteins showing (–log<sub>10</sub> P value values were represented by red dots (–log<sub>10</sub> P value, increase 0.05) or decrease (–log<sub>10</sub> P value less than –0.05). Protein abundances in increasing order on the X-axis. Plot (d) represents no. of total proteins that were up and down-regulated in characterized and uncharacterized</p>	72

	forms	
<b>Figure 4.3</b>	STRING analysis of differentially regulated proteins involved in the biological processes of <i>C. albicans</i> due to the expression of ubiquitin mutants. Protein-protein functional interaction prediction map of downregulated proteins of <i>C. albicans</i> obtained using STRING v.11 database (confidence score 0.700).	78
<b>Figure 4.4</b>	STRING analysis of differentially regulated proteins involved in the metabolic processes of <i>C. albicans</i> due to the expression of ubiquitin mutants. Protein-protein functional interaction prediction map of downregulated proteins of <i>C. albicans</i> obtained using STRING v.11 database (confidence score 0.700). Filled protein nodes that signify the availability of protein 3D structural information either known or predicted.	79
<b>Figure 4.5</b>	KEGG pathway map of glycolysis/ gluconeogenesis. Mappable enzymes within the pathway are highlighted in green and teal denoted as the pathways identified with the aid of the following database: <a href="https://www.kegg.jp/entry/cal00010">https://www.kegg.jp/entry/cal00010</a> through <a href="http://www.cytoscape">http://www.cytoscape</a>	83
<b>Figure 4.6</b>	Gene ontology and STRING analysis of differentially regulated proteins involved in the metabolic processes of <i>C. albicans</i> due to the expression of ubiquitin mutants. Protein-protein functional interaction prediction map of upregulated proteins of <i>C. albicans</i> obtained using STRING v.11 database (confidence score 0.700). Red color indicates the interaction of MET3 with MET10.	84
<b>Figure 4.7</b>	KEGG pathway map of cysteine methionine pathway. Mappable enzymes within the pathway are highlighted in green and were identified with the aid of the following database <a href="https://www.kegg.jp/pathway/map=cal00270&amp;keyword=">https://www.kegg.jp/pathway/map=cal00270&amp;keyword=</a> through <a href="http://www.cytoscape">http://www.cytoscape</a>	87
<b>Figure 5.1</b>	Hypothetical illustration for possibility of polyubiquitin chain	92

	formation	
<b>Figure 5.2</b>	Vector map of P-TET25MC-H for ubiquitin Di- glycine mutants cloning and screening.	93
<b>Figure 5.3</b>	Plasmid PTET25M-CH isolation and PCR for amplification of mutants. (A) 1% agarose gel showing (A) PTET25M-CH 7.5 kb plasmid isolation. lane 1 shows lambda HindIII DNA marker, Lane 2-5 is the plasmid PTET25M-CH containing UbWT, UbEP42, UbL50P, UbI61T respectively (B 1% agarose gel showing shows PCR amplicon 228 bp of ubiquitin mutants, lane 1 contains lambda HindIII 1 kb DNA marker, Lane 2-5 showing PCR amplified ubiquitin mutants UbWT, UbEP42, UbL50P, UbI61T using forward and reverse primers listed in Table5.1	97
<b>Figure 5.4</b>	PTET25M-CH Di glycine ubiquitin mutants cloning. 3% agarose gel showing Plasmid PTET25M-CH double digested (Lane 3,5) with <i>Bgl</i> II and <i>Xma</i> I restriction enzyme. Lane 6 shows UbWT amplified without Di glycine deletion at the C- terminal and lane 7 shows di glycine deletion from UbWT. The difference in their migration pattern is visualized. Similarly, lanes 8, and 9 for UbEP42, lanes 10 and 11 for UbL50P, and lanes 12 and 13 for UbI61T show migration difference. (B) The electropherogram obtained through sequence analysis showed the deletion of Di-glycine in all ubiquitin mutants	98
<b>Figure 5.5</b>	<b>Ubiquitination and K48 and K63 linkage under Diglycine ubiquitin mutants.</b> (A) Coomassie-stained SDS-PAGE gel (12%) of yeast stage lysate. Lane 1, SDS-PAGE marker; lane 2-5 total yeast extract proteome profile after induction of UbWtGG-, UbEP42GG-, UbL50PGG- and UbI61TGG- respectively. (B) Coomassie-stained 12% SDS-PAGE gel of hyphal stage lysate. Lane 1, SDS-PAGE marker; lane 2-5 total <i>C. albicans</i> hyphal stage extract proteome profile after induction of UbWt, UbEP42, UbL50P, and ubI61T respectively. (C) Western blot showing ubiquitination profile. <i>Candida albicans</i> strains harbouring the ubiquitin mutants with Di-glycine (+) and without Di-glycine (-)	100

	<p>constructions were grown in YPD media doxycycline inducer for 3 h. Whole cell extract was separated by SDS-PAGE and probed with an anti ub antibody, allowing the detection of the ubiquitin protein. Lane 2,3 UbWT, Lane 4,5 UbL50P, and Lane 6,7 Ubi61T representing ubiquitination profile with or without Di-glycine (-/+)</p> <p>(D) UbEP42 comparative ubiquitination profile under di-glycine deletion condition. Indicating nondegradative substrate with polyubiquitin chain accumulation decline. (E) K48 and K63 linkage in UbEP42GG- showing a decline in polyubiquitination chain aggregation.</p>	
<b>Figure 5.6</b>	<p>Prediction of Di- glycine Ubiquitin mutants inter-residue 3D geometries by trRosetta and align the structure with ubiquitin mutants having Di-glycine motifs at C-terminal using Mol* 3D viewer from protein data bank (RCSB PDB). The predicted structure models With Di-glycine shown in color cartoons (Orange) and without Di-glycine (Green). TM- scores 0.1 for all the constraints model. (B) Representing combine all Di-glycine deleted ubiquitin mutants structures align model using RPSB PDB Mol* 3D viewer.</p>	102
<b>Figure 5.7</b>	<p>Prediction of Di- glycine Ubiquitin mutants inter-residue 2D geometries by trRosetta The predicted secondary structure showing that all ubiquitin mutants have the same no. Helix, Strand, and Coil.</p>	103

### LIST OF TABLES

Sr. No.	TITLE	Page No.
<b>Table 2.1</b>	Table 1. Primers used in this study.	28
<b>Table 4.1</b>	Common upregulated and downregulated protein under UbEP42, UbL50P, Ubi61T mutants overexpression	73
<b>Table 4.2</b>	Upregulated and Downregulated protein associated with biological process and metabolic pathway under UbEP42, UbL50P, Ubi61T mutant's expression	74

<b>Table 4.3</b>	Primers used for transcription factor analysis	68
<b>Table 5.1</b>	Primers used for this study	94
<b>Table 5.2</b>	<i>C. albicans</i> strains used in this study	95